

North Atlantic summers have warmed more than winters since 1353, and the response of marine zooplankton

Nicholas A. Kamenos¹

School of Geographical and Earth Sciences, University of Glasgow, Glasgow G12 8QQ, Scotland

Edited by Edward A. Boyle, Massachusetts Institute of Technology, Cambridge, MA, and approved November 2, 2010 (received for review May 3, 2010)

Modeling and measurements show that Atlantic marine temperatures are rising; however, the low temporal resolution of models and restricted spatial resolution of measurements (i) mask regional details critical for determining the rate and extent of climate variability, and (ii) prevent robust determination of climatic impacts on marine ecosystems. To address both issues for the North East Atlantic, a fortnightly resolution marine climate record from 1353–2006 was constructed for shallow inshore waters and compared to changes in marine zooplankton abundance. For the first time summer marine temperatures are shown to have increased nearly twice as much as winter temperatures since 1353. Additional climatic instability began in 1700 characterized by ~5–65 year climate oscillations that appear to be a recent phenomenon. Enhanced summer-specific warming reduced the abundance of the copepod *Calanus finmarchicus*, a key food item of cod, and led to significantly lower projected abundances by 2040 than at present. The faster increase of summer marine temperatures has implications for climate projections and affects abundance, and thus biomass, near the base of the marine food web with potentially significant feedback effects for marine food security.

coralline algae | rhodolith | maerl | seasonal

Prior to instrumentally derived temperature records that began in the 1850s, knowledge of seasonal and regional-scale marine temperature changes at high latitudes is limited. Thus at present, hemispheric Atlantic climatic patterns are deduced from multiproxy and instrumental networks using statistical (1, 2) and modeling (3) approaches. Whereas such approaches show that the Atlantic Multidecadal Oscillation (AMO), the North Atlantic Oscillation (NAO), and Atlantic Meridional Overturning Circulation (AMOC) have a substantial influence on Atlantic climate (e.g., ref. 4) they are still unable to capture details that are critical for determining the spatiotemporal distribution of climate oscillations (5, 6) and their associated impacts (7). For example, whereas the AMO, which models suggest reflects multidecadal variation in the AMOC (8), is the dominant influence on summer climate in the United Kingdom (9), at higher European latitudes, seasonal terrestrial tree-ring climate records (6, 10) are not representative of high-latitude sea surface temperatures (11). Limited spatiotemporal resolution, absent proxy validation and organism effects have restricted the distribution of current inshore proxy-derived sea surface temperature (SST) records (12–16). Therefore to date, we have a poor understanding of season specific temperature variability. Similarly, SST, AMO, and AMOC variations are not well represented at high spatiotemporal resolution beyond the instrumental record in North East Atlantic shallow inshore waters (for example 55°50'N 05°50'W). This information is needed to better understand rates of regional climate variability (5).

Marine responses to such climate fluctuations are reflected in the productivity of marine ecosystems from phytoplankton to fish population dynamics (17). Therefore a key requirement for preserving biodiversity and supporting sustainable fishing is

to understand the climatic effects in the lower levels of food webs (18). At present, general circulation models treat the marine environment in a simplified way, lacking the detail required for investigating the impacts of climate on marine ecosystems (7) beyond phytoplanktonic trophic levels. Thus the utility of high resolution marine temperature histories coupled with seasonal records of productivity provide a unique tool for probing climate-marine ecosystem relationships more accurately. With models suggesting that Atlantic SST will continue to rise (19) we need to understand how such changes are likely to affect food security.

The first biweekly resolution record of high-latitude, North East Atlantic, inshore, shallow-water temperature from 1353–2006 is presented and related to seasonal resolution inshore zooplankton changes. Mg/Ca ratios obtained from red coralline algae (*Lithothamnion glaciale*) were converted to in situ marine temperatures using Mg-temperature calibrations performed here and in ref. 20. Red coralline algae are validated (14) archives of palaeoclimate information at subannual time scales (20, 21), which band annually (20, 22) and have a worldwide shallow-water distribution (23). As zooplankton amplify hydroclimatological forcing (24) this investigation targeted historic relationships between North East Atlantic algae-derived in situ temperatures and abundances of the copepod *Calanus finmarchicus*, the most abundant copepod species higher than 50° N (25) and an important food source for cod (26). East Atlantic *C. finmarchicus* abundance was related to algae-derived in situ temperature and that relationship along with model-derived temperature projections (19) were used to project *C. finmarchicus* abundance to 2040.

Results and Discussion

Since the Little Ice Age (LIA) summer marine temperatures in North East Atlantic inshore shallow waters have increased nearly twice as much as those in winter. From the LIA [1400–1700 (27)] to the 1900–2006 mean, summer temperatures rose by $2.7 \pm 0.5^\circ\text{C}$ while winter temperatures rose by $1.4 \pm 0.5^\circ\text{C}$ (Fig. 1). This seasonal difference in change may be attributed either to (i) more suppressed summer than winter temperatures during the LIA or (ii) that since the LIA summers have warmed more than winters. In both cases, summers have warmed more than winters during the last 6 centuries.

Temperature peaks near ~1350, 1700, 1870, 1920, and since the 1970s are similar to present day SSTs. Distinct temperature troughs occurred near ~1540, 1840, 1880, and 1960. In fact, since 1930 the mean temperatures have been significantly higher than the 1353–1929 summer ($F_1 = 50.62$, $p < 0.001$) and winter ($F_1 = 78.39$, $p < 0.001$) temperatures respectively.

Author contributions: N.A.K. designed research, performed research, analyzed data, and wrote the paper.

The author declares no conflict of interest.

This article is a PNAS Direct Submission.

E-mail: nick.kamenos@glasgow.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1006141107/-DCSupplemental.




Fig. 4. Relationship between annual log (natural) *C. finmarchicus* abundance and summer algae-derived in situ temperature on the west coast of Scotland between 1956 and 2001 with 0 years lag.

ton (*C. finmarchicus* and *Calanus helgolandicus*) abundance due to reduced food (zooplankton availability and quality) (26). Since 1958 increasing temperatures have reduced the quality of zooplanktonic food available for larval/juvenile cod [progressive substitution of *C. finmarchicus* by *C. helgolandicus* at higher temperatures (25)], mediated by changes in water column stratification (49, 50), leading to poor recruitment to adult populations in the North Sea (26). In addition, decline in food availability has exacerbated the decline of European cod stocks due to overfishing (7). Added variability in cod recruitment also exists as a consequence of stock specific responses to: (i) temperature [e.g., stocks at different positions in cod's distributional range responding differently (51) and/or changing their spawning locations (52)], (ii) wind and current induced changes in the distribution of cod eggs and larvae (53) and (iii) variable responses to NAO (46, 54). Thus whereas in certain areas warming climate will lead to increased fishery yields (18) it appears that cod yields in the East Atlantic may be altered as a consequence of enhanced summer warming of waters at the time when their food, *C. finmarchicus*, is most sensitive to those changes. Such fluctuations in fish stocks have major economic and social consequences (7) and thus a detailed understanding of climate-induced changes, such as the inclusion of ecological knock-on effects (55), is required.

Conclusions

As global warming begins to represent a risk to global food security on land (56), marine fish stocks are also starting to dwindle (7). Enhanced increases in shallow-water marine summer temperatures observed here, have a negative effect on *C. finmarchicus* abundance in summer when it is most sensitive to temperature. As *C. finmarchicus* is a key prey item for cod this may have detrimental effects higher up the food web. We thus need to further understand if summers will continue to warm more than winters with respect to seasonal climate projections, the stability of the marine climate system and the knock-on effects these synergistic factors will have on fisheries and marine food security.

Materials and Methods

Algal Collection. SCUBA (self-contained underwater breathing apparatus) was used to collect a 20 × 46 cm core through a *L. glaciale* deposit in Loch Sween (56°01.99'N 05°36.13'W) on the west coast of Scotland in January 2007. Live *L. glaciale* on the surface of the deposit protected the core top from disturbance. The core was frozen and sectioned longitudinally; each half was sectioned into 2 cm horizons. To avoid edge effects of the coring process each horizon was recored using a 6 cm diameter core. *L. glaciale* thalli were picked from each horizon and washed. The longest thalli were selected for further analysis.

Rangefinder Dates. Five thalli were selected for radiocarbon rangefinder dating. For each thallus, a branch tip attached directly to the section used in Mg/Ca determination, and containing approximately 20 bands (years of growth), was selected for analysis. Each sample was sonicated to remove any ground-in material and analyzed whole. Accelerator mass spectrometry radiocarbon dating was conducted at the Scottish Universities Environmental Research Centre and adjusted by 405 ¹⁴C years for the marine reservoir effect (57) (Table 1).

Sample Analysis. All analyses were conducted on a Cameca SX100 electron microprobe at the School of Geosciences, University of Edinburgh. Parallel transects ($n = 2$) were made along the length of each thallus. Along each transect, spot analyses (20 μm diam.) simultaneously recording Mg, Ca, and Sr were made at 20 μm steps [electron microprobe conditions: acceleration voltage: 10 kV, beam current: 4 nA, dwell time: 20 sec (see (20) for detailed conditions)]. Samples were imaged using SEM (FEI Quanta 200 F) and any spot analyses that were taken on growth irregularities were rejected (82 of the 12,888 analyses). Remaining values from the 2 transects were averaged to generate a Mg/Ca (molar) time-series. As *L. glaciale* band annually (20), band counting/imaging was used to correspond individual bands to annual peaks and troughs in the Mg/Ca time series that corresponded to maximum and minimum temperatures in individual years (e.g., as in Fig. S1). For the live collected specimens, this process enabled absolute dates to be assigned to each peak/trough. For validation of analytical technique, see SI Text and Fig. S2.

Combined Chronology. Six thalli were used as anchor points in construction of the combined chronology; a core top thallus provided the 0 year (as it was live on collection) and 5 other thalli provided radiocarbon dates. Twenty seven (including anchors) Mg/Ca time series were available; each from an individual thallus. Each time series was presented as both summer maximum and winter minimum molar Mg/Ca [maximum Mg is deposited during summer (20)]. These were detrended (least-squares regression) to remove any effects of band width on the number of Mg/Ca readings yr^{-1} , and hence resolution, when determining the chronology. The anchor point thalli were used to give all other thalli an estimated age (using interpolation). To further refine the age of each thallus, and thus the accuracy of the chronology, the detrended summer Mg/Ca time series determined from each thallus was cross correlated (average overlap = 64 years) with Mg/Ca time series for thalli that grew at similar times (estimated from interpolated age). Relative lagged ages determined by the cross correlation were only accepted if all the following criteria were met (i) correlation coefficient $r >= 0.36$ ($p < 0.001$) (58), (ii) Gleichlaufigkeit, GLK $>= 65\%$ (59), and (iii) Baillie–Plicher t value, $t_{BP} >= 3.6$ (60).

The combined use of radiocarbon dating and dendrochronological techniques allows for a chronology in which no ages are based only on interpolation thus reducing uncertainty introduced by interpolation as well as error associated with the radiocarbon dates. Thus the maximum error on any ages determined will be $t \pm 40$ y (the error on radiocarbon dates) plus/minus any interpolation error (≤ 14 y) whereas the minimum error will be 0 years (the best age determined using cross correlation). As three filtering criteria were used to determine cross correlation significance, it is likely that age model error falls well below the 54 year maximum and is probably nearer 0 years. It is unlikely there is error on the 1857–2006 ages as thalli that grew during these years were live on collection thus are absolutely dated, this is important as these are the years during which comparisons with absolutely dated zooplankton abundance are made.

Mg-Temperature Relationship. The Mg-temperature relationship: $\text{Mg/Ca (molar)} = 0.013 \text{ IST}(\text{°C}) + 0.04$ (IST = in situ temperature, Mg/Ca = molar Mg/Ca ratios in *L. glaciale*, $R^2 = 0.93$, $p < 0.001$, $SE_b \pm 6.5^{-4}$,

Table 1. Radiocarbon data for *L. glaciale* samples from Loch Sween, Scotland

Sample code	Depth (cm)	¹³ C relative to VPDB (‰)	Age	Year
SUERC-16011	8–10	–6.7	Modern	Pre 1950 AD
SUERC-16013	14–16	–6.9	Modern	Pre 1950 AD
SUERC-16005	26–28	–4.1	320 ± 40	1668 ± 40
SUERC-16006	30–32	–12.0	355 ± 40	1651 ± 40
SUERC-16012	44–46	–3.9	630 ± 40	1376 ± 40

Depth denotes the section depth down from top of live surface thalli, sediment depth begins ~8 cm below thallus surface. Ages are given as conventional ¹⁴C yr BP with analytical error of $\pm 1\sigma$.

$SE_{\pm} \pm 6^{-3}$, equivalent error on reconstruction = $\pm 0.5^{\circ}\text{C}$) was determined using measured in situ temperature, the calibration site, technique, and validation in ref. 20 (Fig. S1). Two temperature time series were constructed, for (i) maximum summer temperature and (ii) minimum winter temperature. Fieller's Theorem was used to propagate errors. A gap is present in the time series during 1388–1405 due to the absence of multiple thalli growing during that period. Occasional individual gaps are also present when analyses in all thalli representing a particular time happened to fall on a growth irregularity and were rejected. ANCOVA (assumptions met) was used to compare temperatures between temporal bins.

Spectral Analysis. Multitaper method (MTM) spectral estimation (37, 61, 62) was conducted to estimate the winter algae-derived in situ temperature variability period. Winter temperatures were used as these gave a longer uninterrupted time series than summer temperatures. The MTM bandwidth parameter is $p = 2$ and $K = 3$ Slepian eigentapers were used. Confidence intervals were determined relative to a red noise AR(1) background. Comparison of algae-derived in situ temperature time series with the winter AMO and winter NAO time series was conducted using Magnitude-Squared Coherence—the frequency domain analogue of correlation (62, 63). For both the AMO and NAO the longest suitable time series available were used. The same bandwidth parameter and eigentaper were used as in MTM analysis. In all cases only signals significant at the 99% level were considered. Analyses were conducted on detrended time series (spline/linear model residuals) that were detrended separately.

Temperature-Zooplankton Relationships Data sources. Historic sea water temperatures were algae-derived in situ temperature time-series developed during the current investigation. Projected SSTs from 2000–2040 for $57^{\circ}07'53''\text{N}$ $08^{\circ}12'60''\text{W}$ were obtained from (19) and were developed under an SRES-A2 scenario. *Calanus finmarchicus* abundance (semiquantitative relative scale; e.g., number of zooplankton per unit volume of water sampled) data were collected using a Continuous Plankton Recorder in Area C4 (inshore west coast Scotland) and provided by The Sir Alister Hardy Foundation for Ocean Science (www.sahfos.ac.uk).

Data preparation and relationships. Residuals of each time series were obtained from linear models (variable ~ time). Remaining autocorrelation

was removed by fitting successively higher order Autoregressive Integrated Moving Average models to each input variable (64). All analyses were conducted on the log stabilized residuals.

Model validation and projection. Temperature-zooplankton relationships were determined using models constructed from ~66% of the available time-series data (1958–1987). Those relationships were used to project the zooplankton abundance over the same period as the remaining 33% of the time series (1988–2001) providing projected and actual abundance time series over the period 1988–2001. Projective capacity was validated by regressing the difference between actual zooplankton abundance and projected zooplankton abundance on sums of actual and projected zooplankton abundance (65) between 1988–2001. *C. finmarchicus* abundance was projected to 2040 using relationships [1] and [2]. The projection is useful for detecting changes in trend at decadal scales, however, is not suitable for detecting subannual patterns as error propagation makes subdecadal patterns hard to interpret. This can be observed in the small step change between proxy-derived and modeled temperatures at annual scales (Fig. 3) whereas the linear trendlines, representative of temperature/abundance at decadal scales, are temporally coherent. In addition, the annual step change is not of concern at decadal scales as the variability of modeled temperature is well within the variability of proxy-derived temperature (Fig. 3). Repeated measures ANOVA (assumptions met) was used to compare *C. finmarchicus* abundances at different temporal points.

Statistical analysis. Mg/Ca cross dating was conducted using CDendro (Cybis Elektronik & Data), spectral estimation was conducted using kSpectra Toolkit (Spectraworks, www.spectraworks.com) and all other analyses were conducted using R v2.10 (66).

ACKNOWLEDGMENTS. Thanks to Peter Chung, Chris Hayward, John Faithful, Maggie Cusack, Marian Scott, and Vernon Phoenix for help with sample collection, analysis, and constructive suggestions. Thanks also to the University Marine Biological Station Millport dive team, the Scottish Universities Environmental Research Centre, and The Sir Alister Hardy Foundation for Ocean Science. This research was made possible by Grants NERC NE/D008727 and Royal Society of Edinburgh/Scottish Government RSE 48704/1.

- Luterbacher J, Dietrich D, Xoplaki E, Grosjean M, Wanner H (2004) European seasonal and annual temperature variability, trends, and extremes since 1500. *Science* 303:1499–1503.
- Mann ME, Bradley RS, Hughes MK (1998) Global-scale temperature patterns and climate forcing over the past six centuries. *Nature* 392:779–787.
- Goosse H, Renssen H, Timmermann A, Bradley RS, Mann ME (2006) Using paleoclimate proxy-data to select optimal realisations in an ensemble of simulations of the climate of the past millennium. *Clim Dynam* 27:165–184.
- Luterbacher J, et al. (2002) Extending North Atlantic Oscillation reconstructions back to 1500. *Atmos Sci Lett* 10.1006/asle.2001.0044.
- Mann ME, et al. (2003) On past temperatures and anomalous late-20th century warmth. *EOS Trans Am Geophys Union* 84:256–257.
- Casty C, Wanner H, Luterbacher J, Esper J, Bohm R (2005) Temperature and precipitation variability in the European Alps since 1500. *Int J Climatol* 25:1855–1880.
- Brander K (2010) Impacts of climate change on fisheries. *J Marine Syst* 79:389–402.
- Dong BW, Sutton RT (2005) Mechanism of interdecadal thermohaline circulation variability in a coupled ocean-atmosphere GCM. *J Climate* 18:1117–1135.
- Sutton RT, Hodson DLR (2005) Atlantic Ocean forcing of North American and European summer climate. *Science* 309:115–118.
- Delworth TL, Mann ME (2000) Observed and simulated multidecadal variability in the Northern Hemisphere. *Clim Dynam* 16:661–676.
- D'Arrigo R, Wilson R, Liepert B, Cherubini P (2008) On the "Divergence Problem" in Northern Forests: A review of the tree-ring evidence and possible causes. *Glob Planet Change* 60:289–305.
- Schone BR, et al. (2005) Climate records from a bivalved Methuselah (*Arctica islandica*, Mollusca; Iceland). *Palaeogeog Palaeoclimatol Palaeoecol* 228:130–148.
- Weidman CR, Jones GA, Lohmann KC (1994) The long-lived mollusc *Arctica islandica*: A new palaeoceanographic tool for the reconstruction of bottom temperatures for the continental shelves of the northern North Atlantic Ocean. *J Geophys Res* 99:18,305–18,314.
- Kamenos NA, Cusack M, Huthwelker T, Lagarde P, Scheibling RE (2009) Mg-lattice associations in red coralline algae. *Geochim Cosmochim Acta* 73:1901–1907.
- Cohen AL, Owens KE, Layne GD, Shimizu N (2002) The effect of algal symbionts on the accuracy of Sr/Ca paleotemperatures from coral. *Science* 296:331–333.
- Shirai K, et al. (2005) Deep-sea coral geochemistry: Implication for the vital effect. *Chem Geol* 224:212–222.
- Cushing DH (1982) *Climate and Fisheries* (Academic Press, London) p 294.
- Brown CJ, et al. (2010) Effects of climate-driven primary production change on marine food webs: Implications for fisheries and conservation. *Glob Change Biol* 16:1194–1212.
- Stendel M, Schmith T, Roeckner E, Cubasch U (2004) IPCC-ECHAM4OPYC_SRES_A2_MM. *World Data Centre for Climate* 10.1594/WDC/IPCC_EH4_OPYC_SRES_A2_MM.
- Kamenos NA, Cusack M, Moore PG (2008) Red coralline algae are global paleothermometers with bi-weekly resolution. *Geochim Cosmochim Acta* 72:771–779.
- Halfar J, Zack T, Kronz A, Zachos JC (2000) Growth and high resolution palaeoenvironmental signals of rhodoliths (coralline red algae): A new biogenic archive. *J Geophys Res* 105:22,107–122,116.
- Kamenos NA, Law A (2010) Temperature controls on coralline algal skeletal growth. *J Phycol* 46:331–335.
- Foster MS (2001) Rhodoliths: Between rocks and soft places. *J Phycol* 37:659–667.
- Taylor AH, Allen JI, Clark PA (2002) Extraction of a weak climatic signal by an ecosystem. *Nature* 416:629–632.
- Planque B, Fromentin JM (1996) *Calanus* and environment in the eastern North Atlantic. 1. Spatial and temporal patterns of *C. finmarchicus* and *C. helgolandicus*. *Mar Ecol Prog Ser* 134:101–109.
- Beaugrand G, Brander KM, Lindley JA, Souissi S, Reid PC (2003) Plankton effect on cod recruitment in the North Sea. *Nature* 426:661–664.
- Mann ME, et al. (2009) Global signatures and dynamical origins of the Little Ice Age and Medieval Climate Anomaly. *Science* 326:1256–1260.
- Kreutz KJ, et al. (1997) Bipolar changes in atmospheric circulation during the Little Ice Age. *Science* 277:1294–1296.
- Saenger C, Cohen AL, Oppo DW, Halley RB, Carilli JE (2009) Surface-temperature trends and variability in the low-latitude North Atlantic since 1552. *Nat Geosci* 2:492–495.
- Cook ER, D'Arrigo RD, Mann ME (2002) A well-verified, multiproxy reconstruction of the winter North Atlantic Oscillation index since AD 1400. *J Climate* 15:1754–1764.
- Kerr RA (2000) A North Atlantic climate pacemaker for the centuries. *Science* 288:1984–1986.
- Freiwald A, Henrich R (1994) Reefal coralline algal build-ups within the Arctic circle: Morphology and sedimentary dynamics under extreme environmental seasonality. *Sedimentology* 41:963–984.
- Kremer BP (1981) Aspects of carbon metabolism in marine macroalgae. *Oceanogr Mar Biol Annu Rev* 19:41–95.
- Schlesinger ME, Ramankutty N (1994) An oscillation in the global climate system of period 65–70 years. *Nature* 367:723–726.
- Goodkin NF, Hughen KA, Doney SC, Curry WB (2008) Increased multidecadal variability of the North Atlantic Oscillation since 1781. *Nat Geosci* 1:844–848.
- Mann ME, Park J (1994) Global-scale modes of surface temperature variability on interannual to century timescales. *J Geophys Res* 99:25819–25833.

