



## Microarray analysis of juvenile hormone response in *Drosophila melanogaster* S2 cells

David K. Willis<sup>1a</sup>, Jun Wang<sup>2,4b</sup>, Joliene R. Lindholm<sup>2c</sup>, Anthony Orth<sup>3d</sup> and Walter G. Goodman<sup>2e</sup>

<sup>1</sup>USDA/ARS Vegetable Crops Research Unit, University of Wisconsin – Madison, Madison WI 53706

<sup>2</sup>Department of Entomology, University of Wisconsin – Madison, Madison WI 53706

<sup>3</sup>Genomics Institute of the Novartis Research Foundation, San Diego, CA 92121

<sup>4</sup>Current address: INVITROGEN Corporation, 501 Charmany Drive, Madison, WI 53719

### Abstract

A microchip array encompassing probes for 14,010 genes of *Drosophila melanogaster* was used to analyze the effect of juvenile hormone (JH) on genome-wide gene expression. JH is a member of a group of insect hormones involved in regulating larval development and adult reproductive processes. Total RNA was isolated from *Drosophila* S2 cells after 4 hours treatment with 250 ng/ml (10R) JH III or 250 ng/ml methyl linoleate. A collection of 32 known or putative genes demonstrated a significant change with JH III treatment ( $r > 2.0$ ,  $P \leq 0.005$ ). Of these, the abundance of 13 transcripts was significantly increased and 19 decreased. The expression of a subset of these loci was analyzed by real-time quantitative reverse transcription polymerase chain reaction (RT-qPCR). Three loci that exhibited constant expression in the presence and absence of JH III (RP49 [FBgn0002626], FBgn0023529, and FBgn0034354) were evaluated and found to be reliable invariant reference transcripts for real-time RT-qPCR analysis using *BestKeeper* and *geNorm* software. Increased expression in presence of JH III was confirmed by real-time RT-qPCR analysis. However, only one of five loci that exhibited reduced expression on microarrays could be confirmed as significantly reduced ( $P \leq 0.05$ ). Among the confirmed JH III up-regulated genes were two loci of unknown function (FBgn0040887 and FBgn0037057) and *Epac*, an exchange protein directly activated by cyclic AMP, a guanine nucleotide exchange factor for Rap1 small GTPase.

**Keywords:** real-time RT-qPCR, *Epac*

**Abbreviations:** **JH**, juvenile hormone; **Epac**, exchange protein directly activated by cyclic AMP; **MLA**, methyl linoleate; **ORF**, open reading frame; **RT-qPCR**, quantitative reverse transcription polymerase chain reaction; **RER**, relative expression ratio

**Correspondence:** <sup>a</sup> dkwillis@wisc.edu, <sup>b</sup> jun.wang01@invitrogen.com, <sup>c</sup> jlindholm@entomology.wisc.edu, <sup>d</sup> aorth@gnf.org, <sup>e</sup> goodman@entomology.wisc.edu

**Associate Editor:** Zhijian Jake Tu was editor of this paper.

**Received:** 3 January 2008, **Accepted:** 11 October 2008

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 10, Number 66

#### Cite this paper as:

Willis DK, Wang J, Lindholm JR, Orth A, Goodman WG. 2010. Microarray analysis of juvenile hormone response in *Drosophila melanogaster* S2 cells. *Journal of Insect Science* 10:66 available online: [insectscience.org/10.66](http://insectscience.org/10.66)

## Introduction

---

The insect juvenile hormones (JH) represent a family of acyclic sesquiterpenoids that regulate a diversity of processes in the insect life cycle (Nijhout 1994; Riddiford 1996; Gade et al. 1997; Lafont 2000; Goodman et al. 2005). JH affects insect development by maintaining the larval stage and inhibiting metamorphosis. In adults, JH is involved in regulating reproductive physiology (Riddiford 1996). Although well-studied from the physiological standpoint, the molecular mechanisms underlying JH action remain largely unknown (Jones 1995).

Several molecular mechanisms for JH action have been proposed (Wheeler et al. 2003; Goodman et al. 2005). It has been suggested that JH acts through a specific nuclear receptor complex that modulates gene expression at the level of transcription (Riddiford 1996). This hypothesis is supported by the lipophilic nature of JH and its chemical similarity to the retinoids, compounds known to activate specific nuclear transcription factors, including the vertebrate retinoid X receptor (Mangelsdorf et al. 1995).

Due to its lipophilicity, one might expect the hormone to easily pass through the cellular membrane and interact with cytosolic or nuclear transcription factors; however, there is increasing evidence that suggests JH may act at the membrane level triggering a membrane-receptor-mediated signal transduction pathway. In the male accessory glands of *Drosophila melanogaster*, it has been demonstrated that JH acts via protein kinase C and calcium to stimulate protein synthesis (Yamamoto et al. 1988). This interaction with protein kinase C is a classical signal transduction pathway that involves membrane

receptors and G-coupled proteins (Sevala et al. 1989; Pszczolkowski et al. 2005; Kethidi et al. 2006). Thus, JH may regulate gene expression at multiple levels and through multiple mechanisms.

Genome-wide gene expression analysis by microarray is the method of choice to identify insect genes that are affected by JH treatment. A *Drosophila* microarray chip is currently available that contains probes for 14,010 putative open reading frames (ORF) within the genomic DNA of this model insect (Affymetrix, Inc). While microarray technology is widely used for expression analysis, the technique exhibits problems that are becoming increasingly apparent. Microarray is a reliable method to detect changes in expression of high abundance genes but the accuracy of identifying changes in low abundance gene transcripts is somewhat problematic (Beckman et al. 2004; Morey et al. 2006). Of particular concern is the ability of microarray analysis to correctly identify changes in low abundance genes or down-regulation of medium abundance genes. Both of these problems arise from the interference of background fluorescence with the low intensity signal from low abundance genes or lower expression of medium abundance genes (Beckman et al. 2004). The accuracy of microarray can be optimized by defining a threshold of reliability based on fold-change and P-value from the chip analysis software, but problems with false positives and false negatives remain (Morey et al. 2006).

Real-time quantitative reverse-transcription quantitative PCR (real-time RT-qPCR) has become the standard technology to verify microarray gene expression profiling. Real-time RT-qPCR has many advantages over

microarray for the quantification of specific gene transcripts such as the affordability of performing multiple biological replications and normalizing expression to validated reference RNAs that are known to be invariant under experimental conditions. A major advantage of real-time RT-qPCR is a greatly expanded dynamic range. Microarray analysis can reliably detect expression differences over a three-order of magnitude range (1000-fold) while the dynamic range of real-time RT-qPCR extends over seven orders of magnitude (10 million-fold) (Beckman et al. 2004). This enables the accurate measurement of differences over a much larger range of gene expression levels including medium and low abundant transcripts. Two strategies are commonly employed to enumerate the results obtained by real-time RT-qPCR; the standard curve method (absolute quantification) and the comparative threshold method (relative quantification). Absolute quantification relies on the inclusion of a standard curve on each reaction plate and results in determination of the actual quantity of the target transcript expressed in copy number or weight. This method has the advantage of correcting differences in primer efficiencies. The disadvantage of absolute quantification is the significant reduction in the number of experimental samples that can be run on a single plate. Relative quantification determines changes in steady-state mRNA levels of a gene across multiple samples and biological replicates by determining the change in gene expression relative to a control RNA that is designated as the calibrator (Pfaffl 2001; Rasmussen 2001). With this method, target transcript amounts are expressed as a relative expression ratio (RER) relative to the calibrator. Both methods require the normalization of target gene expression using multiple stably expressed internal control mRNAs. These reference gene

mRNAs must be shown to be stable under the experimental conditions being examined and are evaluated using software programs such as *BestKeeper* or *geNorm* (Vandesompele et al. 2002; Pfaffl et al. 2004). As with any quantitative measure, care must be taken with real-time RT-qPCR to insure that the necessary controls and evaluations have been performed. These include: assessment of RNA quality, assessment of DNA contamination, determination of primer efficiencies and sensitivities, and the use of multiple stable reference RNAs. A recent survey of real-time RT-qPCR publications revealed that only 30% of the published analyses examined satisfied all of these criteria (Bustin 2005).

In this work, we analyzed genome-wide JH III induced expression changes in *Drosophila* S2 cells by microarray. As a control for the lipid component of JH III, methyl linoleate (MLA) was used, as it is a lipid with physical characteristics similar to JH III but is not hormonally active in insects. Microarray expression differences of a select set of genes using real-time RT-qPCR were validated using several reference transcripts that were stably expressed in S2 cells under experimental conditions.

## Materials and Methods

### Purification and quantification of JH homologs

JH III and MLA were purchased from Sigma Chemicals. The biologically active enantiomer (10*R*) of JH III was isolated from a racemic mixture by chiral HPLC chromatography (Cusson et al. 1997).

### Cell culture

*Drosophila* S2 cells (Invitrogen) were maintained in SF900 serum-free medium (Invitrogen) at 27° C. Cells (5X10<sup>5</sup>/ml) were

seeded into 60 mm petri dishes (Nunc) containing 3 ml of medium and allowed to grow for 36 h. Cells at approximately 80% confluency were challenged with 250 ng/ml (10R) JH III using charcoal-stripped 0.1% bovine serum albumin (BSA) as a carrier for 4 h. Control cells were treated with 0.1% BSA alone or 0.1% BSA with 250 ng/ml MLA and harvested after 4 h of treatment.

### Isolation of RNA

Cells were lysed directly in the culture dishes and total RNA extracted using the RNeasy mini kit (Qiagen). RNA was exhaustively treated with 2 U of Turbo DNAase (Ambion) for 1 h at 37°C and quantified by UV spectrophotometry (NanoDrop, Inc.). RNA quality was determined by electrophoresis of samples on denaturing agarose gels. Residual DNA contamination was quantified using real-time RT-qPCR and primers specific for the *Drosophila rp49* gene (Table 1). Those RNA samples showing threshold cycle ( $C_q$ ) values  $\geq 33$  cycles were deemed to be free of DNA contamination.

### Microarray analysis

Total RNA was made into cRNA using standard reagents (Affymetrix, Inc). Duplicate total RNA pools from two independently treated S2 cultures were taken for each sample, and resulting single-dye labeled cRNAs were hybridized to *Drosophila* Genome Arrays (Affymetrix, Inc). Arrays were washed with a custom array washer, and scanned with an Affymetrix 3000 scanner. Cell intensity files were analyzed using the Rosetta Resolver algorithm (Rosetta Biosoftware) and comparisons were performed using Resolver's ratio ANOVA function. Resolver ANOVA analysis is similar to standard ANOVA but uses two inputs, expression measurement quantity and estimated error of measurement quantity. This additional input provides more reliable variance measurements, a necessity when the number of replicates is small (Rajagopalan 2003). This error estimate also brings extra degrees of freedom to the analysis, allowing for fewer false positives and false negatives.

**Table 1.** Primers for real-time RT-qPCR

FlyBase ID (Function)	Primer Sequence (Forward/Reverse)	Amplification Efficiency <sup>a</sup>
<a href="#">FBgn0033102/CG34392 (Epac Guanine Nucleotide Exchange Factor)</a>	CTTACTCAAGGGACTGGTAGAC/ CTCCTTACAGTACGATGGTAGC	1.97
<a href="#">FBgn0002626/CG7939 (Ribosomal protein rp49/RpL32)</a>	CCAAGATCGTGAAGAAGCG/ GTTGGGCATCAGATACTGTC	1.99
<a href="#">FBgn0034354/CG5224 (glutathione transferase)</a>	GAGAAACTGCTTGTACTATGTC/ AGAAACTCGGCGGATTTG	2
<a href="#">Fbgn0037818/CG6465 (aminoacylase)</a>	CCTTGCTAATGGTGTGCTTCG/ CGGGACTTTAGGTTGCTTCTG	2.1
<a href="#">FBgn0023529/CG2918 (ATP binding protein)</a>	CATTGGAAGAAGCATTGGC/ CATAGTGGCGGATTCATCG	1.98
<a href="#">FBgn0040887/CG7418 (Unknown)</a>	AGTGAAACAGGAGTGAGG/ TTGTGGTCAACTTGTAGC	1.98
<a href="#">FBgn0037057/CG10512 (Oxidoreductase)</a>	TGCTTTGCTCCCACCTTC/ CGTCTTCAACTGATTCTCC	1.78
<a href="#">FBgn0039923/CG1793 (RNA polymerase II transcription mediator)</a>	GGAACCTACTACTCACCTATC/ TCATTCTTAGTCCTTCTTCTG	1.89
<a href="#">FBgn0037087/CG7519 (Unknown)</a>	ACTCTATCTTCGTGTTTCG / GATTACTGTTCCCTCTTCGG	1.85
<a href="#">FBgn0036887/CG9231 (Unknown)</a>	TCAGATGTTGCGAGAGAC / AGTGCGATCATTATGTTGG	1.92
<a href="#">FBgn0034614/CG9752 (Unknown)</a>	TACATCCCAATCCAAATCAGC/ ACATCCACGCCTTCCTTG	1.97
<a href="#">FBgn0034199/CG15917 (Unknown)</a>	GCTGGTCTTCCTCCTCAC / GCTGGTATCGTCGTCATCC	1.85

<sup>a</sup>Primer efficiencies were determined from a serial dilution of target DNA using the formula:  $E = 10^{(-1/\text{slope})}$  (Pfaffl 2001; Rasmussen 2001), with the slope determined by the iCycler iQ software.

### Real-time RT-qPCR primer design

Primers were designed based on *D. melanogaster* mRNA sequences obtained from FlyBase that were imported into Beacon Designer software (Premier Biosoft International); a program designed to generate primer pairs suitable for real-time RT-qPCR. The SYBR Green module with program setting 'avoid template structure' was chosen to limit primer sequences to regions of little secondary template structure. Primers were obtained from IDT (Integrated DNA Technologies) and their sequences are shown in Table 1. Both reference and target primers exhibited comparable efficiencies as determined using a dilution series of target DNA. Primer efficiencies were determined from dilution curves using the formula:  $E = 10^{(-1/\text{slope})}$  (Pfaffl 2001; Rasmussen 2001), with the slope determined by the iCycler iQ software (Table 1).

### cDNA synthesis and real-time RT-qPCR

First-strand cDNA synthesis was performed using the iScript cDNA synthesis kit according to the manufacturer's instructions (Bio-Rad Laboratories). Briefly, the reaction was performed with 1.0 µg total RNA in 15 µl RNase-free water, 4 µl 5X iScript reaction mix with a blend of oligo dT and random hexamer primers, and 1 µl iScript reverse transcriptase. The reaction conditions were performed at 25°C for 5 m, 42°C for 30 m, 85°C for 5 m, and the cDNA was stored at 4°C.

Expression of mRNA was analyzed by real-time RT-qPCR using the iCycler iQ detection system (Bio-Rad Laboratories). Samples were performed in triplicate in 25 µl reactions; 12.5 µl iQ SYBR Green Supermix (Bio-Rad Laboratories), 0.2 µM forward and reverse primer, and 11.5 µl of 1:10 diluted cDNA sample. The threshold cycle ( $C_q$ ) is the PCR

cycle at which the fluorescence of the PCR product exceeds an arbitrary threshold. The  $C_q$  of the target transcript in RNA from JH III-challenged S2 cells was compared with the target transcript  $C_q$  generated by RNA from S2 cells treated with MLA. Target gene abundance was normalized to three internal reference transcripts that were shown to be invariant using *BestKeeper* (Pfaffl et al. 2004) and *geNorm* (Vandesompele et al. 2002) software. The RER was calculated as the difference between the  $C_q$  values and was determined using the equation  $2^{-\Delta\Delta C_t}$  as previously modified (Rotenberg et al. 2006). PCR conditions were: 95° C for 3 m, 40 cycles of 95° C, 10 s; 50° C, 45 s and 1 cycle of 95° C, 1 m; 55° C 1 m then followed by a dissociation curve with 80 cycles of 55° C, 10 s with a 0.5° C increase per cycle. To assure PCR accuracy, PCR reaction products were sequenced directly and compared to the expected target sequence.

Statistical analysis of RER values was performed with GraphPad Prism software using the unpaired two-tailed t-test function (GraphPad Software, Inc).

## Results


### Analysis of JH III effect on genomic expression in *Drosophila* S2 cells.

*Drosophila* microarray chips were challenged with RNA from three treatments of S2 cells: (10R) JH III treatment, MLA treatment, or no treatment (no-treatment control). Introduction of lipophilic compounds such as JH III or MLA to culture medium devoid of serum poses a dispersal problem. JH III and MLA are surface active and bind nonspecifically to hydrophobic surfaces (Kramer et al. 1976; Giese et al. 1977). To overcome this problem, we used 0.1% BSA that serves as a carrier molecule to reduce nonspecific binding in all

treatments. MLA is a lipid with physical similarity to JH III but lacks hormonal activity. As shown in Figure 1, MLA has a molecular structure comparable to JH III containing two double bonds and an O-methyl ester. In preliminary experiments, MLA was used as a lipid control for JH I due to the related chemical structure and identical molecular weights (WG Goodman, unpublished data). MLA demonstrated no hormonal activity in a *Manduca sexta* bioassay. In addition, JH I but not MLA induced the expression of hemolymph juvenile hormone binding protein mRNA in *M. sexta* when analyzed by real-time RT-qPCR. MLA was found to have no effect on *D. melanogaster* eclosion success (JR Lindholm and WG Goodman, unpublished). In the present work, MLA was used to control for any potential effects on gene expression caused by the non-specific cellular metabolism of the JH III added to the S2 cells. Comparing genomic expression from JH III treated cells to control cells (Appendix 1 available online) resulted in numerous putative ORFs showing differences. The following criteria were used to identify potentially significant changes between the two treatments: differences in expression  $\geq 2$ -

-fold and P-value  $\leq 0.01$ . Using these criteria, 120 of 14,010 (0.86 %) putative *Drosophila* genes demonstrated differences in expression with 14 ORFs showing an increased expression and 106 ORFs a reduced expression (Appendix 2 available online). Comparing MLA-treated S2 cells to the no-treatment control (Appendix 3 available online) revealed that 63 of 14,010 (0.45%) ORFs displayed significant changes including 10 up-regulated genes and 53 down-regulated genes (Table 2). Comparing RNA from JH III-treated S2 cells to MLA-treated cells (Appendix 4 available online) reduced the number of ORFs demonstrating significant differences as only 32 of 14,010 (0.23%) putative *Drosophila* genes exhibited significant differences with 13 genes showing a  $> 2$ -fold increase and 19 ORFs showing a decreased expression (Table 3).

Most of the JH III up-regulated genes relative to MLA were loci of unknown function (Table 3). However, *Epac* (FBgn0033102), a gene encoding a guanine nucleotide exchange factor of Rap1 small GTPase, showed a  $>3$ -fold increase in expression. FBgn0036313 (a serine/threonine kinase) was induced  $\sim 3$ -fold as were several ORFs with unknown function



**Figure 1.** High quality figures are available online.

**Table 2.** Changes in gene expression between MLA-treated and no-treatment control<sup>a</sup>

Affymetrix Probe Set ID	Sequence Description	Fold Change, MLA to control	P-value, MLA to control	control Raw Data	MLA Raw Data
I41933_at	FBgn0025111 Adenine nucleotide translocase 2 and FBgn0003360 stress-sensitive B	4.3	1.94E-06	0.15	0.67
I42083_at	FBgn0028554 splicing factor 9G8 and FBgn0011737 wee	3.61	3.50E-10	0.37	1.35
I41852_at	FBgn0039809 nucleoside-diphosphate kinase-like and FBgn0000416 Saposin-related	3.21	1.69E-12	0.52	1.71
I41937_at	FBgn0031675 ankyrin and FBgn0027910	3.08	9.65E-04	0.16	0.49
I45883_at	FBgn0031793 protein kinase	2.8	7.93E-03	0.06	0.18
I42630_at	FBgn0036400 DNA repair protein	2.79	9.24E-03	0.05	0.13
I42100_at	FBgn0036448 protein phosphatase-like and FBgn0036447 alpha-1,3-mannosyl-glycoprotein beta-1,4-N-acetylglucosaminyltransferase	2.4	3.54E-04	0.2	0.48
I46000_at	FBgn0031973 serpin	2.33	8.77E-03	0.11	0.25
I42008_at	FBgn0037199 and FBgn0010348 ADP ribosylation factor 79F	2.16	7.60E-04	0.56	1.24
I41868_at	FBgn0032308 voltage-dependent anion-selective channel-like and FBgn0004363 porin	2.05	1.51E-03	0.71	1.49
I50999_at	FBgn0039936	-2.03	1.42E-03	2.01	1.01
I51599_at	FBgn0035128 cell cycle regulator	-2.05	8.26E-09	0.52	0.26
I52387_at	FBgn0027544	-2.07	8.27E-10	0.82	0.4
I45770_at	FBgn0031618	-2.1	6.28E-06	1.08	0.52
I47150_at	FBgn0033833 attacin A-like	-2.11	2.97E-04	0.59	0.29
I52151_at	FBgn0036824 enzyme	-2.13	6.90E-03	2.04	0.98
I43099_i_at	FBgn0000279 Cecropin C	-2.14	1.29E-09	1	0.48
I52312_at	FBgn0034048 glycerol-3-phosphate dehydrogenase-like	-2.16	1.41E-04	1.05	0.5
I54049_at	FBgn0036444	-2.19	8.20E-04	1.18	0.55
I47473_at	FBgn0034407 defense/immunity protein	-2.2	8.79E-03	29.2	13.52
I43770_at	FBgn0014865 Metchnikowin	-2.21	4.52E-05	1.71	0.79
I43164_at	FBgn0000723 Fps oncogene analog	-2.23	5.13E-04	1.02	0.47
I42538_at	FBgn0034341 glutathione transferase	-2.26	8.84E-03	0.63	0.28
I47631_at	FBgn0034647	-2.29	1.37E-05	2.59	1.16
I53684_at	FBgn0000250 cactus	-2.3	8.04E-03	2.02	0.9
I41784_at	FBgn0040997	-2.31	3.45E-04	0.36	0.16
I48352_at	FBgn0035806 peptidoglycan recognition protein	-2.37	6.95E-04	1.76	0.76
I51806_at	FBgn0036732 organic anion transporter	-2.37	3.58E-08	0.92	0.4
I55081_at	FBgn0034108	-2.38	7.55E-04	0.42	0.18
I51937_at	FBgn0037387 sugar transporter-like	-2.42	5.89E-06	0.54	0.23
I54078_at	FBgn0014469 Cytochrome P45-4e2	-2.43	2.03E-03	0.47	0.2
I52188_at	FBgn0020278 locomotion defects	-2.51	3.85E-03	1.22	0.5
I49495_at	FBgn0037583 transcription factor	-2.52	9.77E-03	1.19	0.48
I53942_at	FBgn0030710 transcription factor	-2.53	9.92E-03	0.36	0.15
I43100_f_at	FBgn0000279 Cecropin C	-2.55	5.28E-24	2.78	1.11
I53930_at	FBgn0034354 enzyme	-2.55	3.15E-03	0.52	0.21
I41571_at	FBgn0039211 nucleic acid binding	-2.62	3.30E-05	0.83	0.32
I51520_i_at	FBgn0032854	-2.62	1.19E-06	0.8	0.31
I49552_s_at	FBgn0037682 arylalkylphosphatase	-2.83	5.79E-03	1.17	0.42
I54767_at	FBgn0034491	-2.84	4.20E-03	0.72	0.26
I52918_at	FBgn0037308	-2.88	7.75E-14	1.93	0.68
I43299_at	FBgn0003067 Phosphoenolpyruvate carboxykinase (GTP)	-2.89	1.08E-07	1.86	0.66
I41583_at	FBgn0033980 Cyp6a20	-2.9	4.03E-06	1.46	0.51
I51834_at	FBgn0036371	-2.92	1.86E-04	0.25	0.09
I53894_at	FBgn0039644 ATP-binding cassette transporter	-3.11	1.23E-12	1.38	0.45
I42836_at	FBgn0001223 Heat shock protein 22	-3.22	3.67E-17	1.15	0.36
I43443_at	FBgn0004240 Dipteracin	-3.55	2.40E-05	1.6	0.46
I43098_at	FBgn0000278 Cecropin B	-3.67	3.29E-06	2.15	0.6
I53583_at	FBgn0001224 Heat shock protein 23	-3.81	1.56E-06	1.14	0.31
I48458_at	FBgn0035977 peptidoglycan recognition protein-like	-4.22	3.22E-07	0.85	0.21
I43782_at	FBgn0015037 Cytochrome P45-4p1	-4.23	2.80E-05	0.22	0.05
I51641_at	FBgn0032890 glycerate dehydrogenase-like	-4.44	3.88E-03	0.3	0.07
I43197_at	FBgn0001230 Heat shock protein 68	-4.89	5.77E-04	0.4	0.08
I43096_f_at	FBgn0000276 Cecropin A1	-5.24	5.81E-40	28.57	5.56
I52534_at	FBgn0032889 glycerate dehydrogenase-like	-5.3	4.17E-04	0.25	0.05
I42657_at	FBgn0036658 peptidoglycan recognition protein-like	-5.77	8.73E-11	2.05	0.36
I41374_at	FBgn0012042 Attacin-A	-5.88	9.40E-25	3.86	0.67
I50090_at	FBgn0038530 defense/immunity protein	-5.96	6.68E-11	1.12	0.19
I43097_f_at	FBgn0000277 Cecropin A2	-5.99	9.04E-24	21.86	3.73
I49782_at	FBgn0038059 chaperone	-7.03	2.99E-09	0.77	0.11
I53432_at	FBgn0022073 eukaryotic-initiation-factor-4E binding protein	-8.03	7.72E-11	5.49	0.7
I47220_s_at	FBgn0033959	-9.34	1.22E-25	20.48	2.24
I53731_at	FBgn0001225 Heat shock protein 26	-15.02	3.28E-21	0.72	0.05

<sup>a</sup>ORFs showing an expression difference  $\geq 2.00$  and a P-value  $< 0.01$ .

**Table 3.** Changes in gene expression between (10R) JH III-treated vs. MLA-treated S2 cells<sup>a</sup>

Affymetrix Probe Set ID	Sequence Description	Fold Change (10R) JH III to MLA	P-value, JH III to MLA	MLA Raw Data	(10R) JH III Raw Data
I51791_at	FBgn0033102 Epac: guanyl-nucleotide exchange factor	3.81	4.18E-03	0.12	0.46
I51360_at	FBgn0040887	3.36	2.92E-05	0.92	3.12
I52050_at	FBgn0037057	2.96	3.48E-04	0.27	0.81
I51088_at	FBgn0040603	2.89	5.32E-04	0.17	0.5
I53938_at	FBgn0036313 protein serine/threonine kinase	2.37	1.90E-09	0.58	1.39
I53408_at	FBgn0039414	2.36	2.10E-06	0.71	1.69
I45546_i_at	FBgn0031277	2.31	5.17E-03	1.86	4.34
I52918_at	FBgn0037308	2.31	2.12E-05	0.68	1.59
I52205_at	FBgn0014135 branchless	2.17	3.57E-05	0.4	0.88
I42599_at	FBgn0039011	2.11	1.43E-04	0.61	1.29
I43299_at	FBgn0003067 Phosphoenolpyruvate carboxykinase	2.11	5.70E-03	0.66	1.4
I42434_at	FBgn0035083 calcium binding protein-like	2.06	9.23E-14	2.59	5.39
I53051_at	FBgn0032622	2.04	5.37E-03	0.4	0.82
I53887_at	FBgn0016701 Rab-protein 4	-2.02	3.21E-03	0.84	0.42
I49591_at	FBgn0037759 asparaginase	-2.03	9.24E-03	0.3	0.15
I42100_at	FBgn0036448 protein phosphatase-like and FBgn0036447 alpha-1,3-mannosyl-glycoprotein beta-1,4-N-acetylglucosaminyltransferase	-2.06	2.21E-03	0.48	0.23
I42008_at	FBgn0037199 and FBgn0010348 ADP ribosylation factor 79F	-2.19	6.34E-04	1.24	0.57
I52376_at	FBgn0023529 heat shock protein 70	-2.2	1.80E-03	1.36	0.62
I50989_at	FBgn0039923 transcription factor	-2.24	1.10E-03	1.65	0.74
I49182_at	FBgn0037087	-2.36	2.20E-03	1.15	0.49
I54531_at	FBgn0036887	-2.67	2.86E-03	1.12	0.42
I41933_at	FBgn0025111 Adenine nucleotide translocase 2 and FBgn0003360 stress-sensitive B	-2.79	1.99E-04	0.67	0.24
I41959_at	FBgn0030703 and FBgn0030702	-2.86	5.63E-03	0.49	0.17
I53439_at	FBgn0031094	-3.08	8.01E-03	0.34	0.11
I42083_at	FBgn0028554 splicing factor 9G8 and FBgn0011737 wee	-3.26	8.71E-08	1.35	0.42
I53271_at	FBgn0034614	-3.72	9.61E-03	0.8	0.22
I46252_at	FBgn0032376	-3.88	2.74E-03	0.24	0.06
I47354_at	FBgn0034199	-3.91	4.99E-04	1.9	0.49
I43097_f_at	FBgn0000277 Cecropin A2	-4.25	2.59E-05	3.73	0.88
AFFX-Dros-18S_rRNA_M_at*	D.melanogaster 18S, 5.8S 2S and 28S rRNA genes, complete	-5.11	2.56E-03	12.79	2.52
AFFX-Dros-18S_rRNA_3_at*	D.melanogaster 18S, 5.8S 2S and 28S rRNA genes, complete	-7.8	6.89E-04	13.87	1.79
AFFX-Dros-18S_rRNA_5_at*	D.melanogaster 18S, 5.8S 2S and 28S rRNA genes, complete	-15.94	2.14E-04	5.76	0.36

<sup>a</sup>ORFs showing an expression difference  $\geq 2.00$  and a P-value  $\leq 0.01$ .

\*Accession number M21017



(FBgn0040887, FBgn0037057, and FBgn0040603). Among the down-regulated genes were heat *shock protein 70* (FBgn0023529), a transcription factor (FBgn0039923), and *cecropin A2* (FBgn0000277). A large (-5 to -15-fold) down-regulation of the *Drosophila* 18S rRNA was apparently affected by JH III.

### Verification of microarray analysis

To verify the expression levels derived from the microarray analyses, *Drosophila* S2 cells were treated with (10R) JH III (250 ng/ml) or MLA (250 ng/ml) for 4 h and analyzed for target transcript RER using real-time RT-qPCR. Suitable internal reference gene primers were chosen based on genes that were unaffected by the addition of JH III in the microarray analyses (Appendix 4 available online). These were FBgn0023529, which has ATP binding activity and is involved in response to stress, FBgn0034354, which is a glutathione transferase involved in a toxin defense response, and FBgn0002626 (ribosomal protein *49/RpL32*), which is one of the most commonly employed standards used to normalize gene expression in *Drosophila*. Two criteria were used for reference gene characterization: i) primer efficiencies close to 2.00 (100% efficient); and ii) stable expression in total RNA from MLA-treated and JH III-treated S2 cells. Both reference and target primers exhibited comparable efficiencies as determined using a dilution series of target DNA derived from *D. melanogaster* (Table 1). Reference transcript stability was determined using the *BestKeeper* (Pfaffl et al. 2004) and *geNorm* (Vandesompele et al. 2002) programs. *BestKeeper* is an Excel-based tool designed to determine the correlation between the raw values of real-time RT-qPCR for a particular internal reference gene of interest and the geometric mean (the *BestKeeper Index*) of all of the reference genes tested under various

treatments. The program performs pairwise Pearson correlations between the  $C_q$  values of a candidate gene and the *BestKeeper Index* and reports the measure of the strength of the relationship as an r-value. Ultimately, a strong and significant ( $P < 0.05$ ) correlation (r-value) between the index and the reference RNA candidate determines its stability. The *BestKeeper Index* values were determined from a data set consisting of  $C_q$  values of potential reference transcripts from both treatments (i.e. multiple RNA samples from MLA-treated and JH III-treated S2 cells). The stability of the reference genes *rp49/RpL32*, FBgn0023529, FBgn0034354 and transcripts (as defined by their respective primer sets) was high ( $0.96 > r > 0.73$ ). However, the FBgn0034354 transcript consistently exhibited the lowest correlation to the *BestKeeper Index* and therefore was the least consistent of these three reference RNAs. The raw expression data from each internal reference gene was also analyzed using *geNorm* software (Vandesompele et al. 2002). Average expression stability (M) for the reference genes was less than 0.14, which indicates a high degree of constancy under our experimental conditions. Further, *geNorm* analysis of the optimal number of potential internal reference genes suggested that three reference genes were appropriate for data normalization. We have empirically confirmed the stability and utility of these reference transcripts by calculating the RER of several JH III induced genes individually with all three reference genes and found no statistically significant difference in the RERs (data not shown).

The RER was calculated for several of the loci that were significantly changed upon treatment with JH III (Table 3). For this analysis, RNA was isolated from three independent (10R) JH III- or MLA- treated

cell cultures. Target gene RNA abundances were normalized to the *rp49/RpL32* reference transcript. The RER was calculated using control (MLA treatment) RNAs as a calibrator. The mean of normalized target gene abundance from all three MLA-treated samples was calculated and this value was designated as the ‘calibrator’. Next, the individual target gene abundances from both the JH III- and MLA-treated samples were divided by the calibrator. This allowed the calculation of a RER for each sample replication which was used to compare the means of the treatment RERs statistically. Relative to the MLA-treated cells, the RER of all three JH III up-regulated transcripts were confirmed to be significantly increased (Table 4). *Epac* and two loci of unknown function (FBgn0040887 and FBgn0037057) were tested. In contrast, only one of six genes (FBgn0034199) indicated to be down regulated by the microarray

data was confirmed to be significantly reduced by real-time RT-qPCR (Table 4). *Heat shock protein 70* (FBgn0023529) was predicted to be reduced 2.2-fold by JH III action from microarray data but was found to be increased 1.35-fold by real-time RT-qPCR analysis. The remaining four predicted down-regulated transcripts were not found to be statistically different ( $P > 0.05$ ) from the MLA control by two-tailed t-test (Table 4)

## Discussion

Microarray analyses indicated that the expression of a number of genes was modified positively or negatively in response to a JH III challenge of *Drosophila* S2 cells (Appendix 1); however, when MLA was tested under identical conditions, expression levels for many genes displayed a similar profile for both JH III and MLA (Appendices 1 and 3).

**Table 4.** Analysis of changes in gene expression between (10R) JH III treated and MLA-treated control<sup>a</sup>

Affymetrix Probe Set ID	Sequence Description	Array Fold Change (10R) JH III to MLA	Array P-value, (10R) JH III to MLA	Real-time RT-qPCR Fold Change	Real-time RT-qPCR P-value (two tailed t-test)
151791_at	FBgn0033102 Epac: guanyl-nucleotide exchange factor	3.81	4.18E-03	2.15	2.48E-02
151360_at	FBgn0040887	3.36	2.92E-05	1.46	3.60E-03
152050_at	FBgn0037057	2.96	3.48E-04	1.95	6.00E-03
150989_at	FBgn0023529 heat shock protein 70	-2.24	1.10E-03	-1.2	4.40E-01
149182_at	FBgn0039923 transcription factor	-2.36	2.20E-03	1.23	6.19E-02
154531_at	FBgn0037087	-2.67	2.86E-03	-1.22	5.63E-02
153271_at	FBgn0036887	-3.72	9.61E-03	-1.11	1.17E-01
147354_at	FBgn0034614	-3.91	4.99E-04	-1.23	1.43E-02

<sup>a</sup>ORFs showing a significant expression difference ( $P \leq 0.05$ ) as initially indicated by microarray and confirmed by real-time RT-PCR. Numbers in red indicate a significant increase in expression and numbers in green indicate a significant down-regulation. Numbers in black indicate no significant change.

Less than a 0.5% of the genes examined by microarray analyses demonstrated a highly significant altered level of expression in the presence of JH III but not in the presence of MLA (Appendix 3, Table 4). The up-regulated genes included *Epac*, several loci of unknown function, a serine/threonine kinase, *branchless*, and *phosphoenolpyruvate carboxykinase*. Down-regulated genes included several genes of unknown function, a pre-mRNA splicing factor, *cecropin A2* and *18S rRNA*.

Despite the advantages of microarray technology, real-time RT-qPCR remains the most accurate method to analyze mRNA expression and to verify key relationships identified by microarray analysis (Hembruff et al. 2005). For real-time RT-qPCR to quantitatively assess expression levels of target mRNAs, the selection of appropriate internal reference genes is critical (Vandesompele et al. 2002; Pfaffl et al. 2004; Bustin et al. 2009). It is becoming increasingly evident that reliance on a single reference transcript may lead to significant errors in the analysis of target gene expression (Vandesompele et al. 2002; Pfaffl et al. 2004). Three internal reference RNAs (rp49/RpL32 [FBgn003461 4], FBgn0034354, FBgn0023529) were selected due to their stable expression in S2 cells treated with either JH III or MLA in the microarray analysis. Using three reference genes, as little as a 17% difference in the mean transcript relative expression ratios was shown to be statistically significant (Rotenberg et al. 2006). All three up-regulated genes that we analyzed (*Epac*, FBgn0040887 and FBgn0037057) were confirmed to be up-regulated by real-time RT-qPCR (Table 4). However, only one of six genes predicted to be down-regulated by the addition of JH III was confirmed to be statistically reduced by real-time RT-qPCR. This result illustrates the

absolute need to confirm microarray data by real-time RT-qPCR analysis before time and resources are expended on further analysis (Morey et al. 2006). The relative inability of microarrays to identify down-regulated genes has been noted previously (Beckman et al. 2004; Morey et al. 2006). This phenomenon relates to the decreased reliability and increased variability in the detection of spots on the microarray exhibiting reduced fluorescence (Beckman et al. 2004).

An important consideration with any expression analysis is the choice of the conditions that will be used as the control for the microarray or real-time RT-qPCR analysis. In this study, we used MLA, a lipid physically similar to JH III but without known hormonal activity, as our control treatment of S2 cells. Another recent microarray analysis of genes that are JH III induced in both *Drosophila* and honey bee, used dimethyl sulfoxide (DMSO), the carrier for JH III in the experiments, as a control treatment (Li et al. 2007). It has been previously shown that DMSO significantly increases juvenoid activity when used as a solvent in a bioassay on *Dysdercus cingulatus* (Sláma 1974). A search of online data (Appendices 1, 2, and 3) revealed that all of the genes described in Li et al. (2007) as JH III inducible were also induced by MLA in our microarray analysis (Appendix 5 available online). This raises the possibility that the induction of these loci may be influenced by the metabolism of the JH III lipid backbone. Due to the increased expression in the MLA control, this set of genes was not identified as JH III inducible (Appendix 5 available online). Since the raw microarray data and the conditions used for microarray analysis in Li et al. (2007) have not as yet been published, the reason for the failure of the previous analysis to identify the genes that were found to be JH III inducible in

S2 cells (Table 3, Table 4) is not evident. It may be that DMSO treatment simulates JH III induction to some extent in S2 cells thereby masking the induction, or that this set of genes was induced in S2 cells but not in honey bee by JH III treatment (induction in both insects was a criterion for analysis in the Li et al. 2007 paper). Preliminary real-time RT-qPCR data confirming the specific induction of *Epac* by JH III in both S2 cells and *Drosophila* third instars (Wang et al. 2009) confirms this aspect of our microarray analysis.

In summary, this work details the analysis by microarray of genome-wide gene expression alterations induced by treatment of *Drosophila* S2 cells with JH III. The comparison of JH III treatment to treatment with MLA, a structurally similar but hormonally inactive lipid, revealed only 32 of 14,010 loci responded differentially by microarray analysis. Up-regulated genes were confirmed by real-time RT-qPCR but most predicted down-regulated genes failed confirmation. This indicates that a remarkably small number of genes were specifically affected by JH III. The most intriguing gene that was confirmed to increase expression following JH III treatment was *Epac* that demonstrated highly significant up-regulation in the presence of JH III ( $\geq 3$ -fold) but was refractory to MLA. *Epac*, an exchange factor directly activated by cAMP, is a direct target for cAMP and a guanine-nucleotide exchange factor for the small GTPase, Rap1 (Bos 2005). This suggests that induction of *Epac* expression may be a major component of the JH III hormone's activity in insect development.

### Acknowledgements

This work was supported in part by the Wisconsin Agricultural Experiment Station

(WIS0122) (WGG) and the Alex and Lillian Fier Distinguished Graduate Fellowship (JW).

### References

- Beckman KB, Lee KY, Golden T, Melov S. 2004. Gene expression profiling in mitochondrial disease: assessment of microarray accuracy by high-throughput Q-PCR. *Mitochondrion* 4(5-6): 453-70.
- Bos JL. 2005. Linking *Rap* to cell adhesion. *Current Opinion in Cell Biology* 17(2): 123-128.
- Bustin SA. 2005. Real-time, fluorescence-based quantitative PCR: a snapshot of current procedures and preferences. *Expert Review of Molecular Diagnostics* 5(4): 493-498.
- Bustin, S.A., Benes, V., Garson, J.A., Hellems, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M.W., Shipley, G.L., Vandesompele, J. and Wittwer, C.T., 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.* 55, 611-22.
- Cusson M, Miller D, Goodman WG. 1997. Characterization of antibody 444 using chromatographically purified enantiomers of juvenile hormones I, II, and III: implications for radioimmunoassays. *Analytical Biochemistry* 249(1): 83-87.
- Gade G, Hoffmann KH, Spring JH. 1997. Hormonal regulation in insects: facts, gaps, and future directions. *Physiological Reviews* 77(4): 963-1032.
- Giese C, Spindler KD, Emmerich H. 1977. The solubility of insect juvenile hormone in aqueous solutions and its adsorption by

glassware and plastics. *Zeitschrift Naturforschungen* 32(1): 158-160.

Goodman WG, Granger NA. 2005. The Juvenile Hormones. In: Gilbert LI, Iatrou K, Gill SS, editors. *Comprehensive Molecular Insect Science*. pp. 319-409. Elsevier/Pergamon.

Hembruff SL, Villeneuve DJ, Parissenti AM. 2005. The optimization of quantitative reverse transcription PCR for verification of cDNA microarray data. *Analytical Biochemistry* 345(2): 237-249.

Jones G. 1995. Molecular mechanisms of action of juvenile hormone. *Annual Review of Entomology* 40: 147-169.

Kethidi DR, Li Y, Palli SR. 2006. Protein kinase C mediated phosphorylation blocks juvenile hormone action. *Molecular and Cellular Endocrinology* 247(1-2): 127-134.

Kramer KJ, Dunn PE, Peterson R, Law JH. 1976. Interaction of juvenile hormone with binding proteins in insect hemolymph. In: Gilbert LI, editor. *The Juvenile Hormones*. pp. 327-341. Plenum Press.

Lafont R. 2000. Understanding insect endocrine systems: Molecular approaches. *Entomologia Experimentalis et Applicata* 97(2): 123-136.

Li Y, Zhang Z, Robinson GE, Palli SR. 2007. Identification and characterization of a juvenile hormone response element and its binding proteins. *Journal of Biological Chemistry* 282(52): 37605-37617.

Mangelsdorf DJ, Evans RM. 1995. The RXR heterodimers and orphan receptors. *Cell* 83(6): 841-850.

Morey JS, Ryan JC, Van Dolah FM. 2006. Microarray validation: factors influencing correlation between oligonucleotide microarrays and real-time PCR. *Biological Proceedings Online* 8: 175-93.

Nijhout HF. 1994. *Insect Hormones*. Princeton University Press.

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research* 29(9): 2002-2007.

Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP. 2004. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper-Excel-based tool using pair-wise correlations. *Biotechnology Letters* 26(6): 509-515.

Pszczolkowski MA, Peterson A, Srinivasan A, Ramaswamy SB. 2005. Pharmacological analysis of ovarian patency in *Heliothis virescens*. *Journal of Insect Physiology* 51(4): 445-453.

Rajagopalan D. 2003. A comparison of statistical methods for analysis of high density oligonucleotide array data. *Bioinformatics* 19(12): 1469-1476.

Rasmussen R. 2001. Quantification on the LightCycler instrument. In: Meurer S, Wittwer C, Nakagawara K, editors. *Rapid Cycle Real-Time PCR: Methods and Applications*. pp. 21-34. Springer-Verlag Press.

Riddiford LM. 1996. Molecular aspects of juvenile hormone action in insect metamorphosis. In: Gilbert LI, Tata JR,

Atkinson BG, editors. *Metamorphosis*. pp. 223-251. Academic Press.

Rotenberg D, Thompson TS, German TL, Willis DK. 2006. Methods for effective real-time RT-PCR analysis of virus-induced gene silencing. *Journal of Virological Methods* 138(1-2): 49-59.

Sevala VL, Davey KG. 1989. Action of juvenile hormone on the follicle cells of *Rhodnius prolixus* - Evidence for a novel regulatory mechanism involving protein kinase C. *Experientia* 45(3): 355-356.

Sláma K. 1974. The chemistry and physiology of juvenoids. In: Sláma K, Romanuk M, Sorm F, editors. *Insect Hormones and Bioanalogues*. pp. 90-302. Springer-Verlag.

Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* 3(7): Research0034.

Wheeler DE, Nijhout HF. 2003. A perspective for understanding the modes of juvenile hormone action as a lipid signaling system. *Bioessays* 25(10): 994-1001.

Wang J, Lindholm JR, Willis DK, Orth A, Goodman WG. 2009. Juvenile hormone regulation of *Drosophila* Epac--a guanine nucleotide exchange factor. *Molecular and Cellular Endocrinology* 305(1-2): 30-7.

Yamamoto K, Chadarevian A, Pellegrini M. 1988. Juvenile hormone action mediated in male accessory glands of *Drosophila* by calcium and kinase C. *Science* 239(4842): 916-919.