



Amplified fragment length polymorphism mapping of quantitative trait loci for economically important traits in the silkworm, *Bombyx mori*

Seyed Z Mirhoseini^{1a}, Babak Rabiei^{2b}, Payam Potki^{3c}, and Seyed B Dalirsefat^{4d*}

¹Department of Animal Science, Faculty of Agricultural Sciences, University of Guilan, PO Box 41635-1314, Rasht, Guilan, Iran

²Department of Agronomy & Plant Breeding, Faculty of Agricultural Sciences, University of Guilan, PO Box 41635-1314, Rasht, Guilan, Iran

³Department of Genomics, Agricultural Biotechnology Research Institute of Iran (Rasht), PO Box 41635-4115, Rasht, Guilan, Iran

⁴Department of Sericulture, Faculty of Agricultural Sciences, University of Guilan, PO Box 41635-1314, Rasht, Guilan, Iran

Abstract

Cocoon related characteristics are economically important traits in the silkworm, *Bombyx mori* L. (Lepidoptera: Bombycidae). In this study a genetic linkage map was developed that identified QTL controlling the cocoon weight, cocoon shell weight, and cocoon shell percentage using 161 amplified fragment length polymorphism (AFLP) markers. Twenty *Pst*I/*Taq*I primer combinations were employed to genotype 78 F₂ progenies derived from a cross between P107 Japanese inbred line and Khorasan Lemon Iranian native strain. Among polymorphic markers, 159 AFLP markers were assigned to 24 linkage groups at the LOD threshold of 2.5 that varied in length from 4 to 299 cM. The total length of the linkage map was 2747 cM, giving an average marker resolution of 19.31 cM. A total of 21 AFLP markers were identified that were distributed over the ten linkage groups linked to the three studied traits using the composite interval mapping method. The explained variation rate by QTL controlling cocoon weight, cocoon shell weight, and cocoon shell percentage ranged from 0.02% to 64.85%, 0.2% to 49.11%, and 0.04% to 84.20%, respectively. These QTL controlled by different actions as well as under dominance, additive, partial dominance, dominance, and over dominance.

Keywords: AFLP markers, cocoon traits, QTL mapping

Abbreviations: **ABRII**, Agricultural Biotechnology Research Institute of Iran; **AFLP**, Amplified Fragment Length Polymorphism; **AREO**, Agricultural Research and Education Organization; **BC**, Backcross; **CIM**, Composite Interval Mapping; **cM**, centimorgans; **ESTs**, Expressed Sequence Tags; **IM**, Interval Mapping; **ISRC**, Iran Silkworm Research Center; **LG**, Linkage Group; **LOD**, Logarithm of Odds; **LRT**, Likelihood Ratio Test; **NPV**, Nuclear Polyhedrosis Virus; **PCR**, Polymerase Chain Reaction; **QTL**, Quantitative Trait Loci; **RAPD**, Random Amplified Polymorphic DNA; **RFLP**, Restriction Fragment Length Polymorphism; **SADF**, Selective Amplification of DNA Fragments; **SNP**, Single Nucleotide Polymorphism

Correspondence: ^a mirhosin@guilan.ac.ir, ^b rabiei@guilan.ac.ir, ^c payam_572003@abrii.ac.ir,

^d* bendalir@guilan.ac.ir, *Corresponding author

Associate Editor: Craig Coates was editor of this paper.

Received: 7 April 2009, **Accepted:** 2 May 2010

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 10, Number 153

Cite this paper as:

Mirhoseini SZ, Rabiei B, Potki P, Dalirsefat SB. 2010. Amplified fragment length polymorphism mapping of quantitative trait loci for economically important traits in the silkworm, *Bombyx mori*. *Journal of Insect Science* 10:153 available online: insectscience.org/10.153

Introduction

The silkworm, *Bombyx mori* L. (Lepidoptera: Bombycidae), domesticated for silk production for about 5000 years, is an agriculturally important insect and comprises a large number of geographical races and inbred lines that show substantial variation in their qualitative and quantitative traits (Mirhoseini et al. 2007). With the establishment of stable transformation (Yamao et al. 1999; Tamura et al. 2000), *B. mori* has shown the potential to produce pharmaceutically important proteins in high yield (Tomita et al. 2003), opening up new applications for sericulture in medical, agricultural, and industrial fields (Yamamoto et al. 2006). Currently, it is the major economic resource for nearly 30 million families in countries such as China, India, Vietnam, and Thailand (Miao et al. 2005). In order to make sericulture economically viable, genes affecting growth rate, yield, fiber quality, and virus resistance can be tagged with molecular markers for rapid construction of genetically improved strains. Taking the exclusive investigational advantages of this organism into account, as well as its economic importance, an International Consortium on Lepidopteran Genomics was formed a few years ago to support international cooperation to sequence the genome of *B. mori* and to initiate comparative genomics of other economically important Lepidoptera (Nagaraju and Goldsmith 2002).

Most traits in nature and of importance to agriculture are quantitatively inherited and therefore are difficult to study due to the complex nature of their inheritance. However, recent advances of genomic technologies have led to revolutionary means for unraveling the secrets of genetic variation in quantitative

traits. Genomic technologies allow the molecular characterization of polymorphic markers throughout the entire genome that are then used to identify and map the genes or quantitative trait loci (QTL) underlying a quantitative trait based on linkage analysis (Wu et al. 2007).

A complete linkage map is necessary to efficiently carry out molecular-based analyses such as molecular marker-assisted selection, quantitative trait loci (QTL) mapping of agronomically important traits, prediction of heterosis, and comprehensive investigations of genomic evolution between lineages (Tan et al. 2001). Presently, genome studies in *B. mori* have generated genetic linkage maps based on morphological markers (Doira et al. 1992) and molecular markers including RFLP (Goldsmith 1991; Shi et al. 1995; Nguu et al. 2005), RAPD (Promboon et al. 1995; Yasukochi 1998; Li et al. 2000), SADF and RAPD (He et al. 2001), AFLP (Tan et al. 2001; Lu et al. 2004; Sima et al. 2006), microsatellites (Miao et al. 2005), and SNP (Yamamoto et al. 2006).

Projects have been initiated to find molecular markers that are tightly linked to traits relevant for sericulture, with the related goals of developing tools for marker assisted selection and positional cloning. RAPD or cDNA markers have been associated with the four known densovirus nonsusceptibility loci, *nsd-1* (Abe et al. 1998), *nsd-2* (Abe et al. 2000), *Nid-1* (Kadono-Okuda et al. 2003), and *nsd-Z* (Li et al. 2001). Two large contigs on chromosome 17 that encompass cDNAs closely linked to *Nid-1* and *nsd-2* have been isolated and sequenced (Kadono-Okuda et al. 2003) and are being examined for candidate genes in susceptible and nonsusceptible strains (Goldsmith et al.

2005). A similar strategy was used to screen for RAPD markers linked to resistance to NPV, a potentially devastating pathogen (Yao et al. 2003), and fluoride resistance (Chen et al. 2003). Progress has also been made in assigning RAPDs (Chatterjee and Pradeep 2003), inter-simple sequence repeat markers (Chatterjee and Mohandas 2003), and AFLP (Lu et al. 2004; Li et al. 2006; Sima et al. 2006) to QTL for characters such as larval growth rate and pupal and cocoon weight. A suite of additional fingerprinting tools has been developed for these applications (Nagaraju and Goldsmith 2002). In addition, a collection of about 8500 expressed sequence tags (ESTs) is now available in GenBank and provides an additional source of important anchors in the ongoing *Bombyx* genome study (Nguu et al. 2005).

Although for genome mapping, the ideal genetic marker is codominant, multiallelic, and hypervariable (i.e., segregates in almost every family), some dominant markers are also very useful and powerful in particular situations (Wu et al. 2007). The amplified fragment length polymorphism (AFLP) technique (Zabeau and Vos 1992; Vos et al. 1995) has demonstrated to be a convenient and reliable tool to generate highly polymorphic molecular markers that greatly facilitate building linkage maps (Qi et al. 1997; Waugh et al. 1997). AFLP markers do allow one to construct linkage maps with wide genome coverage without engaging in extensive sequencing or marker development programs. AFLP markers are also faster than individual codominant marker types because a single polymerase chain reaction (PCR) can derive multiple loci simultaneously (Erickson et al. 2004). Because of these features, AFLP has been widely employed for genetic mapping in various organisms.

In this study, significant molecular markers and a large segregating population size were employed to detect QTL linked to economically important traits relevant to the *B. mori* cocoon and to better identify the genome regions of these QTL. Since high-resolution QTL mapping is critical for positional cloning and gene isolation (Zhong et al. 2006), a high resolution AFLP-based genetic linkage map and the results of QTL mapping for economically important cocoon traits are reported.

Materials and Methods

Insect materials and crosses

One F₂ segregating family from mating between a Japanese inbred line (P107) as female parent and an Iranian native strain (Khorasan Lemon) as male parent were used in the study. These inbred line and strain exhibit high phenotype diversity for economically important characters such as whole cocoon weight, cocoon shell weight, and cocoon shell percentage, suggesting that considerable polymorphism exists at the DNA level (Dalirsefat and Mirhoseini, 2007). Indeed, the highest and the least quantities of these traits corresponded to P107 and Khorasan Lemon, respectively. These inbred line and strain have undergone a high degree of inbreeding and are relatively homozygous. A total number of 78 progenies, including 39 males and 39 females from F₂ population, were used to construct the genetic linkage map and QTL detection. The parents and F₁ progenies were used to establish the segregation pattern of the molecular markers. The crossing experiments were done in the Iran Silkworm Research Center (ISRC) located in Rasht, Iran.

AFLP analysis

Genomic DNAs were isolated individually from all the parents, F₁, and F₂ populations, in the moth stage following the phenol/chloroform method (Suzuki et al. 1972) and as modified by Nagaraja and Nagaraju (1995). DNAs were quantified using a known standard (DNA lambda, Roche, www.roche.com) on agarose gels.

All individuals were subjected to genotyping with AFLP markers according to Vos et al. (1995) with some modifications. Briefly, genomic DNA was double digested with *Pst*I and *Taq*I restriction enzymes, which produce polymorphic DNA fragments in *B. mori* (Tan et al. 2001; Mirhoseini et al. 2007). The DNA fragments were ligated with *Pst*I and *Taq*I adaptors, generating template DNA for PCR amplification. Two primers were designed on the basis of adaptor sequences and restriction site sequences to use in PCR amplification. Selective nucleotide sequences were added to the 3' end of each primer. PCR amplification was conducted in two steps: a pre-amplification and a selective amplification. For the selective amplification, a total of 81 primer combinations obtained from two sets of *Pst*I and *Taq*I selective primers (Table 1) were screened. Among them, 20 primer pairs that produced fragments with clear dominance inheritance patterns and reproducibility were used for the linkage analysis. Polymorphism screening of AFLP products was conducted on a 6% polyacrylamide gel using SequiGen 38×30 cm gel apparatus (BioRad Laboratories Inc., www.bio-rad.com). Bands were detected by the silver staining procedure (Promega, www.promega.com, Technical manual No.023), and gel images were scanned and saved as jpeg files for scoring and further analysis.

Linkage analysis and map construction

Using genotype information of 81 AFLP primer combinations, 20 primer combinations that produced clearly readable and polymorphic fragments among parents were employed to analyze linkage mapping. Twenty polymorphic primer combinations generating 161 polymorphic AFLP fragments with a clear dominance inheritance pattern were employed to construct the linkage map and detect QTL; that is, the suitable fragments must show complete dominance expression in one parent and complete recessive expression in the other, and all F₁ individuals must be heterozygous. The AFLP fragments were scored based on 0 and 1 and then converted to A, B, C, and D letters according to the Map manager QTX (Manly et al. 2001) instruction manual. The data were analyzed using the Kosambi map function (Kosambi 1944) of Map manager QTX (Manly et al. 2001) to develop a linkage map for the population. By genotyping 78 progenies from the F₂ population using 161 polymorphic bands, a genotypic data matrix in a dimension of 78 × 161 was constructed and used for linkage mapping. First recombination rates among markers were evaluated, and then recombination rates converted to the map distance based on centiMorgan using the Kosambi map function (Kosambi 1944). Computer software QTL cartographer version 2.5 (Wang et al. 2007) was used to determine the QTL positions, the expected additive and dominance effects, and the phenotypic variance explained by individual QTL. The LOD threshold value for declaring the presence of a QTL was determined by a permutation test ($n = 1000$) (Churchill and Doerge 1994). Genome-wide threshold levels were used to declare significant QTL based at the 5% significance level. Average levels of dominance (h) were estimated using the ratio dominance/additive effects (Stuber et al. 1987).

All molecular experiments were established in the genomics laboratory of the Agricultural Biotechnology Research Institute of North Region (Rasht) under the supervision of the Agricultural Biotechnology Research Institute of Iran (ABRII).

Results

Linkage map construction

Among the 81 AFLP primer combinations screened, approximately one-third of the primer combinations ($n = 28$) produced polymorphic fragments between the P107 inbred line and the Khorasan Lemon native strain. Twenty pairs of AFLP primer combinations were selected for segregation analysis on the F2 population based on reproducibility and the degree of polymorphism. Only the polymorphic fragments that segregated in a dominant

manner and could be scored unambiguously were used for linkage map construction. An example of AFLP gel electrophoresis and polymorphism screening related to the P_{ta}-T_{ta} primer combination is shown in Figure 1.

Twenty *Pst*I/*Taq*I primer combinations produced 845 clearly detected bands, of which 161 qualified polymorphic fragments showing good agreement of 3:1 segregation (for a dominant marker, the segregation ratio is 3:1 in the F2 population) were analyzed for linkage mapping. The frequency of polymorphic AFLP markers derived from the clearly detected bands in the P107 × Khorasan Lemon cross in the silkworm was 19.35%. This frequency was close to that obtained in the Dazao × C₁₀₀ cross of the silkworm (25.7%) (Lu et al. 2004) but it was dramatically lower than in the no. 782 × od100 cross (60.7%) of the silkworm (Tan et al. 2001).

Table 1. Adapters and primers used in AFLP analysis.

	Name	Sequence	
Adapters <i>Pst</i> I	<i>Pst</i> top strand	5'-GACGTGACGGCCGTCATGCA	
	<i>Pst</i> bottom strand	5'-TGACGGCCGTCACG	
Adapters <i>Taq</i> I	<i>Taq</i> top strand	5'-GACGATGAGTCCTGAG	
	<i>Taq</i> bottom strand	5'-CGCTCAGGACTCAT	
Primers <i>Pst</i> I ^a	P01	5'-GACGGCCGTCATGCAG	
	P21	5'-GACGGCCGTCATGCAG TA	
	P22	5'-GACGGCCGTCATGCAG AT	
	P23	5'-GACGGCCGTCATGCAG TC	
	P24	5'-GACGGCCGTCATGCAG AC	
	P31	5'-GACGGCCGTCATGCAG AAC	
	P32	5'-GACGGCCGTCATGCAG AGA	
	P33	5'-GACGGCCGTCATGCAG ATG	
	P34	5'-GACGGCCGTCATGCAG AAG	
	P35	5'-GACGGCCGTCATGCAG TAT	
	Primers <i>Taq</i> I	T01	5'-GATGAGTCCTGAGCGA
		T21	5'-GATGAGTCCTGAGCG ATA
		T22	5'-GATGAGTCCTGAGCG AAT
		T23	5'-GATGAGTCCTGAGCG ATC
T24		5'-GATGAGTCCTGAGCG ATG	
T31		5'-GATGAGTCCTGAGCG AAAT	
T32		5'-GATGAGTCCTGAGCG ACA	
T33		5'-GATGAGTCCTGAGCG AAG	
T34		5'-GATGAGTCCTGAGCG AGC	
T35		5'-GATGAGTCCTGAGCG ATAC	

^a Selective nucleotides shown as bold letters

91 fragments of 161 polymorphic fragments (56.52%) were detected in the male parent (Khorasan Lemon strain), and 70 fragments (43.48%) were observed in the female parent (P107 inbred line). On average, each primer combination generated 8.05 polymorphic fragments that could be used for linkage mapping. The number of polymorphic bands produced using the 20 primer combinations ranged from 3 bands (7.32%) corresponding to P33-T32 to 14 bands (23.33%) for P22-T31 (Table 2).

The linkage map generated from the P107 × Khorasan Lemon cross contained 159 AFLP markers (two markers were unlinked) that were assigned to 24 linkage groups at the

LOD threshold of 2.5 (Figure 2). Average distance between markers was 19.31 cM. The total recombination distance over 24 linkage groups was 2747 cM, which was longer than previous estimates in *B. mori*, i.e. 1800 cM for the dense RAPD map (Yasukochi 1998), 1868.10 cM and 2677.50 cM for the AFLP maps in two F₂ subgroups (Sima et al. 2006), and 1305 cM for SNPs based linkage map (Yamamoto et al. 2006). However, it was shorter than 6512 cM (Tan et al. 2001) and 3676.7 cM (Li et al. 2006) for the AFLP maps and 3431.9 cM (Miao et al. 2005) for the SSR markers reported in backcrossed populations of *B. mori*. Miao et al. (2005) suggested that although many conditions influence map length, including differences in mating




Figure 1. An example of AFLP gel electrophoresis and polymorphism screening corresponding to the P_{at}-T_{tac} primer combination. GM-107 and GF-KL represent female and male parents, respectively. M stands for standard molecular size marker. Polymorphic bands are shown by arrows. High quality figures are available online.

strategy and strains used, the distribution of markers is a possible causative aspect, and increased marker density should converge on a more realistic map length value. As Tan and Ma (1998) demonstrated theoretically, with additional markers typed, the map length may increase when marker density is not saturated or may decrease when marker density is in a saturation state (Tan et al. 2001). For example, Causse et al. (1994) constructed a rice map with 762 markers covering 4026.3 cM, whereas Harushima et al. (1998) obtained a 2275-marker genetic map of rice covering 1521.6 cM. This may explain why the length of our AFLP map is more than that of the *B. mori* linkage map studies mentioned above except for the maps of Tan et al. (2001), Miao et al. 2005, and Li et al. (2006).

Considering that the estimated genome size of *B. mori* is 530 Mbp (Gage 1974), the average physical distance per recombination distance

is about 193 kb/cM. It seems that the AFLP markers did not exhibit significant clustering near centromeres or the distal region of chromosomes, suggesting that they provide good coverage of the genome (Li et al. 2006, Figure 2).

Phenotypic values

The average phenotypic values of cocoon weight, cocoon shell weight, and cocoon shell percentage traits corresponding to each parent and F₂ offspring are shown in Table 3. An extremely high significant difference ($p < 0.01$) for these traits was revealed as a result of comparing the mean phenotypic values between parents using the *t*-test. The mean cocoon weight in line P107 as female parent and Khorasan Lemon as male parent was 1.479 and 1.404 g, respectively. The mean cocoon shell weight in line P107 was 0.324 g, approximately 0.113 g more than that in Khorasan Lemon (0.211 g). In addition, the

Table 2. The observed polymorphisms of twenty *TaqI* and *PstI* primer combinations used in the parents and F₂ population.


Primer name ^a	Total number of bands	Number of polymorphic bands in parents	Number of polymorphic bands in each parents		Observed polymorphism rate
			P107	Khorasan Lemon	
P31-T34	43.00	11.00	5.00	6.00	25.58
P31-T24	44.00	5.00	0.00	5.00	11.36
P22-T31	60.00	14.00	4.00	10.00	23.33
P33-T32	41.00	3.00	2.00	1.00	7.32
P35-T24	38.00	5.00	3.00	2.00	13.16
P32-T33	46.00	7.00	2.00	5.00	15.22
P21-T32	42.00	10.00	6.00	4.00	23.81
P31-T23	50.00	6.00	3.00	3.00	12.00
P24-T33	40.00	8.00	5.00	3.00	20.00
P23-T32	50.00	8.00	5.00	3.00	16.00
P35-T21	45.00	10.00	4.00	6.00	22.22
P35-T35	45.00	7.00	5.00	2.00	15.56
P35-T34	53.00	11.00	7.00	4.00	20.75
P33-T34	45.00	4.00	0.00	4.00	8.89
P31-T22	32.00	11.00	4.00	7.00	34.38
P34-T32	42.00	13.00	4.00	9.00	30.95
P22-T32	35.00	8.00	2.00	6.00	22.86
P22-T34	33.00	8.00	3.00	5.00	24.24
P35-T33	32.00	6.00	2.00	4.00	18.75
P21-T33	29.00	6.00	4.00	2.00	20.69
Total	845.00	161.00	70.00	91.00	387.07
Average	42.25	8.05	3.50	4.55	19.35

^a Primer combinations and sequences shown in Table 1.

cocoon shell percentage in the line P107 was estimated to be 22.23%, which was seven percent more than that in Khorasan Lemon (15.23%) as the male parent. Except for cocoon weight value in the F₂ population, which was higher than both parent values, both the mean cocoon shell weight and cocoon shell percentage traits in the F₂

generation were closely equal to the mean parent values (Table 3).

The frequency distribution of phenotypic values related to each trait studied in the F₂ offspring is illustrated in Figure 3. As it is shown, a number of F₂ offspring demonstrated out of range parent values, especially in the



line P107 for the three studied traits. The least and the highest phenotypic values of cocoon weight in the F₂ population were 1.13 g and 1.83 g, respectively, and the highest value was about 0.35 g more than that in the line P107 (1.479 g). The least and the highest phenotypic values of cocoon shell weight in the F₂ population were 0.22 g and 0.37 g, respectively; consequently, a number of the F₂ offspring had almost 0.046 g more than that in the line P107 phenotypic value (0.324 g). In addition, the least and the highest phenotypic values of cocoon shell percentage in the F₂ progenies were 15.72% and 22.6%, respectively; the highest value was nearly 0.4% higher than that in the line P107 (22.2%) (Figure 3).

QTL analysis

A total of 21 different loci, including QTL, controlling cocoon weight, cocoon shell weight, and cocoon shell percentage traits were detected in the linkage map using a composite interval mapping method at the LOD threshold of 2.5 (Table 4). The selected LOD score plots at the threshold of 3 for linkage groups with the identified QTL provided a basis for identifying the molecular markers most closely linked to the QTL (Figure 4).

In particular, 12 QTL controlling cocoon weight were identified on the LG1, LG5, LG6, LG8, LG16, LG17, and LG19. The additive effects of these QTL ranged from -0.1581 (*cw8*) to +0.0887 (*cw19b*), and their dominance effects ranged from -0.3852 (*cw19a*) to +0.3881 (*cw1a*). Two QTL for

cocoon shell weight were identified and located on the LG16 and LG22 with additive effects ranging from -0.0164 (*cshw16c*) to +0.0459 (*cshw22b*) and dominance effects




Figure 3. Frequency distribution of phenotypic values related to the studied traits in the F₂ segregating *Bombyx mori* population derived from mating between the P107 inbred line and the Khorasan Lemon native strain (the mean phenotypic values of the parents for the three traits are shown in Table 3). High quality figures are available online.

Table 3. Mean phenotypic values of traits studied in the parents and F₂ population.

Trait	Parent		F ₂	Difference between parents (t-value)
	Line P107	Khorasan Lemon		
Cocoon weight (g)	1.479±0.230	1.404±0.195	1.707±0.249	6.06*
Cocoon shell weight (g)	0.324±0.029	0.211±0.027	0.296±0.033	43.02*
Cocoon shell percentage (%)	22.23±0.02	15.23±0.02	17.54±1.95	41.99*

* statistical difference at the significant level of P<0.01

Table 4. Genetic parameters as estimated by composite interval mapping of the QTL affecting the cocoon related traits in the silkworm.

Trait	Linkage group	QTL	Neighboring markers	Position (cM)	LOD	R ² (%)	Additive effect	Dominance effect	h ²	
Cocoon weight (C.W.)	1	cw1a	Patg/Taca1-Paac/Ttg7	16.01	2.5217	4.29	-0.0735	0.3881	-5.278	
	1	cw1c	Ptc/Taca11-Paga/Taag8	77.94	2.7645	0.03	-0.0060	-0.3802	62.944	
	5	cw5a	Pat/Taca7-Ptat/Ttg6	22.01	2.6030	0.89	0.0352	-0.3142	-8.913	
	5	cw5b	Ptat/Ttg6-Patg/Taca6	42	2.6020	1.44	0.0447	-0.2625	-5.880	
	6	cw6	Ptat/Taag14-Pat/Taca10	79.1	3.8932	2.01	-0.0559	-0.3835	6.858	
	8	cw8	Paac/Ttg8-Ptat/Ttg7	22.01	3.0872	18.33	-0.1581	0.3690	-2.334	
	16	cw16c	Paac/Ttc4-Pat/Ttaci1	36.55	2.8329	1.20	-0.0409	-0.0559	1.367	
	16	cw16d	Paac/Tat8-Ptat/Ttac2	172.57	2.6349	0.97	0.0385	-0.3771	-9.802	
	16	cw16f	Ptc/Taca7-Ptat/Ttac3	242.79	4.9963	1.01	-0.0347	-0.3775	10.886	
	17	cw17	Ptat/Ttg1-Ptat/Taag10	30.49	3.5747	3.72	-0.0723	-0.3846	5.318	
	19	cw19a	Paac/Tagc4-Paac/Tat7	49.29	3.7596	1.43	0.0433	-0.3852	-8.891	
	19	cw19b	Paac/Ttc2-Pat/Tagc16	179.81	3.4050	6.53	0.0887	-0.0262	-0.296	
	Cocoon shell weight (C.Sh.W.)	16	cshw16c	Ptc/Taca7-Ptat/Ttac3	234.79	4.1126	10.04	-0.0164	-0.0314	1.917
		22	cshw22b	Ptat/Ttac7-Ptat/Taag1	156.61	3.7283	49.12	0.0459	0.0377	0.820
		8	cshp8	Paac/Ttg8-Ptat/Ttg7	20.01	3.5878	14.64	0.0118	-0.0361	-3.059
		9	cshp9	Ptat/Tagc4-Paac/Tagc5	0.01	2.9946	8.14	-0.0086	-0.0269	3.115
Cocoon shell percentage (C.Sh.P.)	16	cshp16	Paac/Tat8-Ptat/Ttac2	154.57	2.5384	1.53	0.0039	-0.0134	-3.468	
	19	cshp19a	Paac/Tagc4-Paac/Tat7	49.29	3.5142	0.64	-0.0025	0.0352	-14.047	
	23	cshp23a	Pat/Taat1-Ptat/Taag3	2.01	4.5747	61.68	-0.0272	0.0255	-0.936	
	23	cshp23b	Ptat/Taag3-Pat/Taca14	21.41	4.6637	69.18	-0.0347	0.0007	-0.020	
	23	cshp23d	Pac/Taag11-Paag/Taca1	71.99	4.7863	47.64	-0.0309	0.0210	-0.678	

^a The ratio dominance/additive effects. Under dominance or recessive if $h < 0$, additive if $h=0-0.20$, partial dominance if $h=0.21-0.80$, dominance if $h=0.81-1.20$, and over dominance if $h > 1.20$ (Stuber et al. 1987).
P<0.01

ranging from -0.0314 (*cshw16c*) to +0.037 (*cshw22b*). Finally, fourteen QTL were identified for cocoon shell percentage located on the LG8, LG9, LG16, LG19, and LG23. The additive effects of these QTL ranged from -0.0347 (*cshp23b*) to +0.0118 (*cshp8*), and their dominance effects ranged from -0.0361 (*cshp8*) to + 0.0352 (*cshp19a*) (Table 4).

The explained variation rate by QTL controlling cocoon weight, cocoon shell weight, and cocoon shell percentage ranged from 0.02% to 64.85%, 0.2% to 49.11%, and 0.04% to 84.20%, respectively (Table 4).

Discussion

In the present study, an AFLP-based linkage map containing 159 AFLP markers in a total

length of 2747 cM and an average marker resolution of 19.31 cM was developed for *B. mori*. Using this map, a total of 21 AFLP markers linked to cocoon weight, cocoon shell weight, and cocoon shell percentage were identified using a composite interval mapping method (Table 4). Recently, 11 QTL (Lu et al. 2004) and 40 QTL (Li et al. 2006) for whole cocoon weight, cocoon shell weight, ratio of shell weight and weight of pupae have been reported, and Javadi Taklimi (2006) accounted 5 QTL controlling ratios of shell weight in a backcrossed population (BC_1) of *B. mori*. These differences may be due to sample size, number and type of primer combinations used, and crosses established. Though apart from employing different primer pairs, Lu et al. (2004) and Li et al. (2006) used 44 BC_1 progenies, and Javadi Taklimi (2006)




Figure 4.




Figure 4.




Figure 4.




Figure 4.




Figure 4. Composite interval mapping of the cocoon related traits in *Bombyx mori*. Significance thresholds are indicated by horizontal lines, with LOD = 3.0 (genome wide $P < 0.001$) as determined by 1000 permutations of the mapping data (Churchill and Doerge 1994). High quality figures are available online.

applied only seven polymorphic primer combinations. Practically, factors such as the number of molecular markers used, types of crosses, sample size of segregating population, number of genes controlling the traits, and existence of gene interaction may influence the statistical power of QTL mapping (Zhong et al. 2006).

It seems that some QTL had pleiotropic effects on the traits. One QTL controlling both cocoon weight and cocoon shell weight traits located on the LG16 between neighboring markers Ptc/Taca7-Ptat/Ttac3 was observed. In addition, three QTL were observed controlling both cocoon weight and cocoon shell percentage traits located on the LG8, LG16, and LG19 between neighboring

markers Paac/Ttg8-Ptat/Ttg7, Paac/Tat8-Ptat/Ttac2 and Paac/Tagc4-Paac/Tat7.

A total of 19.35% of clearly readable and qualified AFLP bands were polymorphic between the P107 inbred line and the Khorasan Lemon native strain of *B. mori*. A higher level (61%) of polymorphic AFLP marker has been reported by Tan et al. (2001) in a single backcross (no. 782 and od100) family of *B. mori*. To explain this approach, they discussed several factors:

1. Employing two distinct *B. mori* strains in the present study, P107 and Khorasan Lemon are two examples of distinct silkworm inbred lines and strains. The former is from the Japanese bivoltine

system and the latter is from the Iranian native monovoltine system.

2. Detecting high levels of polymorphisms by the AFLP technique (Huys et al. 1996; Latorra et al. 1996; Mackill et al. 1996; Wan et al. 1999)
3. A large fraction of the silkworm genome consists of families of transposable elements such as *Bm1*, *BMCI* (a member of the *LINE1* family), *mariner*, *mariner*-like elements (*Bmmar1*), long terminal repeat transposons (LTRs), non-long terminal transposons (nonLTRs), and others (Ueda et al. 1986; Herrero and Wang 1991; Xiong and Eickbush 1993; Xiong et al. 1993; Robertson and Asplund 1996; Tomita et al. 1997; Shimizu et al. 2000; Wang et al. 2000).

Among the 20 pairs of AFLP primer combinations applied in this study, an average of 9.2 polymorphic AFLP markers per primer combination for linkage analysis and QTL mapping was recognized. This rate was considerably lower than two other AFLP linkage and QTL mapping studies on silkworm with 35.7 (Tan et al. 2001) and 36.4 (Lu et al. 2004) fragments per primer. This may be due to the degree of differences between parental lines and strains and the primer combinations used.

The present AFLP map consisted of 24 linkage groups, whereas the haploid genome of *B. mori* has 28 chromosomes. As Promboon et al. (1995), Young et al. (1998), and He (1998) reported, this may be due to nonequivalence between the number of linkage groups and the number of chromosomes. In the RFLP based linkage map by Goldsmith (1991), 15 linkage groups were reported. However, by using morphological (Doira et al. 1992), RAPD (Yasukochi et al. 1998), RFLP (Nguu et al. 2005), AFLP (Sima

et al. 2006), and SNP (Yamamoto et al. 2006) markers, 28 linkage groups and using SSR (Miao et al. 2005) markers, 29 linkage groups have been recognized in *B. mori*. It has also been shown that the large number of chromosomes in the haploid *B. mori* genome ($n = 28$), typical of Lepidoptera, makes it difficult to construct maps without missing some chromosomes (Yasukochi 1998).

The whole cocoon weight, cocoon shell weight, and cocoon shell percentage are the major economic traits in *B. mori* that are controlled by a polygene (Li et al. 2006). In the present study, a single F_2 population derived from a cross between P107 Japanese inbred line and Khorasan Lemon Iranian native strain was used as the mapping population. Among the 21 QTL for the traits studied, one QTL had dominance effect, 13 QTL had under dominance or recessive effects, and seven QTL had over dominance effects (Table 4). Li et al. (2006) in a backcrossed population (BC_1) derived from a cross between C_{100} and Dazao detected 40 QTL for whole cocoon weight and related traits, of which 19 were additive effect QTL and 21 were reduced effects QTL.

In summary, 159 AFLP markers were employed to construct a linkage map for *B. mori*, with an average marker resolution of 19.31 cM. We identified 21 QTL ($n = 21$) using the composite interval mapping method that affects whole cocoon weight and related traits. The effects of these QTL were under dominance, dominance, and over dominance. Since AFLP amplification is highly reproducible, the development of an AFLP linkage map and subsequently the identification of strain-specific markers for tracking allele frequency changes and quantitative trait loci (QTL) analysis for economically important traits provides an

invaluable tool for improving *B. mori* breeds, strains, and hybrids in order to enhance the silk production.

Acknowledgments

This work was supported by funds from Agricultural Biotechnology Research Institute of Iran (ABRII) under the supervision of Agricultural Research and Education Organization (AREO) (an organization of the Ministry Jihad-e Agriculture) and Agricultural Biotechnology Research Institute of North Region (Rasht). We thank Mr. Moeineddin Mavvajpour (the director) and Dr. Alireza Seydavi (the adviser) from ISRC (Iran Silkworm Research Centre) for establishing silkworm crosses and phenotypic information and for their invaluable advice. The technical assistance of the staff of ISRC is gratefully acknowledged. We are also grateful to Mohammad Naserani for contribution of data and typing of phenotypic records in Excel, for editing and inputting data in the linkage map and QTL software for further analysis.

References

- Abe H, Harada T, Kanehara M, Shimada T, Ohbayashi F, Oshiki T. 1998. Genetic mapping of RAPD markers linked to the denonucleosis refractoriness gene, *nsd-1*, in the silkworm, *Bombyx mori*. *Genes and Genetic Systems* 73: 237-42.
- Abe H, Sugasaki T, Kanehara M, Shimada T, Gomi ST, Ohbayashi F. 2000. Identification and genetic mapping of RAPD markers linked to the denonucleosis refractoriness gene, *nsd-2*, in the silkworm, *Bombyx mori*. *Genes and Genetic Systems* 75: 93-96.
- Causse MA, Fulton TM, Cho YG, Ahn SN, Chumwongse J, Wu K, Xiao J, Yu Z, Ronald PC, Hanington SE, Second G, McCouch SR, Tanksley SD. 1994. Saturated molecular map of the rice genome based on an interspecific backcross population. *Genetics* 138: 1251-1274.
- Chatterjee SN, Pradeep AR. 2003. Molecular markers (RAPD) associated with growth, yield and origin of the silkworm, *Bombyx mori* L. in India. *Genetika* 39: 1612-24.
- Chatterjee SN, Mohandas TP. 2003. Identification of ISSR markers associated with productivity traits in silkworm, *Bombyx mori* L. *Genome* 46: 438- 47.
- Chen K, Yao Q, Li M, Wang Y. 2003. Screening of RAPD markers for fluoride resistance in *Bombyx mori* L. *International Journal of Industrial Entomology* 7: 11-14.
- Churchill GA, Doerge RW. 1994. Empirical threshold values for quantitative trait mapping. *Genetics* 138: 963-971.
- Dalirsefat SB, Mirhoseini SZ. 2007. Assessing genetic diversity in Iranian native silkworm (*Bombyx mori* L.) strains and Japanese commercial lines using AFLP markers. *Iranian Journal of Biotechnology* 5(1): 25-33.
- Doira H, Fujii H, Kawaguchi Y, Kihara H, Banno Y. 1992. Important Genetic Resources. In: *Genetic Stocks and Mutations of Bombyx mori*. p 73. Isseido Press.
- Erickson DL, Fenster BC, StenØien HK, Price D. 2004. Quantitative trait locus analyses and the study of evolutionary process. *Molecular Ecology* 13: 2505-2522.
- Gage LP. 1974. The *Bombyx mori* genome: Analysis by DNA reassociation kinetics. *Chromosoma* 45: 27-42.

- Goldsmith MR. 1991. The *Bombyx mori* genome-mapping project. *Sericologia* 31: 145-155.
- Goldsmith MR, Shimada T, Abe H. 2005. The genetics and genomics of the silkworm, *Bombyx mori*. *Annual Review of Entomology* 50: 71-100.
- Harushima Y, Yano M, Shomura A, Sato M, Shimano T, Kuboki Y, Yamamoto T, Lin SY, Antonio BA, Parco A, Kajiya H, Huang N, Yamamoto K, Nagamura Y, Kurata N, Khush GS, Sasaki T. 1998. High-density rice genetic linkage map with 2275 markers using a single F2 population. *Genetics* 148: 479-494.
- He NJ. 1998. *Construction of the molecular linkage map of the silkworm (Bombyx mori L.)*. Ph.D. Thesis, Southwest Agricultural University, Chongqing, China.
- He LJ, Lu C, Li B, Zhou ZY, Xiang ZH. 2001. The construction of linkage map of *Bombyx mori* by combination of SADF and RAPD. *Acta Entomologica Sinica* 44(4): 476-482. (in Chinese)
- Herrer RJ, Wang J. 1991. Evidence for a relationship between the *Bombyx mori* middle repetitive Bm1 sequence family and U1 snRNA. *Genetica* 84: 31-37.
- Huys G, Coopman R, Janssen P, Kersters K. 1996. High-resolution genotypic analysis of the genus *Aeromonas* by AFLP fingerprinting. *International Journal of Systematic Bacteriology* 46: 572-580.
- Javadi Taklimi SA. 2006. *Identification QTL controlling cocoon shell percentage in silkworm*. M.Sc. dissertation, Faculty of Agricultural Science, University of Guilan.
- Kadono-Okuda K, Ogoyi DO, Nohata J, Sasanuma S, Sasanuma M, Eguchi R, Hara W, Yamamoto K, Mita K. 2003. Sequence analysis of BAC contigs covering a densovirus-nonsusceptibility gene, *nsd-2* in *Bombyx mori*. *Lepidoptera Abstracts*. Presented at the Sixth International Workshop *Molecular Biology and Genetics of Lepidoptera*, Kolymbari, Crete.
- Kosambi DD. 1944. The estimation of map distance from recombination values. *Annals of Eugenics* 12: 172-175.
- Latorra D, Schanfield MS. 1996. Analysis of human specificity in AFLP systems APOB, PAH, and D1S80. *Forensic Science International* 83: 15-25.
- Li B, Lu C, Zhou ZY. 2000. The Construction of RAPD linkage map of *Bombyx mori*. *Acta Genetica Sinica* 27(2): 127-132. (in Chinese)
- Li B, Lu C, Zhao AC, Xiang ZH. 2006. Multiple interval mapping for whole cocoon weight and related economically important traits QTL in silkworm (*Bombyx mori*). *Agricultural Sciences in China* 5(10): 798-804.
- Li M, Yao Q, Hou C, Cheng L, Chen K. 2001. Studies on RAPD markers linked to the denonucleosis refractoriness gene, *nsd-Z*, in the silkworm, *Bombyx mori*. *Sericologia* 41: 409-415.
- Lu C, Li B, Zhao AC, Xiang ZH. 2004. QTL mapping of economically important traits in Silkworm (*Bombyx mori*), *Science in China Series C Life Sciences* 47(5): 477-484.
- Mackill DJ, Zhang Z, Redona ED, Colowit PM. 1996. Level of polymorphism and genetic mapping of AFLP markers in rice. *Genome* 39: 969-977.

- Manly KF, Cudmore JR, Meer JM. 2001. Map Manager QTX: Cross-platform software for genetic mapping. *Mammalian Genome* 12: 930-932.
- Miao XX, Xu SJ, Lia MH, Lia MW, Huang JH, Dai FY, Marino SW, Mills DR, Zeng P, Mita K, Jia SH, Zhang Y, Liu WB, Xiang H, Guo QH, Xu AY, Kong XY, Lin HX, Shi YZ, Lu G, Zhang X, Huang W, Yasukochi Y, Sugasaki T, Shimada T, Nagaraju J, Xiang ZH, Wang SY, Goldsmith MR, Lu C, Zhao GP, Huang YP. 2005. Simple sequence repeat-based consensus linkage map of *Bombyx mori*. *Proceedings of the National Academy of Sciences U.S.A.* 102: 16303-16308.
- Mirhoseini SZ, Dalirsefat SB, Pour Khairandish M. 2007. Genetic characterization of Iranian native *Bombyx mori* strains by using amplified fragment length polymorphism markers. *Journal of Economic Entomology* 100(3): 939-945.
- Nagaraja GM, Nagaraju J. 1995. Genome fingerprinting in silkworm, *Bombyx mori*, using random arbitrary primers. *Electrophoresis* 16: 1633-1638.
- Nagaraju J, Goldsmith MR. 2002. Silkworm genomics-progress and prospects. *Current Science* 83: 415-425.
- Nguu EK, Kadono-Okada K, Mase K, Kosegawa E, Hara W. 2005. Molecular linkage map for the silkworm, *Bombyx mori*, based on restriction fragment length polymorphism of cDNA clones. *Journal of Insect Biotechnology and Sericology* 74: 5-13.
- Promboon A, Shimada T, Fujiwara H, Kobayashi M. 1995. Linkage map of RAPDs in the silkworm, *Bombyx mori*. *Genetics Research* 66: 1-7.
- Qi X, Stam P, Lindhout P. 1997. Use of the locus specific AFLP markers to construct a high density molecular map in barley. *Theoretical and Applied Genetics* 96: 376-384.
- Robertson HM, Asplund ML. 1996. Bmmar1: A basal lineage of the mariner family of transposable elements in the silkworm moth, *Bombyx mori*. *Insect Biochemistry and Molecular Biology* 26: 945-954.
- Shi J, Heckel DG, Goldsmith MR. 1995. A genetic linkage map for the domesticated silkworm, *Bombyx mori*, based on restriction fragment length polymorphisms. *Genetics Research* 66: 109-126.
- Shimizu K, Kamba M, Sonobe H, Kanda T, Klinakis AG, Savakis C, Tamura T. 2000. Extrachromosomal transposition of the transposable element minos occurs in embryos of the silkworm *Bombyx mori*. *Insect Molecular Biology* 9: 277-281.
- Sima YH, Li B, Chen DX, Sun DB, Zhao AC, Zhang L, Lu C, He SM, Xiang ZH. 2006. Construction and analysis of an AFLP molecular linkage map of the silkworm (*Bombyx mori*). *Chinese Journal of Agricultural Biotechnology* 3(1): 25-31.
- Stuber CW, Edwards MD, Wendel JF. 1987. Molecular markers facilitated investigations of quantitative trait loci in maize. II. Factors influencing yield and its component traits. *Crop Science* 27: 639-648.
- Suzuki Y, Gage LP, Brown DD. 1972. The genes for fibroin in *Bombyx mori*. *Journal of Molecular Biology* 70: 637-649.
- Tamura T, Thibert C, Royer C, Kanda T, Eappen A, Kamba M, Kômoto N, Thomas JL, Mauchamp B, Chavancy G, Shirk P, Fraser M,

- Prudhomme JC, Couble P. 2000. Germline transformation of the silkworm, *Bombyx mori* L. using a piggyBac transposon-derived vector. *Nature Biotechnology* 18: 81-84.
- Tan YD, Ma RL. 1998. Estimates of lengths of genome and chromosomes of rice using molecular markers. *Journal of Biomathematics* 13:1022-1027. (in Chinese)
- Tan YD, Wan CL, Zhu YF, Lu C, Xiang Z, Deng HW. 2001. An amplified fragment length polymorphism map of the silkworm, *Genetics* 157(3): 1277-1284.
- Tomita M, Munetsuna H, Sato T, Adachi T, Hino R, Hayashi M, Shimizu K, Nakamura N, Tamura T, Yoshizato K. 2003. Transgenic silkworms produce recombinant human type III procollagen in cocoons. *Nature Biotechnology* 21: 52-56.
- Tomita S, Sohn BH, Tamura T. 1997. Cloning and characterization of a mariner-like element in the silkworm, *Bombyx mori*. *Genes and Genetic Systems* 72: 219-228.
- Ueda H, Mizuno S, Shimura K. 1986. Transposable genetic element found in the 5'-flanking region of the fibroin H-chain gene in a genomic clone from the silkworm *Bombyx mori*. *Journal of Molecular Biology* 190: 319-327.
- Vos P, Hogers R, Bleeker M, Reijans M, Lee TVD, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M. 1995. AFLP: A new technique for DNA fingerprinting. *Nucleic Acids Research* 23: 4407-4414.
- Wan CL, Zhu YF, Tan YD, Lu C. 1999. Application of AFLP markers to detection of genetic polymorphic loci in the silkworm (*Bombyx mori*, L.). *Biotechnology* 9: 4-9. (in Chinese)
- Wang S, Basten CJ, Zeng ZB. 2007. *Windows QTL cartographer* Version 2.5. *Statistical Genetics*, North Carolina State University, USA.
- Wang W, Swevers L, Iatrou K. 2000. Mariner (Mos1) transposase and genomic integration of foreign gene sequences in *Bombyx mori* cells. *Insect Molecular Biology* 9: 145-155.
- Waugh R, Bonar N, Baird E, Thomas B, Graner A, Thos WTB, Powell W. 1997. Homology of AFLP products in three mapping populations of barley. *Molecular Genetics and Genomics* 255: 311-321.
- Wu R, Ma CX, Casella G. 2007. *Statistical Genetics of Quantitative Traits-Linkage, Maps, and QTL*. Springer.
- Xiong Y, Eickbush TH. 1993. Dong, a non-long terminal repeat (non-LTR) retrotransposable element from *Bombyx mori*. *Nucleic Acids Research* 21: 1318.
- Xiong Y, Burke WD, Eickbush TH. 1993. Pao, a highly divergent retrotransposable element from *Bombyx mori* containing long terminal repeats with tandem copies of the putative R region. *Nucleic Acids Research* 21: 2117-2123.
- Yamamoto K, Narukawa J, Kadono-Okuda K, Nohata J, Sasanuma M, Suetsugu Y, Banno Y, Fujii H, Goldsmith MR, Mita K. 2006. Construction of a single nucleotide polymorphism linkage map for the silkworm, *Bombyx mori*, based on bacterial artificial chromosome end sequences. *Genetics* 173: 151-161.
- Yamao M, Katayama N, Nakazawa H, Yamakawa M, Hayashi Y, Hara S, Kamei K, Mori H. 1999. Gene targeting in the silkworm by use of a baculovirus. *Genes and Development* 13: 511-516.

Yao Q, Li MW, Wang Y, Wang WB, Lu J, Dong Y, Chen KP. 2003. Screening of molecular markers for NPV resistance in *Bombyx mori* L. (*Lep.*, *Bombycidae*). *Journal of Applied Entomology* 127(3): 134-136.

Yasukochi Y. 1998. A dense genetic map of the silkworm, *Bombyx mori*, covering all chromosomes based on 1018 molecular markers. *Genetics* 150: 1513-1525.

Young WP, Wheeler PA, Coryell VH, Keim P, Thorgaard GH. 1998. A detailed linkage map of rainbow trout produced using doubled haploids. *Genetics* 148: 839-850.

Zabeau M, Vos P. 1992. Selective restriction fragment amplification: A general method for DNA fingerprinting, European Patent Application no. 92402629.7, publication no. 0 534 858 A1.

Zhong D, Menge DM, Temu EA, Chen H, Yan G. 2006. Amplified fragment length polymorphism mapping of quantitative trait loci for malaria parasite susceptibility in the Yellow Fever mosquito *Aedes aegypti*. *Genetics* 173: 1337-1345.

!