

NADPH Oxidase Deficiency in X-Linked Chronic Granulomatous Disease

DAVID C. HOHN and ROBERT I. LEHRER

From the Department of Surgery, University of California, San Francisco, 94143, and the Department of Medicine, University of California, Los Angeles, California 90024

ABSTRACT We measured the cyanide-insensitive pyridine nucleotide oxidase activity of fractionated resting and phagocytic neutrophils from 11 normal donors, 1 patient with hereditary deficiency of myeloperoxidase, and 7 patients with X-linked chronic granulomatous disease (CGD). When measured under optimal conditions (at pH 5.5 and in the presence of 0.5 mM Mn^{++}), NADPH oxidase activity increased four-fold with phagocytosis and was six-fold higher than with NADH. Phagocytic neutrophils from patients with CGD were markedly deficient in NADPH oxidase activity.

INTRODUCTION

X-linked chronic granulomatous disease (CGD),¹ a hereditary disorder of male children, is characterized by serious susceptibility to infection (2). Although neutrophils (PMN) from patients with CGD ingest microorganisms effectively, they fail to kill many bacteria and fungi at a normal rate (3, 4). Phagocytosis by normal PMN is associated with major changes in oxidative metabolism, including increased cyanide-insensitive consumption of oxygen (5-7), stimulation of glucose oxidation via the hexose monophosphate shunt (HMPS) (7, 8), increased generation of hydrogen peroxide and superoxide (9-11), and augmented reduction of the redox dye, nitroblue tetrazolium (NBT)

This work was presented in part at the National Meeting of the American Federation for Clinical Research, Atlantic City, N. J., May, 1974, and published in abstract form (1).

Received for publication 21 June 1974 and in revised form 14 November 1974.

¹ Abbreviations used in this paper: CGD, chronic granulomatous disease; GRF, granule-rich fraction; HMPS, hexose monophosphate shunt; MPO, myeloperoxidase; NBT, nitroblue tetrazolium; PBS, phosphate-buffered saline; PMN, polymorphonucleated neutrophils.

(12). All of these features of increased oxidative metabolism are deficient in CGD PMN (12-13).

The primary enzyme of the phagocytic oxidative burst is thought to be a cyanide-insensitive pyridine nucleotide-linked oxidase that catalyzes the conversion of oxygen to hydrogen peroxide and superoxide. Major controversy has arisen over which nucleotide is oxidized, and evidence for both NADH and NADPH has been advanced (14-19). There is agreement that the controlling factor for HMPS activity is the availability of $NADP^+$ (20). If the primary oxidase were NADPH-linked, the $NADP^+$ produced would directly stimulate the HMPS. An NADH-linked oxidase would require at least one additional metabolic pathway for the provision of $NADP^+$ (15, 17).

In view of the controversy attending this aspect of neutrophil function, we decided to examine the oxidase activity of human PMN, employing resting and phagocytic PMN from normal donors, a group of patients with X-linked CGD, and a patient with myeloperoxidase (MPO) deficiency.

METHODS

Control PMN were obtained from nine adults and two children without CGD, and from one patient with hereditary MPO deficiency. Five of the seven male patients with CGD ranged from 4 to 12 yr of age, and the others were 18 and 28 yr old. Two of the patients were brothers, and all but one, who had *Pseudomonas* sepsis, were free of serious infection when studied. The diagnosis of CGD was established by measurements of the oxygen consumption, HMPS activity, quantitative iodination, quantitative NBT reduction, and microbicidal activity (*Staphylococcus aureus* 502 A and *Candida albicans*) by PMN.

Leukocytes were separated by the Hypaque-Ficoll technique of Böyum (21), and granulocytes were harvested with less than 2% contamination by mononuclear cells. The mean percentages of eosinophils in these preparations were 5.3% (normal controls), 6.8% (MPO-deficient subject), and 6.8% (CGD patients). Erythrocytes were sedimented in dextran, and residual contaminating erythrocytes were lysed by sus-

pending the cell button in cold 0.22% saline for 45 sec and then restoring isotonicity. Granulocytes were centrifuged at 250 *g* for 10 min and resuspended in Ca⁺⁺ and Mg⁺⁺-free phosphate-buffered saline (PBS)² at a concentration of 1 × 10⁸ cells/ml. Zymosan³ was opsonized by adding AB serum to achieve a concentration of 5 mg/ml, incubating the mixture for 30 min at 37°C, and washing and resuspending the zymosan in PBS at a final concentration of 15 mg zymosan/ml. Phagocytosis was induced by adding 15 mg of zymosan/10⁸ PMN, providing approximately 10 zymosan particles/PMN. The cell-zymosan suspension was rotated (30 rpm) at 37°C for 3 min, so that virtually all the neutrophils contained three or more ingested zymosan particles. The reaction was stopped by addition of ice-cold PBS, and the resting and phagocytic cells were immediately centrifuged at 400 *g* for 3 min at 4°C. The cells were resuspended at a concentration of 5 × 10⁷ PMN/ml in 1 or 2 ml of ice-cold 0.34 M sucrose adjusted to pH 7.4 with NaHCO₃ and homogenized immediately in a motor-driven Teflon-glass homogenizer at 0°C until 95% cell breakage had been achieved. With our homogenizer, 5 min of homogenization was required. After centrifugation at 250 *g* for 10 min, the supernate was centrifuged at 27,000 *g* for 20 min to yield a granule-rich sediment and a clear supernate. The granule-rich sediment was resuspended at a concentration of 2 × 10⁸ cell equivalents/ml in 0.34 M sucrose, pH 7.4.

Measurements of oxygen consumption were made at 1 mM nucleotide concentration at 37°C with a Gilson model KM oxygraph⁴ fitted with a Clark-Yellow Springs Instruments oxygen electrode⁵. NADPH oxidase activity was linear with time during the 1st min after addition of enzyme. Reaction rates were calculated from the maximum slope achieved during this period. Nucleotide consumption was monitored as the decrease in absorbance at 340 nm at room temperature in a Beckman Kintrac VII recording spectrophotometer⁶ with 0.16 mM nucleotide concentrations and 10 mm path-length cuvettes. NADH³ (98% pure) and NADPH³ (95-99% pure) were weighed and dissolved in phosphate buffer immediately before use.

The effect of manganese and hydrogen ion concentrations on NADH and NADPH oxidase activity in the 27,000 *g* granule-rich fraction (GRF) of resting and phagocytic neutrophils was determined polarographically at 1 mM nucleotide concentration, and the data were expressed as nanomoles O₂ consumed per minute per milligram protein. Both studies were made in 0.065 M Na⁺-K⁺-phosphate buffer with 125 mM sucrose and 2 mM KCN. Studies of the effect of manganese were conducted at pH 5.5 over a concentration range of 0-0.5 mM manganese chloride. Determinations at higher manganese concentration were limited by the formation of insoluble salts. The influence of hydrogen ion concentration was determined in the presence of 0.5 mM Mn⁺⁺ as the pH was varied from 7.0 to 5.5. Spontaneous oxidative degradation of both nucleotides increased as the pH was lowered. At pH 5.5 in the phosphate-MnCl₂-KCN buffer system, the mean spontaneous oxidation rates for NADH and NADPH were 4.1 and 40.5 nmol O₂/min measured polarographically at 1 mM NAD(P)H and 2.1 and 6.1 nmol nucleotide/min measured spectrophotometrically at 0.16 mM NAD(P)H. In each determination, the corresponding spontaneous oxidation rate was subtracted from the value ob-

tained in the presence of enzyme. After determining the basal rate of O₂ consumption, enzymatic activity was measured by adding from 80 to 400 μg of granule protein; the higher amounts were employed as the concentrations of manganese and hydrogen ion were decreased. There was no spontaneous O₂ consumption by granule preparations in the absence of pyridine nucleotides.

Oxidase activity was present in the 250 *g* and 27,000 *g* sediments but absent in the 27,000 *g* supernate. Subsequent studies were made in 0.065 M phosphate buffer at pH 5.5 with 0.5 mM manganese chloride, 125 mM sucrose, and 2 mM KCN with the 27,000 *g* GRF as the enzyme source. For NADPH oxidase determinations phagocytic (80 μg protein) or resting (160 μg protein) GRF was added after the basal rate of spontaneous oxidation of nucleotide was established. Similar amounts of protein were used in our initial spectrophotometric determinations of NADH consumption. Because of the low levels of activity found, later spectrophotometric and all polarographic measurements of NADH oxidase activity employed 320 μg of granule protein. Blank values for NADPH oxidation were 27.5% (polarographic assay) and 23.8% (spectrophotometric assay) of the mean enzymatic oxidation by GRF from phagocytic PMN. The corresponding blank values for NADH auto-oxidation were 29% (polarographic assay) and 78% (spectrophotometric assay). Oxidase activity was expressed as nanomoles of O₂ or nucleotide consumed per minute per milligram protein or per 10⁷ PMN equivalents. Zymosan had no intrinsic NAD(P)H oxidase activity, and addition of zymosan to homogenized nonphagocytic PMN caused no increase in NAD(P)H oxidase activity. When PMN were incubated with unopsonized zymosan, there was little microscopic evidence of particle ingestion, and no increase in NAD(P)H oxidase activity was measured.

Protein was determined by the method of Lowry, Rosebrough, Farr, and Randall (22), with bovine serum albumin as the standard. The mean total protein in GRF obtained from normal phagocytic PMN (1.31 mg/10⁸ PMN) was 12% lower than that obtained from normal resting PMN (1.49 mg/10⁸ PMN). The corresponding mean values for GRF protein content of patients with CGD were 1.34 mg/10⁸ phagocytic PMN and 1.45 mg/10⁸ resting PMN.

RESULTS

Effect of manganese and pH on oxidase activity.

The effects of manganese and pH on oxidase activity in the 27,000 *g* sediment of phagocytic PMN are shown in Figs. 1 and 2. Oxidase activity with both nucleotides increased 25-fold as the manganese concentration was raised from zero to 0.5 mM. NADPH oxidation was three to six times higher than NADH oxidation at all manganese concentrations and nearly six times higher in the absence of manganese.

When measured in the presence of 0.5 mM manganese, lowering the pH from 7.0-5.5 resulted in a four-fold increase in oxidase activity with both nucleotides; oxidation of NADPH was at least 4.8 times higher than NADH oxidation at all pH values. NAD(P)H oxidase activity of GRF from nonphagocytic cells also varied with Mn⁺⁺ concentration and pH; these activities were optimal at pH 5.5 and at a concentration of 0.5 mM Mn⁺⁺. The NAD(P)H oxidase activity of GRF

² Grand Island Biological Co., Berkeley, Calif.

³ Sigma Chemical Co., St. Louis, Mo.

⁴ Gilson Medical Electronics, Inc., Middleton, Wis.

⁵ Yellow Springs Instrument Co., Yellow Springs, Ohio.

⁶ Beckman Instruments, Inc., Fullerton, Calif.




FIGURE 1 Effect of manganese concentration on NAD(P)H oxidase activity measured polarographically at 1 mM NAD(P)H concentration in pH 5.5 phosphate buffer. Data are from a representative experiment using the 27,000 g granule-rich sediment of phagocytic PMN as the enzyme source.

varied linearly with the protein concentration over the range employed in our polarographic assays.

In three experiments, we found the apparent K_m for NADPH to be 3.1, 3.3, and 5.7×10^{-3} M for GRF from resting PMN and 2.4, 2.5, and 4.0×10^{-3} M for GRF from corresponding phagocytic PMN.

Localization of oxidase activity in PMN fractions. The localization of oxidase activity in normal PMN fractions, measured polarographically and expressed as total O₂ consumed per minute per 10⁷ PMN equivalents, is shown in Fig. 3. No oxidase activity was detectable with either nucleotide in the whole homogenates of resting PMN, although low levels with both NADPH and NADH were detected in the isolated 27,000 g sediment. This NAD(P)H oxidase activity in GRF from resting PMN was eliminated when granules were suspended in the corresponding 27,000 g supernate rather than in 0.34 M sucrose. Homogenates of phagocytic PMN contained high NADPH oxidase activity; this was localized entirely to the 250 g and 27,000 g sediments, and activity in these fractions was maintained in the presence of the corresponding 27,000 g supernate. NADH oxidase activity in homogenate of phagocytic PMN was substantially lower and also localized




FIGURE 2 Effect of pH on NAD(P)H oxidase activity measured polarographically at 1 mM NAD(P)H concentration with 0.5 mM Mn²⁺. Data are from an experiment with the same 27,000 g sediment of phagocytic PMN used in Fig. 1.

to these sediments. We detected no oxidase activity with either NADPH or NADH in the 27,000 g supernate of either resting or phagocytic PMN.

Under the conditions employed, with 1 mM NADPH as the substrate, oxygen consumption by homogenate




FIGURE 3 NAD(P)H oxidase activities in homogenate and fractions of resting (R) and phagocytic (Ø) PMN. Oxidase activity was measured polarographically. Data are from a representative experiment. The dashed line indicates the level of O₂ consumption by 10⁷ phagocytic PMN (10 zymosan particles/PMN).




FIGURE 4 NADPH oxidase activity in 27,000 *g* sediment of resting and phagocytic PMN from controls (11) and CGD patients (7). Oxidase activity was measured polarographically (1 mM NADPH) and spectrophotometrically (0.16 mM NADPH). Vertical brackets represent the range of 10 control determinations, open circles the mean values, and solid dots the individual data points of CGD patients.

or the combined 250 *g* and 27,000 *g* sediments of phagocytic PMN exceeded the increment in oxygen consumption of an equivalent number of intact phagocytic PMN.

NADPH oxidase activities of controls and patients with CGD. The specific NADPH oxidase activity of the 27,000 *g* GRF of resting and phagocytic PMN from controls and the patients with CGD are compared in Fig. 4. In all cases, the 27,000 *g* fraction of normal




FIGURE 5 NADH oxidase activity in 27,000 *g* sediment of resting and phagocytic PMN from controls (11) and patients with CGD (7). Oxidase activity was measured polarographically (1 mM NADH) and spectrophotometrically (0.16 mM NADH). Vertical brackets represent the control range, open circles the mean values, and solid dots the individual data points of CGD patients.

phagocytic PMN had higher oxidase activity than the corresponding fraction from resting PMN. Mean NADPH oxidase values from resting and phagocytic PMN, respectively, were 180 and 696 nmol O₂/min per mg protein measured polarographically and 103 and 251 nmol NADPH/min per mg protein measured spectrophotometrically. As dissimilar substrate concentrations and temperatures were used in the two assay methods, these values cannot be used to establish the exact stoichiometry of the oxidase reaction.

NADPH oxidase activity was measured in the 27,000 *g* fraction of PMN from two children, ages 5 and 10, without CGD or serious infection. Resting levels were 85 and 248, and phagocytic levels were 492 and 769. NADPH oxidase activity of comparable fractions from MPO-deficient PMN was 49 (resting) and 367 (phagocytic) nmol O₂/min per mg protein. These polarographic values of NADPH oxidase activity were well within the range obtained with PMN from normal adult donors.

Fig. 4 clearly indicates a substantial overlap in NADPH oxidase values between the CGD and control groups when resting PMN were compared. When NADPH oxidase activities of phagocytic PMN were compared, all CGD values were well below the control range, both polarographically and spectrophotometrically. This difference between mean control and mean CGD values was highly significant ($P < 0.001$) when the data was analyzed by standard *t*-test. In contrast to the consistent increase in NADPH oxidase activity induced by phagocytosis in normal PMN, leukocytes from all of the patients with CGD lacked significant phagocytic activation.

NADH oxidase activity of controls and patients with CGD. The NADH oxidase activity in the 27,000 *g* sediment of PMN from control donors and patients with CGD is shown in Fig. 5. When measured polarographically in nonphagocytic leukocytes at 1 mM NADH concentration, mean specific NADH oxidase activity was 27 in control PMN and 17.2 in CGD PMN. Although the mean NADH oxidase activity of phagocytic PMN from patients with CGD was lower than the control values (17.7 vs. 87 nmol O₂/min per mg protein), there was overlap between individual control and CGD data points. As was the case with NADPH oxidase activity, NADH oxidase activity in PMN from patients with CGD was not activated by phagocytosis. Phagocytic PMN from all controls showed some increase over resting levels of NADH oxidation; the mean increment in NADH oxidase activity was substantially less than that in NADPH oxidase activity (60 vs. 516 nmol O₂/min per mg protein).

Low levels of enzyme activity, necessitating large amounts of enzyme, caused difficulties in measuring

NADH oxidase activity spectrophotometrically at 0.16 mM NADH concentration. There were individual control determinations where no phagocytic increase in NADH oxidase activity was measured; in some of these cases NADH oxidase activity was too low to allow accurate measurement without prohibitively large amounts of enzyme.

Oxidase activity of PMN from mothers of CGD patients. NADPH oxidase determinations were made on PMN from mothers of three of the CGD patients. Quantitative NBT reduction and O₂ consumption by PMN from two of these subjects were well below the normal control range; spectrophotometric oxidase activities with NADPH were 83 and 84 (resting PMN) and 145 and 279 (phagocytic PMN). NBT reduction and O₂ consumption by PMN from the third mother were normal; polarographic oxidase activities with NADPH were 11 (resting) and 306 (phagocytic). Additional studies of heterozygotes are required; at present, we are unable to conclude that the GRF from PMN of heterozygous subjects is significantly deficient in NADPH oxidase activity.

DISCUSSION

Phagocytosis by mammalian neutrophils is accompanied by marked augmentation of oxidative metabolism. The basic observations of increases in O₂ consumption, glucose oxidation via the hexose monophosphate pathway, and hydrogen peroxide production have been readily confirmed by many investigators. Recognition that PMN from patients with CGD do not show these features of increased oxidative metabolism after ingestion of particles or microorganisms has led to a general appreciation that these metabolic responses are related to microbicidal function. However, controversy has arisen over the particular enzyme(s) primarily responsible for this burst in oxidative metabolism. Clearly, elucidation of the enzymatic basis of the defect in CGD neutrophils could contribute to resolution of the controversy over the primary oxidase of normal human neutrophils.

Cagan and Karnovsky (14), and Baehner, Gilman, and Karnovsky (15) have provided evidence for an NADH-linked primary oxidase in the nonphagocytic leukocytes of guinea pigs and man. Although the levels measured were low, these authors reported sufficient oxidase activity in disrupted PMN to account for the burst in oxidative metabolism by intact phagocytic cells. Although Baehner and Karnovsky (23) found abnormally low levels of NADH oxidase activity in the KCl extracts of nonphagocytic leukocytes from five children with CGD, others have been unable to confirm these observations (13).

Evidence for an NADPH-linked primary oxidase has been advanced by a number of investigators, including

Iyer and Quastel (18), Rossi et al. (16, 17, 24), and Paul, Strauss, Jacobs, and Sbarra (19). Although Rossi's group has not reported studies of patients with CGD, they have found a five-fold increase in NADPH oxidase activity with phagocytosis by normal human PMN and have stressed the importance of Mn⁺⁺ and H⁺ concentrations for obtaining optimal enzyme activity (24).

Our own studies have led us to conclude that many of the discrepancies between previous studies of neutrophil oxidase activity are attributable to methodologic differences with major consequences. These include differences related to the H⁺ and Mn⁺⁺ concentrations of the assay mixtures, use of nonphagocytic or phagocytic cells as enzyme sources, and the method of fractionation of the cells to be tested. Each will be commented on briefly.

Our studies offer additional confirmation to the generally held belief that pyridine nucleotide oxidase activity is optimal at acid pH levels (9, 17, 19), yet some investigators have chosen to perform measurements at suboptimal pH levels (13–15, 23). This has reduced the sensitivity of their measurements and necessitated extrapolation to presumed optimal pH values to determine whether the measured enzyme levels would suffice to account for the magnitude of the metabolic stimulation of an equivalent number of phagocytic cells (23).

Considerable importance attends the use of Mn⁺⁺ in assays of oxidase activity. We found that addition of a variety of other cations, including Mg⁺⁺, Ca⁺⁺, Fe⁺⁺, Fe⁺⁺⁺, Cu⁺⁺, and Zn⁺⁺, caused no increase in NAD(P)H oxidase activity. Iyer and Quastel noted the Mn⁺⁺-stimulated oxidation of NADH and NADPH by oxidases of guinea pig PMN (18); Roberts and Quastel later attributed this effect, in part, to a Mn⁺⁺-dependent oxidation of these nucleotides by MPO (25, 26). However, several considerations indicated that Mn⁺⁺ exerts an effect on NAD(P)H oxidase activity that is completely independent of MPO. Rossi, Romeo, and Patriarca found that Mn⁺⁺ stimulated the oxidase activity of MPO-deficient neutrophils (24), a finding also confirmed in the present study. We have found that neutrophils from subjects with CGD have low levels of oxidase activity in the presence of Mn⁺⁺, despite their normal levels of MPO. The failure of MPO to obscure these measurements of NAD(P)H oxidase activity may be attributed to the presence of 2 mM KCN in the mixtures used to assay oxidase activity. We observed that this concentration sufficed to inhibit completely the Mn⁺⁺-dependent oxidation of both nucleotides by purified human MPO.⁷

We do not yet know if the Mn⁺⁺ effect on NAD(P)H oxidase activity is expressed in intact cells. Although

⁷ Generously provided by Dr. J. Schultz.

neutrophils are rich in zinc, we are aware of only one report dealing with their content of manganese (27). Atomic absorption spectrophotometric studies obtained on neutrophils fractionated by the methods employed in this study revealed Mn content in the 27,000 g GRF of normal neutrophils, where the metal was found to be 7.9% as abundant on a molar basis as zinc. The 27,000 g supernate had no detectable Mn. Although additional studies will be required to ascertain the functional significance of human neutrophil Mn^{++} , its presence and relative abundance in the oxidase-containing fraction suggest that its role in NAD(P)H oxidase activity may be more than an interesting *in vitro* phenomenon.

The selection of phagocytic versus nonphagocytic cells for measurements of oxidase activity is of critical significance. We found no consistent difference between normal and CGD neutrophils when we examined nonphagocytic cells. Yet clear and consistent differences in NADPH oxidation were apparent when phagocytic cells were compared. In preparing phagocytic PMN for our studies, we selected conditions that induced a rapid and maximal stimulation of O_2 consumption and made our enzyme measurements on cells fractionated soon after attaining rates of O_2 consumption some 15–20 times their basal levels. In the only other study that has employed phagocytic PMN from patients with CGD, Baehner and Nathan stimulated the cells with polystyrene spheres or bacteria for 15 min, and granules were lysed by repeated cycles of freezing and thawing (12). In our hands, both prolonged incubation (more than 7 min) of cell-particle suspensions and freeze-thaw treatment resulted in major losses of enzyme activity.

A final important methodologic point relates to the fraction in which enzyme activity was measured. We found essentially no activity in the 27,000 g cytosol after homogenization of the cells in 0.34 M sucrose. Activity was localized to the 250 g and 27,000 g sediments, both of which contained large numbers of granules as shown by morphology and by measurements of marker enzymes; oxidase activity of GRF was completely abolished by boiling and by incubation with pronase and trypsin. Our attempts to solubilize the oxidase by extraction of GRF with dilute acids or isotonic or hypertonic KCl resulted in considerable loss of activity. We preferred to make our measurements on the 27,000 g GRF, a reasonably consistent enzyme source that was not unduly influenced by variations in the extent of successful cell disruption. It should be noted that Rossi et al. had earlier recommended the use of this fraction.

Baehner and Karnovsky reported that NADH oxidase activity in 30,000 g alkaline KCl supernate of

nonphagocytic neutrophils was lower in five children with CGD than in controls (23). It appears, however, that the controls were patients who may have had leukocytosis associated with infection (28). Holmes, Page, and Good did not confirm these differences with normal uninfected subjects as controls (13). Baehner and Karnovsky reported that NADPH oxidase in neutrophil granules was approximately one-sixth as active as the NADH oxidase of normal neutrophils. Their measurements were made on nonphagocytic PMN, at pH 7, in the absence of manganese. NADPH oxidase activity was not reported in phagocytic cells, in patients with CGD, or under optimal conditions of assay.

Baehner and Nathan found lower levels of NADH oxidase activity in supernates of phagocytic cells from two children with CGD than in normal controls (12). They found little NADH oxidase activity in the granule fraction of normal cells and found no NADPH oxidase activity in either granules or supernate. Their granules, however, were subjected to repeated freezing and thawing, a treatment that would markedly decrease NADPH oxidase activity. Furthermore, NADPH oxidase was measured at pH 7.0 and in the absence of Mn^{++} , resulting in further enzyme attenuation.

Although we maintain reservations concerning the validity of comparing oxidase activity in disrupted cell fractions with levels of O_2 consumption by intact leukocytes, our studies indicate that, when NADPH served as the substrate, O_2 consumption by homogenized cells considerably exceeds the level of O_2 utilization by maximally phagocytic PMNs. These high levels of total and specific NADPH oxidase activity were observed only in homogenates of normal phagocytic neutrophils; nonphagocytic PMNs had much lower oxidase activity.

Our studies revealed that nonphagocytic PMN from patients with CGD had mean oxidase activities with both NADPH and NADH that were somewhat lower than corresponding mean values of the control group; however, overlap between data points was substantial, and separation of control and CGD groups was incomplete. The mean values of NADH oxidase activity of phagocytic PMN from patients with CGD were lower than the corresponding control values, but the ranges overlapped. Although none of the aforementioned measurements allowed us to separate CGD from normal PMN, our data show a marked difference when the NADPH oxidase activities of phagocytic PMN are compared. Phagocytic neutrophils from seven patients with CGD were deficient in NADPH oxidase activity, with wide separation between patient and control groups. Moreover, PMN from none of the CGD patients showed the activation of NADPH oxidase ac-

tivity by phagocytosis that was consistently observed with control PMN. We believe that these data indicate that absence of a cyanide-insensitive NADPH oxidase or deficiency in its activation is the primary metabolic defect in the X-linked CGD syndrome.

Attention should now be directed to a number of important problems that remain. These include determining whether NADPH and NADH are oxidized by the same or separate oxidases, precise localization of the oxidase(s), isolation and purification of the enzyme(s), and elucidation of the mechanism and effectors involved in the activation of oxidase activity by the phagocytic event.

ACKNOWLEDGMENTS

We thank Kathryn I. Mitchell and Dr. Franz E. Hohn for their skillful technical assistance.

The research was supported in part by grants from the National Institutes of Health (AI 10547, CA 11067, and GMO 1474-08 SUR), the American Cancer Society (CF-3258), and by a grant from the Research Corporation (Brown-Hazen Fund).

REFERENCES

- Hohn, D. C., and R. I. Lehrer. 1974. Identification of the defect in X-linked chronic granulomatous disease. *Clin. Res.* **22**: 394A. (Abstr.)
- Bridges, R. A., H. Berendes, and R. A. Good. 1959. A fatal granulomatous disease of childhood. The clinical, pathological, and laboratory features of a new syndrome. *Am. J. Dis. Child.* **97**: 387-408.
- Holmes, B., P. G. Quie, D. B. Windhorst, and R. A. Good. 1966. Fatal granulomatous disease of childhood. An inborn abnormality of phagocytic function. *Lancet.* **1**: 1225-1228.
- Quie, P. G., J. G. White, B. Holmes, and R. A. Good. 1967. In vitro bactericidal capacity of human polymorphonuclear leukocytes: diminished activity in chronic granulomatous disease of childhood. *J. Clin. Invest.* **46**: 668-679.
- Baldrige, C. W., and R. W. Gerard. 1933. The extra respiration of phagocytosis. *Am. J. Physiol.* **103**: 235-236.
- Cohn, Z. A., and S. I. Morse. 1960. Functional and metabolic properties of polymorphonuclear leukocytes. I. Observations on the requirements and consequences of particle ingestion. *J. Exp. Med.* **111**: 667-687.
- Sbarra, A. J., and M. L. Karnovsky. 1959. The biochemical basis of phagocytosis. I. Metabolic changes during the ingestion of particles by polymorphonuclear leukocytes. *J. Biol. Chem.* **234**: 1355-1362.
- Stähelin, H., M. L. Karnovsky, A. E. Farnham, and E. Suter. 1957. Studies on the interaction between phagocytes and tubercle bacilli. III. Some metabolic effects in guinea pigs associated with infection with tubercle bacilli. *J. Exp. Med.* **105**: 265-277.
- Iyer, G. Y. N., M. F. Islam, and J. H. Quastel. 1961. Biochemical aspects of phagocytosis. *Nature (Lond.)*. **192**: 535-541.
- Paul, B., and A. J. Sbarra. 1968. The role of the phagocyte in host-parasite interactions. XIII. The direct quantitative estimation of H₂O₂ in phagocytizing cells. *Biochim. Biophys. Acta.* **156**: 168-178.
- Babier, B. M., R. S. Kipnes, and J. T. Curnutte. 1973. Biological defense mechanisms: the production by leukocytes of superoxide, a potential bactericidal agent. *J. Clin. Invest.* **52**: 741-744.
- Baehner, R. L., and D. G. Nathan. 1967. Quantitative nitroblue tetrazolium test in chronic granulomatous disease. *N. Engl. J. Med.* **278**: 971-976.
- Holmes, B., A. R. Page, and R. A. Good. 1967. Studies of the metabolic activity of leukocytes from patients with a genetic abnormality of phagocytic function. *J. Clin. Invest.* **46**: 1422-1432.
- Cagan, R. H., and M. L. Karnovsky. 1964. Enzymatic basis of the respiratory stimulation during phagocytosis. *Nature (Lond.)*. **204**: 255-257.
- Baehner, R. L., N. Gilman, and M. L. Karnovsky. 1970. Respiration and glucose oxidation in human and guinea pig leukocytes: comparative studies. *J. Clin. Invest.* **49**: 692-700.
- Rossi, F., and M. Zatti. 1964. Biochemical aspects of phagocytosis in polymorphonuclear leukocytes: NADH and NADPH oxidation by the granules of resting and phagocytic cells. *Experientia (Basel)*. **20**: 21-23.
- Patriarca, P., R. Cramer, S. Moncalvo, F. Rossi, and D. Romeo. 1971. Enzymatic basis of metabolic stimulation of leukocytes during phagocytosis: the role of activated NADPH oxidase. *Arch. Biochem. Biophys.* **145**: 255-262.
- Iyer, G. Y. N., and J. H. Quastel. 1963. NADPH and NADH oxidation by guinea pig polymorphonuclear leukocytes. *Can. J. Biochem. Physiol.* **41**: 427-434.
- Paul, B. B., R. R. Strauss, A. A. Jacobs, and A. J. Sbarra. 1972. Direct involvement of NADPH oxidase with the stimulated respiratory and hexose monophosphate shunt activities in phagocytizing leukocytes. *Exp. Cell Res.* **73**: 456-462.
- Beck, W. S. 1958. Occurrence and control of the phosphogluconate oxidation pathway in normal and leukemic leukocytes. *J. Biol. Chem.* **232**: 271-283.
- Böyum, A. 1968. Isolation of mononuclear cells and granulocytes from human blood. Isolation of mononuclear cells by one centrifugation, and of granulocytes by combining centrifugation and sedimentation at 1 g. *Scand. J. Clin. Lab. Invest. Suppl.* **97**: 77-89.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**: 265-275.
- Baehner, R. L., and M. L. Karnovsky. 1968. Deficiency of reduced nicotinamide-adenine dinucleotide oxidase in chronic granulomatous disease. *Science (Wash., D. C.)*. **162**: 1277-1279.
- Rossi, F., D. Romeo, and P. Patriarca. 1972. Mechanism of phagocytosis-associated oxidative metabolism in polymorphonuclear leukocytes and macrophages. *RES, J. Reticuloendothel. Soc.* **12**: 127-149.
- Roberts, J., and J. H. Quastel. 1964. Oxidation of reduced triphosphopyridine nucleotide by guinea pig polymorphonuclear leukocytes. *Nature (Lond.)* **202**: 85-86.
- Akazawa, T., and E. E. Conn. 1958. The oxidation of reduced pyridine nucleotides by peroxidase. *J. Biol. Chem.* **232**: 403-415.
- Seitz, J. F. 1969. *The Biochemistry of the Cells of Blood and Bone Marrow*. Charles C. Thomas, Publisher, Springfield, Ill. 141.
- Karnovsky, M. L. 1973. Chronic granulomatous disease—pieces of a cellular and molecular puzzle. *Fed. Proc.* **32**: 1527-1533.