



# Cardiac gene expression profiling – the quest for an atrium-specific biomarker

A.H. Maass, A-M.R. De Jong, M.D. Smit, L. Gouweleeuw, R.A de Boer, W.H. Van Gilst, I.C. Van Gelder

Biomarkers are gaining increasing interest to predict risk but also to aid in diagnostics. Tissue-specific biomarkers are of utmost importance to detect diseases of respective organs. As of yet there are no atrium-specific biomarkers for risk stratification of atrial disease, such as atrial fibrillation. Bioinformatics such as mRNA microarrays can help to detect tissue-enriched and possibly tissue-specific expressed genes that can be targets for biomarkers. We describe an approach to identify genes preferably expressed in atrial cardio-

myocytes compared with ventricular cardiomyocytes by RNA microarray and confirmed by quantitative real-time polymerase chain reaction. By this approach we identified several atrium-enriched genes but also ventricle-enriched genes. As expected atrial natriuretic peptide (ANP) mRNA showed higher expression in atrial cardiomyocytes while with adrenergic stimulation expression was almost as high in ventricular as in atrial cells. Brain-type natriuretic peptide (BNP), however, was not different between atrial and ventricular cells giving a possible explanation for increased levels of NT-proBNP in atrial fibrillation patients. Interesting identified candidates are *serpine1* and *Itbp2* as atrium-enriched genes whereas *alpha-adrenergic receptor subtype 1b* and *S100A1* expression was significantly higher in ventricular cells. The identified genes need to be confirmed in human tissue and might ultimately be tested as potential biomarkers for atrial stress. (Neth Heart J 2010;18:610-4.)

**A.H. Maass**

**R.A de Boer**

*Departments of Cardiology and Experimental Cardiology, Thoraxcenter, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands*

**A-M.R. De Jong**

**W.H. Van Gilst**

*Department of Experimental Cardiology, Thoraxcenter, University Medical Center Groningen, University of Groningen, and Interuniversity Cardiology Institute of the Netherlands, the Netherlands*

**M.D. Smit**

*Department of Cardiology, Thoraxcenter, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands*

**L. Gouweleeuw**

*Department of Experimental Cardiology, Thoraxcenter, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands*

**I.C. Van Gelder**

*Department of Cardiology, Thoraxcenter, University Medical Center Groningen, University of Groningen, Groningen, and Interuniversity Cardiology Institute of the Netherlands, the Netherlands*

Keywords: Atrial, Ventricular; Cardiac Myocytes; Fibroblasts; Atrial Fibrillation; Microarray.

**B**iomarkers are increasingly used to predict the risk for certain diseases but also to monitor disease progression. For heart diseases, biomarkers have long been used to measure the degree of myocardial damage with creatine kinase and its isoforms. In recent years, troponin T and troponin I as more sensitive markers for myocardial damage and n-terminal pro-brain type natriuretic peptide (NT-proBNP) are routinely used for risk stratification and monitoring of the clinical situation and success of therapy. Heart failure seems to be a prototypical target for biomarkers as decompensation is difficult to predict with clinical parameters only.<sup>1</sup>

Correspondence to: A.H. Maass

Department of Cardiology, Thoraxcenter, University Medical Center Groningen, University of Groningen, PO Box 30.001, 9700 RB Groningen, the Netherlands

E-mail: a.h.maass@thorax.umcg.nl

Atrial fibrillation (AF) is often a consequence of underlying heart disease such as hypertension and heart failure.<sup>2</sup> Not all patients with these diseases, however, develop atrial arrhythmias.<sup>3</sup> The occurrence of AF is of prognostic relevance. Once AF occurs treatment is often difficult as structural remodelling such as fibrosis has progressed so far that it is irreversible.<sup>4,6</sup> To identify (a set of) biomarkers measuring atrial stress could help to predict patients at risk for atrial arrhythmias and could aid in initiating more aggressive therapy and to test efficacy of new or known treatment regimens.<sup>7</sup>



### Methods

Neonatal rat atrial and ventricular myocytes (NRAMs, NRVMs) and fibroblasts were isolated from one- to two-day-old Sprague Dawley rats by trypsin digestion, as described previously.<sup>8</sup> For microarray analysis total RNA was isolated using TRIzol reagent (Invitrogen Corporation), and cleaned using the Nucleospin II kit (Macherey-Nagel). Processing of RNA and microarray analysis was performed in the core lab (Department of Genetics, University Medical Center Groningen, University of Groningen, the Netherlands). RNA was processed using the Ambion RNA kit (Applied Biosystems) and Illumina Rat Ref12 Expression Platform was used for the microarray. Quantitative real-time polymerase chain reaction (qRT-PCR) was used to confirm microarray data of selected genes and to investigate effects of stimulation with 20  $\mu$ M of the  $\alpha$ -adrenergic agonist phenylephrine for 24 hours on both NRAM and NRVM.<sup>9</sup> Total RNA was isolated using the Nucleospin II kit (Macherey-Nagel) and converted to cDNA by QuantiTect Reverse Transcription (Qiagen). Gene ex-

pression was measured with Absolute QPCR SYBR Green ROX Mix (Abgene) in the presence of 7.5 ng cDNA and 200 nM forward and reverse primers. qRT-PCR was conducted on the Biorad CFX384 (Biorad). Gene expression levels were corrected for 36b4 reference gene expression, and values were expressed relative to NRVM control levels. Primers used were ANP forward atgggctctctctcatcac, ANP reverse tctaccgcatcttctcctc, BNP forward acaatccagatgcagaagct, BNP reverse gggccttggtcctttgaga,  $\alpha$ -myosin heavy chain ( $\alpha$ MHC) forward gacaactctcccgtttgg,  $\alpha$ MHC reverse aagatcaccgggacttctc, Adra1b forward aaccttgggcattgtagtctg, Adra1b reverse tgaggcagctgttgaagtag, serpine1 forward cggcagcagatccaagatg, serpine1 reverse ggtccc-gctggacaagatg, S100a1 forward ggagaccctcatcaatgtg, S100a1 reverse cagcatctgcatccttctg, Wisp2 forward ttctggccacttctctcctc, Wisp2 reverse ttacagcagccacagc-catc, pln forward ttgtcttcttgccatcatgg, pln reverse cagcttgtcacagaagcatcac, 36b4 forward gttgctcagtgctcactc, 36b4 reverse gcagccgcaatgcagatgg. Microarray analysis was performed using GeneSpring GX (Version 10.0, Agilent Technologies). Statistical significance was tested using unpaired t-test with Benjamini-Hochberg correction. qRT-PCR results are expressed as mean values  $\pm$  standard error of the mean (SEM) and statistical analysis was performed using Mann-Whitney u-test, using SPSS (Version 16, SPSS Inc). Values of  $p < 0.05$  were considered statistically significant.

### Results

Atrial and ventricular cells showed no significant morphological difference besides a slightly larger cell size of NRAM (figure 1).



*Figure 1. Primary neonatal rat atrial (1A) and ventricular (1B) myocytes and fibroblasts were stained for  $\alpha$ -actinin with specific antibodies (green) and total actin was stained with Texas red-phalloidin. Cardiac myocytes stain both red and green, whereas fibroblasts stain red for actin exclusively. Nuclei are stained blue with DAPI.*

**Table 1A.** Atrium-enriched genes.

Gene symbol	Fold A/V microarray	P value	Fold A/V PCR	P value	Protein
Myl7-pred	7.9	0.005	nd		Regulatory myosin light chain7
Ltb2	4.5	0.03	2.92	0.01	Latent TGF- $\beta$ binding protein2
Serpine1	4.37	0.01	4.76	<0.01	Serine peptidase inhibitor
Nppa	3.33	0.03	5.98	<0.01	Atrial natriuretic peptide
Wisp2	2.52	0.08	2.47	0.07	WNT1 inducible signalling pathway
Myh6	2.02	0.03	1.63	0.02	$\alpha$ -myosin heavy chain

A/v= ratio atrial vs. ventricular expression, nd= not determined.


**Table 1B.** Ventricle-enriched genes.

Gene symbol	Fold V/A microarray	P value	Fold V/A PCR	P value	Protein
Adra1b	6.19	0.01	4.68	0.03	Adrenergic $\alpha$ -1b receptor
Ckm	3.62	0.03	nd		Muscle creatine kinase
Pln	2.49	0.06	1.59	0.01	Phospholamban
S100A1	2.43	0.02	2.31	0.02	S100 calcium binding protA1

V/A= ratio ventricular vs. atrial expression, nd= not determined.

We excluded genes with low abundance and the ones that showed less than twofold difference from our further analysis. This yielded 80 differentially expressed genes with statistical significance. Furthermore, we focused on genes with a known function and possible physiological or pathophysiological roles in cardiac diseases. Not surprisingly, atrial natriuretic peptide (ANP) mRNA showed a sixfold higher expression in NRAM. Table 1 lists the most important differentially expressed genes. In table 1A atrial-enriched genes are shown whereas table 1B lists important ventricle-enriched genes. There were some differences in known cardiac structural genes such as MHC, creatine kinase, phospholamban, and regulatory myosin light chain. As further interesting candidates, we investigated serpine1 and wisp2, which showed fivefold and 2.5-fold higher expression in NRAMs, respectively. The alpha-adrenergic receptor subtype 1b and S100A1 showed fivefold and 2.3-fold higher expression in NRVMs, respectively. Latent transforming growth factor beta-binding protein 2 (ltb2) was expressed threefold higher in atrial cells, but further analysis showed that it is exclusively expressed in cardiac fibroblasts. Even though our analysis was done in isolated cardiac myocytes the cultures are always contaminated with 10 to 20% non-myocytes, whereas in the heart 90% of cells are non-myocytes.

To determine if there were differences between natriuretic peptides under basal conditions and stress, we treated the cells with the strong hypertrophic stimulus phenylephrine. This led to a fivefold increase of ANP and BNP in NRVM. Levels of ANP under these pathological conditions were almost as high as in NRAM (figure 2).



*Figure 2. Comparison of mRNA expression of atrial (ANP) and brain-type (BNP) natriuretic peptides in primary neonatal rat atrial (NRAM) and ventricular (NRVM) myocytes. Cells were also treated with 20  $\mu$ M phenylephrine (PE) to induce hypertrophy which leads to increases in both ANP and BNP in NRVM only to levels comparable with NRAM at baseline.*

### Discussion

We demonstrate differentially expressed genes in atrial and ventricular cardiac myocytes via RNA microarray that are confirmed via qRT-PCR. These results must clearly be confirmed in human tissue before we think about using them as biomarkers of atrial stress or disease. We identified, however, several interesting candidates including serpine1 and ltb2.



Serpine1, also known as plasminogen activator inhibitor 1 (PAI1), is a pleiotropic factor involved in multiple processes, such as inflammation, fibrinolysis, and insulin resistance.<sup>10</sup> So far atrial-enriched expression had not been described. PAI-1 has been implicated in the development of atrial fibrillation. Elevated levels of PAI-1 independently predict development of AF after cardiac surgery.<sup>11</sup>

Ltbp2 is a fibroblast-specific factor and the differences we demonstrate with the microarrays are caused by differences in atrial and ventricular fibroblasts (data not shown). It has been suggested that cardiac fibroblasts can also produce natriuretic peptides.<sup>12</sup> In our hands, however, expression in fibroblasts was 100-fold lower than in cardiac myocytes (data not shown). There seems to be a profound difference between atrial and ventricular fibroblasts. Atrial fibroblasts seem to respond stronger to transforming growth factor  $\beta$  (TGF- $\beta$ ).<sup>13</sup> Ltbp2 could be involved in this mechanism, as it has been suggested to enhance TGF- $\beta$  function.

There are obviously no optimal biomarkers. Individual markers or panels for certain situations can be envisioned. If a biomarker for tissue damage is to be developed, this protein needs to be tissue-specific and once it is released into the circulation, half-life should not be too short. To detect stress instead of damage, the biomarker needs to be a secreted protein, again with a decent half-life in plasma or serum.

A problem with biomarkers might be regulation through pathological signals leading to 'aberrant' expression. A good example is the increase in ANP mRNA in hypertrophic NRVM as shown in our study. Levels can reach the levels in NRAM with adrenergic stimulation. NT-proBNP is a prototypical marker for ventricular stress. We show here that NRAM express as much BNP mRNA as NRVM. Stress can lead to an increase in BNP mRNA in NRAM. This might add to the high levels of NT-proBNP in patients with AF. NT-proBNP has been shown to be an independent predictor of the development of AF.<sup>14</sup> In advanced heart failure, however, we showed that AF affects (NT-pro)ANP levels, but not (NT-pro)BNP levels, although NT-proBNP was an independent determinant of prognosis in advanced heart failure, irrespective of the rhythm, AF or sinus rhythm.<sup>15</sup> Nevertheless, the previous concept of ventricular stress only leading to increase in serum NT-pro-BNP needs to be revised.

### Outlook

Further experiments are aimed at utilising human cardiac tissue to look for differentially expressed genes in patients. This is troubled by cell composition but our approach has shown that this approach does not preclude identification of differentially expressed genes from certain cell types as we detected cardiomyocyte-specific as well as fibroblast-specific genes. We thus envision finding structural markers for atrial myocytes but also genes that are specific for atrial versus ventricular fibroblasts. Microarray experiments can obviously be combined with the candidate approach. As an example for this approach, a recent publication has

suggested endothelin-1 as an important player in atrial fibrillation in patients with structural heart disease.<sup>16</sup>

A novel approach will be to look for atrial specific microRNAs (miRNAs). Pinto's group has recently demonstrated that circulating miRNAs can be detected in patient blood and that certain miRNAs are enriched in patients with heart failure.<sup>17</sup> It is possible that there are atrium-specific miRNAs that can also be detected in blood.

To identify biomarkers for atrial stress, we will use a secretion trap screen with human atrial cDNA expressed in secretion-deficient yeast to identify known or novel secreted proteins.<sup>18</sup> We hope to find a secreted factor with sufficient atrial enrichment that is abundant and stable enough to be measured in human serum or plasma.

Treatment of patients with AF is currently undergoing a shift of paradigm. The most relevant outcomes are related to patient well being and objective measures of cardiovascular morbidity. In symptomatic patients this includes abolishment of AF. To improve outcome in patients with AF, patient-tailored therapy could be aided by biomarkers. In addition to identifying patients at risk of AF they could help in stratifying therapy. We still do not know which patients will fail rhythm control therapy. On the other hand, in patients where we chose for rate control it might be necessary to identify patients with an adverse outcome who need more aggressive therapy. ■

### References

- 1 Braunwald E. Biomarkers in heart failure. *N Engl J Med*. 2008;358:2148-59.
- 2 Camm AJ, Kirchhof P, Lip GYH, Schotten U, Savelieva I, Ernst S, et al. Guidelines for the management of atrial fibrillation. *Eur Heart J*. 2010; 31:2369-429.
- 3 Buck S, Rienstra M, Maass AH, Nieuwland W, Van Veldhuisen DJ, Van Gelder IC. Cardiac resynchronization therapy in patients with heart failure and atrial fibrillation: importance of new-onset atrial fibrillation and total atrial conduction time. *Europace*. 2008;10:558-65.
- 4 Cosio FG, Aliot E, Botto GL, Heidebuchel H, Geller CJ, Kirchhof P, et al. Delayed rhythm control of atrial fibrillation may be a cause of failure to prevent recurrences: reasons for change to active antiarrhythmic treatment at the time of the first detected episode. *Europace*. 2008;10:21-7.
- 5 Van Gelder IC, Crijns HJ, Tieleman RG, Brugemann J, De Kam PJ, Gosselink AT, et al. Chronic atrial fibrillation. Success of serial cardioversion therapy and safety of oral anticoagulation. *Arch Intern Med*. 1996;156:2585-92.
- 6 Ahmed S, Rienstra M, Crijns HJ, Links TP, Wiesfeld AC, Hillege HL, et al. Continuous vs episodic prophylactic treatment with amiodarone for the prevention of atrial fibrillation: a randomized trial. *JAMA*. 2008;300:1784-92.
- 7 Smit MD, Van Gelder IC. Will we be able to predict in which atrial fibrillation patients a rhythm control strategy will be successful? *Europace*. 2009;11:846-7.
- 8 Maass A, Langer SJ, Oberdorf-Maass S, Bauer S, Neyses L, Leinwand LA. Rational promoter selection for gene transfer into cardiac cells. *J Mol Cell Cardiol*. 2003;35:823-31.
- 9 Maass AH, Ikeda K, Oberdorf-Maass S, Maier SK, Leinwand LA. Hypertrophy, fibrosis, and sudden cardiac death in response to pathological stimuli in mice with mutations in cardiac troponin T. *Circulation*. 2004;110:2102-9.
- 10 Cesari M, Pahor M, Incalzi RA. Plasminogen Activator Inhibitor-1 (PAI-1): A Key Factor Linking Fibrinolysis and Age-Related Subclinical and Clinical Conditions. *Cardiovasc Ther*. 2010;28:e72-9.
- 11 Pretorius M, Donahue BS, Yu C, Greelish JP, Roden DM, Brown NJ. Plasminogen activator inhibitor-1 as a predictor of postoperative atrial fibrillation after cardiopulmonary bypass. *Circulation*. 2007;116(11 Suppl):I1-7.

- 12 Jarai R, Kaun C, Weiss TW, Speidl WS, Rychli K, Maurer G, et al. Human cardiac fibroblasts express B-type natriuretic peptide: fluvastatin ameliorates its up-regulation by interleukin-1alpha, tumour necrosis factor-alpha and transforming growth factor-beta. *J Cell Mol Med.* 2009;13:4415-21.
- 13 Burstein B, Libby E, Calderone A, Nattel S. Differential behaviors of atrial versus ventricular fibroblasts: a potential role for platelet-derived growth factor in atrial-ventricular remodeling differences. *Circulation.* 2008;117:1630-41.
- 14 Patton KK, Ellinor PT, Heckbert SR, Christenson RH, DeFilippi C, Gottdiener JS, et al. N-terminal pro-B-type natriuretic peptide is a major predictor of the development of atrial fibrillation: the Cardiovascular Health Study. *Circulation.* 2009;120:1768-74.
- 15 Rienstra M, Van Gelder IC, Van den Berg MP, Boomsma F, Van Veldhuisen DJ. Natriuretic peptides in patients with atrial fibrillation and advanced chronic heart failure: determinants and prognostic value of (NT-) ANP and (NT-pro)BNP. *Europace.* 2006;8:482-7.
- 16 Mayyas F, Niebauer M, Zurick A, Barnard J, Gillinov AM, Chung MK, et al. Association of left atrial endothelin-1 with atrial rhythm, size, and fibrosis in patients with structural heart disease. *Circ Arrhythm Electrophysiol.* 2010;3:369-79.
- 17 Tijssen AJ, Creemers EE, Moerland PD, de Windt LJ, van der Wal AC, Kok WE, et al. MiR423-5p as a circulating biomarker for heart failure. *Circ Res.* 2010;106:1035-9.
- 18 Frost RJ, Engelhardt S. A secretion trap screen in yeast identifies protease inhibitor 16 as a novel antihypertrophic protein secreted from the heart. *Circulation.* 2007;116:1768-75.