

Magnesium citrate with a single dose of sodium phosphate for colonoscopy bowel preparation

Yong Sung Choi, Jung Pil Suh, Jong Kyu Kim, In Taek Lee, Eui Gon Youk, Doo Seok Lee, Do Sun Kim, Doo Han Lee

Yong Sung Choi, Jung Pil Suh, Jong Kyu Kim, Department of Gastroenterology, Daehang Hospital, Seoul 137-820, South Korea
In Taek Lee, Eui Gon Youk, Doo Seok Lee, Do Sun Kim, Doo Han Lee, Department of Surgery, Daehang Hospital, Seoul 137-820, South Korea

Author contributions: Choi YS performed data analysis and edited the manuscript; Suh JP and Kim JK were involved in editing the manuscript; Lee IT, Youk EG and Lee DS were involved in data collection and analysis; Kim DS and Lee DH provided financial support for publishing this work.

Correspondence to: Jung Pil Suh, MD, Department of Gastroenterology, Daehang Hospital, Seoul 137-820, South Korea. yschoi71@dreamwiz.com

Telephone: +82-2-63888114 Fax: +82-2-63888115

Received: July 20, 2010 Revised: August 17, 2010

Accepted: August 24, 2010

Published online: January 14, 2011

Abstract

AIM: To evaluate the efficacy and acceptability of magnesium citrate and a single dose of oral sodium phosphate (45 mL) solution for morning colonoscopy bowel preparation.

METHODS: A total of 159 patients were randomly assigned to receive two split doses of 90 mg of sodium phosphate (Group I, $n = 79$) or magnesium citrate (250 mL, the day before the procedure) followed by 45 mL of sodium phosphate (the day of procedure, Group II, $n = 80$). The quality of bowel cleansing and the acceptability of each regimen were compared, including the satisfaction, taste, willing to repeat and adverse effects of each regimen.

RESULTS: The quality of bowel cleansing of Group II was as good as that of Group I (An Aronchick scale score of good or excellent: 70.9% vs 81.0%, respectively, $P = 0.34$; the Ottawa system score: 4.4 ± 2.6 vs 3.8 ± 3.0 , respectively, $P = 0.76$). There was no statistically

significant difference between both groups with regard to acceptability, including the satisfaction, taste and willingness to repeat the regimen. A significantly greater number of older patients (over 65 years old) in Group II graded the overall satisfaction as satisfactory (48.1% vs 78.1%, respectively; Group I vs Group II, $P = 0.01$). There were no significant adverse reactions.

CONCLUSION: Magnesium citrate and a single dose of sodium phosphate was as effective and tolerable as the conventional sodium phosphate regimen and is a satisfactory option.

© 2011 Baishideng. All rights reserved.

Key words: Colonoscopy; Bowel preparation; Efficacy; Acceptability; Magnesium citrate; Sodium phosphate

Peer reviewer: Josep M Bordas, MD, Department of Gastroenterology IMD, Hospital Clinic, Llusanes 11-13 at, Barcelona 08022, Spain

Choi YS, Suh JP, Kim JK, Lee IT, Youk EG, Lee DS, Kim DS, Lee DH. Magnesium citrate with a single dose of sodium phosphate for colonoscopy bowel preparation. *World J Gastroenterol* 2011; 17(2): 242-248 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v17/i2/242.htm> DOI: <http://dx.doi.org/10.3748/wjg.v17.i2.242>

INTRODUCTION

The worldwide introduction of screening colonoscopy programs has led to a dramatic increase in the number of colonoscopies. Despite the development of techniques and instruments, the difficulties in bowel preparation for a complete colonoscopic examination still remain a challenge to the physician and the examinee. These difficulties may result in significant noncompliance and thus subop-

timal preparation, which subsequently may lead to missed lesions^[1], an increased procedure time and the need for repeat colonoscopy^[2], as well as reluctance to undergo follow-up examinations.

Sodium phosphate is a low-volume laxative, and it is known to be superior to 4l polyethylene glycol (PEG) solution, both for acceptability by the patient and for the quality of bowel cleansing^[3-6]. However, a significant proportion of patients complain that its bad taste is unacceptable and there are adverse reactions such as nausea, vomiting and headache. Moreover, more potential adverse reactions associated with hyperphosphatemia may theoretically develop under any conditions that increase absorption or decreased elimination of phosphate^[7]. Unfortunately, poor compliance with the instructions about the dosage or interval of oral sodium phosphate and undetected chronic illness, particularly in the elderly, may result in a higher possibility of unexpected adverse events. Recent reports of renal failure associated with sodium phosphate have raised increasing concerns^[8-12] in spite of some reports demonstrating its safety in a series of clinical trials^[13,14].

In this study, we wanted to find a better and easier preparation regimen for a complete colonoscopic examination. We have experienced that the usual bowel preparation for sigmoidoscopy in our hospital using magnesium citrate the day before the procedure scheduled in the next morning was able to debulk the colon of solid fecal material without much patient discomfort. Therefore, we hypothesized that taking magnesium citrate the day before the procedure and then taking a single dose of sodium phosphate (45 mL) on the day of procedure would be as effective a regimen as the conventional two doses of sodium phosphate and more tolerable for morning colonoscopy.

MATERIALS AND METHODS

Study design, patient enrollment and the end outcomes

This was a prospective, randomized, endoscopist-blinded, controlled study. All consecutive patients referred to the Outpatient Department of Gastroenterology and Surgery at Daehang Hospital (Seoul, Korea) between September 2009 and December 2009 for colonoscopy in the morning were enrolled in this clinical study. The exclusion criteria were as follows: ileus or suspected bowel obstruction, significant gastroparesis or gastric outlet obstruction, the presence of serious medical conditions such as severe cardiac, renal, hepatic or metabolic diseases, pregnancy, lactation and a history of prior colon or rectal surgery. Written informed consent was obtained from each patient. No patient refused to participate. The primary end outcomes in this study were the efficacy of bowel preparation according to the Aronchick scale and the Ottawa Bowel Preparation Scale (OBPS), the patients' tolerance, the overall satisfaction, the taste and the willingness to repeat the same preparation regimen, if necessary. The sec-

ondary end outcomes included assessment of the OBPS components (i.e. the right vs middle vs left colon, and the fluid score) and a subgroup assessment according to age, gender, a prior bowel cleansing regimen and so forth.

Colon bowel preparation

A total of 159 patients were randomly assigned to receive two split doses of 90 mg of oral sodium phosphate (Group I, $n = 79$) or 250 mL of magnesium citrate (magnesium carbonate 4.3 g and anhydrous citric acid 7.8 g/100 mL) (the day before the procedure) and this was followed by 45 mL of sodium phosphate (the day of the procedure, Group II, $n = 80$). Group I ingested a single dose (45 mL) of sodium phosphate starting at 8:00 PM on the day before the procedure and then they ingested the remaining 45 mL of sodium phosphate at 5:00 AM on the day of colonoscopy. Group II ingested magnesium citrate (one bottle; 250 mL) at 8:00 PM on the preceding day and a single dose (45 mL) of sodium phosphate at 5:00 AM on the procedure day. Both groups took one tablet of bisacodyl as pretreatment 30 min before each laxative on the day before the procedure. All the patients who had taken oral sodium phosphate were instructed to drink an additional 1-1.5 L of water. Both groups had a thick liquid diet for dinner on the day before the procedure, and they took nothing further by mouth after 6:00 PM. All the colonoscopy procedures were performed between 8:30 AM and 11:30 AM. The quality of bowel cleansing, the tolerability, the satisfaction, the willing to repeat the procedure and the adverse effects of each regimen were compared.

Evaluation of bowel preparation

Efficacy of bowel cleansing: The endoscopists, who were blinded to the form of preparation, performed the colonoscopy exams. To standardize and minimize interobserver variation, we had a meeting before the start of this study and an investigator gave a standardized explanation for assessing the quality of bowel preparation during the meeting. A poster with endoscopic photos that demonstrated the scoring system was placed in each endoscopy room as a reminder. The endoscopists were also requested to take pictures of each colon segment before cleansing the segments with saline during the procedure.

Bowel cleansing was assessed at the end of the colonoscopy using the OBPS and the Aronchick scale. The OBPS has a potential score ranging from 0 (excellent preparation, no fluid) to 14 (inadequate in all segments with a large amount of fluid) (Table 1). The Aronchick scale was as follows: excellent (a small volume of clear liquid or greater than 95% of the surface was seen); good (a large volume of clear liquid covering 5% to 25% of the surface but greater than 90% of the surface was seen); fair (some semisolid stool that could be suctioned or washed away but greater than 90% of the surface was seen); and poor (semisolid stool that could not be suctioned or washed away and less than 90% of the surface was seen).

There were ten colonoscopists who performed colonos-

Table 1 Summary of the scoring system for the Ottawa Bowel Preparation Scale

Cleanliness	
Excellent (0): mucosal detail clearly visible. If fluid present, it is clear. Almost no stool residue	
Good (1): some turbid fluid or stool residue but mucosal detail still visible. Washing and suctioning not necessary	
Fair (2): turbid fluid or stool residue obscuring mucosal detail and contour. However, mucosal detail becomes visible with suctioning. Washing not necessary	
Poor (3): presence of stool obscuring mucosal detail and contour. However, with suctioning and washing, a reasonable view is obtained	
Inadequate (4): solid stool obscuring mucosal detail and contour despite aggressive washing and suctioning	
Fluid: small (0), moderate (1), large (2)	

Table 2 Baseline characteristics (mean ± SD) n (%)

Variables	Group I (n = 79)	Group II (n = 80)	P value
Age (yr), median (range)	59 (46-77)	60 (48-78)	NS
Male	40 (50.6)	50 (62.5)	NS
Cecal intubation time (s)	300.6 ± 171.9	314 ± 190.5	NS
Weight (kg)	61.5 ± 15.6	64.1 ± 15.3	NS
Prior colonoscopy	45 (56.9)	44 (55.0)	NS
With PEG	11 (13.9)	14 (17.5)	
With Sodium phosphate	34 (43.0)	30 (37.5)	
Hypertension	12 (15.2)	15 (18.8)	NS
Diabetes	8 (10.1)	5 (6.3)	NS
Constipation (< 3/wk)	12 (15.2)	14 (17.6)	NS

PEG: Polyethylene glycol; NS: Not significant.

copies in this study. All the colonoscopists were experts who had experienced at least 1000 cases of colonoscopy after their gastrointestinal or coloproctologic fellowship. To check the interobserver variation for rating the efficacy of bowel preparation, all the colonoscopy pictures were reviewed by two experienced endoscopists (Choi YS and Suh JP) who were blinded to the preparation methods.

Patient acceptability

All the patients completed a questionnaire before colonoscopy. It was used to assess the past medical history, compliance, discomfort symptoms during preparation and the overall satisfaction. The investigated symptoms were nausea or vomiting (present = 1, absent = 0), abdominal pain or discomfort (present = 1, none = 0) and abdominal distension (present = 1, none = 0). The overall satisfaction with the preparation was rated on a 3 scale (satisfactory, fair and unsatisfactory). The willingness to repeat the same preparation method in the future, if necessary, was also evaluated with a “yes” or “no” question.

Statistical analysis

All the continuous variables are presented as means and standard deviations or as medians and ranges, as appropriate. The χ^2 test was used for comparisons between categorical variables and Student’s *t* test was used for com-




Figure 1 Quality of bowel cleansing. A: Quality of bowel cleansing. The patients are graded as good or excellent quality vs fair or poor quality with the conventional sodium phosphate regimen (Group I) or the magnesium citrate and a single dose sodium phosphate regimen (Group II); B: Segmental quality of bowel cleansing. The patients are scored with colon segment grades of poor or inadequate with the conventional sodium phosphate regimen (Group I) or the magnesium citrate and a single dose sodium phosphate regimen (Group II). NS: Not significant.

parisons between continuous variables. *P* values less than 0.05 were regarded as statistically significant.

RESULTS

Baseline characteristics

A total of 180 consecutive patients were enrolled and randomized into two groups (90 patients in each group). Twenty-one patients were excluded because the procedure was postponed or canceled. Finally, 159 patients were included in the analysis. There was no significant difference in the baseline characteristics between both groups (Table 2). In all cases, cecal intubation was successful. There were no serious complications or adverse events immediately after the procedure.

Assessment of the quality of bowel cleansing

There was no significant difference in bowel cleansing quality as evaluated by the Aronchick scale between both bowel preparation regimens. Group I had 56 satisfactory preparations (defined as excellent or good) of 79 (70.9%) compared with 62 of 80 (81.0%) for Group II (*P* = 0.34) (Figure 1A).




Figure 2 Patient acceptability. A: For overall satisfaction, 65.8% in Group I and 82.0% in Group II replied that they were “satisfied” or it was “fair”; B: For assessing taste, the percentage of patients who responded “good or tolerable” was not significantly different between Group I (69.6%) and Group II (73.4%); C: For the willingness to repeat the regimen, the percentage of patients who responded “yes” was not significantly different between Group I (83.5%) and Group II (80.8%). NS: Not significant.

When bowel preparation was assessed by the Ottawa scale, the preparation quality was not significantly different as well (Table 3). The mean score was not significantly different between both groups for the right colon ($P = 0.54$), the mid colon ($P = 0.47$), the rectosigmoid colon ($P = 0.97$) or the total amount of fluid ($P = 0.48$). In both groups, the patients with colon segment grades of poor or inadequate were 12 (15.2%) *vs* 15 (18.8%), 5 (6.3%) *vs* 6 (8.0%) and 9 (11.4%) *vs* 7 (9.0%) for the right colon, the mid colon and the rectosigmoid colon, respectively (Figure 1B).

Table 3 Efficacy of bowel cleansing by Ottawa scoring system (mean \pm SD)

Variables	Group I ($n = 79$)	Group II ($n = 73$)	P value
Right colon	1.4 \pm 1.1	1.3 \pm 1.1	NS
Mid colon	1.0 \pm 0.9	0.8 \pm 0.9	NS
Rectosigmoid colon	1.3 \pm 0.9	1.2 \pm 1.0	NS
Fluid present	0.7 \pm 0.6	0.4 \pm 0.6	NS
Overall score	4.4 \pm 2.6	3.8 \pm 3.0	NS

NS: Not significant.




Figure 3 Overall satisfaction with bowel preparation in the elderly (> 65 years). 48.1% in Group I and 78.1% in Group II replied that they were “satisfied” or it was “fair” ($P = 0.017$).

There was no significant difference of the segmental bowel preparation quality either.

Assessment of patient acceptability (overall satisfaction, taste and willing to repeat)

The percentage of patients with symptoms such as abdominal distension with the two preparation regimens was 23% *vs* 34%, respectively, nausea or vomiting 45% *vs* 41%, respectively, and abdominal pain 15% *vs* 13%, respectively.

When patients were asked whether they were satisfied with the bowel preparation regimen, 65.8% (52 of 79) in Group I and 82.0% (62 of 80) in Group II replied that they were “satisfied” or it was “fair” (Figure 2A). When patients were asked how the laxatives tasted, the patients who responded “good or tolerable” were not significantly different between Group I (69.6%) *vs* Group II (73.4%) (Figure 2B). When patients were asked whether they were willing to repeat the same preparation regimen if necessary, the percentage of patients who responded “yes” was not significantly different between Group I (83.5%) *vs* Group II (80.8%) (Figure 2C).

Subgroup assessment of the bowel cleansing quality and patient acceptability

When the patients over 65 years old were asked whether they were satisfied with the bowel preparation regimen, 48.1% (13 of 27) in group I and 78.1% (25 of 32) in group II replied that they were “satisfied” or it was “fair” ($P = 0.017$) (Figure 3). The question about taste and will-

ingness to repeat revealed no significant differences between both groups.

When only the patients who had previously undergone colonoscopy and had taken PEG or oral sodium phosphate were asked about patient acceptability, the prior sodium phosphate users generally had a preference for this new regimen rather than the new PEG users concerning overall satisfaction (80.0% *vs* 64.3%, respectively, $P = 0.26$), taste tolerance (80% *vs* 71.4%, respectively, $P = 0.52$) and willing to repeat (90.0% *vs* 61.3%, respectively, $P = 0.04$).

DISCUSSION

The early detection of colonic neoplasm *via* colonoscopy and its removal by polypectomy is popular and increasing, yet many people avoid screening colonoscopy because they fear the preparation required for colonoscopy. The two most commonly-used regimens for colonoscopic bowel preparation are currently a large volume of osmotically balanced PEG solution and a small volume of an osmotically active agent, oral sodium phosphate. Four liters of PEG solution is known to be safe and effective, but poorly tolerated. In contrast, sodium phosphate is known to be well-tolerated and as effective as PEG solution. Oral sodium phosphate has been shown to be very safe in a large series of clinical trials^[13,14]. Its administration is known to be associated with only an asymptomatic, mild increase in serum phosphate (mean increase: 3 mg/dL) and a minimal decrease in serum calcium (mean decrease: 0.3 mg/L); these changes are readily reversible^[10]. Clinically significant adverse reactions, including severe hyperphosphatemia, hypocalcemia, seizure, tetany, hypovolaemia and acute renal failure are thought to be extremely rare, with a reported incidence of 1.44 per million patients^[10,15]. Under normal conditions, phosphate is absorbed in the small intestine and eliminated by the kidney as calcium phosphate^[15]. The conventional dosage of oral phosphate for bowel cleansing is one bottle (45 mL) the evening before the procedure and another bottle the next morning. Any condition that can increase absorption or decrease elimination of phosphate such as repeated dosing, dehydration, the use of diuretics and ACE inhibitor, or a preexisting illness that may impair phosphate homeostasis, for example, hyperparathyroidism, renal insufficiency or delayed intestinal transit can be a risk factor. Indeed, fatal cases have been observed among the patients with a history of renal dysfunction^[16-18], ischemic colitis^[17], cirrhosis^[18] and in elderly patients with normal renal function^[19,20]. The maximum safe dose of sodium phosphate is 90 mL^[21]. Several studies on the adverse effects of high doses of sodium phosphate have suggested that these high doses should be avoided^[13-15]. If the recommended dose of 60 g (90 mL) is surpassed, or if the interval between doses is < 5 h, severe hyperphosphatemia could develop^[10,15,18,22,23]. Fine *et al*^[24] have found that the mortality rate was 33%, and that the risk of death was high if the serum phosphate level increased beyond 32.69 mg/dL (10.56 mmol/L). Most of the deaths reported in the lit-

erature have been caused by arrhythmia or heart attack associated with electrolyte changes and dehydration^[25]. Therefore, adequate fluid intake is necessary. Markowitz *et al*^[10] has suggested that patients must be encouraged to drink eight cups of fluids (1920 mL) and Rex *et al*^[2] have promoted taking 3.6 L of clear fluids. However, it was reported that the standard dose of sodium phosphate could induce the development of hyperphosphatemia even in low-risk and well-hydrated patients^[7].

In this study, we attempted to debulk the colon of solid fecal material with magnesium citrate at first and thereby decrease the volume of oral sodium phosphate required for bowel cleansing without compromising the quality of the bowel preparation. Rather, we achieved superior overall satisfaction in the elderly patients who were pretreated with magnesium citrate. It is practically significant to find that this new regimen is better tolerated by elderly patients since the potential for complications associated with bowel preparation may be relatively increased in elderly patients due to poor compliance with the recommended fluid intake. In addition, elderly patients have a greater possibility of having an underlying chronic illness that may be undetected. Therefore, we think a lower volume of sodium phosphate can be safer and better suited for everybody. The patients who had previously taken two doses of sodium phosphate for bowel preparation had a tendency to prefer this new regimen rather than the PEG solution, suggesting that this regimen can be an alternative option for the intolerant patients.

We found that overall rates of adequate bowel cleansing in both groups were relatively high (Group I, 70.9% and Group II, 81%), which suggested that bisacodyl as pretreatment and procedure time scheduled for the morning had influenced the overall quality of bowel cleansing. Both regimens were scheduled for morning because afternoon colonoscopies have higher failure rates than morning colonoscopies, and the suggested scheduling of all outpatient colonoscopies is preferentially in the morning to avoid suboptimal procedures^[26]. The dosing time of magnesium citrate was determined as 9 h before taking the oral sodium phosphate. In a previous study that used magnesium citrate as pretreatment to decrease the volume of PEG solution, magnesium citrate was taken at 1-2 h before ingesting 2 L of PEG solution^[27,28]. However, we expect that a 9 h time interval would be optimal, with reference to the usual interval time for the split-dosed PEG solution regimen or the oral sodium phosphate regimen.

Our study has several limitations. First, the safety profiles, including electrolytes, hemoglobin and the BUN/creatinine change, were not evaluated to prove the safety advantages of this new regimen compared to the conventional sodium phosphate regimen, although the volume of sodium phosphate was reduced by half. Second, we excluded the patients with medical conditions that are currently contraindicated for conventional sodium phosphate use such as heart failure, cirrhosis, renal disease, very old age patients or those taking diuretics or ACE inhibitors and so forth. Thus, the findings in our study need to be

confirmed in a large study with various groups of patients. Third, there were ten colonoscopists involved in this study. Although all the endoscopists were trained in using the scale with photographs illustrating each point on the OBPS, and the degree of agreement was acceptable according to a κ coefficient of greater than 0.4, the study could have been more powerful if all the colonoscopies had been performed by a single colonoscopist. Fourth, there was a discrepancy between “better” overall satisfaction and a “similar” propensity of willingness to repeat the bowel preparation regimen among the elderly patients. Further tailored evaluation of a large group of elderly patients is needed to verify this.

As mentioned above, the use of the conventional dose of sodium phosphate is theoretically limited for patients with any condition that can increase the absorption or decrease the elimination of phosphate. If we think a lower volume of sodium phosphate might be safer, then this must be verified in individual cases since the mechanism of an adverse reaction to sodium phosphate and its risk factors has not been fully discovered. Further evaluation of the safety profile is mandatory.

In conclusion, a regimen of magnesium citrate and a single dose of oral sodium phosphate was as effective as the conventional two doses of sodium phosphate, and our new regimen was well tolerated. Therefore, this regimen could be a good option to routinely prepare for morning colonoscopy.

COMMENTS

Background

Oral sodium phosphate is a hyperosmolar, low-volume laxative for colonoscopy bowel preparation, and it is known to be as effective as polyethylene glycol solution. It is generally better tolerated, but the bad taste and uncomfortable abdominal symptoms such as nausea and vomiting frequently lead to poor compliance with the oral sodium phosphate regimen, and all this can subsequently cause incomplete bowel cleansing. Moreover, potential adverse reactions associated with hyperphosphatemia may develop.

Innovations and breakthroughs

This is the first clinical study to evaluate the efficacy and tolerability of magnesium citrate and a single dose of oral sodium phosphate for bowel preparation prior to morning colonoscopy. The findings demonstrated this modified regimen, in which the volume of sodium phosphate was reduced by half was as effective and tolerable as the conventional sodium phosphate regimen.

Applications

The study results suggest that magnesium citrate and a single dose of oral sodium phosphate can be an effective and satisfactory option to routinely prepare for morning colonoscopy.

Terminology

Magnesium citrate acts as an osmotic laxative which is often used for bowel preparation

Peer review

In this well designed paper the author has proposed a combination of magnesium citrate the day before the procedure and sodium phosphate early in the morning of the day of the procedure, associated with one tablet of bisacodyl as compared to standard cleansing using two doses of sodium phosphate, with one tablet of bisacodyl. The paper is interesting and the design is adequate, as are the results and comments.

- copy preparation quality on detection of suspected colonic neoplasia. *Gastrointest Endosc* 2003; **58**: 76-79
- 2 **Rex DK**, Imperiale TF, Latinovich DR, Bratcher LL. Impact of bowel preparation on efficiency and cost of colonoscopy. *Am J Gastroenterol* 2002; **97**: 1696-1700
- 3 **Cohen SM**, Wexner SD, Binderow SR, Noguera JJ, Daniel N, Ehrenpreis ED, Jensen J, Bonner GF, Ruderman WB. Prospective, randomized, endoscopic-blinded trial comparing precolonoscopy bowel cleansing methods. *Dis Colon Rectum* 1994; **37**: 689-696
- 4 **Frommer D**. Cleansing ability and tolerance of three bowel preparations for colonoscopy. *Dis Colon Rectum* 1997; **40**: 100-104
- 5 **Hsu CW**, Imperiale TF. Meta-analysis and cost comparison of polyethylene glycol lavage versus sodium phosphate for colonoscopy preparation. *Gastrointest Endosc* 1998; **48**: 276-282
- 6 **Golub RW**, Kerner BA, Wise WE Jr, Meesig DM, Hartmann RF, Khanduja KS, Aguilar PS. Colonoscopic bowel preparations--which one? A blinded, prospective, randomized trial. *Dis Colon Rectum* 1995; **38**: 594-599
- 7 **Casais MN**, Rosa-Diez G, Pérez S, Mansilla EN, Bravo S, Bonfiglio FC. Hyperphosphatemia after sodium phosphate laxatives in low risk patients: prospective study. *World J Gastroenterol* 2009; **15**: 5960-5965
- 8 **Sica DA**, Carl D, Zfass AM. Acute phosphate nephropathy--an emerging issue. *Am J Gastroenterol* 2007; **102**: 1844-1847
- 9 **Desmeules S**, Bergeron MJ, Isenring P. Acute phosphate nephropathy and renal failure. *N Engl J Med* 2003; **349**: 1006-1007
- 10 **Markowitz GS**, Stokes MB, Radhakrishnan J, D'Agati VD. Acute phosphate nephropathy following oral sodium phosphate bowel purgative: an underrecognized cause of chronic renal failure. *J Am Soc Nephrol* 2005; **16**: 3389-3396
- 11 **Carl DE**, Sica DA. Acute phosphate nephropathy following colonoscopy preparation. *Am J Med Sci* 2007; **334**: 151-154
- 12 **Markowitz GS**, Radhakrishnan J, D'Agati VD. Towards the incidence of acute phosphate nephropathy. *J Am Soc Nephrol* 2007; **18**: 3020-3022
- 13 **Hookey LC**, Depew WT, Vanner S. The safety profile of oral sodium phosphate for colonic cleansing before colonoscopy in adults. *Gastrointest Endosc* 2002; **56**: 895-902
- 14 **Belsey J**, Epstein O, Heresbach D. Systematic review: oral bowel preparation for colonoscopy. *Aliment Pharmacol Ther* 2007; **25**: 373-384
- 15 **Tan HL**, Liew QY, Loo S, Hawkins R. Severe hyperphosphatemia and associated electrolyte and metabolic derangement following the administration of sodium phosphate for bowel preparation. *Anaesthesia* 2002; **57**: 478-483
- 16 **Martin RR**, Lisehora GR, Braxton M Jr, Barcia PJ. Fatal poisoning from sodium phosphate enema. Case report and experimental study. *JAMA* 1987; **257**: 2190-2192
- 17 **Ullah N**, Yeh R, Ehrinpreis M. Fatal hyperphosphatemia from a phosphosoda bowel preparation. *J Clin Gastroenterol* 2002; **34**: 457-458
- 18 **Azzam I**, Kovalev Y, Storch S, Elias N. Life threatening hyperphosphatemia after administration of sodium phosphate in preparation for colonoscopy. *Postgrad Med J* 2004; **80**: 487-488
- 19 **Farah R**. Fatal acute sodium phosphate enemas intoxication. *Acta Gastroenterol Belg* 2005; **68**: 392-393
- 20 **Aydogan T**, Kanbay M, Uz B, Kaya A, Isik A, Bozalan R, Erkman M, Akcay A. Fatal hyperphosphatemia secondary to a phosphosoda bowel preparation in a geriatric patient with normal renal function. *J Clin Gastroenterol* 2006; **40**: 177
- 21 **US Food and Drug Administration**. Science backgrounder: Safety of sodium phosphates oral solution. 2001. Available from: URL: <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm173897.htm>
- 22 **US Food and Drug Administration**. Science background

REFERENCES

- 1 **Harewood GC**, Sharma VK, de Garmo P. Impact of colonos-

- paper: Acute phosphate nephropathy and renal failure associated with the use of oral sodium phosphate bowel cleansing products. Available from: URL: <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm161581.htm>
- 23 **Rex DK.** Phosphate nephropathy. *Am J Gastroenterol* 2008; **103**: 807
- 24 **Fine A,** Patterson J. Severe hyperphosphatemia following phosphate administration for bowel preparation in patients with renal failure: two cases and a review of the literature. *Am J Kidney Dis* 1997; **29**: 103-105
- 25 **Rex DK.** Dosing considerations in the use of sodium phosphate bowel preparations for colonoscopy. *Ann Pharmacother* 2007; **41**: 1466-1475
- 26 **Sanaka MR,** Shah N, Mullen KD, Ferguson DR, Thomas C, McCullough AJ. Afternoon colonoscopies have higher failure rates than morning colonoscopies. *Am J Gastroenterol* 2006; **101**: 2726-2730
- 27 **Sharma VK,** Steinberg EN, Vasudeva R, Howden CW. Randomized, controlled study of pretreatment with magnesium citrate on the quality of colonoscopy preparation with polyethylene glycol electrolyte lavage solution. *Gastrointest Endosc* 1997; **46**: 541-543
- 28 **Sharma VK,** Chockalingham SK, Ugheoke EA, Kapur A, Ling PH, Vasudeva R, Howden CW. Prospective, randomized, controlled comparison of the use of polyethylene glycol electrolyte lavage solution in four-liter versus two-liter volumes and pretreatment with either magnesium citrate or bisacodyl for colonoscopy preparation. *Gastrointest Endosc* 1998; **47**: 167-71

S- Editor Sun H L- Editor O'Neill M E- Editor Zheng XM