

## CX<sub>3</sub>CR1 is critical for Salmonella-induced migration of dendritic cells into the intestinal lumen

Claudio Nicoletti,<sup>1\*</sup> Juan L. Arques<sup>1,2</sup> and Eugenio Bertelli<sup>3</sup>

<sup>1</sup>Laboratory of Mucosal Immunology; Programme of Integrated Biology of the GI Tract; Institute of Food Research; Norwich, UK;

<sup>2</sup>Department Tecnología de Alimentos; INIA; Madrid, Spain; <sup>3</sup>Department of Pharmacology "G. Segre"; University of Siena; Siena, Italy

We have demonstrated that direct antigen sampling of bacteria by intestinal dendritic cells (DCs) is accompanied by a rapid migration of CD11c<sup>+</sup>CX<sub>3</sub>CR1<sup>+</sup>MHCII<sup>+</sup>CD8 $\alpha$ <sup>-</sup>CD11b<sup>-</sup> DCs into the intestinal lumen upon exposure to non-invasive  $\Delta$ SPI1-Salmonella. Importantly, intraluminal DCs internalized Salmonella but were not able to cross the epithelium to return into tissue, thus showing that these DCs do not function as antigen-presenting cells and participate in the conventional regulation of immune responses to intestinal pathogens. Here we show that the presence of the chemokine receptor CX<sub>3</sub>CR1, that plays a vital role in DC-mediated antigen sampling and clearance in the gut, is also instrumental for the transepithelial migration of DCs. The latter observation, along with the notion that CX<sub>3</sub>CR1-deficient mice displayed higher susceptibility to Salmonella infection compared to wild-type mice raises the possibility that Salmonella-induced migration of "bacteria-capturing" DCs into the lumen may be an important mechanism of mucosal defence and clearance.

### Salmonella Induces Flagellin- and MyD88-Dependent Migration of Bacteria-Capturing Dendritic Cells into the Gut Lumen

Intestinal DCs play a critical role in the orchestration of mucosal immune responses,<sup>1</sup> however in the gut-DCs appeared to be also directly involved in sampling luminal antigens by extending

cellular processes between epithelial cells and shuttle them across the epithelial barrier.<sup>2-4</sup> This event is facilitated by the expression of tight junction protein that enables sampling without altering the integrity of the intestinal barrier. It has been hypothesized that direct DC sampling is followed by migration of bacteria-loaded DCs to the mesenteric lymph node (MLN) where they present antigens to T cells.<sup>5</sup> We have recently described that DC direct sampling is not the sole event taking place at mucosal interface in the gut during Salmonella infection.<sup>6</sup> Indeed, using both isolated intestinal loops and oral delivery of Salmonella we observed that a significant number of CD11c<sup>+</sup>CX<sub>3</sub>CR1<sup>+</sup>MHCII<sup>+</sup>CD8 $\alpha$ <sup>-</sup>CD11b<sup>-</sup> DCs traversed the epithelial barrier and moved into the lumen prior or following internalization of GFP-labelled Salmonella. Post Salmonella-challenge analysis of the epithelial barrier, that involved detailed study of tight junctions and intestinal permeability showed that DCs migration was not due to Salmonella-associated damages of the epithelium. The numbers of DCs within the gut lumen at 1.5 h after the introduction of Salmonella were 50-fold greater than after challenge with *E. coli* or PBS and after 3 h, the number of luminal DCs increased another ~3-fold in the lumen of Salmonella-treated mice.

Our work also determined that flagellin is a key signal, although not the only one, for DC migration. This was assessed by the use of Salmonella variant ( $\Delta$ *fljC*  $\Delta$ *fljB*) that lack flagellin and transgenic mice that lack the adaptor molecule MyD88, that is

**Key words:** dendritic cell, mucosal immunity, Salmonella, antigen sampling, cell migration, immune exclusion, mucosal clearance

Submitted: 11/16/09

Revised: 02/26/10

Accepted: 03/04/10


Previously published online:

[www.landesbioscience.com/journals/gut-microbes/article/11711](http://www.landesbioscience.com/journals/gut-microbes/article/11711)

\*Correspondence to:

Claudio Nicoletti; Email: [claudio.nicoletti@bbsrc.ac.uk](mailto:claudio.nicoletti@bbsrc.ac.uk)

Addendum to: Arques JL, Hautefort I, Ivory K, Bertelli E, Regoli M, Clare S, Hinton JCD, Nicoletti C. Salmonella induces flagellin- and MyD88-dependent migration of bacteria-capturing dendritic cells into the gut lumen. *Gastroenterology* 2009; 137:579-87.



**Figure 1.** Isolated loops were carried out in small intestine of wild-type C57BL/6 and transgenic MyD88 and CX<sub>3</sub>CR1-KO mice of same genetic background. The small intestine was isolated by ligatures at the level of terminal duodenum and terminal ileum. Saline suspensions containing bacteria ( $4 \times 10^8$ /ml) was injected in the loop and the intestine was returned to the abdominal cavity for 3 h. Luminal contents were then carefully recovered by gently flushing the intestine with PBS and the numbers of DCs assessed by flow cytometry. DC migration was completely abolished in CX<sub>3</sub>CR1-KO mice and was not different from levels observed in MyD88 where the numbers of DCs did not differ from PBS-treated mice.<sup>6</sup>

required for the activation signals following flagellin/TLR5 engagement.<sup>7</sup> The role of flagellin was also assessed by generating a double SPI1-SPI2 deficient Salmonella double mutant ( $\Delta$ SPI1  $\Delta$ ssrA) and a flagellated variant of *E. coli* K12. In either case the bacterial challenge did not induce DC migration suggesting an important role for the SPI2-mediated vesicular intracellular transport of flagellin,<sup>8</sup> and the requirement for additional pathogen-specific signal that were absent in the flagellated, but not pathogenic variant of *E. coli* used. Parallel experiments also showed that soluble flagellin did not induced DCs migration. It is also interesting to highlight that the lack of migration upon challenge with different variants of *E. coli* showed that LPS is not involved in this event. Also, by challenging different areas of the gut by using the isolated loops technique we determined that DC migration took place exclusively in the small intestine, including jejunum and ileum, but not in the colon. We also demonstrated that intraluminal DCs internalized Salmonellae prior or following migration into the gut lumen and that DC traffic was unidirectional; DCs were not able to cross the epithelial barrier and return into the intestinal tissue.

The latter observations ruled out the possibility that these DCs play a role in antigen-presentation. Taken together, these data demonstrated that the introduction of non-invasive Salmonella in the small intestine triggers at least two different events; DCs can either sample Salmonella and shuttle these bacteria back across the epithelial barrier or they can migrate into the intestinal lumen prior to or following internalization of Salmonella. These data also show that transepithelial DC protrusions do not always represent antigen-sampling devices; instead it appeared that the majority of these structures identify DCs in their migration into the intestinal lumen.

### Salmonella-Induced DC Migration, but Not Bacterial Translocation Is Impaired in CX<sub>3</sub>CR1-Deficient Mice

Conflicting results have been reported on the role of the chemokine receptor CX<sub>3</sub>CR1 on the ability of DCs to form transepithelial dendrites and sample directly luminal bacteria. CX<sub>3</sub>CR1<sup>+</sup> DCs are the target for the CX<sub>3</sub>CL1/fractalkine, a transmembrane chemokine expressed at the surface and basolateral compartment

of intestinal epithelial cells (IEC) predominantly, but not exclusively in the terminal ileum.<sup>3,4,9</sup> One study reporting that CX<sub>3</sub>CR1-deficient mice were unable to sample luminal Salmonella,<sup>4</sup> was followed by another one suggesting that CX<sub>3</sub>CR1 was not directly involved in the formation of DC protrusions across the intestinal epithelium.<sup>3</sup> Important, both studies were done in mice of identical genetic make up (C57BL/6), thus showing that this was not the origin of this discrepancy. Most important, in addressing the role of CX<sub>3</sub>CR1<sup>+</sup> DC it was reported that CX<sub>3</sub>CR1-deficient mice showed increased susceptibility to Salmonella infection, thus suggesting that these antigen-sampling DCs are a major component of defence against pathogenic microorganisms.<sup>4</sup> We then extended our investigation of Salmonella-induced DC migration in CX<sub>3</sub>CR1 knock-out (KO) mice and we observed that following short-term (3 h) challenge, carried out using isolated loops DC migration is drastically reduced (Fig. 1). Numbers of DCs into the lumen were significantly reduced compared to wild-type mice and comparable to levels observed in MyD88 mice. In addition, early translocation (5 h) of non-invasive Salmonella in the jejunum and ileum was not significantly affected in CX<sub>3</sub>CR1-KO mice (Table 1) confirming the finding that the absence of transepithelial dendrites in the terminal ileum of CX<sub>3</sub>CR1-KO did not impair bacteria entry to the lamina propria and suggesting the existence of multiple entry pathways for pathogens and an overall minor role for DC-mediated sampling.<sup>10</sup>

### Protection against Bacteria Infection in the Gut: DC-Sampling vs DC-Migration?

Is it possible that both DC-sampling and migration are required for protection against bacterial infections? Also, do these events complement each other by playing a role at different stages of the immune response to pathogens? In regard to DC sampling, its biological relevance and role in the generation of immune responses remains to be determined. To this end it is interesting to notice that following oral challenge a very small number of *E. coli* was translocated by the lamina propria

(LP)-DCs while that vast majority (<100-fold difference) crossed the epithelial barrier via M cells of Peyer's patch.<sup>4</sup> Interestingly, this result was obtained by comparing levels of CFU within the MLNs and PPs as parameters to evaluate DC and M cell transport, respectively. However, this way of assessing DC-mediated sampling has to be interpreted cautiously. Indeed, determining levels of CFU within the MLN does not take into consideration both the contribution of villous-M cells,<sup>11,12</sup> and the fact that bacteria transported by PPs also end up in MLN.<sup>13,14</sup> In regard to the latter observation, the finding that no bacteria were recovered in the MLN of CX<sub>3</sub>CR1-KO mice despite an intact M cell-mediated transport is particularly intriguing.<sup>4</sup> This would mean that pathogens that invade the host via PP-associated M cells would not reach the MLN and in so doing they would escape, critical immunological check points associated to the intestinal immune system. However, this is not the case. Indeed, it has been demonstrated that also pathogens invading the host exclusively via the PPs are subsequently transported to the MLN;<sup>15-17</sup> thus showing that both PP and LP drain to the MLN. This notion is also supported by a detailed anatomical investigation of the mouse intestine;<sup>18</sup> this study confirmed that the entire small intestine, including PPs drained into the MLN and did not detect any alternative pathway for PP-derived lymph and its cellular contents.

Furthermore, another study reported that orally delivered fluorescence-labelled *Enterobacter cloacae* were recovered only within the PPs but not within the LP suggesting that, at least in this case DCs did not sample luminal bacteria.<sup>19</sup> These results taken together showed that PP-M cells are the major site of antigen up-take, a critical step for the induction of mucosal and systemic immune responses and that DC-sampling, as a whole represents a small scale event, thus making it difficult to envisage its biological relevance. In addition, other relevant features of DC sampling are the subject of debate. For example, in one case high numbers of DC extensions were detected in the jejunum and a small number in the terminal ileum, where their numbers increased following

**Table 1.** Transepithelial transport of non-invasive salmonella after oral challenge CFU/gr. of tissue

	Jejunum	Ileum	Colon
Wild-type	680 ± 230	895 ± 380	257 ± 98
CX <sub>3</sub> CR1-KO	590 ± 332	1012 ± 426	390 ± 57

CX<sub>3</sub>CR1-KO (C57BL/6 background) and wild-type mice and (n = 6/group) were orally administrated with 5 × 10<sup>8</sup>-10<sup>9</sup> non-invasive ΔSPI1-Salmonella. Tissues were removed after 5 h, washed in antibiotics and subsequently homogenized in HBSS buffer. Serial dilutions of the homogenates were plated on LB agar plates overnight at 37°C and then colonies were counted.

challenge with non-invasive Salmonella;<sup>3</sup> in contrast others have reported that transepithelial dendrites were observed only in the terminal ileum and not in the remaining areas of the small intestine.<sup>4</sup> Also the underlying molecular mechanism appeared to vary according to the different areas of the gut with DC-sampling being TLR-dependent in the small intestine,<sup>3</sup> but TLR-independent in the colon.<sup>20</sup> Interestingly, the formation of DC extensions is also strain-specific.<sup>10</sup> These were observed in C57BL/6 but were absent in BALB/c mice and this led to suggest that DC-mediated antigen sampling is not a universal phenomenon.<sup>13</sup> To this end it would be interesting to dissect in detail immune responses and determine the rate of mortality in these two mouse strains after challenge with invasive Salmonella.

At present much less is known on the potential role of DCs that migrate rapidly into the intestinal lumen. Indeed, this event poses several key questions. Firstly, do sampling and migration involve different DC subpopulations? The possibility that different subpopulations of DCs may actively sample luminal antigens is strongly suggested by the observation that DC protrusions in CX<sub>3</sub>CR1-GFP and MHC II-GFP mice only partially overlapped.<sup>21</sup> Also, it would be interesting to determine the expression of CD103 on DCs involved in both antigen-sampling and migration into the intestinal lumen.<sup>22,23</sup> Secondly, and most compelling, what is the function, if any, of these cells into the gut lumen? The observation that CX<sub>3</sub>CR1-KO mice are not able to send DCs into the intestinal lumen upon Salmonella infection may enable us to formulate some hypothesis on their role. Indeed, as discussed above, it was reported that CX<sub>3</sub>CR1 deficient mice showed higher rate of mortality following Salmonella infection and albeit direct

evidence was lacking this was attributed to their inability to form intraepithelial DC protrusions and sample luminal antigens.<sup>4</sup> Although a quantitative analysis of Salmonella-induced recruitment of DCs in the LP of CX<sub>3</sub>CR1-KO mice is currently lacking, in light of our previous paper,<sup>6</sup> and the present work on the role of CX<sub>3</sub>CR1 receptor in DC migration and bacterial transport to the LP we would like to propose that the increased susceptibility of these CX<sub>3</sub>CR1-deficient mice to Salmonella infection could also reflect the absence of "Salmonella-capturing" DCs into the lumen at the early stage of infection. Indeed, the lack of DC migration, along with other defects in monocyte (GR1<sup>+</sup>) recruitment may contribute to an overall diminished capability to cope with mucosal infection.<sup>24</sup> As we have previously discussed,<sup>6</sup> at present we can not rule out the possibility that DC migration is simply a physical consequence of the mechanical pressure of cells that migrate within the LP towards the intestinal lumen to perform antigen-sampling following certain bacterial stimuli. However, we feel it is unlikely that DC migration into the lumen would represent an accidental event as it would inevitably lead to the loss of important regulatory cells. Alternatively, these cells may be part of a defence mechanism of cell (phagocyte)-mediated immune exclusion that limits the number of pathogens crossing the epithelial barrier that would complement and potentiate the mucous and IgA antibody secretion-mediated system. A similar hypothesis was first proposed by Bellamy and Nielsen.<sup>25</sup> The authors observed that a large number of phagocytes moved into the lumen following exposure to enteric antigen and suggested that antigen "escaping" the immune exclusion barrier provided by both specific IgA and mucous and

penetrating the epithelial barrier triggered a rapid migration of cells with the task of removing and eliminating the offending pathogen. Furthermore, the observation that DC migration is restricted to the small bowel also would lend support to this hypothesis; indeed, in the past it was observed that small intestine juices upregulated phagocytosis;<sup>26</sup> this feature along with the known opsonising properties of IgA would make the small intestine the ideal environment for the implementation of such a defence strategy.

### Concluding Remarks

This work demonstrates that CX<sub>3</sub>CR1 plays a critical role in Salmonella-induced migration into the intestinal lumen but it does not significantly affect early transport of bacteria to the LP. This would suggest that intraluminal DC have an important role in mucosal clearance during the early phase of bacterial infection in the gut; the lack of this defence mechanism, as seen in CX<sub>3</sub>CR1-KO mice may determine higher susceptibility to Salmonella infection. Finally, it is worth to highlight that the co-existence of DC-sampling and migration in response to the same antigenic stimulus, in this case non-invasive Salmonella, illustrates the highly dynamic nature of both gut epithelium and immune system in response to bacterial infections and the complexity of the signalling network that governs host-pathogen interaction at mucosal interface. The challenge now is to identify the host molecular mechanisms and the bacterial signals triggering these two events and to determine their biological relevance in regard to immunity and mucosal clearance of intestinal pathogens.

### Acknowledgements

The authors wish to thank L. Zuidmeer for helpful discussions and suggestions.

This work was funded by the Biotechnology and Biological Sciences Research Council, UK (to Claudio Nicoletti) and University of Siena, Italy (to Eugenio Bertelli).

### References

1. Coombes JL, Powrie F. Dendritic cells in intestinal immune regulation. *Nature Rev Immunol* 2008; 8:435-46.
2. Rescigno M, Urbano M, Valzasina B, Francolini M, Rotta G, Bonasio R, et al. Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to sample bacteria. *Nat Immunol* 2001; 2:361-7.
3. Chieppa M, Rescigno M, Huang AY, Germain RN. Dynamic imaging of dendritic cell extension into the small bowel lumen in response to epithelial cell TLR engagement. *J Exp Med* 2006; 203:2841-52.
4. Niess JH, Brand S, Gu X, Landsman L, Jung S, McCormick BA, et al. CX<sub>3</sub>CR1-mediated dendritic cell access to the intestinal lumen and bacterial clearance. *Science* 2005; 307:248-54.
5. Niedergang F, Didierlaurent A, Kraehenbuhl JP, Sirard JC. Dendritic cells: the host Achilles's heel for mucosal pathogens? *Trends Microbiol* 2004; 12:79-88.
6. Arques JL, Hautefort I, Ivory K, Bertelli E, Regoli M, Claire S, et al. Salmonella induces flagellin- and MyD88-dependent migration of bacteria-capturing dendritic cells into the gut lumen. *Gastroenterology* 2009; 137:579-87.
7. Hayashi F, Smith KD, Ozinsky A, Hawn TR, Yi EC, Goodlett DR, et al. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature* 2001; 410:1099-103.
8. Lyons S, Wang L, Casanova JE, Sitaraman SV, Merlin D, Gewirtz AT. *Salmonella typhimurium* transcytoses flagellin via an SPI2-mediated vesicular transport pathway. *J Cell Sci* 2004; 117:5771-80.
9. Muehlhoefer A, Saubermann LJ, Gu X, Heckenkamp KL, Xavier R, Blumberg RS, et al. Fraktalkine is an epithelial and endothelial cell-derived chemoattractant for intraepithelial lymphocytes in the small intestinal mucosa. *J Immunol* 2000; 164:3368-76.
10. Vallon-Eberhard A, Landsman L, Yogev N, Verrier B, Jung S. Trans epithelial pathogen uptake into the small intestinal lamina propria. *J Immunol* 2006; 176:2465-9.

11. Yamamoto M, Terahara K, Sasakawa C, Suzuki T, Nochi T, Yokota Y, et al. Intestinal villous M cells: an antigen entry site in the mucosal epithelium. *Proc Natl Acad Sci USA* 2004; 101:6110-5.
12. Knoop KA, Kumar N, Butler BR, Sakthivel SK, Taylor RT, Nochi T, et al. RANKL is necessary and sufficient to initiate development of antigen-sampling M cells in the intestinal epithelium. *J Immunol* 2009; 183:5738-47.
13. Iwasaki A. Mucosal dendritic cells. *Annu Rev Immunol* 2007; 25:381-418.
14. Johansson-Lindbom B, Agace WW. Generation of gut-homing T cells and their localization to the small intestinal mucosa. *Immunol Rev* 2007; 215:226-42.
15. McDonald TT, Carter PB. Cell-mediated immunity to intestinal infection. *Infect Immun* 1980; 28:516-32.
16. Marco AJ, Prats N, Ramos JA, Briones V, Blanco M, Dominguez L, et al. A microbiological, histopathological and immunohistological study of the intragastric inoculation of *Listeria monocytogenes* in mice. *J Comp Pathol* 1992; 107:1-9.
17. Marco AJ, Altamira J, Prats N, López S, Dominguez L, Domingo M, et al. Penetration of *Listeria monocytogenes* in mice infected by the oral route. *Microb Pathog* 1997; 23:255-63.
18. Carter PB, Collins FM. The route of enteric infection in normal mice. *J Exp Med* 1974; 134:1189-203.
19. Macpherson AJ, Uhr T. Induction of protective IgA by intestinal dendritic cells carrying commensal bacteria. *Science* 2004; 303:1662-5.
20. Hapfelmeier S, Müller AJ, Stecher B, Kaiser P, Barthel M, Endt K, et al. Microbe sampling by mucosal dendritic cells is a discrete, MyD88-independent step in *DeltainvG S. typhimurium* colitis. *J Exp Med* 2008; 205:437-50.
21. Rescigno M. Before they were gut dendritic cells. *Immunity* 2009; 31:454-6.
22. Bogunovic M, Ginhoux F, Helft J, Shang L, Hashimoto D, Greter M, et al. Origin of the lamina propria dendritic cell network. *Immunity* 2009; 31:513-25.
23. Jung S, Varol C, Vallon-Eberhard A, Elinav E, Aychek T, Shapira Y, et al. Intestinal lamina propria dendritic cell subsets have different origin and functions. *Immunity* 2009; 31:502-12.
24. Auffray C, Fogg DK, Narni-Mancinelli E, Senechal B, Trouillet C, Saederup N, et al. CX<sub>3</sub>CR1<sup>+</sup> CD115<sup>+</sup> CD135<sup>+</sup> common macrophage/DC precursors and the role of CX<sub>3</sub>CR1 in their response to inflammation. *J Exp Med* 2009; 206:595-606.
25. Bellamy JEC, Nielsen NO. Immune-mediated emigration of neutrophils into the lumen of the small intestine. *Infect Immun* 1974; 9:615-9.
26. Girard JP, de Kalbermatten A. Antibody activity in human duodenal fluid. *Eur J Clin Invest* 1970; 1:188-95.