



Published in final edited form as:

*Biochemistry*. 2010 September 21; 49(37): 7969–7971. doi:10.1021/bi101098f.

## Direct Evidence for Conformational Heterogeneity in Human Pancreatic Glucokinase from High-Resolution NMR

Mioara Larion<sup>§,\*</sup>, Roberto K. Salinas<sup>§,‡,\*</sup>, Lei Bruschiweiler-Li<sup>§,‡</sup>, Rafael Brüschiweiler<sup>§,‡</sup>, and Brian G. Miller<sup>§</sup>

<sup>§</sup> Department of Chemistry and Biochemistry, The Florida State University, Tallahassee, Florida 32306-4390

<sup>‡</sup> National High Magnetic Field Laboratory, The Florida State University, Tallahassee, Florida, 32310

### Abstract

High-resolution NMR is used to investigate the conformational dynamics of human glucokinase, a 52 kDa monomeric enzyme that displays kinetic cooperativity. <sup>1</sup>H-<sup>15</sup>N TROSY spectra of uniformly labeled glucokinase, collected in the absence and presence of glucose, reveal significant cross-peak overlap and heterogeneous peak intensities that persist over a range of temperatures. <sup>15</sup>N-specific labeling of isoleucines and tryptophans, reporting on backbone and side-chain dynamics respectively, demonstrate that both unliganded and glucose-bound enzymes sample multiple conformations, although glucose stabilizes certain conformations. These results provide the first direct evidence of glucokinase conformational heterogeneity and hence shed light on the molecular basis of cooperativity.

Glucokinase is a 52 kDa enzyme that catalyzes the rate-limiting step of glucose metabolism in pancreatic  $\beta$ -cells (1,2). Glucokinase activity is allosterically regulated by its sugar substrate, displaying a sigmoidal response to glucose concentrations spanning the physiological range. The kinetic cooperativity of glucokinase is mechanistically interesting because the enzyme functions exclusively as a monomer (3). Current models suggest that cooperativity relies upon a slow, glucose-dependent conformational change that occurs with a rate constant slower than  $k_{cat}$  (4). Consistent with that postulate, crystal structures of glucokinase confirm that large-scale structural alterations accompany substrate binding (5). To date, essentially no experimental work has been directed toward understanding the functional dynamics of the enzyme. For this reason, we initiated an investigation of the solution state dynamics of glucokinase using high-resolution NMR.

Initially, we collected the <sup>1</sup>H-<sup>15</sup>N TROSY spectrum of uniformly labeled glucokinase (120  $\mu$ M) at 25°C using an 800 MHz NMR spectrometer equipped with a cryogenic probe. This spectrum displayed very weak peak intensities and considerable cross-peak overlap. Increasing protein concentrations up to 520  $\mu$ M did not remove heterogeneous peak intensities or alleviate cross-peak overlap. Analytical ultracentrifugation experiments at 40  $\mu$ M indicated the presence of a monomeric species. 1D <sup>1</sup>H and 2D <sup>1</sup>H-<sup>15</sup>N HSQC NMR spectra obtained over a range from 50  $\mu$ M to 500  $\mu$ M showed no change in the spectral

<sup>§</sup>To whom correspondence should be addressed. Phone: (850)645-6570 Fax (850) 644-8281 miller@chem.fsu.edu, Phone: (850) 644.1768 Fax (850) 644-8281 rbruschw@magnet.fsu.edu.

\*Both authors contributed equally to this work.

#### SUPPORTING INFORMATION AVAILABLE

Detailed labeling approaches and experimental procedures and Figures S1–S6. This material is available free of charge via the Internet at <http://pubs.acs.org>.

characteristics with the increase in protein concentration (Figures S1–S2). Together, these data support retention of a monomeric state and preclude aggregation as the cause for the poor spectral resolution. Next, we collected  $^1\text{H}$ - $^{15}\text{N}$  TROSY spectra over a range of temperatures from 5–45°C. At temperatures above 40°C the protein rapidly precipitated. Below this temperature, we observed a consistent increase in spectral resolution and peak intensities with increasing temperatures (Figure S3). Despite this fact, the 40°C  $^1\text{H}$ - $^{15}\text{N}$  TROSY spectrum still displayed substantial cross-peak overlap and a heterogeneous distribution of peak intensities (Figure 1A). The size of glucokinase (52 kDa) does not fully explain its unusual NMR behavior, as indicated by comparison with the  $^1\text{H}$ - $^{15}\text{N}$  TROSY spectrum of monomeric arginine kinase (42 kDa), which displays a much weaker temperature dependence than glucokinase and a consistently higher spectral resolution (Figure S3) (6). Interestingly, the addition of glucose to glucokinase did not alleviate cross-peak overlap, especially in the  $^1\text{H}$  region near 8 ppm, although several new cross-peaks appeared (Figure 1B). Together these observations suggest that glucokinase is an intrinsically mobile enzyme whose structure and dynamics are modulated by temperature and substrate binding.

To probe glucokinase dynamics via a different approach, we used NMR to detect the exchange of amide protons with  $\text{D}_2\text{O}$ . H/D exchange was initiated by dissolving a lyophilized sample of uniformly  $^{15}\text{N}$  labeled enzyme in 100%  $\text{D}_2\text{O}$ . Successive 2D  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectra were acquired over a period of 20 hours at pH 7.0 and 25°C (Figure S4). The  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectra show a substantial loss of signal within the first two hours of exchange. During this period, approximately 75% of the total signal was lost when compared to the 1D spectrum in 100%  $\text{H}_2\text{O}$ . The rapidity of this loss prevented us from determining the effect of glucose addition upon the exchange rate of these protons. The remaining signal (~25%) underwent slower exchange, the rate of which was unaffected by the presence of glucose. The observation that a significant fraction of the amide protons undergoes relatively rapid exchange suggests that the backbone of one or more glucokinase conformational states is highly solvent exposed, both in the absence and presence of glucose.

To simplify the  $^1\text{H}$ - $^{15}\text{N}$  TROSY spectrum, and to detect discrete conformational states, we incorporated site-specific  $^{15}\text{N}$  labels in the backbone atoms of isoleucine residues and in the side chain atoms of tryptophan. Glucokinase contains 17 isoleucine residues that are intrinsically dispersed throughout the structure and have the potential to globally probe functional dynamics (Figure 2A). Glucokinase contains three tryptophan residues, two of which appear to experience significant environmental changes upon glucose association (5). Isoleucine and tryptophan labeling was achieved as previously described (7, 8) and site-specific  $^{15}\text{N}$  incorporation was confirmed via MALDI-TOF mass spectrometry (Figure S5). The  $^1\text{H}$ - $^{15}\text{N}$  TROSY spectrum of unliganded, isoleucine-labeled glucokinase shows two strong cross-peaks and 8 additional peaks that slightly exceed the noise level (Figure 2B). The addition of glucose leads to the emergence of extra cross-peaks and produces more homogenous peak intensities (Figures 2C and S6). Similarly, the  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectrum of unliganded, tryptophan-labeled glucokinase displays one strong cross-peak with a  $^1\text{H}$  chemical shift at 10 ppm and a second, very weak cross-peak near 9.5 ppm (Figure 3A). In the presence of saturating glucose, the  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectrum displays three prominent tryptophan cross peaks, two of which possess weak extra peaks in close vicinity (Figure 3B). The differential number of cross-peaks observed in the absence and presence of glucose in both the isoleucine and tryptophan  $^{15}\text{N}$ -labeled spectra demonstrates that glucose binding stabilizes distinct enzyme conformations. Moreover, the appearance of extra tryptophan side chain signals in the  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectrum suggests that glucose-bound glucokinase samples multiple conformational states on a relatively slow time scale.

Several possible explanations exist for the unique spectral characteristics of human glucokinase. The differential number of cross-peaks observed in the absence and presence of glucose could be interpreted as unliganded glucokinase populating a large number of states. The resonances of these states would have to display a distribution of chemical shifts with peak intensities that are below the detection limit. From the signal-to-noise ratio of the tryptophan-labeled  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectrum, we estimate an upper population limit for each of these states around 2%. A second possibility is that unliganded glucokinase undergoes conformational exchange between a smaller number of discrete states on a time scale that approaches the inverse of the chemical shift difference. The strong temperature dependence of the  $^1\text{H}$ - $^{15}\text{N}$  TROSY spectrum favors such a coalescence phenomenon (Figure S3). A third possibility, very rapid amide proton exchange with the solvent, is unlikely to be responsible for cross-peak disappearance; even in the  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectra of small highly flexible peptides, cross-peaks can be observed (9).

In summary, the results of our preliminary NMR investigation suggest that glucokinase is capable of sampling multiple conformational states, both in the absence and presence of glucose. The detection of additional cross-peaks in the glucose-bound spectrum suggests that glucose binding alters the dynamics of glucokinase conformational exchange. Indeed, the ability of glucose to modulate enzyme dynamics may provide a general mechanism by which this substrate can confer allosteric regulation. Using the sample conditions established here, quantitative NMR investigations seem feasible to establish the rate of glucokinase conformational exchange in the presence of various cooperative and non-cooperative ligands. Such information is critical to our understanding of the link between enzyme functional dynamics and kinetic cooperativity.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

This work was supported, in part, by grants from the Florida Biomedical Research Program (09KN08), the NIH (1R01DK081358) and the NSF (MCB-0918362). R.K.S. is recipient of a postdoctoral fellowship from the American Heart Association. We thank Dr. Xiaogang Niu for providing an NMR sample of arginine kinase.


## Abbreviations

<b>NMR</b>	nuclear magnetic resonance
<b>TROSY</b>	transverse relaxation optimized spectroscopy
<b>HSQC</b>	heteronuclear single quantum coherence spectroscopy
<b>HEPES</b>	4-(2-hydroxyethyl)-1-piperazine ethanesulfonic acid
<b>TCEP</b>	tris(2-carboxyethyl)phosphine


## References

1. Wilson JE. Hexokinases. *Rev Biochem, Pharmacol.* 1995; 126:65–198.
2. Ferre T, Riu E, Bosch F, Valera A. Evidence from transgenic mice that glucokinase is rate limiting for glucose utilization in the liver. *FASEB J.* 1996; 10:1213–1218. [PubMed: 8751724]
3. Cornish-Bowden, A.; Cardenas, M. Glucokinase: A monomeric enzyme with positive cooperativity. *Glucokinase and glycemic disease: From basics to novel therapeutics.* In: Matschinsky, FM.; Magnuson, MA., editors. *Front Diabetes.* Vol. 16. Karger, Basel: 2004. p. 125-134.
4. Neet KE, Ainslie GR Jr. Hysteretic enzymes. *Methods Enzym.* 1980; 64:192–226.

5. Kamata K, Mitsuya M, Nishimura T, Eiki J, Nagata Y. Structural basis for allosteric regulation of the monomeric allosteric enzyme human glucokinase. *Structure*. 2004; 12:429–438. [PubMed: 15016359]
6. Davulcu O, Flynn PF, Chapman MS, Skalicky JJ. Intrinsic domain and loop dynamics commensurate with catalytic turnover in an induced-fit enzyme. *Structure*. 2009; 17:1356–1367. [PubMed: 19836335]
7. Muchmore DC, McIntosh LP, Russell CB, Anderson DE, Dahlquist FW. Expression and nitrogen-15 labeling of proteins for nitrogen-15 Nuclear Magnetic Resonance. *Methods in Enzymology*. 1989; 177:44–73. [PubMed: 2691846]
8. Rodriguez-Mias RA, Pellicchia M. Use of selective Trp side chain labeling to characterize protein-protein and protein-ligand interactions by NMR spectroscopy. *J Am Chem Soc*. 2003; 125:2892–2893. [PubMed: 12617653]
9. Corrêa R, Farah CS, Salinas RK. Mg<sup>2+</sup> ions bind at the C-terminal region of skeletal muscle  $\alpha$ -tropomyosin. *Biopolymers*. 2009; 91:583–590. [PubMed: 19280641]
10. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. UCSF Chimera—a visualization system for exploratory research and analysis. *J Comput Chem*. 2004; 13:1605–1612. [PubMed: 15264254]




**FIGURE 1.**  $^1\text{H}$ - $^{15}\text{N}$  TROSY NMR spectrum at 800 MHz of uniformly  $^{15}\text{N}$ -labeled glucokinase ( $520\ \mu\text{M}$  in 50 mM HEPES pH 7.0, 50 mM KCl, and 4 mM TCEP) in the absence (A) and presence (B) of 50 mM glucose. The spectra were recorded at 40 °C, as matrices of  $2048(N_2) \times 290(N_1)$  real points and accumulation of 512 scans (in the absence of glucose) or 128 scans (in the presence of glucose), multiplied by a cosine apodization function and zero-filled up to  $2048(N_2) \times 512(N_1)$  data points. Negative peaks, colored in green, are due to the leakage of anti-TROSY contributions.



**FIGURE 2.**

(A) Crystal structure of human glucokinase depicting the location of isoleucine (green spheres) and tryptophan (blue spheres) residues. Glucose is colored red and white. Mesh plots of  $^1\text{H}$ - $^{15}\text{N}$  TROSY spectra of  $^{15}\text{N}$  isoleucine labeled glucokinase in the absence (B) and presence (C) of glucose, collected at 40 °C. Image was created with Chimera (10) and PDB entry 1V4S (5). For clarity, the allosteric activator is not depicted in the glucokinase structure.

**FIGURE 3.**

$^1\text{H}$ - $^{15}\text{N}$  HSQC NMR spectra of glucokinase with  $^{15}\text{N}$  tryptophan side chain labels in the absence (A) and presence (B) of glucose. The spectra were acquired as matrices of  $2048(N_2) \times 128(N_1)$  or  $2048(N_2) \times 64(N_1)$  real points and accumulation of 128 or 64 scans in the absence or presence of glucose, respectively. The indirect dimension of the HSQC spectrum of the glucose-bound sample was extended up to 128 points by linear prediction. All dimensions were multiplied by a cosine square apodization function and zero filled up to  $2048(N_2) \times 256(N_1)$  data points. The spectrum in the absence of glucose was acquired using a smaller spectral window in the  $^{15}\text{N}$  dimension (813 Hz compared to 2433 Hz in the presence of glucose).