

REVIEW

Open Access

Potassium channels as a potential therapeutic target for trigeminal neuropathic and inflammatory pain

Mamoru Takeda^{1*}, Yoshiyuki Tsuboi², Junichi Kitagawa³, Kazuharu Nakagawa⁴, Koichi Iwata², Shigeji Matsumoto¹

Abstract

Previous studies in several different trigeminal nerve injury/inflammation models indicated that the hyperexcitability of primary afferent neurons contributes to the pain pathway underlying mechanical allodynia. Although multiple types of voltage-gated ion channels are associated with neuronal hyperexcitability, voltage-gated K⁺ channels (Kv) are one of the important physiological regulators of membrane potentials in excitable tissues, including nociceptive sensory neurons. Since the opening of K⁺ channels leads to hyperpolarization of cell membrane and a consequent decrease in cell excitability, several Kv channels have been proposed as potential target candidates for pain therapy. In this review, we focus on common changes measured in the Kv channels of several different trigeminal neuropathic/inflammatory pain animal models, particularly the relationship between changes in Kv channels and the excitability of trigeminal ganglion (TRG) neurons. We also discuss the potential of Kv channel openers as therapeutic agents for trigeminal neuropathic/inflammatory pain, such as mechanical allodynia.

Introduction

Pain caused by a lesion of the peripheral or central nervous system is commonly termed neuropathic pain, and this type of pain frequently persists, even following normal repair of the injured tissue [1-3]. In a clinically significant proportion of cases, the neuropathic pain becomes chronic, severely debilitating, and extremely difficult to treat. Although several different types of neuropathic pain animal models have been developed and extensively studied [4,5], no common therapeutic molecular target has been identified for neurons located in the nociceptive pathway.

Multiple types of voltage-gated ion channels are associated with neuronal excitability. Among these, voltage-gated K⁺ (Kv) channels are important physiological regulators of membrane potentials, action potential shape, and firing adaptation in excitable tissues including nociceptive sensory neurons [6-8]. Dorsal root ganglion (DRG) and trigeminal ganglion (TRG) neurons expressed three distinct classes of K⁺ currents in varying quantities: slow-inactivating sustained (K-current; I_K),

fast-inactivating transient (A-current; I_A), and slow-inactivating transient (D-current; I_D) currents based on their inactivation of kinetics and sensitivities to tetraethylammonium (TEA), 4-aminopyridine (4-AP) and α -dendrotoxin (α -DTX), respectively [9-13]. Peripheral nerve injury and inflammation markedly reduces the densities of Kv channels, implicating them in the development of neuropathic/inflammatory pain [14-17]. Since the opening of K⁺ channels leads to hyperpolarization of cell membrane and a resultant decrease in cell excitability, several types of Kv channels have been proposed as potential target candidates for such pain pathways [18]. Recent studies in various pain models identified the voltage-gated KCNQ/Kv7 channel (M-current) opener [19-21] and alterations in either calcium-activated (K_{Ca}) or ATP-sensitive potassium (K_{ATP}) channels as potential therapeutic targets for neuropathic/inflammatory pain [22,23].

We recently reported on the hyperexcitability of primary afferent neurons in several different trigeminal nerve injury and inflammation animal models, including chronic constriction nerve injury (CCI), axotomy, and inflammatory models [17,24-26]. In these models, we systemically investigated the mechanism of mechanical allodynia underlying changes in the TRG neuronal

* Correspondence: m-takeda@tokyo.ndu.ac.jp

¹Department of Physiology, School of Life Dentistry at Tokyo, Nippon Dental University, 1-9-20, Fujimi-cho, Chiyoda-ku, Tokyo, 102-8159, Japan
Full list of author information is available at the end of the article

excitability due to several types of neuropathic/inflammatory conditions in rats, using behavioral analysis as well as extracellular single-unit and whole-cell patch-clamp recordings.

This review therefore focuses on changes in the Kv channels studied in different trigeminal neuropathic/inflammatory pain animal models, and particularly the relationships between Kv channels and TRG neuron excitability. We also discuss potential therapeutic targets identified thus far for the prevention of pathological pain, such as those targeting mechanical allodynia.

Pathological pain models

Peripheral nerve injury/inflammation produces sensory abnormalities associated with chronic pain. Peripheral tissue inflammation can alter the property of somatic sensory pathways, resulting in behavioral hypersensitivity due to the increased responses to pain caused by noxious stimulation (hyperalgesia) as well as the normally non-noxious stimulation (allodynia). Several different types of animal nerve injury models have been developed to elucidate the neuronal mechanisms underlying these abnormal pain sensations (allodynia and hyperalgesia) [27-29]. Among these models, the CCI model shows inflammation and nerve injury, and can mimic neuropathic pain in human [27,30]. CCI of the sciatic nerve is indeed one of the most reliable models of neuropathic pain [31]. Following peripheral nerve injury, primary afferent neurons exhibit high frequency background activity or irregular burst discharges [27]. On the other hand, in the axotomy of primary afferent neuron model (*axotomy model*), the hyperexcitability of primary afferent neurons induced by peripheral nerve injury is thought to result from extensive changes in the ionic currents of DRG neurons [32,33]. Moreover, changes in the excitability of primary afferent neurons were also observed at the adjacent intact primary afferent neurons after nerve injury (*axotomy spared neuron models*) [34,35].

Any discussion of trigeminal neuropathic pain should consider the following points. First, the trigeminal nerve has some unique characteristics that may influence its response to injury, such as its embryological origin, and the proportion of myelinated and unmyelinated nerves, and sympathetic fibers [36-38]. In addition, the location of some branches of the nerve within bony canals protects them from exogenous stimuli, but makes them vulnerable to pressure (e.g., edema or displacement of fractured bone fragments) [36]. To elucidate the mechanism underlying trigeminal mechanical allodynia in different neuropathic pain/inflammatory models, we recently investigated four such animal models, as described in the following sections.

Changes in potassium currents and trigeminal neuropathic and inflammatory pain

Figure 1 summarizes the changes in TRG neuronal excitability associated with the changes in Kv channels using whole-cell patch-clamp recording in our trigeminal neuropathic/inflammatory pain animal models [17,24-26]. In each animal model, mechanical allodynia was determined by withdrawal threshold for the mechanical stimulation of the whisker pad area.

CCI-model

Trigeminal nerve has a predominant sensory function compared with that of other peripheral nerves such as sciatic nerve because it is predominantly composed of sensory afferents, and only contains a small number of efferent fibers. It can be assumed that the infraorbital nerve (ION) does not contain any motor fibers, thus the trigeminal CCI model provides evidence to demonstrate the neuropathic pain mechanism following a purely sensory nerve injury [25]. In our study, mechanical allodynia appeared at three days after ION-CCI and lasted for 14 days. Whole-cell patch-clamp recordings from medium-/large-diameter CCI-TRG neurons revealed that both I_K and I_A in rats with ION-CCI were significantly smaller than in naïve rats (Figure 1A). Changes in the duration of the depolarization phase (DDP) and repolarization phase (DRP) contribute to the I_A and I_K , respectively [12], thus suppression of these current in TRG neurons following CCI correlated with a decrease in DDP and an increase in DRP, respectively. Also, the TRG neuronal action potentials (and consequently, the total action potential duration) in rats with ION-CCI were significantly prolonged than those in naïve rats. Under current-clamp experiments, TRG neurons following ION-CCI showed a significant decrease in threshold currents for generating spike and a depolarizing effect of resting membrane potentials. In addition, depolarizing pulses following ION-CCI significantly increased the discharge frequencies of the TRG neuronal action potentials. These changes in medium-/large-diameter CCI-TRG neurons could contribute to the development of mechanical allodynia.

Axotomy-spared neuron model

Uninjured TRG neurons show similar alterations to adjacent TRG neurons following injury [24,34]. In this same model, we found that mechanical allodynia in the territory of ION was observed at 3 days after inferior alveolar nerve (IAN) transection. As shown in Figure 1B, voltage-clamp recording indicated that both I_K and I_A in the medium-/large-diameter spared (uninjured) TRG neurons innervating second branches of the trigeminal nerve (ION) in rats with IAN were significantly smaller than those in naïve rats, as for the CCI model. Corresponding with these changes, we also measured





Types of neuropathic/ inflammatory animal models (Ref.)	Changes in TRG neuronal excitability Whole cell patch-clamp	
	V-clamp	I-clamp
A Chronic constriction nerve Injury (Kitagawa et al., 2006) 	<ul style="list-style-type: none"> ● Medium-/large-diameter <u>CCI</u> neuron • I_A ↓ • I_K ↓ 	<ul style="list-style-type: none"> • Spike frequency ↑ • Spike duration ↑ • Threshold ↓ • Vm (Depo) ↑
B Axotomy-neighboring neurons (Tsuboi et al., 2004) 	<ul style="list-style-type: none"> ● Medium-/large-diameter <u>uninjured</u> neuron • I_A ↓ • I_K ↓ 	<ul style="list-style-type: none"> • Spike frequency ↑ • Spike duration ↑ • Threshold ↓ • Vm (Depo) ↑
C Axotomy- regenerated neurons (Nakagawa et al., 2010) 	<ul style="list-style-type: none"> ● Medium-/large-diameter <u>regenerated</u> neuron • I_A ↓ • I_K ↓ 	<ul style="list-style-type: none"> • Spike frequency ↑ • Spike duration ↑ • Threshold ↓ • Vm (Depo) ↑
D Inflammation -inflamed neurons (Takeda et al., 2006) 	<ul style="list-style-type: none"> ● Small-diameter <u>inflamed</u> neuron • I_A ↓ • I_K ▼ 	<ul style="list-style-type: none"> • Spike frequency ↑ • Threshold ↓ • Vm (Depo) ▲

Figure 1 Trigeminal neuropathic/inflammatory pain models and changes in TRG neuronal activities. **A:** *Chronic constriction nerve injury (CCI) model.* Three days after infraorbital nerve (ION) chronic constriction. Target trigeminal ganglion (TRG) neurons were labeled by fluorogold (FG). Whole-cell patch-clamp recordings under current-/voltage-clamp configurations were performed on medium-/large-diameter CCI neurons. **B:** *Axotomy-neighboring neuron model.* In this model, we found mechanical allodynia in the territory of ION at 2 days after inferior alveolar nerve (IAN) transection. Uninjured TRG neurons innervating the territory of ION were labeled by FG and patch-clamp recordings under current-/voltage-clamp configuration was performed on medium-/large-diameter neurons. **C:** *Axotomy-regenerated neuron model.* In this model, we observed mechanical allodynia at 14 days after IAN transection. FG-injection at 14 days after IAN transection showed massive labeling of trigeminal ganglion containing FG-labeled small-/large-diameter neurons and the patch-clamp recordings under current-/voltage-clamp configuration indicated axotomy-regenerated neurons. **D:** *Inflammation model.* Inflammation was induced by injection of CFA into the rat temporomandibular joint (TMJ). We found mechanical allodynia in the territory of ION at 2 days after CFA injection. In FG-labeled small-diameter TRG neurons innervating TMJ, the whole-cell patch-clamp recordings under current-/voltage-clamp configurations were performed on inflammatory neurons. Blackened circles indicate the trigeminal ganglion neurons. I-III: trigeminal nerves, I_A : transient K current, I_K : sustained K currents, ↑: increase, ↓: decrease, ▼: tendency for decrease in I_K , but not significant, ▲: tendency for depolarization of membrane potential (Vm), but not significant, Depo: Depolarization, V-clamp : Voltage-clamp, I-clamp: Current-clamp.

the following changes under current-clamp conditions: 1) frequencies of the spike discharge induced by depolarizing pulses and duration of uninjured TRG neurons in rats with IAN were significantly increased; and 2) significant decrease in threshold current for generation of spike and a depolarizing effect of resting membrane potentials of spared TRG neurons after IAN transection. These changes of medium-/large-diameter spared TRG neurons might contribute to the development of mechanical allodynia through paracrine signaling (e.g., brain-derived neurotrophic factor) from adjacent injured TRG neurons [24].

Axotomy-regenerated neuron model

Injured nerves regenerate several weeks after nerve damage [39-41]. Some clinical reports have noted that areas innervated by the regenerated nerve show an altered sensitivity to a variety of stimuli compared to those innervated by the intact nerve activity [42-45]. The regenerated neurons also exhibit ectopic discharges [46,47]. Recently we investigated the neural mechanisms underlying abnormal pain following regeneration of the injured IAN [26]. In this model, we observed mechanical allodynia at 11-14 days after IAN transection. Injection of the retrograde tracer, fluorogold (FG), at 14 days after IAN transection showed extensive labeling of TRG neurons, suggesting that most of the TRG neurons present at that stage were regenerated. Whole-cell patch-clamp recording of FG-labeled small-/medium-diameter reinnervated TRG neurons showed a significant decrease in both I_A and I_K in the TRG neurons, a response that was associated with an increase in the spike generation, resulting in the hyperexcitability of reinnervated IAN-TRG neurons (Figure 1C). Taken together with previous results from intracellular single-unit recording [26], these changes may be due to the mechanical allodynia.

Inflammation-model


Complete Freund's adjuvant (CFA) models of inflammation in the orofacial region have been developed in rats to study the trigeminal nervous system [48,49]. Temporomandibular joint (TMJ) inflammation is associated with spreading pain and hyperalgesia [50] and TMJ disorder patients complain of pain from innocuous vibrotactile stimulation [51]. We recently reported that TMJ inflammation modulates the excitability of $A\beta$ -TRG neurons innervating the facial skin via a paracrine mechanism due to the release of substance P (SP) from the small-diameter TRG neurons. Such a release may be important in determining the trigeminal inflammatory allodynia associated with TMJ disorders [52,53]. Using this TMJ inflammatory model, we found that mechanical allodynia in the whisker skin region and I_A densities in the small-diameter TRG neurons innervating TMJ were both significantly decreased in inflamed rats compared to naïve rats [17]. TMJ inflammation significantly

reduced the threshold current and significantly increased action potential firing evoked by depolarizing current pulses. In this model there is a tendency for decreases in I_K density in TMJ inflamed rats compared to naïve rats, although the change is not significant. Since application of 4-aminopyridine (A-type K channel blocker) to naïve rats TRG neurons mimicked the changes in the firing properties observed after CFA treatment [17], I_A suppression of the TRG neurons innervating the TMJ could affect trigeminal inflammatory allodynia in TMJ disorders via paracrine mechanisms of SP in the trigeminal ganglia.

Potassium channels as a potential target for trigeminal neuropathic/inflammatory pain

Lawson et al. [18] reported that several types of Kv channels have been proposed as potential target candidates for pain therapy. It was recently demonstrated that a selective KCNQ/Kv7 channel opener (retigabine), which mediates M-currents, selectively reduces the activity of axotomized $A\delta/C$ -fibers, but not uninjured axons [19], and human C-fiber axons [21]. Xu et al. [20] also demonstrated that this KCNQ/Kv7 channel opener could attenuate allodynia due to TMJ inflammation in rats. Similarly, axotomy reduces K_{Ca} channel activity in the small to medium sized DRG neurons, which would also increase membrane excitability [21]. Thus, these channels may be involved in the development of hyperalgesia and allodynia. On the contrary, spinal nerve ligation suppressed K_{ATP} channels only in large-diameter neurons, but not in small-diameter neurons, from hyperalgesic rats [23].

As described in the preceding chapter, experiments in our trigeminal neuropathic/inflammatory pain model indicated that the common changes in the Kv channels, such as I_A , I_K in rats with neuropathic/inflammatory pain, were significantly suppressed compared to those in naïve rats. Since previous reports suggested that I_A and I_K are important for regulating the firing frequency and duration of action potentials in the TRG neuron, respectively [12,13], these Kv changes caused an incremental spike discharge and prolongation of duration of action potentials in the neuropathic/inflammatory pain rats. The relationship between depression of Kv and excitability of TRG neurons under trigeminal neuropathic/inflammatory pain conditions are summarized in Figure 2. It is well established that the duration of action potentials in primary afferents influences the amount of neurotransmitter released from the soma and the peripheral and central terminals [10,11]. Increased action potential duration and firing frequencies prolong the opening of voltage-gated Ca^{2+} channels, probably potentiating Ca^{2+} influx and causing an increase in the neurotransmitter released from the cell bodies and/or



peripheral and central terminals (Figure 2B). Thus, enhancing the amount of neurotransmitter released from central terminals probably contributes to the hyperexcitability of second-order nociceptive and wide dynamic-range neurons in the trigeminal spinal nucleus (central sensitization) [24,25,49-54]. In addition, increasing the neurotransmitters/neuromodulators (e.g., SP) released from the cell body of TRG neurons through a paracrine or autocrine mechanism may activate the neighboring TRG neurons [52,53]. Since previous studies suggested that peripheral inflammation can depolarize trigeminal satellite-glia cells via activation of

neurokinin 1 receptor (SP release from small diameter TRG neurons) [55,56], such changes may further promote the interaction between satellite-glia cells and TRG neurons associated with pathological pain within the trigeminal ganglia [55-57].

The precise mechanism by which Kv depression of TRG neurons affects the neuropathic/inflammatory condition remains to be determined. Recent studies demonstrated that small- to medium-diameter TRG neurons express glial cell line-derived neurotrophic factor (GDNF) and its receptor components, such as GFR α 1 [58,59]. To this end, we recently found that acute

application of GDNF enhances the neuronal excitability of adult rat small-diameter TRG neurons, which innervate the facial skin in the absence of neuropathic and inflammatory conditions [60]. This potentiation of small-diameter TRG neuronal excitability is mediated by inhibition of Kv channels, and transection of sciatic nerve in rat upregulated the expression of GDNF and GFR α -1 mRNA in the proximal nerve trunk [61,62]. Thus, GDNF-induced potentiation of sensory neuronal excitability may account for enhanced pain sensitivity, resulting from increased levels of GDNF after tissue injury, and further studies are needed to elucidate possible mechanisms underlying this effect.

Finally, opening of K⁺ channels leads to hyperpolarization of the cell membrane, and a resultant decrease in cell excitability. We thus propose the potential development of I_K and/or I_A channel openers as therapeutic agents for preventing trigeminal neuropathic pain, such as mechanical allodynia. Kv channel subunits, Kv2.1 and Kv2.2, contribute to the I_K channel and play a distinct role in regulating neuronal excitability [63,64]. However, Kv1.4 and 4.1-4.3 channels may also be involved in forming A-type Kv channels in sensory neurons [65-67]. Among these Kv channel α subunits, Kv1.4, 2.2, and 4.2 mRNAs are downregulated under neuropathic conditions involving the transection and CCI of sciatic nerves [15,16]. Rasband et al. [65] demonstrated that Kv1.4 is the sole Kv α subunit expressed in smaller diameter DRG neurons, and not Kv4-family channels. Since the pharmacology and gating properties of DRG A-type Kv channels are similar to those of heterogeneously expressed Kv4 channels [66,67], we could not completely rule out a potential role for Kv4.1, 4.2, and 4.3 in neuropathic pain. Previously, we reported that TMJ inflammation decreased the expression of Kv1.4 subunits in small-/medium-diameter (A δ -/C-) TRG neurons and that this may contribute to trigeminal inflammatory allodynia associated with TMJ disorders [68]. Nerve injury also induces accelerated reduction in Kv 1.4 expression in the small-diameter nociceptive DRG neurons [65]. Taken together, these findings suggest Kv 1.4 channel as candidate molecular targets, and further studies are needed to explore this possibility.

Conclusion

Several different animal models of trigeminal nerve injury/inflammation showed depression of both I_A and I_K in TRG neurons, compared to naïve rats. These common changes contribute to the incremental spike discharge and action potential prolongation in these neurons, and may alter the properties of trigeminal pain pathways. Our results therefore would suggest that Kv channel (I_A and I_K) openers have potential as

therapeutic agents against trigeminal neuropathic/inflammatory pain, such as mechanical allodynia.

List of abbreviations used

4-AP: 4-aminopyridine; CCI: chronic constriction nerve; CFA: Complete Freund's adjuvant; DDP: duration of depolarization; DRG: dorsal root ganglion; DRP: duration of repolarization; α -DTX: α -dendrotoxin; FG: fluorogold; GDNF: Glial cell-derived neurotrophic factor; I_A: fast inactivating transient current (A-current); I_K: dominant sustained current (K-current); IAN: Inferior alveolar nerve; ION: infraorbital nerve; K_{ATP}: ATP-sensitive potassium; K_{Ca}: Calcium activated potassium; Kv: voltage-gated K⁺ channels; SP: Substance P; TEA: Tetraethylammonium; TMJ: temporomandibular joint; TRG: trigeminal ganglion.

Author details

¹Department of Physiology, School of Life Dentistry at Tokyo, Nippon Dental University, 1-9-20, Fujimi-cho, Chiyoda-ku, Tokyo, 102-8159, Japan.

²Department of Physiology, School of Dentistry, Nihon University, 1-8-13, Kandasurugadai, Chiyoda-ku, Tokyo, 101-8310, Japan.

³Division of Oral Physiology, Department of Oral Biological Science, Niigata University Graduate School of Medical and Dental Sciences, 2-5274, Gakkocho-dori, Niigata, 951-8514, Japan. ⁴Department of Hygiene and Oral Health, Showa University School of Dentistry, 1-5-8, Hatanodai, Shinagawa-ku, Tokyo, 142-8555, Japan.

Authors' contributions

MT conceived and wrote the manuscript. YT, JK, and KN carried out the experiments in the main part of references [242526]. KI designed the main part of experiments [242526]. SM conceptualized the manuscript and helped to finalize the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 2 November 2010 Accepted: 10 January 2011

Published: 10 January 2011

References

- Devor M: Strategies for finding new pharmacological targets for neuropathic pain. *Clin Pain Head Reports* 2004, **8**:187-191.
- Niv D, Devor M: Refractory neuropathic pain: the nature and extent of the problem. *Pain Prac* 2006, **6**:3-9.
- Schwartz J, Naff N: The management of neuropathic pain. *Neurosurg Clin N Am* 2004, **15**:231-239.
- Scholtz J, Woolf CJ: Can we conquer pain? *Nat Neurosci* 2002, **5**:1062-1067.
- Costigas M, Scholz J, Woolf CJ: Neuropathic pain: A maladaptive response of nervous system to damage. *Annu Rev Neurosci* 2009, **32**:1-32.
- Ficker E, Heinemann U: Slow and fast transient potassium currents in cultured rat hippocampus cells. *J Physiol (Lond)* 1992, **445**:431-455.
- Wu LG, Saggau P: Presynaptic inhibition of elicited neurotransmitter release. *Trend Neurosci* 1997, **20**:204-212.
- Hille B: Potassium channels and chloride channels. In *Ion channels of excitable membranes*. 3 edition. Edited by: Hille BD. Sunderland: Sinauer Associates; 2001:134-167.
- Everill B, Rizzo MA, Kocsis JD: Morphologically identified cutaneous afferent DRG neurons express three different potassium currents in varying properties. *J Neurophysiol* 1998, **79**:1814-1824.
- Takeda M, Tanimoto T, Ikeda M, Kadoi J, Matsumoto S: Activation of GABA_B receptor inhibits the excitability of rat small diameter trigeminal root ganglion neurons. *Neuroscience* 2004, **123**:491-505.
- Takeda M, Tanimoto T, Ikeda M, Kadoi J, Nasu M, Matsumoto S: Opioidergic modulation of excitability of rat trigeminal root ganglion neuron projections to the superficial layer of cervical dorsal horn. *Neuroscience* 2004, **125**:995-1008.
- Yoshida S, Matsumoto S: Effects of α -Dendrotoxin on K⁺ currents and action potentials in tetrodotoxin resistant adult rat trigeminal ganglion neurons. *J Pharmacol Exp Ther* 2005, **314**:437-445.

13. Matsumoto S, Yoshida S, Takahashi M, Saiki C, Takeda M: **The roles of I_D, I_A and I_K in the electrophysiological functions of small-diameter rat trigeminal ganglion neurons.** *Curr Mol Pharmacol* 2010, **3**:30-36.
14. Everill B, Kocsis JD: **Reduction in potassium currents in identified cutaneous afferent dorsal root ganglion neurons after axotomy.** *J Neurophysiol* 1999, **82**:700-708.
15. Kim DS, Choi JO, Rim HD, Cho HJ: **Down regulation of voltage gated potassium channels a gene expression in dorsal root ganglion following chronic constriction injury of rat sciatic nerve.** *Mol Brain Res* 2002, **105**:146-152.
16. Park SY, Choi JY, Lee YS, Cho H, Kim DS: **Down regulation of voltage-gated potassium channel a gene expression by axotomy and neurotrophin in rat dorsal root ganglia.** *Mol Cell* 2003, **16**:256-259.
17. Takeda M, Tanimoto T, Ikeda M, Nasu M, Kadoi J, Yoshida S, Matsumoto S: **Enhanced excitability of rat trigeminal root ganglion neurons via decrease in A-type potassium currents following temporomandibular joint inflammation.** *Neuroscience* 2006, **138**:621-630.
18. Lawson K: **Potassium channel as targets for the management of pain.** *Cent Nerv Syst Agents Med Chem* 2006, **6**:119-128.
19. Rosa C, Lopez-Garcia JA: **Retigabine, the specific KCNQ channel opener, blocks ectopic discharges in axotomized fibers.** *Pain* 2008, **138**:537-545.
20. Xu W, Wu Ym, Bi Y, Tan L, Gan Y, Wang KW: **Activation of voltage-gated KCNQ/Kv7 channels by anticonvulsant retigabine attenuates mechanical allodynia of inflammatory temporomandibular joint in rats.** *Mol Pain* 2010, **6**:49.
21. Lang PM, Fleckenstein J, Passmore GM, Brown DA, Grafe P: **Retigabine reduces the excitability of unmyelinated peripheral human axons.** *Neuropharmacology* 2008, **54**:1271-1278.
22. Sarantopoulos CD, McCallum JB, Ringaud M, Fuchs A, Kwok WM, Hogan QH: **Opposing effects of spinal nerve ligation on calcium-activated potassium currents in axotomized and adjacent mammalian primary afferent neurons.** *Brain Res* 2007, **1132**:84-99.
23. Kawano T, Zoga V, McCallum JB, Wu HE, Gemes G, Liang MY, Abram S, Kwok WM, Hogan QH, Sarantopoulos CD: **ATP-sensitive potassium currents in rat primary afferent neurons: Biophysical, pharmacological properties, and alterations by painful nerve injury.** *Neuroscience* 2009, **162**:431-443.
24. Tsuboi Y, Takeda M, Tanimoto T, Ikeda M, Matsumoto S, Kitagawa J, Teramoto K, Shimizu K, Yamazaki Y, Shima A, Ren K, Iwata K: **Alternation of the second branch of the trigeminal nerve activity following inferior alveolar nerve transection in rats.** *Pain* 2004, **111**:323-334.
25. Kitagawa J, Takeda M, Suzuki I, Kadoi J, Tsuboi Y, Honda K, Matsumoto S, Nakagawa H, Tanabe A, Iwata K: **Mechanisms involved in modulation of trigeminal primary afferent activity in rats with peripheral mononeuropathy.** *Eur J Neurosci* 2006, **24**:1976-1986.
26. Nakagawa K, Takeda M, Tsuboi Y, Kondo M, Kitagawa J, Matsumoto S, Kobayashi A, Sessle BJ, Shinoda M, Iwata K: **Mechanisms involved in modulation of trigeminal primary afferent activity in rats with peripheral mononeuropathy.** *Mol Pain* 2010, **6**:9.
27. Bennett GJ, Xie YK: **A peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man.** *Pain* 1988, **33**:87-107.
28. Decosterd I, Allchorne A, Woolf CJ: **Progressive tactile hypersensitivity after a peripheral nerve crush: non-noxious mechanical stimulus induced neuropathic pain.** *Pain* 2002, **100**:155-162.
29. Seltzer Z, Dubner R, Shir Y: **A novel behavioral model of neuropathic pain disorders produced in rats partial sciatic nerve injury.** *Pain* 1990, **43**:205-218.
30. Stussle SL, Crisp T, McBunney DL, Schechter JB, Lovell JA, Cruce WL: **Neuropathic pain in aged rats: Behavioral responses and astrocytic activation.** *Exp Brain Res* 2001, **137**:219-227.
31. Guilbaud G, Benoist JM, Jazat F, Gauton M: **Neuronal responsiveness in the ventrobasal thalamic complex of rats with an experimental peripheral mononeuropathy.** *J Neurophysiol* 1990, **64**:1537-1554.
32. Abdula FA, Smith PA: **Axotomy- and autotomy-induced changes in the excitability of rat dorsal root ganglion neurons.** *J Neurophysiol* 2001, **85**:630-643.
33. Abdula FA, Smith PA: **Changes in Na⁺ channels currents of rat dorsal root ganglion neurons following autotomy and axotomy-induced autotomy.** *J Neurophysiol* 2002, **88**:2518-2529.
34. Fukuoka T, Noguchi K: **Contribution of the spared primary afferent neurons to the pathomechanisms of neuropathic pain.** *Mol Neurobiol* 2002, **26**:57-67.
35. Fukuoka T, Tokunaga A, Tachibana T, Dai Y, Yamanaka H, Noguchi K: **VR1, but not P2X(3) increase in the spared L4 DRG in rats with L5 spinal nerve ligation.** *Pain* 2002, **99**:111-120.
36. Fried K, Bongenhillem FM, Boissonade FM, Robinson PP: **Nerve injury-induced pain in the trigeminal system.** *Neuroscientist* 2001, **7**:155-165.
37. Sessle BJ: **Acute and chronic, craniofacial pain: brainstem mechanisms of nociceptive transmission and neuroplasticity, and their clinical correlates.** *Crit Rev Oral Biol Med* 2000, **11**:57-91.
38. Hoffmann KD, Matthews MA: **Comparison of sympathetic neurons in orofacial and upper extremity nerve: implications for causalgia.** *J Oral Maxillofac Surg* 1990, **48**:720-726.
39. Cain P, Frank ME, Barry MA: **Recovery of chorda tympani nerve function following injury.** *Exp Neurol* 1996, **141**:337-346.
40. Lago N, Uchida E, Ramchandran A, Navarro Z: **Neurological assessment of regenerative electrodes for bidirectional interfacing injured peripheral nerves.** *IEEE Trans Biomed Eng* 2007, **54**:1129-1137.
41. Lee YS, Lin CY, Robertson RT, Hisao I, Lin VW: **Motor recovery and anatomical evidence of axonal regrowth in spinal cord-repaired adult rats.** *J Neuropathol Exp Neurol* 2004, **63**:233-245.
42. deMedinacell L, Hurpeau J, Merle M, Bergorre H: **Cold and post-traumatic pain: modeling of the peripheral nerve message.** *BioSystems* 1989, **49**:228-240.
43. Ochs G, Schenk M, Struppler A: **Painful dysaesthesia following peripheral nerve injury: clinical and electrophysiological study.** *Brain Res* 1989, **496**:228-240.
44. Saito T, Schbamori Y, Manabe Y, Yamaguchi T, Yamamoto T, Ohtsubo T, Saito H: **Morphological and functional study of regenerated corda tympani nerves in humans.** *Ann Otol Rhinol Laryngol* 2000, **109**:703-709.
45. Sawai S, Kanai K, Nakata M, Hiraga A, Misawa S, Isose S, Hattori T, Kuwabara S: **Changes in excitability properties associated with axonal regeneration in human neuropathy and mouse Wallerian degeneration.** *Clin Neurophysiol* 2008, **119**:1097-1105.
46. Gorodetskaya N, Constantin C, Jägg W: **Ectopic activity in cutaneous regenerating afferent nerve fibers following nerve lesion in the rat.** *Eur J Neurosci* 2003, **18**:2487-2497.
47. Jägg W, Grossmann L, Gorodetskaya N: **Mechano- and thermosensitivity of regenerating cutaneous afferent nerve fibers.** *Exp Brain Res* 2009, **196**:101-114.
48. Imbe H, Iwata K, Zhou QQ, Zou S, Dubner R: **Orofacial deep and cutaneous tissue inflammation and trigeminal neuronal activation.** *Cells Tissue Organs* 2001, **169**:238-247.
49. Iwata K, Tashiro A, Tsuboi Y, Imai T, Sumino R, Morimoto T, Dubner R, Ren K: **Medullary dorsal horn neuronal activity in rats with persistent temporomandibular joint and perioral inflammation.** *J Neurophysiol* 1999, **82**:1244-1253.
50. Sessle BJ, Hu JW: **Mechanism of pain arising from articular tissues.** *Can J Physiol Pharmacol* 1999, **69**:617-626.
51. Takeda M, Tanimoto T, Ikeda M, Nasu M, Kadoi J, Shima Y, Ohta H, Matsumoto S: **Temporomandibular joint inflammation potentiates the excitabilities of trigeminal root ganglion neurons innervating the facial skin in rats.** *J Neurophysiol* 2005, **93**:2723-2738.
52. Takeda M, Tanimoto T, Nasu M, Ikeda M, Kadoi J, Matsumoto S: **Activation of NK1 receptor of trigeminal root ganglion via substance P paracrine mechanism contributes to the mechanical allodynia in the temporomandibular joint inflammation in rats.** *Pain* 2005, **116**:375-385.
53. Iwata K, Imai T, Tsuboi Y, Tashiro A, Ogawa A, Morimoto T, Masuda Y, Tachibana Y, Hu JW: **Alternation of medullary dorsal horn neuronal activity following inferior alveolar nerve transection in rats.** *J Neurophysiol* 2001, **86**:2868-2877.
54. Nomura H, Ogawa A, Tashiro A, Morimoto T, Hu JW, Iwata K: **Induction of Fos protein-like immunoreactivity in the trigeminal spinal nucleus caudalis and upper cervical cord following noxious and non-noxious mechanical stimulation of whisker pad of the rat with a inferior alveolar nerve transection.** *Pain* 2002, **95**:225-238.
55. Takeda M, Tanimoto T, Kadoi J, Nasu M, Takahashi M, Kitagawa J, Matsumoto S: **Enhanced excitability of nociceptive trigeminal ganglion neurons by satellite glial cytokine following peripheral inflammation.** *Pain* 2007, **29**:155-166.
56. Takeda M, Takahashi M, Matsumoto S: **Contribution of activated interleukin receptors in trigeminal ganglion neurons to hyperalgesia via**

- satellite glial interlekin-1 β paracrine mechanism. *Brain Behav Immun* 2008, **22**:1016-1023.
57. Takeda M, Takahashi M, Matsumoto S: Contribution of activation of satellite glia in the sensory ganglia to pathological pain. *Neurosci Biobehav Rev* 2009, **33**:784-792.
 58. Quartu M, Sera MP, Bachis A, Lai ML, Ambu R, Del Fiacco M: Glial cell line-derived neurotrophic factor-like immunoreactivity in human trigeminal ganglion and nucleus. *Brain Res* 1999, **847**:196-202.
 59. Quartu M, Serra MP, Mascia F, Boi M, Lai ML, Spano A, Del Fiacco M: GDNF family ligand receptor components Ret and GFR alpha-1 in the human trigeminal ganglion and sensory nuclei. *Brain Res Bull* 2006, **69**:393-403.
 60. Takeda M, Kitagawa J, Nasu M, Takahashi M, Iwata K, Matsumoto S: Glial cell line-derived neurotrophic factor acutely modulates the excitability of rat small diameter trigeminal ganglion neurons innervating facial skin. *Brain Behav Immun* 2010, **24**:72-82.
 61. Hoke A, Cheng C, Zochodne DW: Expression of glial cell line-derived neurotrophic factor family of growth factors in peripheral nerve injury rats. *Neuroreport* 2000, **11**:1651-1654.
 62. Bennett DL, Boucher TJ, Armanini MP, Poulsen KT, Michael GJ, Priestley JV, Phillips HS, McMahon SB, Shelton DL: The glial cell line-derived neurotrophic factor family receptor components are differentially regulated within sensory neurons after nerve injury. *J Neurosci* 2000, **20**:427-443.
 63. Murakoshi H, Trimmer JS: Identification of Kv2.1 K⁺ channel as a major component of the delayed rectifier K⁺ current in rat hippocampal neurons. *J Neurosci* 1999, **19**:1728-1735.
 64. Malin SA, Nerbonne JM: Delayed rectifier K⁺ currents, IK are encoded by Kv2 alpha-subunits and regulate tonic firing in mammalian sympathetic neurons. *J Neurosci* 2002, **22**:10094-10105.
 65. Rasband MN, Park EW, Vanderah TW, Lai J, Porreca F, Trimmer JS: Distinct potassium channels on pain-sensing neurons. *Proc Natl Acad Sci USA* 2001, **98**:13373-13378.
 66. Mathie A, Woolton JRA, Watkins CS: Voltage-activated potassium channels in mammalian neurons and their block by novel pharmacological agents. *Gen Pharmacol* 1998, **30**:13-14.
 67. Winkleman DL, Beck CL, Ypey DL, O'Leary ME: Inhibition of the A-type K channels of dorsal root ganglion neurons by long duration of anesthetic butamben. *J Pharmacol Exp Ther* 2005, **314**:1177-1186.
 68. Takeda M, Tanimoto T, Nasu M, Matsumoto S: Temporomandibular joint inflammation decreases the voltage-gated K⁺ channel subtype Kv1.4-immunoreactivity of in trigeminal ganglion neurons in rats. *Eur J Pain* 2008, **12**:189-195.

doi:10.1186/1744-8069-7-5

Cite this article as: Takeda *et al.*: Potassium channels as a potential therapeutic target for trigeminal neuropathic and inflammatory pain. *Molecular Pain* 2011 **7**:5.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

