

Comment

## Availability of new Bayesian-delimited gecko names and the importance of character-based species descriptions

Leaché & Fujita [1] present an empirical example of Bayesian species delimitation (BSD; [2]) to recognize three new species of African geckos from within the range of the widespread taxon *Hemidactylus fasciatus*, Gray 1842. As with any new method, BSD will undoubtedly generate questions and discussions about its practicality and assumptions. However, the use of this method of delimiting species does not obviate the need to adhere to the practice of describing or defining species on the basis of intrinsic characters.

In order for any species name in zoology to be available, it must satisfy the applicable provisions of Articles 11–20 of the *International Code of Zoological Nomenclature* (1999) (hereafter ‘the Code’). Relevant to this issue, ‘To be available, every new name published after 1930... must be accompanied by a description or definition that states in words characters that are purported to differentiate the taxon (Article 13.1.1, p. 17)’ where a character is ‘any attribute of organisms used for recognizing, differentiating, or classifying taxa’ (Glossary, p. 101). Leaché & Fujita [1] do not provide descriptions or definitions for their new taxa, only putative diagnoses (recommended but not required under the Code (Recommendation 13A)), and these are based not upon characters but upon degree of support under their species delimitation model (e.g. ‘This species includes all populations that cluster with those from the Togo Hills included in this study with strong support in the Bayesian species delimitation model’). Although geographical location could be argued to be an organismal attribute, (i) Article 12.3 and, by implication, Article 13.1 [3] clarify that the mention of a locality does not in itself constitute a description, and (ii) Leaché & Fujita [1] have not actually used locality as a character, but as an indicator of the cluster to which populations belong. The new names they propose thus lack definitions or descriptions of *organismal* attributes (characters) as required by the Code and their diagnoses consist solely of extrinsic relational statements about *populations*. The new names proposed for members of the *H. fasciatus* group (*Hemidactylus coalescens*, *Hemidactylus eniangii*, *Hemidactylus kyaboboensis*) fail to conform to Article 13 and are *nomina nuda*, and thus unavailable under the Code.

The diagnoses of their new species are similar to the definitions of clade names under phylogenetic taxonomy, but their species descriptions are also unavailable under

the *PhyloCode* [4], which defers to ‘provisions of the appropriate rank-based code (e.g. ... ICZN)’ for species availability.

The unavailability of Leaché and Fujita’s gecko names rests on more than a technicality. We emphasize this point to highlight *the very purpose* of the taxonomic protocols inherent in the Code’s adherence to character-based species definitions. The application of Leaché & Fujita’s approach would no doubt facilitate the description of many new species, and in light of the ongoing biodiversity crisis facing the planet [5] and the need to describe new species before they are lost to extinction [6], this might be viewed positively. However, the deleterious impact of premature taxonomic inflation to conservation efforts is well documented [7–9]. Thus, our concern is that the conservation benefit from a flood of superficial new species descriptions might be outweighed by the cost of taxonomic instability and confusion. For example, unambiguous allocation of a specimen to one of Leaché & Fujita’s new species of *Hemidactylus* could only be accomplished by its incorporation into their species delimitation model. This would necessitate the collection of DNA sequence data and replication of their analytical protocol for each unknown sample. Alternatively, specimens could be compared directly to the cited type material but, in the absence of informative diagnostic characters, this would effectively entail a re-analysis of the entire *Hemidactylus fasciatus* group.

The Code’s requirement for a character-based description or definition forces taxonomists to make comparisons with similar or related forms and to identify features intrinsic to the organism. Character differences not only reflect lineage independence, but also they make the associated species names the very currency of communication in systematic biology. Although we suspect few biologists believe either that all descriptions under the Code employ a uniform degree of rigor, or that the biological reality of a species is dependent upon the enumeration of a set of diagnostic characters, the focus on characters, no matter how artificial, allows for comparability. This decreases taxonomic ambiguity and instability, and results in a system of names that can be used both by taxonomists and by the much larger community of consumers of taxonomic information. Characters must be included by all authors publishing new species descriptions, whether or not they use new or traditional means of species delimitation.

We submit that there are at least three alternative approaches that are superior to creating unavailable names. First, genetic characters can be used to satisfy

Article 13.1.1 of the *Code* [10], just as morphological characters do (e.g. [11]). Second, evolutionarily distinct lineages (i.e. 'candidate species' *sensu* [12]) may be left unnamed until additional data allow for proper descriptions (e.g. [13,14]). Finally, morphology can be studied in conjunction with application of the Bayesian model.

Leaché & Fujita [1] stated 'In terms of *H. fasciatus*, we are not aware of any morphological or ecological characteristics that differentiate these lineages,' but it is unclear whether a morphological survey or analysis was undertaken. Although some organisms exhibit limited interspecific external variation owing to extreme morphological conservativeness and/or homoplasy [15], descriptions of many 'cryptic' gecko species have successfully used tree-based data (gene tree monophyly and distance data) in conjunction with explicit character-based descriptions and diagnoses (e.g. [16,17]). Corroborating their new method with a set of differentiating characters would have been an excellent way to bolster Leaché & Fujita's [1] test of BSD. Any single method of inference can be misleading, but the agreement between independent classes of data can result in increased confidence in hypotheses such as species delimitations [18].

We do not deny that the populations of West African *Hemidactylus* named by Leaché & Fujita [1] represent valid taxa under any of several lineage-based species concepts [19]. The Bayesian approach to species delimitation may well provide objective criteria for discerning species boundaries using genetic data and we favour further exploration of this approach, especially if used in combination with surveys of characters that can be used to recognize species. If, however, as in the paper by Leaché & Fujita, Bayesian species descriptions have as their sole underpinnings the degree of support for a pattern of clustering under a particular model, we regard them as both practically unworkable and philosophically inadvisable.

Aaron M. Bauer<sup>1,\*</sup>, James F. Parham<sup>2</sup>, Rafe M. Brown<sup>3</sup>, Bryan L. Stuart<sup>4</sup>, Lee Grismer<sup>5</sup>, Theodore J. Papenfuss<sup>6</sup>, Wolfgang Böhme<sup>7</sup>, Jay M. Savage<sup>8</sup>, Salvador Carranza<sup>9</sup>, Jesse L. Grismer<sup>1,3</sup>, Philipp Wagner<sup>7</sup>, Andreas Schmitz<sup>10</sup>, Natalia B. Ananjeva<sup>11</sup> and Robert F. Inger<sup>12</sup>

<sup>1</sup>Department of Biology, Villanova University, 800 Lancaster Avenue, Villanova, PA 19085, USA

<sup>2</sup>Alabama Museum of Natural History, University of Alabama, Box 870340, Tuscaloosa, AL 35487, USA

<sup>3</sup>Natural History Museum and Biodiversity Research Center and Department of Ecology and Evolutionary Biology, 1345 Jayhawk Blvd, Dyche Hall, University of Kansas, Lawrence, KS 66045, USA

<sup>4</sup>North Carolina Museum of Natural Sciences, 11 West Jones Street, Raleigh, NC 27601, USA

<sup>5</sup>Department of Biology, La Sierra University, 4500 Riverwalk Parkway, Riverside, CA 92515, USA

<sup>6</sup>Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA

<sup>7</sup>Zoologisches Forschungsmuseum Alexander Koenig, Adenauerallee 160, 53113 Bonn, Germany

<sup>8</sup>Department of Biology, San Diego State University, San Diego, CA 92182-4614, USA

<sup>9</sup>Institute of Evolutionary Biology (CSIC-UPF), 08003 Barcelona, Spain

<sup>10</sup>Department of Herpetology and Ichthyology, Muséum d'Histoire Naturelle, C.P. 6434, CH-1211, Genève 6, Switzerland

<sup>11</sup>Division of Herpetology and Ornithology, Zoological Institute, Russian Academy of Sciences, Universitetskaya nab., 1, St Petersburg 199034, Russia

<sup>12</sup>Division of Amphibians and Reptiles, The Field Museum of Natural History, Chicago, IL, USA

\*Author for correspondence (aaron.bauer@villanova.edu).

## REFERENCES

- 1 Leaché, A. D. & Fujita, M. K. 2010 Bayesian species delimitation in West African forest geckos (*Hemidactylus fasciatus*). *Proc. R. Soc. B* **277**, 3071–3077. (doi:10.1098/rspb.2010.0662)
- 2 Yang, Z. & Rannala, B. 2010 Bayesian species delimitation using multilocus sequence data. *Proc. Natl Acad. Sci. USA* **107**, 9264–9269. (doi:10.1073/pnas.0913022107)
- 3 Čermák, S. 2009 The Plio-Pleistocene record of *Hypolagus* (Lagomorpha, Leporidae) from the Czech and Slovak Republics with comments on systematics and classification of the genus. *Bull. Geosci.* **84**, 497–524. (doi:10.3140/bull.geosci.1104)
- 4 Cantino, P. D. & de Queiroz, K. 2010 PhyloCode. International code of phylogenetic nomenclature version 4c. See <http://www.ohio.edu/phylocode/>.
- 5 Pimm, S. L., Russell, G. R., Gittleman, J. L. & Brooks, T. M. 1995 The future of biodiversity. *Science* **269**, 347–350. (doi:10.1126/science.269.5222.347)
- 6 Erwin, T. L. & Johnson, P. J. 2000 Naming species, a new paradigm for crisis management in taxonomy: rapid journal validation of scientific names enhanced with more complete descriptions on the internet. *Coleopt. Bull.* **54**, 269–278.
- 7 Isaac, N. J. B., Mallet, J. & Mace, G. M. 2004 Taxonomic inflation: its influence on macroecology and conservation. *Trends Ecol. Evol.* **19**, 464–469. (doi:10.1016/j.tree.2004.06.004)
- 8 Mace, G. M. 2004 The role of taxonomy in species conservation. *Phil. Trans. R. Soc. Lond. B* **359**, 711–719. (doi:10.1098/rstb.2003.1454)
- 9 Parham, J. F., Türkozan, O., Stuart, B. L., Arakelyan, M., Shafei, S., Macey, J. R. & Papenfuss, T. J. 2006 Genetic evidence for premature taxonomic inflation in Middle Eastern tortoises. *Proc. Calif. Acad. Sci.* **57**, 955–964.
- 10 ICZN (International Commission on Zoological Nomenclature) 1999 *International code of zoological nomenclature*, 4th edn. London, UK: International Trust for Zoological Nomenclature.
- 11 Chippindale, P. T., Hillis, D. M., Wiens, J. J. & Price, A. H. 2000 Phylogenetic relationships and systematic revision of central Texas hemidactyliine plethodontid salamanders. *Herpetol. Monogr.* **14**, 1–80. (doi:10.2307/1467045)
- 12 Morando, M., Avila, L. J. & Sites Jr, J. W. 2003 Sampling strategies for delimiting species: genes, individuals, and populations in the *Liolaemus elongatus-kriegi* complex (Squamata; Liolaemidae) in Andean–Patagonian South America. *Syst. Biol.* **52**, 159–185. (doi:10.1080/10635150390192717)
- 13 Inger, R. F., Stuart, B. L. & Iskandar, D. T. 2009 Systematics of a widespread Southeast Asian frog, *Rana chalconota* (Amphibia: Anura: Ranidae). *Zool. J. Linn. Soc.* **155**, 123–147. (doi:10.1111/j.1096-3642.2008.00440.x)
- 14 Oliver, P. M., Adams, M., Lee, M. S. Y., Hutchinson, M. N. & Doughty, P. 2009 Cryptic diversity in

- vertebrates: molecular data double estimates of species diversity in a radiation of Australian lizards (*Diplodactylus*, Gekkota). *Proc. R. Soc. B* **276**, 2001–2007. (doi:10.1098/rspb.2008.1881)
- 15 Bickford, D., Lohman, D. J., Sodhi, N. S., Ng, P. K. L., Meier, R., Winker, K., Ingram, K. K. & Das, I. 2007 Cryptic species as a window on diversity and conservation. *Trends Ecol. Evol.* **22**, 148–155. (doi:10.1016/j.tree.2006.11.004)
- 16 Bauer, A. M., Jackman, T., Sadler, R. A. & Whitaker, A. H. 2006 A revision of the *Bavayia validiclavis* group (Squamata: Gekkota: Diplodactylidae), a clade of New Caledonian geckos exhibiting microendemism. *Proc. Calif. Acad. Sci.* **57**, 503–547.
- 17 Welton, L. J., Siler, C., Diesmos, A. C. & Brown, R. M. 2010 Phylogeny-based species delimitation of southern Philippine bent-toed geckos and a new species of *Cyrtodactylus* (Squamata: Gekkonidae) from western Mindanao and the Sulu Archipelago. *Zootaxa* **2390**, 49–68.
- 18 de Queiroz, K. 2007 Species concepts and species delimitation. *Syst. Biol.* **56**, 879–886. (doi:10.1080/10635150701701083)
- 19 de Queiroz, K. 1998 The general lineage concept of species, species criteria, and the process of speciation. In *Endless forms: species and speciation* (eds D. J. Howard & S. H. Berlocher), pp. 57–75. Oxford, UK: Oxford University Press.