



Published in final edited form as:

Arch Gen Psychiatry. 2010 October ; 67(10): 1012–1024. doi:10.1001/archgenpsychiatry.2010.111.

A Meta-analysis of Depression During Pregnancy and the Risk of Preterm Birth, Low Birth Weight, and Intrauterine Growth Restriction

Dr. Nancy K. Grote, PhD, Dr. Jeffrey A. Bridge, PhD, Dr. Amelia R. Gavin, PhD, Dr. Jennifer L. Melville, MD, Dr. Satish Iyengar, PhD, and Dr. Wayne J. Katon, MD

School of Social Work (Drs Grote and Gavin), Departments of Obstetrics and Gynecology (Dr Melville) and Psychiatry and Behavioral Sciences (Dr Katon), University of Washington, Seattle; Department of Pediatrics and The Research Institute at Nationwide Children's Hospital, The Ohio State University, Columbus (Dr Bridge); and Department of Statistics, University of Pittsburgh, Pittsburgh, Pennsylvania (Dr Iyengar)

Abstract

Context—Maternal depressive symptoms during pregnancy have been reported in some, but not all, studies to be associated with an increased risk of preterm birth (PTB), low birth weight (LBW), and intrauterine growth restriction (IUGR).

Objective—To estimate the risk of PTB, LBW, and IUGR associated with antenatal depression.

Data Sources and Study Selection—We searched for English-language and non-English-language articles via the MEDLINE, PsycINFO, CINAHL, Social Work Abstracts, Social Services Abstracts, and Dissertation Abstracts International databases (January 1980 through December 2009). We aimed to include prospective studies reporting data on antenatal depression and at least 1 adverse birth outcome: PTB (<37 weeks' gestation), LBW (<2500 g), or IUGR (<10th percentile for gestational age). Of 862 reviewed studies, 29 US-published and non-US-published studies met the selection criteria.

Data Extraction—Information was extracted on study characteristics, antenatal depression measurement, and other biopsychosocial risk factors and was reviewed twice to minimize error.

Data Synthesis—Pooled relative risks (RRs) for the effect of antenatal depression on each birth outcome were calculated using random-effects methods. In studies of PTB, LBW, and IUGR that used a categorical depression measure, pooled effect sizes were significantly larger (pooled RR [95% confidence interval]=1.39 [1.19–1.61], 1.49 [1.25–1.77], and 1.45 [1.05–2.02], respectively) compared with studies that used a continuous depression measure (1.03 [1.00–1.06], 1.04 [0.99–1.09], and 1.02 [1.00–1.04], respectively). The estimates of risk for categorically defined antenatal

Correspondence: Nancy K. Grote, PhD, School of Social Work, 4101 15th Ave NE, University of Washington, Seattle, WA 98105 (ngrote@u.washington.edu).

Additional Contributions: We gratefully acknowledge the contribution of additional data from some of the investigators of the original studies included in this meta-analysis. We also thank all those who responded to our inquiries, even if data from their studies could not be made available. Finally, we thank Kimberly A. Yonkers, MD, Yale University, for her thoughtful review of the manuscript, for which no compensation was received.

Financial Disclosure: Dr Bridge has received honoraria for participation in a suicidality consensus conference supported by independent educational or other unrestricted grant support from AstraZeneca, Dainippon Sumitomo, Eli Lilly, Forest Research Institute, Ortho-McNeil-Janssen, Pfizer Inc, Roche Pharmaceuticals, Sanofi-Aventis, Schering-Plough, Sepracor, and United BioSource; and he has presented on suicide prevention in children and adolescents at a conference supported in part by Eli Lilly and Lundbeck. Dr Katon has received honoraria for lectures from Lilly, Forest, Wyeth, and Pfizer and has served on advisory boards for Lilly and Wyeth during the last 2 years.

depression and PTB and LBW remained significant when the trim-and-fill procedure was used to correct for publication bias. The risk of LBW associated with antenatal depression was significantly larger in developing countries (RR=2.05; 95% confidence interval, 1.43–2.93) compared with the United States (RR=1.10; 95% confidence interval, 1.01–1.21) or European social democracies (RR=1.16; 95% confidence interval, 0.92–1.47). Categorically defined antenatal depression tended to be associated with an increased risk of PTB among women of lower socioeconomic status in the United States.

Conclusions—Women with depression during pregnancy are at increased risk for PTB and LBW, although the magnitude of the effect varies as a function of depression measurement, country location, and US socioeconomic status. An important implication of these findings is that antenatal depression should be identified through universal screening and treated.

Preterm birth (PTB), Low birth weight (LBW), and intrauterine growth restriction (IUGR) are the leading causes of neonatal, infant, and childhood morbidity, mortality, and neurodevelopmental impairments and disabilities worldwide.^{1–5} Maternal depression during pregnancy has begun to be recognized as a factor that may adversely alter pregnancy outcomes.^{6–8} Depression also has been linked to known risk factors for adverse pregnancy outcomes such as smoking,^{9,10} substance abuse,¹¹ hypertension,^{12,13} preeclampsia,^{14,15} and gestational diabetes.^{16,17} Recent estimates of the prevalence of major depression during pregnancy show that from 8.3%¹⁸ to 12.7%¹⁹ of US women experience this condition. Moreover, many community-based studies have indicated that poor urban women from minority backgrounds^{20–22} are at least twice as likely as middle-class women^{23–25} to meet diagnostic criteria for major and minor depression during pregnancy and the postpartum period (20%–25% vs 9%–13%, respectively). These findings are congruent with epidemiological data showing higher rates of depression in poor young women²⁶ and with data on prevalence rates of perinatal depression for women in developing countries.^{27–29}

Research findings during the last decades on the links between antenatal depression and PTB, LBW, and IUGR have revealed a relatively inconsistent and inconclusive picture.³⁰ Some evidence indicates that depression during pregnancy may be significantly related to PTB,^{31,32} LBW,^{33,34} and IUGR,³⁵ whereas other studies have reported no direct association.^{36–38} Reasons for these contradictory results are most likely related to differences in (1) study design, methods, sample sizes, and the timing, frequency, and type of antenatal depression measurement; (2) misclassification bias with respect to depression or birth outcomes; (3) the populations studied; and (4) the extent to which studies control for confounding factors of PTB, LBW, or IUGR such as socioeconomic status (SES), race/ethnicity, antidepressant use during pregnancy, smoking, substance abuse, previous preterm birth, or obstetric/medical complications.

We therefore conducted a meta-analysis of all available studies to quantify the strength of the relationships between depression during pregnancy and PTB, LBW, and IUGR and examined potential moderators of negative birth outcomes, such as categorical vs continuous measurement of antenatal depression, race or SES of the sample, and country location of the study (ie, whether the study was from the United States, a developing country, or a European social democracy). A social democracy was defined broadly as a democratic welfare state that incorporates both capitalist and socialist practices and provides universal access to health care.^{39,40} We expected that women with antenatal depression who lived in developing countries, relative to their peers in the United States or social democracies, would show greater disparities in the likelihood of PTB, LBW, and IUGR because of their more limited access to adequate prenatal, health, and mental health care.⁴¹ Similarly, we expected that socioeconomically disadvantaged, depressed, pregnant women within the United States would show a higher probability of negative birth outcomes compared with their middle-or upper-class counterparts because they have less access to ongoing adequate

health and mental health services.^{42–44} Finally, given that categorical measures of antenatal depression more closely approximate clinical diagnoses of major depression than do continuous measures, we hypothesized that the former would show a stronger association with adverse birth outcomes.

METHODS

The methods for conducting and reporting the meta-analysis followed state-of-the-art guidelines.^{45–47}

SEARCH STRATEGY AND STUDY SELECTION

Study investigators (N.K.G., A.R.G., J.L.M., and W.J.K.) retrieved potential studies based on a literature search of English-language and non-English-language articles from January 1980 through December 2009 using the MEDLINE, PsycINFO, CINAHL, Social Work Abstracts, Social Services Abstracts, and Dissertation Abstracts International databases. The time frame ensured that the applied standards for the categorical measures of depression were consistent with the *Diagnostic and Statistical Manual of Mental Disorders* (Third Edition)⁴⁸ or later criteria. We used the following keywords and their combinations: *depression, depressive symptoms, pregnancy, prematurity, antenatal, prenatal, birthweight, birth weight, preterm birth, gestational age, fetal growth restriction, intrauterine growth restriction, and small-for-gestational age*. Relevant articles were also identified through references of retrieved articles and contact with prominent investigators in the field.

Published and unpublished English-language and non-English-language observational studies were included in the meta-analysis if they assessed depressive symptoms or unipolar depression diagnoses by means of a depression-screening questionnaire or structured psychiatric interview at 1 or more times during pregnancy. Studies were included if they reported sufficient data to calculate an effect size between depressive symptoms/diagnoses and at least 1 adverse birth outcome: PTB, LBW, or IUGR. Studies were excluded if they used a retrospective design to measure antenatal depression, did not use a prospective or longitudinal design, combined unipolar and bipolar depression diagnoses to measure antenatal depression,⁴⁹ or reported the same data on antenatal depression and PTB, LBW, or IUGR from a previous article. Of 862 reviewed studies, 29 published studies met the inclusion criteria.

DATA EXTRACTION

Adverse Birth Outcomes—Two investigators (one of whom was N.K.G.) reviewed all the studies. Standardized data collection forms were developed a priori for data extraction.⁴⁷ In cases of disagreement in coding, we reached agreement through consensus. The mean percentage of disagreement in coding across all 29 studies included in the meta-analysis was 2.1% (95% confidence interval [CI], 1.6%–2.6%). When a report did not contain sufficient data to calculate an effect size, we contacted the primary author up to 3 times to obtain this information. Six of 9 authors contacted provided the requested data. The following types of studies^{31–38,50–70}(Table 1) were included in the meta-analysis: PTB (n=20), LBW (n=11), and IUGR (n=12). The latter included studies examining the keyword topics *fetal growth restriction* or *small-for-gestational age*. We extracted information from each study pertinent to each adverse outcome using the authors' definition of clinical significance. Typically, PTB was defined as less than 37 weeks' gestation, LBW was defined as less than 2500 g, and IUGR was defined as a fetal weight lower than the 10th percentile for gestational age as determined through an ultrasound.^{80,81} Alternatively, IUGR was defined in some studies as LBW controlling for gestational age,^{56,60,64} a fetal weight lower than the 15th percentile for gestational age,⁷⁰ or a fetal weight lower than the 10th customized percentile.⁶³

Study Characteristics and Antenatal Depression Measures—Information extracted from each study included year of publication, mean maternal age, mean gestational age at first depression assessment, timing and frequency of antenatal depression measurement, country location, country rating of inequality of income distribution (ie, Gini coefficient⁸²), sample size, and the predominant (ie, >60% of the participants) race/ethnicity, SES, parity, marital status, educational level, and work status of the sample. We also recorded the type of antenatal depression measure used (ie, depression-screening questionnaire or structured psychiatric interview).

Other Biopsychosocial Risk Factors for Adverse Birth Outcomes—We coded the extent to which each study controlled for a group of variables observed in the literature to be risk factors for each adverse birth outcome, including demographic variables such as maternal age, SES, parity, race/ethnicity, educational level, marital status, work status, and sex of the infant; psychiatric variables such as antenatal anxiety, stress, drug/alcohol use, antidepressant medication use, and smoking; and obstetric/medical variables such as history of PTB or LBW, current gestational diabetes or preeclampsia, prepregnancy maternal weight or body mass index, and gestational age. We designated the following as key control variables because they have shown the strongest and most consistent associations with adverse birth outcomes in the literature: (1) smoking or substance abuse,^{10,11,83–85} (2) race/ethnicity or SES,^{42,60,86} (3) previous PTB,^{87–89} and (4) selective serotonin reuptake inhibitor (SSRI) antidepressant use.^{90–94}

Methodologic Quality Assessment—Two of the investigators (one of whom was N.K.G.) rated each study on 6 components of methodologic quality, which we developed by modifying the instrument by Downs and Black⁹⁵ for randomized controlled trials and observational studies. We used a consensus approach in which any differences were resolved before assigning a final rating (intraclass correlation coefficient, 0.97; 95% CI, 0.93–0.98). The 6 components evaluated (1) the size of the sample, (2) the representativeness of the sample, (3) whether the sample was clearly described, (4) the reliability and validity of the measure used to assess antenatal depression, (5) whether the statistical tests were appropriate and controlled for the key variables described previously, and (6) whether the study response rate (ie, those who declined to enter the study) and attrition rate (ie, those who entered the study but dropped out) were reported and taken into account statistically. The 3 levels of quality for each component (not adequate, somewhat adequate, and adequate) received equal weights in scoring. A composite quality score was created for each study (Table 1), which was a sum of the number of the 6 components rated (total score range, 0–12).

DATA ANALYSIS

The association of antenatal depressive symptoms or diagnoses with each adverse birth outcome was examined using relative risks (RRs). To do this, we considered odds ratios (ORs) as surrogates for RRs because when outcomes undergoing study are relatively uncommon, the relative odds approximate RRs. One study⁵⁵ used the correlation coefficient as the measure of effect size between antenatal depression and LBW; in this case, we computed the RR by means of transformation from the Pear-son correlation to the standardized mean difference and then from this difference to the log OR.⁹⁶ Two studies of PTB^{60,62} reported hazard ratios as a measure of association between antenatal depression and time to delivery. Because hazard ratios provide a control for calendar time, we consider an effect size derived from Cox regression as a measure of RR.⁹⁷ For each birth outcome, there were a sufficient number of studies to calculate an effect size with a corresponding 95% CI.⁹⁸ We weighted the study-specific RR by the inverse of its variance to compute a pooled RR using random-effects models. A 2-tailed $P < .05$ was used to determine statistical

significance. Statistical analyses were performed with Comprehensive Meta-analysis version 2.2 (Biostat, Englewood, New Jersey) and SPSS version 17.0 (SPSS Inc, Chicago, Illinois) statistical software.

Heterogeneity of effect size was assessed using the Cochran Q χ^2 statistic ($P \leq .10$) and the I^2 statistic (a transformation of the Cochran Q that indicates the percentage of variation in the effect size estimate attributable to heterogeneity rather than sampling error).⁹⁹ A nonsignificant χ^2 statistic suggests that the obtained pooled RR represents a unitary effect and that any variability in effect sizes is caused by random error rather than the influence of other potential moderator variables. For outcomes in which the test of homogeneity of effect sizes was significant, random-effects meta-regression analyses and moderator analyses were conducted to determine whether 6 study characteristics could explain variability across studies: (1) country location: United States, developing country, or social democracy; (2) the country rating of inequality of income distribution (ie, Gini coefficient); (3) sample SES; (4) sample race (eg, black or white), controlling for SES; (5) study methodologic quality; and (6) categorical vs continuous antenatal depression measurement. Sensitivity analyses¹⁰⁰ (known as “leave-one-out”) were conducted by iteratively deleting each study and calculating the resulting effect sizes.

We followed a group of a priori decision rules for pooling data from each study. First, we used the most typical cutoff value for the validated depression scale in each study. Second, when the cutoff score was trichotomized, we used the typical cutoff value to determine the mean of the scores for the medium- and high-risk groups to comprise the group with depression. Third, when a study examined a depression-only group and a depression plus antidepressant medication group, we pooled the effect sizes for the 2 groups in the primary analysis. We then conducted sensitivity analyses¹⁰⁰ for the studies that stratified for antidepressant medication use during pregnancy to compare the birth outcomes for depressed women treated and not treated with antidepressants. Fourth, for studies that measured depression more than once at different times or trimesters during pregnancy, we used the mean of the effect sizes in the primary analyses. Fifth, we used both categorical and continuous measures of antenatal depression in the primary analyses and then conducted moderator analyses based on the categorical-continuous distinction. Sixth, in the primary analyses we included 28 studies that used a validated measure of antenatal depression and an additional study that did not report validity or reliability data for its antenatal depression measure.⁶⁹ We also conducted sensitivity analyses comparing the birth outcomes for the 28 studies using validated measures with the outcomes for the 29 studies used in the primary analyses.

Publication bias was assessed visually using a funnel plot and quantitatively using an adjusted rank correlation test¹⁰¹ and a regression procedure to measure funnel plot asymmetry.¹⁰² The trim-and-fill method by Duval and Tweedie^{103,104} was used to adjust for potential publication bias. The trim-and-fill method assesses asymmetry in the funnel plot, imputes the number of suspected missing studies, and recalculates the adjusted effect size estimate. The adjusted result can be used as a sensitivity analysis to indicate the extent to which publication bias may affect the pooled estimate.¹⁰⁵

RESULTS

The study retrieval and selection strategy is illustrated in Figure 1. Of 862 citations meeting initial search criteria, 54 articles were retrieved and 3 were identified from the references of the retrieved articles, making a total of 57. Of these, 28 studies were excluded (23 studies met at least 1 of the exclusion criteria, and 5 studies represented duplicate studies in which the same data were reported in a previous article), leaving a total of 29 articles included in

the meta-analysis. Table 1 gives the characteristics of the studies included in the meta-analysis.

ANTENATAL DEPRESSION AND RISK OF ADVERSE BIRTH OUTCOMES

Preterm Birth—Twenty studies evaluated the association between antenatal depression and PTB, with RRs ranging from 1.01 to 4.90 (Table 2). Eleven of the studies found no significant association. Using the random-effects model, depression during pregnancy was significantly associated with PTB (RR=1.13; 95% CI, 1.06–1.21). Significant heterogeneity across studies was noted ($Q_{19} = 49.0$; $P < .001$; $I^2 = 61\%$).

Low Birth Weight—Eleven studies evaluated the association between antenatal depression and LBW with RRs ranging from 1.02 to 4.75 (Table 2). Six of the studies found no significant association. The random-effects meta-analysis showed that antenatal depression was significantly associated with LBW (RR=1.18; 95% CI, 1.07–1.30). Significant heterogeneity across studies was found ($Q_{10} = 33.8$; $P < .001$; $I^2 = 70\%$).

Intrauterine Growth Restriction—Twelve studies evaluated the association between antenatal depression and IUGR, with RRs ranging from 0.69 to 11.28 (Table 2). Only 2 studies reported a significant association. The summary RR calculated from the random-effects model showed that antenatal depression was not significantly associated with IUGR (RR=1.03; 95% CI, 0.99–1.08). Significant heterogeneity across studies was noted ($Q_{11} = 22.4$; $P = .02$; $I^2 = 51\%$).

MODERATORS OF OUTCOME

Moderator analyses were conducted to explore sources of heterogeneity (Table 3). As expected, studies of PTB, LBW, and IUGR that used a categorical depression predictor yielded larger ($P < .05$ for all) pooled RRs (1.39 [95% CI, 1.19–1.61], 1.49 [1.25–1.77], and 1.45 [1.05–2.02], respectively) than studies that used a continuous depression predictor (1.03 [1.00–1.06], 1.04 [0.99–1.09], and 1.02 [1.00–1.04], respectively). In PTB trials, heterogeneity among studies was reduced by the addition of the depression predictor moderator (categorical: $Q_{15} = 24.6$; $P = .06$; $I^2 = 39\%$; continuous: $Q_3 = 4.7$; $P = .20$; $I^2 = 36\%$).

As expected, country location (developing nation, social democracy, or United States) was also a significant moderator of the association between antenatal depression and LBW (Table 3 and Figure 2). In developing nations, 2 studies of antenatal depression and LBW yielded a pooled RR of 2.05 (95% CI, 1.43–2.93).^{34,66} In studies from social democracies^{36, 50, 55} and the United States,^{33,35,37,53,67,69} the resulting summary RRs were 1.16 (95% CI, 0.92–1.47) and 1.10 (95% CI, 1.01–1.21), respectively. Significant heterogeneity was still present in US studies ($Q_5 = 18.7$; $P = .002$; $I^2 = 73\%$). A similar effect ($Q_2 = 7.1$; $P = .03$) was found after excluding 2 US studies^{35,37} that used a continuous depression predictor, which yielded a pooled RR of 1.50 (95% CI, 1.22–1.84), and that eliminated heterogeneity across US trials ($P > .45$; $I^2 = 0\%$). Table 3 reveals a similar pattern for studies evaluating the association between antenatal depression and IUGR, with a significant pairwise difference in RR between developing countries and social democracies ($P = .048$) and a near significant pairwise difference in RR between developing countries and the United States ($P = .05$). These data indicate that depressed pregnant women in the developing world are twice as likely to experience IUGR as their European or US counterparts.

Limiting the analysis to the 10 US studies that used a categorically defined depression predictor showed a trend ($Q_1 = 3.54$; $P = .06$) for antenatal depression to be associated with an increased risk of PTB among women of lower SES (RR=1.69; 95% CI, 1.14–2.50) but not in women of middle- or upper-income status (RR=1.13; 95% CI, 0.99–1.29).

Inequality of income across countries, study quality, type of depression severity measurement, and use of adjusted vs unadjusted effect size estimates were not significant moderators of PTB, LBW, or IUGR. Race was not a significant moderator of PTB in US studies of predominantly low SES populations. Type of IUGR measurement was not a significant moderator of outcome in IUGR studies. In the 5 studies of PTB^{31,57,62,67,68} for which stratification by antidepressant medication treatment during pregnancy was possible, the summary RR was comparable for depressed women treated and not treated with antidepressants.

Leave-One-Out Analyses—Sensitivity analyses revealed that no single study unduly influenced the pooled RR estimates of the association between antenatal depression and PTB, LBW, and IUGR (data available on request). In particular, leaving out the 1 study that did not report validity or reliability data for its measure of antenatal depression did not significantly alter the findings observed for LBW in the primary analyses.

Publication Bias—For all 3 outcome measures (PTB, LBW, and IUGR), visual inspection of funnel plots in which each study's effect size was plotted against the standard error showed marked asymmetry, suggesting that small studies with negative results may not have been published. Formal testing using the regression intercept approach¹⁰² confirmed the possibility of publication bias for PTB ($P < .001$), LBW ($P = .001$), and IUGR ($P = .008$). As indicated in Table 4, the trim-and-fill adjusted RRs for each adverse birth outcome are generally lower than the unadjusted RRs. More important, however, the estimates of RR for categorically defined antenatal depression and PTB and LBW were robust to the effects of publication bias.

COMMENT

This meta-analysis showed that depression during pregnancy, regardless of the type of antenatal depression measurement (ie, categorical or continuous), is associated with modest but statistically significant risks of PTB and LBW. Furthermore, the estimates of risk for PTB and LBW from categorically defined antenatal depression appear resilient to the effects of publication bias. Sensitivity analyses showed that the magnitude of RR for these adverse birth outcomes was consistent within the set of 20 studies examining PTB and the set of 11 studies investigating LBW. Although the sizes of the significant RRs for PTB and LBW posed by antenatal depression are modest, the test of relevance in our study is not statistical significance but public health significance. That is, given the prevalence of antenatal depression in a population of pregnant women, what will be the likely burden of PTB or LBW for their infants? Thus, a relatively small effect size magnified by a large population base can have a considerable and noteworthy effect on public health.

We also found evidence that the type of depression measurement (categorical vs continuous) moderated the strength of the associations between antenatal depression and PTB, LBW, and IUGR, thereby eliminating or reducing the heterogeneity associated with these findings. Results for categorical measures of antenatal depression revealed that having major depression or clinically significant depressive symptoms significantly increased the RR of PTB by 39%, the risk of LBW by 49%, and the risk of IUGR by 45%. As expected, continuous measures of antenatal depression showed a similar albeit weaker pattern, indicating that every 1-point increase in depression severity was associated with a 3% significantly increased risk of PTB and nonsignificantly increased risks of LBW (4%) and IUGR (2%). To place the categorical results for antenatal depression in context, we note that smoking was observed to have a dose-dependent relationship with PTB,^{106,107} with smoking more than 10 cigarettes a day shown to increase the likelihood of PTB between 33 and 36 weeks by 40% and of PTB at 32 weeks or less by 60%. Furthermore, in a cohort

analysis of a large, ethnically diverse population,¹¹ substance use disorders were associated with a 2.4-fold higher risk of PTB and a 3.7-fold higher risk of LBW, and black race increased the likelihood of PTB by 60% and of LBW by 2-fold. Thus, the magnitude of risk for PTB and LBW posed by antenatal depression is comparable to the risk of smoking 10 or more cigarettes a day for PTB but is relatively modest contrasted to the greater risks of black race and substance abuse associated with PTB and LBW.

Moderator analyses for country location revealed that the RR of delivering an infant with LBW or IUGR was higher among women from developing countries who experienced antenatal depression than their counterparts in the United States or social democracies. Moreover, in US studies, categorically defined antenatal depression tended to be associated with an elevated risk of PTB in women of predominantly lower SES but not in women of middle- or upper-income status. Pregnant women of lower SES in the United States are also twice as likely to experience antenatal major and minor depression as are women from middle- to upper-income strata.^{20–22} Whereas cross-cultural variability in the prevalence of perinatal depression certainly exists,²⁹ estimates of the prevalence of perinatal depression in several developing countries^{27,28} are similar to the higher depression rates in pregnant US women of lower SES. Thus, many socioeconomically disadvantaged childbearing women in developing countries and in the United States experience a double-barreled threat: an increased risk of becoming depressed during their pregnancies and an increased likelihood of experiencing adverse birth outcomes once they have antenatal depression. Depressed pregnant women living near or below poverty levels are subject to large amounts of acute and chronic stress, such as living in unsafe neighborhoods, experiencing racial/ethnic or economic discrimination, and confronting food inadequacy in their households.^{22,108,109} At the same time, despite Medicaid or other public insurance coverage during pregnancy, their mental health problems are seldom accurately diagnosed and they often lack access to specialty mental health services.^{110–112}

Several potential direct and indirect causal pathways through which antenatal depression leads to adverse pregnancy outcomes have been proposed. One possibility is that prenatal stress or depression during pregnancy might promote adverse birth outcomes through the dysregulation of the hypothalamic-pituitary-adrenocortical axis, stimulating the release of stress hormones, such as cortisol and catecholamines. These biological changes may result in placental hypofusion and consequent restriction of oxygen and nutrients to the fetus, leading to fetal growth restriction and/or precipitation of PTB.^{113–117} Other mechanisms include the possibility that antenatal depression might compromise immune system functioning,¹¹⁸ which in turn may lead to a reproductive tract infection triggering PTB.¹¹⁷ The harmful public health effect of antenatal depression on birth outcomes is further heightened by evidence that depression during pregnancy is associated with risky but modifiable health practices, such as poor nutrition and hygiene, lack of motivation to obtain prenatal care or to follow medical recommendations, and smoking and/or alcohol and substance abuse, all of which adversely affect pregnancy outcomes.^{11,33,85,119}

Clearly, pregnancy is an important time to universally screen women for depression, especially those who are socioeconomically disadvantaged, and to improve their timely access to evidence-based prenatal and mental health services.¹²⁰ Improved accuracy of diagnosis and treatment of antenatal depression combined with education about harmful but potentially modifiable lifestyle practices could lead to decreased rates of PTB and LBW.

Reducing the rates of these adverse birth outcomes is a critically important public health issue. During childhood, PTB is associated with an increased risk of mortality³; adverse medical outcomes^{4,121–123} including respiratory distress syndrome, cerebral palsy, chronic lung disease, vision and hearing loss, and neurodevelopmental disabilities; cognitive

difficulties¹²⁴, and psychiatric problems,¹²⁴ such as internalizing and externalizing behaviors and attention-deficit/hyperactivity disorder. Long-term consequences of PTB in adulthood include diminished rates of reproduction and, for women born preterm, an increased risk of next-generation PTB, fetal stillbirth, and infant mortality.³ Pernicious child and adult outcomes related to LBW^{125–133} and IUGR^{134–139} reflect patterns similar to that of PTB. Furthermore, in the United States, the annual economic costs of medically managing the consequences of these adverse birth outcomes are enormous.^{140,141} For example, approximately 75% of admissions to the neonatal intensive care unit are related to prematurity.¹⁴² Daily neonatal intensive care unit costs in the United States exceed \$3500 per infant, and it is not unusual for costs to reach \$1 million for a prolonged stay.¹⁴²

A strength of our meta-analysis is that our search included US and non-US English-language studies, as well as a study in French that was translated into English by an expert. Thus, the 29 studies included in our meta-analysis came from 12 non-US countries, indicating significant international representation. Our meta-analysis also included studies that varied in the extent to which they controlled for confounding factors related to PTB, LBW, and IUGR. For example, one-third of the studies (n=10) controlled for SSRI use^{31,57,62,67,68} or reported that SSRI use was unlikely in their sample.^{34,36,51,64,66} An increasing number of women with depression are prescribed antidepressant medications during pregnancy.¹⁴³ These medications, especially SSRIs, have been significantly linked with LBW in some^{49,92,93,144} but not all studies.^{145–147} Wisner et al⁶⁸ recently found that infants who were continuously exposed to either SSRIs or major depression throughout the 3 trimesters of pregnancy were more likely to be born preterm than were infants with partial or no exposure. Differentiating between the effects of depression or depressive symptoms and the effects of antidepressants on birth outcomes is challenging because (1) researchers typically investigate the effects of one without controlling adequately for the other; (2) use of antidepressants during pregnancy occurs at different times, dosages, and durations; (3) recognition and treatment of depression by the physician is often associated with depression severity and persistence; and (4) depression is also associated with the use of additional prescription and nonprescription medications¹⁴⁸ and other potential confounders, such as smoking or substance use disorders.

Finally, prior evidence has shown that key variables in addition to depression, such as substance use or abuse, race/ethnicity or SES, and previous PTB, are strongly and consistently predictive of negative birth outcomes. Most of the studies in our meta-analysis (80%) controlled for at least 2 of these key predictors, but few controlled for all of them. In addition, most of the studies did not control for stressful life events and other psychiatric comorbidities of depression, such as antenatal anxiety, which has been linked with adverse birth outcomes in some studies.⁶ A recent meta-analysis of anxiety symptoms and birth outcomes, however, did not show evidence of this association.¹⁴⁹

Only 5 of 29 studies in our meta-analysis assessed major depression during pregnancy by using diagnostic criteria adhering to or compatible with the *Diagnostic and Statistical Manual of Mental Disorders* (Fourth Edition).¹⁵⁰ Most studies used relatively short screening tools to evaluate levels of depressive symptoms and to set cutoff scores for describing clinically significant depressive symptoms. Although categorizing depressive symptom levels may be related to the diagnosis of clinically significant depression, it is not a substitute nor may it be as accurate as a structured interview. Finally, although we found possible publication bias, the findings of an elevated risk of PTB and LBW associated with categorically defined depression remained robust to trim-and-fill analyses that corrected for this bias.

Limitations of the studies reviewed suggest the need for a large prospective epidemiological study to simultaneously evaluate the RRs during pregnancy of depression, SSRI use, smoking, substance use disorders, key sociodemographic variables, obstetric/medical variables, and important behavioral health practices, including engagement in adequate prenatal and medical care and nutrition. Multiple assessments of major and minor antenatal depression should be performed, using criteria from the *Diagnostic and Statistical Manual of Mental Disorders* (Fourth Edition) as well as measures assessing depression severity. The RRs of anxiety disorders and anxiety severity should also be assessed because these psychopathologic conditions typically accompany depression and were found to be related in some, but not all, studies to adverse birth outcomes. Ideally, this future study would prospectively gather data during the prepregnancy, pregnancy, and postpartum periods, considering that a broader perspective on a woman's health status may be necessary to better understand the risk factors associated with harmful birth outcomes.^{58,151,152}

Our overall pattern of findings in this meta-analysis highlights the salient public health risk of PTB and LBW posed by antenatal depression, particularly for socioeconomically disadvantaged women in developing countries and in the United States. Furthermore, mounting evidence from this meta-analysis and other sources⁶⁸ suggests that untreated major depression during pregnancy is as likely to lead to poor birth outcomes as is treatment with SSRIs. An important implication of these findings is that pregnant women should be universally screened for depression and provided guideline-level treatment before childbirth. Given that untreated antenatal depression is the most robust predictor of postpartum depression and has additional serious adverse consequences for infant and child development beyond harmful birth outcomes, women and their obstetrics professionals will need to weigh the costs and benefits of treating antenatal depression pharmacologically, especially when treatment with evidence-based psychotherapy is not available or desired.

Acknowledgments

Funding/Support: Dr Grote's role in this study was supported by grant K23-MH 67595 from the National Institute of Mental Health (NIMH), National Institutes of Health (NIH). Dr Bridge's role was supported by grant K01-MH069948 from the NIMH. Dr Gavin's role was supported by grant 1KL2RR025015-01 from the National Center for Research Resources, a component of the NIH and NIH Roadmap for Medical Research. Dr Melville's role was supported by grant K23 MH070704 from the NIMH. Dr Katon's role was supported by grant K24 MH069471 from the NIMH.

Role of the Sponsor: The NIMH had no role in the design and conduct of the study; the collection, analysis, and interpretation of the data; or the preparation, review, or approval of the manuscript.

References

1. March of Dimes. Perinatal Profile: Statistics for Monitoring State Maternal and Infant Health, 1999 & 2000 editions. White Plains, NY: Perinatal Data Center; 2000.
2. World Health Organization. Physical Status: The Use and Interpretation of Anthropometry. Geneva, Switzerland: World Health Organization; 1995. WHO Technical Report Series 854
3. Swamy GK, Ostbye T, Skjaerven R. Association of preterm birth with long-term survival, reproduction, and next-generation preterm birth. *JAMA* 2008;299(12):1429–1436. [PubMed: 18364485]
4. Allen MC, Jones MD Jr. Medical complications of prematurity. *Obstet Gynecol* 1986;67(3):427–437. [PubMed: 2935764]
5. Wilson-Costello D, Friedman H, Minich N, Fanaroff AA, Hack M. Improved survival rates with increased neurodevelopmental disability for extremely low birth weight infants in the 1990s. *Pediatrics* 2005;115(4):997–1003. [PubMed: 15805376]

6. Alder J, Fink N, Bitzer J, Hösli I, Holzgreve W. Depression and anxiety during pregnancy: a risk factor for obstetric, fetal and neonatal outcome? a critical review of the literature. *J Matern Fetal Neonatal Med* 2007;20(3):189–209. [PubMed: 17437220]
7. Bonari L, Pinto N, Ahn E, Einarson A, Steiner M, Koren G. Perinatal risks of untreated depression during pregnancy. *Can J Psychiatry* 2004;49(11):726–735. [PubMed: 15633850]
8. Kramer MS. Determinants of low birth weight: methodological assessment and meta-analysis. *Bull World Health Organ* 1987;65(5):663–737. [PubMed: 3322602]
9. Kyrklund-Blomberg NB, Cnattingius S. Preterm birth and maternal smoking: risks related to gestational age and onset of delivery. *Am J Obstet Gynecol* 1998;179(4):1051–1055. [PubMed: 9790397]
10. Ellard GA, Johnstone FD, Prescott RJ, Ji-Xian W, Jian-Hua M. Smoking during pregnancy: the dose dependence of birthweight deficits. *Br J Obstet Gynaecol* 1996;103(8):806–813. [PubMed: 8760712]
11. Kelly RH, Russo J, Holt VL, Danielsen BH, Zatzick DF, Walker E, Katon W. Psychiatric and substance use disorders as risk factors for low birth weight and preterm delivery. *Obstet Gynecol* 2002;100(2):297–304. [PubMed: 12151153]
12. Haelterman E, Bréart G, Paris-Llado J, Dramaix M, Tchobroutsky C. Effect of uncomplicated chronic hypertension on the risk of small-for-gestational age birth. *Am J Epidemiol* 1997;145(8):689–695. [PubMed: 9125995]
13. Sibai BM, Caritis SN, Hauth JC, MacPherson C, VanDorsten JP, Klebanoff M, Landon M, Paul RH, Meis PJ, Miodovnik M, Dombrowski MP, Thurnau GR, Moawad AH, Roberts J. National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Preterm delivery in women with pregestational diabetes mellitus or chronic hypertension relative to women with uncomplicated pregnancies. *Am J Obstet Gynecol* 2000;183(6):1520–1524. [PubMed: 11120521]
14. Kurki T, Hiilesmaa V, Raitasalo R, Mattila H, Ylikorkala O. Depression and anxiety in early pregnancy and risk for preeclampsia. *Obstet Gynecol* 2000;95 (4):487–490. [PubMed: 10725477]
15. Xiong X, Mayes D, Demianczuk N, Olson DM, Davidge ST, Newburn-Cook C, Saunders LD. Impact of pregnancy-induced hypertension on fetal growth. *Am J Obstet Gynecol* 1999;180(1 pt 1):207–213. [PubMed: 9914605]
16. Kozhimannil KB, Pereira MA, Harlow BL. Association between diabetes and perinatal depression among low-income mothers. *JAMA* 2009;301(8):842–847. [PubMed: 19244191]
17. Evers IM, de Valk HW, Visser GH. Risk of complications of pregnancy in women with type 1 diabetes: nationwide prospective study in the Netherlands. *BMJ* 2004;328(7445):915. [published online ahead of print April 5, 2004]. 10.1136/bmj.38043.583160.EE. [PubMed: 15066886]
18. Vesga-López O, Blanco C, Keyes K, Olfson M, Grant BF, Hasin DS. Psychiatric disorders in pregnant and postpartum women in the United States. *Arch Gen Psychiatry* 2008;65(7):805–815. [PubMed: 18606953]
19. Gavin NI, Gaynes BN, Lohr KN, Meltzer-Brody S, Gartlehner G, Swinson T. Perinatal depression: a systematic review of prevalence and incidence. *Obstet Gynecol* 2005;106(5 pt 1):1071–1083. [PubMed: 16260528]
20. Hobfoll SE, Ritter C, Lavin J, Hulsizer MR, Cameron RP. Depression prevalence and incidence among inner-city pregnant and postpartum women. *J Consult Clin Psychol* 1995;63(3):445–453. [PubMed: 7608357]
21. Scholle SH, Haskett RF, Hanusa BH, Pincus HA, Kupfer DJ. Addressing depression in obstetrics/gynecology practice. *Gen Hosp Psychiatry* 2003;25(2):83–90. [PubMed: 12676420]
22. Siefert K, Bowman PJ, Heflin CM, Danziger S, Williams DR. Social and environmental predictors of maternal depression in current and recent welfare recipients. *Am J Orthopsychiatry* 2000;70(4):510–522. [PubMed: 11086529]
23. Gotlib IH, Whiffen VE, Wallace PM, Mount JH. Prospective investigation of postpartum depression: factors involved in onset and recovery. *J Abnorm Psychol* 1991;100(2):122–132. [PubMed: 2040762]
24. O'Hara MW, Neunaber DJ, Zekoski EM. Prospective study of postpartum depression: prevalence, course, and predictive factors. *J Abnorm Psychol* 1984;93(2):158–171. [PubMed: 6725749]

25. O'Hara MW, Swain AM. Rates and risk of postpartum depression: a meta-analysis. *Int Rev Psychiatry* 1996;8(1):18–37.
26. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS. Lifetime and 12-month prevalence of *DSM-III-R* psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry* 1994;51(1):8–19. [PubMed: 8279933]
27. Patel V, Rodrigues M, DeSouza N. Gender, poverty, and postnatal depression: a study of mothers in Goa, India. *Am J Psychiatry* 2002;159(1):43–47. [PubMed: 11772688]
28. Rahman A, Iqbal Z, Harrington R. Life events, social support and depression in childbirth: perspectives from a rural community in the developing world. *Psychol Med* 2003;33(7):1161–1167. [PubMed: 14580070]
29. Halbreich U, Karkun S. Cross-cultural and social diversity of prevalence of postpartum depression and depressive symptoms [published online ahead of print February 7, 2006]. *J Affect Disord* 2006;91(2–3):97–111.10.1016/j.jad.2005.12.051 [PubMed: 16466664]
30. Yonkers K, Wisner K, Stewart D, Oberlander TF, Dell DL, Stotland N, Ramin S, Chaudron L, Lockwood C. Management of depression during pregnancy: a report from the American Psychiatric Association and the American College of Obstetricians and Gynecologists. *Gen Hosp Psychiatry* 2009;31(5):403–413. [PubMed: 19703633]
31. Dayan J, Creveuil C, Marks MN, Conroy S, Herlicoviez M, Dreyfus M, Tordjman S. Prenatal depression, prenatal anxiety, and spontaneous preterm birth: a prospective cohort study among women with early and regular care. *Psychosom Med* 2006;68(6):938–946. [PubMed: 17079701]
32. Orr ST, James SA, Blackmore Prince C. Maternal prenatal depressive symptoms and spontaneous preterm births among African-American women in Baltimore, Maryland. *Am J Epidemiol* 2002;156(9):797–802. [PubMed: 12396996]
33. Neggers Y, Goldenberg R, Cliver S, Hauth J. The relationship between psychosocial profile, health practices, and pregnancy outcomes. *Acta Obstet Gynecol Scand* 2006;85(3):277–285. [PubMed: 16553174]
34. Rahman A, Iqbal Z, Bunn J, Lovel H, Harrington R. Impact of maternal depression on infant nutritional status and illness: a cohort study. *Arch Gen Psychiatry* 2004;61(9):946–952. [PubMed: 15351773]
35. Steer RA, Scholl TO, Hediger ML, Fischer RL. Self-reported depression and negative pregnancy outcomes. *J Clin Epidemiol* 1992;45(10):1093–1099. [PubMed: 1474405]
36. Andersson L, Sundström-Poromaa I, Wulff M, Aström M, Bixo M. Neonatal outcome following maternal antenatal depression and anxiety: a population-based study. *Am J Epidemiol* 2004;159(9):872–881. [PubMed: 15105180]
37. Copper RL, Goldenberg RL, Das A, Elder N, Swain M, Norman G, Ramsey R, Cotroneo P, Collins BA, Johnson F, Jones P, Meier AM. National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. The preterm prediction study: maternal stress is associated with spontaneous preterm birth at less than thirty-five weeks' gestation. *Am J Obstet Gynecol* 1996;175(5):1286–1292. [PubMed: 8942502]
38. Nordentoft M, Lou HC, Hansen D, Nim J, Pryds O, Rubin P, Hemmingsen R. Intrauterine growth retardation and premature delivery: the influence of maternal smoking and psychosocial factors. *Am J Public Health* 1996;86(3):347–354. [PubMed: 8604759]
39. Huber, E.; Stephens, JD. *Development and Crisis of the Welfare State: Parties and Policies in Global Markets*. Chicago, Illinois: University of Chicago Press; 2001.
40. Esping-Andersen, G. *Welfare States in Transition: National Adaptations in Global Economies*. Thousand Oaks, California: Sage Publications; 1996.
41. Prince M, Patel V, Saxena S, Maj M, Maselko J, Phillips MR, Rahman A. No health without mental health. *Lancet* 2007;370(9590):859–877. [PubMed: 17804063]
42. Blackmore-Prince C, Kieke B Jr, Kugaraj KA, Ferré C, Elam-Evans LD, Krule-witch CJ, Gaudino JA, Overpeck M. Racial differences in the patterns of singleton preterm delivery in the 1988 National Maternal and Infant Health Survey. *Matern Child Health J* 1999;3(4):189–197. [PubMed: 10791359]

43. Wang PS, Lane M, Olfson M, Pincus HA, Wells KB, Kessler RC. Twelve-month use of mental health services in the United States: results from the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005;62(6):629–640. [PubMed: 15939840]
44. Murray CJ, Kulkarni S, Ezzati M. Eight Americas: new perspectives on US health disparities. *Am J Prev Med* 2005;29(5 suppl 1):4–10. [PubMed: 16389119]
45. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA, Thacker SB. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA* 2000;283(15):2008–2012. [PubMed: 10789670]
46. Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF. the QUORUM Group. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. *Lancet* 1999;354(9193):1896–1900. [PubMed: 10584742]
47. Lipsey, MW.; Wilson, DB. *Practical Meta-analysis*. Thousand Oaks, California: Sage Publications; 2001.
48. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 3. Washington, DC: American Psychiatric Association; 1980.
49. Oberlander TF, Warburton W, Misri S, Aghajanian J, Hertzman C. Neonatal outcomes after prenatal exposure to selective serotonin reuptake inhibitor antidepressants and maternal depression using population-based linked health data. *Arch Gen Psychiatry* 2006;63(8):898–906. [PubMed: 16894066]
50. Berle JO, Mykletun A, Daltveit AK, Rasmussen S, Holsten F, Dahl AA. Neonatal outcomes in offspring of women with anxiety and depression during pregnancy: a linkage study from The Nord-Trøndelag Health Study (HUNT) and Medical Birth Registry of Norway. *Arch Womens Ment Health* 2005;8(3):181–189. [PubMed: 15959623]
51. Chung TK, Lau TK, Yip AS, Chiu HF, Lee DT. Antepartum depressive symptomatology is associated with adverse obstetric and neonatal outcomes. *Psychosom Med* 2001;63(5):830–834. [PubMed: 11573032]
52. Dayan J, Creveuil C, Herlicoviez M, Herbel C, Baranger E. Antenatal depression, a risk factor for prenatal delivery [in French]. *Presse Med* 1999;28 (31):1698. [PubMed: 10554610]
53. Diego MA, Field T, Hernandez-Reif M, Schanberg S, Kuhn C, Gonzalez-Quintero VH. Prenatal depression restricts fetal growth. *Early Hum Dev* 2009;85(1):65–70. [PubMed: 18723301]
54. Dole N, Savitz DA, Hertz-Picciotto I, Siega-Riz AM, McMahon MJ, Buekens P. Maternal stress and preterm birth. *Am J Epidemiol* 2003;157(1):14–24. [PubMed: 12505886]
55. Elsenbruch S, Benson S, Rütcke M, Rose M, Dudenhausen J, Pincus-Knackstedt MK, Klapp BF, Arck PC. Social support during pregnancy: effects on maternal depressive symptoms, smoking and pregnancy outcome [published online ahead of print November 16, 2006]. *Hum Reprod* 2007;22(3):869–877.10.1093/humrep/del432 [PubMed: 17110400]
56. Evans J, Heron J, Patel RR, Wiles N. Depressive symptoms during pregnancy and low birth weight at term: longitudinal study. *Br J Psychiatry* 2007;191:84–85. [PubMed: 17602131]
57. Gavin AR, Holzman C, Siefert K, Tian Y. Maternal depressive symptoms, depression, and psychiatric medication use in relation to risk of preterm delivery. *Womens Health Issues* 2009;19(5):325–334. [PubMed: 19733802]
58. Haas JS, Fuentes-Afflick E, Stewart AL, Jackson RA, Dean ML, Brawarsky P, Escobar GJ. Prepregnancy health status and the risk of preterm delivery. *Arch Pediatr Adolesc Med* 2005;159(1):58–63. [PubMed: 15630059]
59. Hedegaard M, Henriksen TB, Sabroe S, Secher NJ. Psychological distress in pregnancy and preterm delivery. *BMJ* 1993;307(6898):234–239. [PubMed: 8369684]
60. Hoffman S, Hatch MC. Depressive symptomatology during pregnancy: evidence for an association with decreased fetal growth in pregnancies of lower social class women. *Health Psychol* 2000;19(6):535–543. [PubMed: 11129356]
61. Jesse DE, Seaver W, Wallace DC. Maternal psychosocial risks predict preterm birth in a group of women from Appalachia. *Midwifery* 2003;19(3):191–202. [PubMed: 12946335]

62. Li D, Liu L, Odouli R. Presence of depressive symptoms during early pregnancy and the risk of preterm delivery: a prospective cohort study. *Hum Reprod* 2009;24(1):146–153. [PubMed: 18948314]
63. Paarlberg KM, Vingerhoets AJ, Passchier J, Dekker GA, Heinen AG, van Geijn HP. Psychosocial predictors of low birthweight: a prospective study. *Br J Obstet Gynaecol* 1999;106(8):834–841. [PubMed: 10453835]
64. Patel V, Prince M. Maternal psychological morbidity and low birth weight in India. *Br J Psychiatry* 2006;188:284–285. [PubMed: 16507972]
65. Perkin MR, Bland JM, Peacock JL, Anderson HR. The effect of anxiety and depression during pregnancy on obstetric complications. *Br J Obstet Gynaecol* 1993;100(7):629–634. [PubMed: 8369244]
66. Rondó PH, Ferreira RF, Nogueira F, Ribeiro MC, Lobert H, Artes R. Maternal psychological stress and distress as predictors of low birth weight, prematurity and intrauterine growth retardation. *Eur J Clin Nutr* 2003;57(2):266–272. [PubMed: 12571658]
67. Suri R, Altshuler L, Hellemann G, Burt VK, Aquino A, Mintz J. Effects of antenatal depression and antidepressant treatment on gestational age at birth and risk of preterm birth. *Am J Psychiatry* 2007;164(8):1206–1213. [PubMed: 17671283]
68. Wisner KL, Sit DK, Hanusa BH, Moses-Kolko EL, Bogen DL, Hunker DF, Perel JM, Jones-Ivy S, Bodnar LM, Singer LT. Major depression and antidepressant treatment: impact on pregnancy and neonatal outcomes. *Am J Psychiatry* 2009;166(5):557–566. [PubMed: 19289451]
69. Zimmer-Gembeck MJ, Helfand M. Low birthweight in a public prenatal care program: behavioral and psychosocial risk factors and psychosocial intervention. *Soc Sci Med* 1996;43(2):187–197. [PubMed: 8844923]
70. Goldenberg RL, Cliver SP, Cutter GR, Hoffman HJ, Copper RL, Gotlieb S, Davis RO. Maternal psychosocial characteristics and intrauterine growth retardation. *Pre-Peri-Nat Psychol J* 1991;6(2):129–134.
71. Spitzer RL, Williams JB, Kroenke K, Linzer M, deGruy FV III, Hahn SR, Brody D, Johnson JG. Utility of a new procedure for diagnosing mental disorders in primary care: the PRIME-MD 1000 study. *JAMA* 1994;272(22):1749–1756. [PubMed: 7966923]
72. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale: an updated literature review. *J Psychosom Res* 2002;52(2):69–77. [PubMed: 11832252]
73. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561–571. [PubMed: 13688369]
74. Radloff LS. The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas* 1977;1(3):385–401.
75. Murray D, Cox JL. Screening for depression during pregnancy with the Edinburgh Depression Scale (EPDS). *J Reprod Infant Psychol* 1990;8(2):99–107.
76. First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. Structured Clinical Interview for DSM-IV Axis I Disorders: SCID-I—Clinician Version. Washington, DC: American Psychiatric Press; 1997.
77. Goldberg DP, Hillier VF. A scaled version of the General Health Questionnaire. *Psychol Med* 1979;9(1):139–145. [PubMed: 424481]
78. Luteijn, F.; Hamel, LF.; Bouman, TK.; Kok, AR. HSCL Hopkins Symptom Checklist (Manual). Lisse, the Netherlands: Swets and Zeitlinger; 1984.
79. World Health Organization, Division of Mental Health. SCAN: Schedules for Clinical Assessment in Neuropsychiatry. Geneva, Switzerland: World Health Organization; 1994.
80. Brenner WE, Edelman DA, Hendricks CH. A standard of fetal growth for the United States of America. *Am J Obstet Gynecol* 1976;126(5):555–564. [PubMed: 984126]
81. Robertson, NRC. Textbook of Neonatology. New York, NY: Churchill Livingstone; 1986.
82. United Nations Development Programme. Human Development Report 2006: Beyond Scarcity: Power, Poverty, and the Global Water Crisis: Table 15: Inequality in Income or Expenditure. [Accessed September 1, 2007]. <http://hdr.undp.org/hdr2006/>

83. Cliver SP, Goldenberg RL, Cutter GR, Hoffman HJ, Copper RL, Gotlieb SJ, Davis RO. The relationships among psychosocial profile, maternal size, and smoking in predicting fetal growth retardation. *Obstet Gynecol* 1992;80(2):262–267. [PubMed: 1635741]
84. Lundsberg LS, Bracken MB, Saftlas AF. Low-to-moderate gestational alcohol use and intrauterine growth retardation, low birthweight, and preterm delivery. *Ann Epidemiol* 1997;7(7):498–508. [PubMed: 9349918]
85. Wen SW, Goldenberg RL, Cutter GR, Hoffman HJ, Cliver SP. Intrauterine growth retardation and preterm delivery: prenatal risk factors in an indigent population. *Am J Obstet Gynecol* 1990;162(1):213–218. [PubMed: 2301495]
86. Lieberman E, Ryan KJ, Monson RR, Schoenbaum SC. Risk factors accounting for racial differences in the rate of premature birth. *N Engl J Med* 1987;317(12):743–748. [PubMed: 3627184]
87. Adams MM, Elam-Evans LD, Wilson HG, Gilbertz DA. Rates of and factors associated with recurrence of preterm delivery. *JAMA* 2000;283(12):1591–1596. [PubMed: 10735396]
88. Mercer BM, Goldenberg RL, Moawad AH, Meis PJ, Iams JD, Das AF, Caritis SN, Miodovnik M, Menard MK, Thurnau GR, Dombrowski MP, Roberts JM, McNellis D. National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. The preterm prediction study: effect of gestational age and cause of preterm birth on subsequent obstetric outcome. *Am J Obstet Gynecol* 1999;181(5 pt 1):1216–1221. [PubMed: 10561648]
89. Esplin MS, O'Brien E, Fraser A, Kerber RA, Clark E, Simonsen SE, Holmgren C, Mineau GP, Varner MW. Estimating recurrence of spontaneous preterm delivery. *Obstet Gynecol* 2008;112(3):516–523. [PubMed: 18757647]
90. Davis RL, Rubanowice D, McPhillips H, Raebel MA, Andrade SE, Smith D, Yood MU, Platt R. HMO Research Network Center for Education, Research in Therapeutics. Risks of congenital malformations and perinatal events among infants exposed to antidepressant medications during pregnancy. *Pharmacoepidemiol Drug Saf* 2007;16(10):1086–1094. [PubMed: 17729378]
91. Källén B. Neonate characteristics after maternal use of antidepressants in late pregnancy. *Arch Pediatr Adolesc Med* 2004;158(4):312–316. [PubMed: 15066868]
92. Simon GE, Cunningham ML, Davis RL. Outcomes of prenatal antidepressant exposure. *Am J Psychiatry* 2002;159(12):2055–2061. [PubMed: 12450956]
93. Wen SW, Yang Q, Garner P, Fraser W, Olatunbosun O, Nimrod C, Walker M. Selective serotonin reuptake inhibitors and adverse pregnancy outcomes. *Am J Obstet Gynecol* 2006;194(4):961–966. [PubMed: 16580283]
94. Zeskind PS, Stephens LE. Maternal selective serotonin reuptake inhibitor use during pregnancy and newborn neurobehavior. *Pediatrics* 2004;113(2):368–375. [PubMed: 14754951]
95. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;52(6):377–384. [PubMed: 9764259]
96. Rosenthal, R. Parametric measures of effect size. In: Cooper, H.; Hedges, LV., editors. *The Handbook of Research Synthesis*. New York, New York: Russell Sage Foundation; 1994. p. 231–244.
97. McGettigan P, Henry D. Cardiovascular risk and inhibition of cyclooxygenase: a systematic review of the observational studies of selective and nonselective inhibitors of cyclooxygenase 2. *JAMA* 2006;296(13):1633–1644. [PubMed: 16968831]
98. Rosenthal, R. *Meta-Analytic Procedures for Social Research*. Newbury Park, California: Sage Publications; 1991.
99. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539–1558. [PubMed: 12111919]
100. Greenhouse, JB.; Iyengar, S. Sensitivity analyses and diagnostics. In: Cooper, HM.; Hedges, LV., editors. *The Handbook of Research Synthesis*. New York, New York: Russell Sage Foundation; 1994. p. 383–398.
101. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50(4):1088–1101. [PubMed: 7786990]

102. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315(7109):629–634. [PubMed: 9310563]
103. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56 (2):455–463. [PubMed: 10877304]
104. Sutton AJ, Duval SJ, Tweedie RL, Abrams KR, Jones DR. Empirical assessment of effect of publication bias on meta-analyses. *BMJ* 2000;320(7249):1574–1577. [PubMed: 10845965]
105. Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Performance of the trim and fill method in the presence of publication bias and between-study heterogeneity. *Stat Med* 2007;26(25): 4544–4562. [PubMed: 17476644]
106. Berkowitz GS, Papiernik E. Epidemiology of preterm birth. *Epidemiol Rev* 1993;15(2):414–443. [PubMed: 8174665]
107. Harlow BL, Frigoletto FD, Cramer DW, Evans JK, LeFevre ML, Bain RP, McNellis D. the RADIUS Study Group. Determinants of preterm delivery in low-risk pregnancies. *J Clin Epidemiol* 1996;49(4):441–448. [PubMed: 8621995]
108. Belle D, Doucet J. Poverty, inequality, and discrimination as sources of depression among US women. *Psychol Women Q* 2003;27:101–113.
109. Grote NK, Bledsoe SE, Wellman J, Brown C. Depression in African American and white women with low incomes: the role of chronic stress. *Soc Work Public Health* 2007;23(2–3):59–88. [PubMed: 19306588]
110. Miranda J, Chung JY, Green BL, Krupnick J, Siddique J, Revicki DA, Belin T. Treating depression in predominantly low-income young minority women: a randomized controlled trial. *JAMA* 2003;290(1):57–65. [PubMed: 12837712]
111. Grote NK, Zuckoff A, Swartz H, Bledsoe SE, Geibel S. Engaging women who are depressed and economically disadvantaged in mental health treatment. *Soc Work* 2007;52(4):295–308. [PubMed: 18232240]
112. Kelly R, Zatzick D, Anders T. The detection and treatment of psychiatric disorders and substance use among pregnant women cared for in obstetrics. *Am J Psychiatry* 2001;158(2):213–219. [PubMed: 11156803]
113. Borders AE, Grobman WA, Amsden LB, Holl JL. Chronic stress and low birth weight neonates in a low-income population of women. *Obstet Gynecol* 2007;109(2 pt 1):331–338. [PubMed: 17267833]
114. Lundy BL, Jones NA, Field T, Nearing G, Davalos M, Pietro PA, et al. Prenatal depression effects on neonates. *Infant Behav Dev* 1999;22(1):119–129.
115. Talge NM, Neal C, Glover V. Early Stress, Translational Research and Prevention Science Network: Fetal and Neonatal Experience on Child and Adolescent Mental Health. Antenatal maternal stress and long-term effects on child neurodevelopment: how and why? *J Child Psychol Psychiatry* 2007;48(3–4):245–261. [PubMed: 17355398]
116. van Goozen SHM, Fairchild G, Snoek H, Harold GT. The evidence for a neurobiological model of childhood antisocial behavior. *Psychol Bull* 2007;133 (1):149–182. [PubMed: 17201574]
117. Federenko IS, Wadhwa PD. Women’s mental health during pregnancy influences fetal and infant developmental and health outcomes. *CNS Spectr* 2004;9(3):198–206. [PubMed: 14999160]
118. Herbert TB, Cohen S. Stress and immunity in humans: a meta-analytic review. *Psychosom Med* 1993;55(4):364–379. [PubMed: 8416086]
119. Zuckerman B, Amaro H, Bauchner H, Cabral H. Depressive symptoms during pregnancy: relationship to poor health behaviors. *Am J Obstet Gynecol* 1989;160(5 pt 1):1107–1111. [PubMed: 2729387]
120. Stratton, KR.; Howe, CJ.; Battaglia, FC. *Fetal Alcohol Syndrome: Diagnosis, Epidemiology, Prevention, and Treatment*. Washington, DC: National Academy Press; 1996.
121. Knoches AM, Doyle LW. Long-term outcome of infants born preterm. *Baillieres Clin Obstet Gynaecol* 1993;7(3):633–651. [PubMed: 7504604]
122. Marlow ES, Hunt LP, Marlow N. Sensorineural hearing loss and prematurity. *Arch Dis Child Fetal Neonatal Ed* 2000;82(2):F141–F144. [PubMed: 10685988]

123. Majnemer A, Riley P, Shevell M, Birnbaum R, Greenstone H, Coates AL. Severe bronchopulmonary dysplasia increases risk for later neurological and motor sequelae in preterm survivors. *Dev Med Child Neurol* 2000;42(1):53–60. [PubMed: 10665976]
124. Bhutta AT, Cleves MA, Casey PH, Cradock MM, Anand KJ. Cognitive and behavioral outcomes of school-aged children who were born preterm: a meta-analysis. *JAMA* 2002;288(6):728–737. [PubMed: 12169077]
125. Roussounis SH, Hubley PA, Dear PR. Five-year-follow-up of very low birth-weight infants: neurological and psychological outcome. *Child Care Health Dev* 1993;19(1):45–59. [PubMed: 7678787]
126. Antoniadou L, MacGregor AJ, Andrew T, Spector TD. Association of birth weight with osteoporosis and osteoarthritis in adult twins. *Rheumatology (Oxford)* 2003;42(6):791–796. [PubMed: 12730541]
127. Barker DJ, Lackland DT. Prenatal influences on stroke mortality in England and Wales. *Stroke* 2003;34(7):1598–1602. [PubMed: 12817108]
128. Bhargava SK, Sachdev HS, Fall CHD, Osmond C, Lakshmy R, Barker DJ, Biswas SK, Ramji S, Prabhakaran D, Reddy KS. Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N Engl J Med* 2004;350(9):865–875. [PubMed: 14985484]
129. Kajantie E, Osmond C, Barker DJ, Forsén T, Phillips DI, Eriksson JG. Size at birth as a predictor of mortality in adulthood: a follow-up of 350 000 person-years [published online ahead of print March 11, 2005]. *Int J Epidemiol* 2005;34(3):655–663.10.1093/ije/dyi048 [PubMed: 15764690]
130. Paneth N. The impressionable fetus? fetal life and adult health. *Am J Public Health* 1994;84(9):1372–1374. [PubMed: 8092355]
131. Curhan GC, Willett WC, Rimm EB, Spiegelman D, Ascherio AL, Stampfer MJ. Birth weight and adult hypertension, diabetes mellitus, and obesity in US men. *Circulation* 1996;94(12):3246–3250. [PubMed: 8989136]
132. Ekholm K, Carstensen J, Finnström O, Sydsjö G. The probability of giving birth among women who were born preterm or with impaired fetal growth: a Swedish population-based registry study. *Am J Epidemiol* 2005;161(8):725–733. [PubMed: 15800264]
133. Wang X, Zuckerman B, Coffman GA, Corwin MJ. Familial aggregation of low birth weight among whites and blacks in the United States. *N Engl J Med* 1995;333(26):1744–1749. [PubMed: 7491139]
134. Barker DJ. Early growth and cardiovascular disease. *Arch Dis Child* 1999;80(4):305–307. [PubMed: 10086930]
135. Bernstein IM, Horbar JD, Badger GJ, Ohlsson A, Golan A. the Vermont Oxford Network. Morbidity and mortality among very-low-birth-weight neonates with intrauterine growth restriction. *Am J Obstet Gynecol* 2000;182(1 pt 1):198–206. [PubMed: 10649179]
136. Fattal-Valevski A, Leitner Y, Kutai M, Tal-Posener E, Tomer A, Lieberman D, Jaffa A, Many A, Harel S. Neurodevelopmental outcome in children with intrauterine growth retardation: a 3-year follow-up. *J Child Neurol* 1999;14(11):724–727. [PubMed: 10593549]
137. Ferguson AC. Prolonged impairment of cellular immunity in children with intrauterine growth retardation. *J Pediatr* 1978;93(1):52–56. [PubMed: 77323]
138. Kok JH, den Ouden AL, Verloove-Vanhorick SP, Brand R. Outcome of very pre-term small for gestational age infants: the first nine years of life. *Br J Obstet Gynaecol* 1998;105(2):162–168. [PubMed: 9501780]
139. Malloy MH. Size for gestational age at birth: impact on risk for sudden infant death and other causes of death, USA 2002 [published online ahead of print February 21, 2007]. *Arch Dis Child Fetal Neonatal Ed* 2007;92(6):F473–F478. [PubMed: 17314115]
140. Committee on Understanding Premature Birth and Assuring Healthy Outcomes, Board on Health Sciences Policy. Societal costs of preterm birth. In: Behrman, RE.; Butler, AS., editors. *Preterm Birth: Causes, Consequences, and Prevention*. Vol. chap 12. Washington, DC: National Academies Press; 2007.
141. Cuevas KD, Silver DR, Brooten D, Youngblut JM, Bobo CM. The cost of prematurity: hospital charges at birth and frequency of rehospitalizations and acute care visits over the first year of life:

- a comparison by gestational age and birth weight. *Am J Nurs* 2005;105(7):56–64. [PubMed: 15995395]
142. Muraskas J, Parsi K. The cost of saving the tiniest lives: NICUs versus prevention. *Virtual Mentor* 2008;10(10):655–658.
143. Cooper WO, Willy ME, Pont SJ, Ray WA. Increasing use of antidepressants in pregnancy. *Am J Obstet Gynecol* 2007;196(6):544.e1–544.e5. [PubMed: 17547888]
144. Chambers CD, Johnson KA, Dick LM, Felix RJ, Jones KL. Birth outcomes in pregnant women taking fluoxetine. *N Engl J Med* 1996;335(14):1010–1015. [PubMed: 8793924]
145. Chun-Fai-Chan B, Koren G, Favez I, Kalra S, Voyer-Lavigne S, Boshier A, Shakir S, Einarson A. Pregnancy outcome of women exposed to bupropion during pregnancy: a prospective comparative study. *Am J Obstet Gynecol* 2005;192(3):932–936. [PubMed: 15746694]
146. Einarson A, Fatoye B, Sarkar M, Lavigne SV, Brochu J, Chambers C, Mastroiacovo P, Addis A, Matsui D, Schuler L, Einarson TR, Koren G. Pregnancy outcome following gestational exposure to venlafaxine: a multicenter prospective controlled study. *Am J Psychiatry* 2001;158(10):1728–1730. [PubMed: 11579012]
147. Sivojelezova A, Shuhaiber S, Sarkissian L, Einarson A, Koren G. Citalopram use in pregnancy: prospective comparative evaluation of pregnancy and fetal outcome. *Am J Obstet Gynecol* 2005;193(6):2004–2009. [PubMed: 16325604]
148. Källén BA, Otterblad Olausson P. Maternal use of selective serotonin re-uptake inhibitors in early pregnancy and infant congenital malformations. *Birth Defects Res A Clin Mol Teratol* 2007;79(4):301–308. [PubMed: 17216624]
149. Littleton HL, Breitkopf CR, Berenson AB. Correlates of anxiety symptoms during pregnancy and association with perinatal outcomes: a meta-analysis. *Am J Obstet Gynecol* 2007;196(5):424–432. [PubMed: 17466693]
150. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4. Washington, DC: American Psychiatric Association; 1994.
151. Misra DP, Guyer B, Allston A. Integrated perinatal health framework: a multiple determinants model with a life span approach. *Am J Prev Med* 2003;25 (1):65–75. [PubMed: 12818312]
152. Lu MC, Tache V, Alexander GR, Kotelchuck M, Halfon N. Preventing low birth weight: is prenatal care the answer? *J Matern Fetal Neonatal Med* 2003;13(6):362–380. [PubMed: 12962261]



Figure 1. Identification of independent studies for inclusion in meta-analysis (adapted from QUOROM flowchart guidelines⁴⁶).




Figure 2. Effect of antenatal depression on the risk of low birth weight (LBW) in developing nations, European social democracies, and the United States. CI indicates confidence interval.

Table 1

Characteristics of Studies Included in the Meta-analysis^a

Source	Country/Gini Coefficient	Sample Size	Sample SES	Race/Ethnicity	Depression Measure	Assessment Times/Triesters	RR (95% CI)			Control Variables	Quality Rating
							PTB	LBW	IUGR		
Andersson et al, 36 2004	Sweden/25.0	1465	Mixed	White	PRIME-MD ⁷¹	Once/second	1.19 (0.59–2.40)	1.19 (0.40–3.56)	...	Yes	12
Berle et al, ⁵⁰ 2005	Norway/25.8	680	Mixed	White	HADS-D ⁷²	Once/third	...	1.78 (0.23–13.89)	...	No	8
Chung et al, ⁵¹ 2001	Hong Kong/43.4	642	Mixed	Chinese	BDI ⁷³	Twice/first, second, and third	1.60 (0.69–3.72)	Yes	8
Copper et al, ³⁷ 1996	United States/40.8	2593	Lower	Black	CES-D ⁷⁴	Once/second and third	1.03 (0.99–1.06)	1.02 (1.00–1.04)	100 (0.97–1.03)	Yes	10
Dayan et al, ⁵² 1999	France/32.7	392	Mixed	White	EPDS ⁷⁵	Once/second	2.10 (1.10–4.10) ^b	No	4
Dayan et al, ³¹ 2006	France/32.7	641	Mixed	White	EPDS ⁷⁵	Once/second	4.90 (1.60–14.90) ^c	Yes	10
Diego et al, ⁵³ 2009	United States/40.8	79	Mixed	Mixed	SCID ⁷⁶	Once/second	2.61 (0.73–9.33)	4.75 (0.94–24.00)	11.28 (0.64–197.35)	No	4
Dole et al, ⁵⁴ 2003	United States/40.8	1962	Mixed	Black/white	CES-D ⁷⁴	Once/second and third	1.12 (0.91–1.38)	Yes	11
Eisenbruch et al, 55 2007	Germany/28.3	896	Mixed	White	CES-D ⁷⁴	Once/first	...	1.16 (0.91–1.47)	...	No	8
Evans et al, ⁵⁶ 2007	England/30.0	10 967	Mixed	White	EPDS ⁷⁵	Twice/second and third	1.29 (0.87–1.91)	Yes	9
Gavin et al, ⁵⁷ 2009	United States/40.8	3019	Mixed	White	CES-D ⁷⁴	Once/second	1.26 (0.61–2.61)	Yes	10
Goldenberg et al, 70 1991	United States/40.8	1545	Lower	Black	CES-D ⁷⁴	Once/second	2.00 (0.94–4.26)	Yes	8
Haas et al, ⁵⁸ 2005	United States/40.8	1619	Mixed	Mixed	CES-D ⁷⁴	Twice/second and third	1.05 (0.66–1.67)	Yes	11
Hedegaard et al, 59 1993	Denmark/24.7	5872	Mixed	White	GHQ ⁷⁷	Twice/second and third	1.33 (1.10–1.60) ^c	Yes	11
Hoffman and Hatch, ⁶⁰ 2000	United States/40.8	666	Mixed	White	CES-D ⁷⁴	3/Second and third	1.07 (0.87–1.31)	...	0.69 (0.23–2.07)	Yes	10

Source	Country/Gini Coefficient	Sample Size	Sample SES	Race/Ethnicity	Depression Measure	Assessment Times/Trimesters	RR (95% CI)			Quality Rating	
							PTB	LBW	IUGR		
Jesse et al, ⁶¹ 2003	United States/40.8	119	Lower	White	2-Item validated screening tool	Once/second	3.19 (1.08–9.44) ^b	Yes	5
Li et al, ⁶² 2009	United States/40.8	791	Mixed	Mixed	CES-D ⁷⁴	Once/first and second	1.60 (0.71–3.63)	Yes	8
Neggers et al, ³³ 2006	United States/40.8	3149	Lower	Black	CES-D ⁷⁴	Once/second	1.30 (1.03–1.64) ^b	1.40 (1.09–1.79) ^c	0.99 (0.75–1.31)	Yes	10
Nordentoft et al, 38 1996	Denmark/24.7	2432	Mixed	White	GHQ ⁷⁷	Once/second	1.01 (0.98–1.05)	...	1.01 (0.97–1.05)	No	9
Orr et al, ³² 2002	United States/40.8	1399	Lower	Black	CES-D ⁷⁴	Once/first and second	1.96 (1.04–3.71) ^b	Yes	10
Paarlberg et al, ⁶³ 1999	The Netherlands/30.9	396	Mixed	White	HSCl ⁷⁸	3/First, second, and third	1.03 (0.98–1.08)	Yes	9
Patel and Prince, 64 2006	India/32.5	245	Lower	Indian	GHQ ⁷⁷	Once/third	3.49 (1.48–8.23) ^c	Yes	8
Perkin et al, ⁶⁵ 1993	United States/40.8	1515	Lower	White	GHQ ⁷⁷	3/First, second, and third	1.28 (0.95–1.73)	Yes	10
Rahman et al, ³⁴ 2004	Pakistan/30.6	265	Lower	Pakistani	SCAN ⁷⁹	Once/third	...	2.10 (1.32–3.35) ^c	...	Yes	9
Rondó et al, ⁶⁶ 2003	Brazil/54.0	865	Lower	Brazilian	GHQ ⁷⁷	3/Second and third	2.32 (1.18–4.58) ^b	1.97 (1.12–3.47) ^b	1.58 (0.84–2.96)	Yes	9
Steer et al, ³⁵ 1992	United States/40.8	389 ^d	Lower	Black	BDI ⁷³	Once/third	1.06 (1.01–1.11) ^b	1.07 (1.02–1.12) ^c	1.05 (1.00–1.11) ^b	Yes	5
Suri et al, ⁶⁷ 2007	United States/40.8	90	Mixed	...	SCID ⁷⁶	Once/first, second, and third	1.41 (0.26–7.73)	1.86 (0.18–19.45)	...	No	6
Wisner et al, ⁶⁸ 2009	United States/40.8	238	Mixed	Mixed	SCID ⁷⁶	Continuous/first, second, and third	2.62 (1.09–6.29) ^b	Yes	7
Zimmer-Gembeck and Heland, ⁶⁹ 1996	United States/40.8	3073	Lower	Mixed	Nonvalidated	Once/first and second	...	1.65 (1.12–2.42) ^b	...	Yes	8

Abbreviations: BDI, Beck Depression Inventory; CES-D, Center for Epidemiologic Studies Depression scale; CI, confidence interval; ellipses, not available; EPDS, Edinburgh Depression Scale; GHQ, General Health Questionnaire; HADS-D, Hospital Anxiety and Depression Scale–Depression; HSCl, Hopkins Symptom Checklist; IUGR, intrauterine growth restriction; LBW, low birth weight; PRIME-MD, Primary Care Evaluation of Mental Disorders; PTB, preterm birth; RR, relative risk; SCAN, Schedules for Clinical Assessment in Neuropsychiatry; SCID, Structured Clinical Interview for DSM-IV; SES, socioeconomic status (as defined by income level, educational level, or type of insurance).

^a Each row in this table describes an independent study of antenatal depression and adverse birth outcomes. Gini coefficient refers to the numerical coefficient indicating the degree of income inequality in each country (0=perfect equality; 100=perfect inequality). Effect sizes are represented by RRs and 95% CIs. Control variables include (1) demographic (age, SES, parity, race/ethnicity, educational level, marital status, work status, and sex of infant); (2) psychiatric (anxiety, stress, alcohol and substance abuse, selective serotonin reuptake inhibitor use, smoking, and previous depression or psychiatric illness); and (3) obstetric/medical (gestational age; prepregnancy body mass index; previous PTB, LBW, or IUGR; previous or current hypertension, diabetes, or preeclampsia; or other obstetric complications). Study quality ratings range from 0 to 12.

^b $P < .05$.

^c $P < .01$.

^d Only the adult (≥ 18 years) sample was included in the meta-analysis.

Table 2

Effect of Antenatal Depression on Outcomes of PTB, LBW, and IUGR

Outcome	No. of Studies	Relative Risk (95% CI) ^a	P Value	Q _{adj} Within	Heterogeneity	
					P Value	Variance Explained, %
PTB	20	1.13 (1.06–1.21)	<.001	49.0 ₁₉	<.001	61
LBW	11	1.18 (1.07–1.30)	.001	33.8 ₁₀	<.001	70
IUGR	12	1.03 (0.99–1.08)	.14	22.4 ₁₁	.02	51

Abbreviations: CI, confidence interval; IUGR, intrauterine growth restriction; LBW, low birth weight; PTB, preterm birth.

^aPooled effect size was estimated using the random-effects model.

Table 3
Moderators of Effect of Antenatal Depression on Outcomes of PTB, LBW, and IUGR

Moderator	Within Group						Effect of Moderator		
	No. of Studies	Relative Risk (95% CI) ^a	P Value	Heterogeneity			Q _{df} Between	P Value	Variance Explained, %
				Q _{df} Within	P Value	Variance Explained, %			
PTB									
Depression predictor									
Categorical	16	1.39 (1.19–1.61)	<.001	24.6 ₁₅	.06	39	14.2 ₁	.001	29
Continuous	4	1.03 (1.00–1.06)	.05	4.7 ₃	.20	36			
Study location ^b									
European social democracy	5	1.37 (1.01–1.85)	.04	20.0 ₄	<.001	80			
United States	14	1.10 (1.03–1.19)	.005	22.2 ₁₃	.05	41	1.8 ₁	.18	4
Study quality ^c									
≤6	5	1.70 (0.99–2.92)	.05	10.0 ₄	.04	60			
>6	15	1.14 (1.06–1.24)	.001	37.4 ₁₄	.001	63	2.0 ₁	.15	4
Effect size									
Adjusted	16	1.18 (1.08–1.28)	<.001	38.4 ₁₅	.001	61	0.6 ₁	.45	1
Unadjusted	4	1.46 (0.84–2.53)	.18	7.0 ₃	.07	57			
LBW									
Depression predictor									
Categorical	9	1.49 (1.25–1.77)	<.001	9.8 ₈	.28	18	14.6 ₁	<.001	43
Continuous	2	1.04 (0.99–1.09)	.10	3.3 ₁	.07	70			
Study location ^d									
Developing nation	2	2.05 (1.43–2.93)	<.001	0.0 ₁	.86	0	10.7 ₂	.005	32
European social democracy	3	1.16 (0.92–1.47)	.20	0.2 ₂	.92	0			
United States	6	1.10 (1.01–1.21)	.03	18.7 ₅	.002	73			
Study quality ^c									
≤6	3	1.59 (0.63–4.00)	.33	3.5 ₂	.18	42	0.1 ₁	.78	0
>6	8	1.39 (1.11–1.73)	.004	27.7 ₇	<.001	75			

Moderator	Within Group					Effect of Moderator			
	No. of Studies	Relative Risk (95% CI) ^a	P Value	Q _{adj} Within	Heterogeneity P Value	Variance Explained, %	Q _{adj} Between	P Value	Variance Explained, %
Effect size							0.17 ₁	.68	0
Adjusted	7	1.17 (1.06–1.30)	.003	29.0 ₆	<.001	79			
Unadjusted	4	1.27 (0.88–1.82)	.20	3.1 ₃	.37	4			
IUGR									
Depression predictor							4.5 ₁	.04	20
Categorical	8	1.45 (1.05–2.02)	.03	14.0 ₇	.05	50			
Continuous	4	1.02 (1.00–1.04)	.12	3.7 ₃	.29	20			
Study location ^e							4.0 ₂	.14	19
Developing nation	2	2.22 (1.03–4.79)	.04	2.1 ₁	.14	53			
European social democracy	3	1.02 (0.99–1.05)	.21	1.8 ₂	.41	0			
United States	6	1.03 (0.96–1.10)	.41	9.6 ₅	.09	48			
Study quality ^c							0.5 ₁	.49	2
≤6	2	2.20 (0.26–18.80)	.47	2.6 ₁	.11	62			
>6	10	1.03 (0.98–1.08)	.30	17.3 ₉	.04	48			
Effect size							0.4 ₁	.52	2
Adjusted	10	1.04 (0.99–1.11)	.14	19.4 ₉	.20	54			
Unadjusted	2	2.17 (0.24–19.60)	.49	2.7 ₁	.10	63			

Abbreviations: CI, confidence interval; IUGR, intrauterine growth restriction; LBW, low birth weight; PTB, preterm birth.

^aPooled effect size was estimated using the random-effects model.

^bOnly 1 study was conducted in a developing nation.

^cStudy quality ratings range from 0 to 12.

^dPairwise effect of moderator for LBW: developing nation vs European social democracy, QI=7.0, P=.01; developing nation vs United States, QI=10.7, P=.001; European social democracy vs United States, QI=0.2, P=.68.

^ePairwise effect of moderator for IUGR: developing nation vs European social democracy, QI=3.9, P=.048; developing nation vs United States, QI=3.8, P=.05; European social democracy vs United States, QI=0.0, P=.84.

Table 4

Comparison of Unadjusted Pooled RRs and Trim-and-Fill Adjusted Pooled RRs

Depression Predictor	No. of Studies	Unadjusted Pooled RR (95% CI) ^a	No. of Missing Studies	Trim-and-Fill Adjusted Pooled RR (95% CI) ^b
PTB				
Overall	20	1.13 (1.06–1.21)	10	1.07 (0.99–1.15)
Categorical	16	1.39 (1.19–1.61)	6	1.24 (1.04–1.47)
Continuous	4	1.03 (1.00–1.06)	1	1.03 (1.00–1.07)
LBW ^c				
Overall	11	1.18 (1.07–1.30)	6	1.10 (1.00–1.22)
Categorical	9	1.49 (1.25–1.77)	4	1.34 (1.10–1.64)
IUGR				
Overall	12	1.03 (0.99–1.08)	4	1.03 (0.97–1.09)
Categorical	8	1.45 (1.05–2.02)	3	1.17 (0.82–1.68)
Continuous	4	1.02 (1.00–1.04)	2	1.00 (0.98–1.03)

Abbreviations: CI, confidence interval; IUGR, intrauterine growth restriction; LBW, low birth weight; PTB, preterm birth; RR, relative risk.

^aUsing random-effects models.

^bUsing random-random effects trim-and-fill models.

^cOnly 2 studies used a continuous depression predictor; the trim-and-fill algorithm requires 3 or more studies.