

NDRG2-mediated Modulation of SOCS3 and STAT3 Activity Inhibits IL-10 Production

Eun Byul Lee^{1#}, Aeyung Kim^{2#}, Kyeongah Kang¹, Hyeree Kim¹ and Jong-Seok Lim^{1*}

¹Department of Biological Science and the Research Center for Women's Diseases, Sookmyung Women's University, Seoul 140-742,

²Department of Molecular Science and Technology, Ajou University, Suwon 443-749, Korea

Background: N-myc downstream regulated gene 2 (NDRG2) is a member of the NDRG gene family. Our previous report indicated a possible role for NDRG2 in regulating the cytokine, interleukin-10 (IL-10), which is an important immunosuppressive cytokine. Several pathways, including p38-MAPK, NF- κ B, and JAK/STAT, are used for IL-10 production, and the JAK/STAT pathway can be inhibited in a negative feedback loop by the inducible protein, SOCS3. In the present study, we investigated the effect of NDRG2 gene expression on IL-10 signaling pathway that is modulated via SOCS3 and STAT3. **Methods:** We generated NDRG2-overexpressing U937 cell line (U937-NDRG2) and treated the cells with PMA to investigate the role of NDRG2 in IL-10 production. U937 cells were also transfected with SOCS3- or NDRG2-specific siRNAs to examine whether the knockdown of SOCS3 or NDRG2 influenced IL-10 expression. Lastly, STAT3 and SOCS3 induction was measured to identify the signaling pathway that was associated with IL-10 production. **Results:** RT-PCR and ELISA assays showed that IL-10 was increased in U937-mock cells upon stimulation with PMA, but IL-10 was inhibited by overexpression NDRG2. After PMA treatment, STAT3 phosphorylation was decreased in a time-dependent manner in U937-mock cells, whereas it was maintained in U937-NDRG2 cells. SOCS3 was markedly reduced in U937-NDRG2 cells compared with U937-mock cells. IL-10 production after PMA stimulation was reduced in U937 cells when SOCS3 was inhibited, but this effect was less severe when NDRG2 was inhibited. **Conclusion:** NDRG2 expression modulates SOCS3 and STAT3 activity, eventually leading to the inhibition of IL-10 production.

[Immune Network 2010;10(6):219-229]

INTRODUCTION

N-myc downstream regulated gene 2 (NDRG2), a member of a new family of genes involved in cell differentiation, is normally expressed in the brain, heart, and muscle (1). It has also been reported that NDRG2 is expressed specifically in dendritic cells (DCs) differentiated from monocytes or CD34⁺ progenitor cells (2). NDRG2 expression in breast cancer cells has been shown to inhibit activation of signal transducer and activator of transcription 3 (STAT3) via induction of suppressor of cytokine signaling 1 (SOCS1) (3). NDRG2 has several roles in cell growth, differentiation and apoptosis, but its function remains to be clarified (2,4).

In a previous report, we found a possible role for NDRG2 in regulating the cytokine, interleukin-10 (IL-10). IL-10 is an important anti-inflammatory cytokine with a crucial role in preventing inflammatory and autoimmune pathologies (5). IL-10 functions as an immune suppressor and can also promote the growth of lymphocytes *in vivo* (6). As IL-10 limits the immune response to pathogens, this cytokine thus prevents damage to the host. IL-10 is produced by T cells, B cells, dendritic cells, and monocytes/macrophages, and its immunosuppressive activity has been documented in IL-10 knockout mice, which have increased autoimmune disease and increased resistance to infection (7-9). Because of its pro-

Received on November 9, 2010. Revised on November 15, 2010. Accepted on November 18, 2010.

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Corresponding Author. Tel: 82-2-710-9560; Fax: 82-2-2077-7322; E-Mail: jslim@sookmyung.ac.kr

#These authors contributed equally to this work.

Keywords: NDRG2, IL-10, SOCS3, STAT3, U937

duction by many immune cells, IL-10 clearly plays a crucial role as a feedback regulator of diverse immune responses. IL-10 mediates its effect by binding to the heterodimeric IL-10 receptor (IL-10R); Jak1 and Tyk2 are then transphosphorylated and activated, and these molecules subsequently activate the downstream transcription factor, STAT3, and in some cases, STAT1 and STAT5 (10-12). There are several mechanisms of regulation of IL-10 expression, and this regulation is different depending on the type of cell or location. Currently, studies investigating immunosuppressive processes of diverse cells or formation and maintenance of regulatory T cells by IL-10 are being performed (13).

p38 mitogen-activated protein kinase (p38-MAPK), nuclear factor- κ B (NF- κ B) and JAK/STAT pathways are the known signaling pathways for innate IL-10 production (14,15). In particular, activation of STAT3 is essential for the effects of IL-10 (16). Macrophages increase their production of IL-10 by activating STAT3 in an autocrine fashion, and the activated STAT3 then plays an important role in inhibiting the activation of macrophages and neutrophils (7). In mouse T cells, the induction of IL-10 by IL-27 depends on both STAT1 and STAT3, and STAT3 is also involved in IL-6-mediated IL-10 expression (17-19). By contrast, a recent study in human monocytes describes an inhibitory role for IL-27 on IL-10 production through STAT1 (20). Another study suggests that IL-10 induces its own expression by human monocyte-derived macrophages in an autocrine manner through the activation of STAT3 (7).

Activated STATs dimerize and migrate to the nucleus where they activate the transcription of specific genes among; some of these genes are members of the suppressors of cytokine signaling (SOCS) family (21). SOCS proteins (SOCS1-7 and cytokine-inducible SH2-containing protein (CIS)) negatively regulate cytokine signaling pathways (22). The inducible SOCS3 protein inhibits the JAK/STAT pathway in a negative feedback loop using a variety of mechanisms (23). In addition, it is known that SOCS3 plays an important role in the negative regulation of inflammatory responses via the induction of IL-10. Among SOCS family members, SOCS1, SOCS3 and SOCS5 are mainly involved in the regulation of IL-4 and IL-10 pathways. Both SOCS1 and SOCS3 inhibit JAK tyrosine kinase activity via their N-terminal kinase inhibitory region. SOCS1 directly binds to the activation loop of JAK molecules through its Src-homology 2 domain, whereas SOCS3 inhibits the action of JAK molecules only in the presence of a receptor that it binds (24). However, both SOCS1 and SOCS3 are activated

by IL-4 or IL-10 and function as potent inhibitors of IL-4/STAT6 or IL-10/STAT3 activation (25). It has been reported that targeted deletion of SOCS3 in macrophages results in markedly enhanced IL-6-induced STAT3 activation (26).

Monocytes are known to be one of the major sources of IL-10. In this study, we used the human monocytic cell line, U937, which was derived from malignant cells obtained from the pleural effusion of a patient with histiocytic lymphoma in 1974. U937 cells display many characteristics of monocytes and have thus served as a model for the differentiation of monocytes and macrophages *in vitro*. This cell line matures and differentiates in response to a number of soluble stimuli and adopts the morphology and characteristic features of mature macrophages. Herein, we generated NDRG2-overexpressing U937 cells and treated them with a stimulus to investigate the role of NDRG2 in IL-10 production. We also investigated whether the IL-10 signaling pathway was modulated via SOCS3 and STAT3.

MATERIALS AND METHODS

Cell culture and treatment

U937 cells were grown in RPMI (Gibco/Invitrogen, Carlsbad, CA) supplemented with 10% heat-inactivated fetal bovine serum (Gibco). Cells were maintained in an atmosphere of 5% CO₂ in a 37°C humidified incubator. Phorbol 12-myristate 13-acetate (PMA) was purchased from Sigma-Aldrich (St. Louis, MO) and cucurbitacin I, a selective inhibitor of JAK/STAT3, and AG490, an inhibitor of constitutive STAT3-DNA binding, were purchased from Calbiochem (Merck KGaA, Darmstadt, Germany).

Transfection

The NDRG2 gene was cloned into the pcDNA3.1 vector (Invitrogen, Carlsbad, CA) and the resulting pcDNA3.1-NDRG2 plasmid was introduced into U937 cells by electroporation with a Gene Pulser (Bio-Rad, Hercules, CA) at 960 μ F and 300 V. Transfectants resistant to 1 mg/ml G418 (Sigma) were screened for NDRG2 expression. NDRG2 expression was confirmed by RT-PCR and western blotting. NDRG2-expressing cells were expanded in culture for further study.

RNA interference

The SMARTpool siRNA specific for human SOCS3 was purchased from Dharmacon (Lafayette, CO), and NDRG2-specific siRNA and control non-targeting siRNA were purchased from

Santa Cruz Biotechnology (Santa Cruz, CA). U937-mock and NDRG2-overexpressing cells were transfected with 100 nM of control, SOCS3- or NDRG2-specific siRNAs using an Amaxa Nucleofector™ (program V-001, Amaxa, Cologne, Germany). After transfection, cells were immediately placed in a 37°C humidified incubator.

Reverse transcription (RT)

Total RNA from harvested cells was isolated using TRI reagent (Molecular Research Center, Cincinnati, OH) and 5 µg of total RNA was reverse transcribed (RT) into cDNA. For the RT reaction, RNA was first incubated with an oligo-(dT) primer at 70°C to denature the RNA secondary structure and then incubated at room temperature for 10 min to allow the primer to anneal to the RNA. Other components of the RT assay, including dNTPs (Bioneer, Daejeon, Korea), M-MLB reverse transcriptase, and RT buffer (Promega, Madison, WI), were then added to the reaction. The RT reaction was performed at 37°C for 1 h and then at 100°C for 5 min. The RT product was then used as a template for a 20 µl PCR reaction.

Polymerase chain reaction (PCR)

Oligonucleotide primers for PCR were designed according to the published sequences (GenBank). PCR primers were purchased from Bioneer, or GENOTECH (Daejeon, Korea). The following primer sequences were used: β -actin: 5'-cca cac ctt cta caa tga gc-3' (sense), 5'-tga ggt agt cag tca ggt cc-3' (antisense); NDRG2: 5'-gga ttc atg gcg gag ctg cag gag g-3' (sense), 5'-gaa ttc tca aca gga gac ctg cat ggt-3' (antisense); IL-10: 5'-gat gcc ttc agc aga gtg aag a-3' (sense), 5'-cat ggc ttt gta gat gcc ttt c-3' (antisense); SOCS1: 5'-aga ccc ctt ctg acc tct tg-3' (sense), 5'-ctg cac agc aga aaa taa agc-3' (antisense); SOCS2: 5'-gga tgg tac tgg gga agt atg act-3' (sense), 5'-agt gga tca gat gaa cca cac tct-3' (antisense); SOCS3: 5'-cac atg gca caa gca caa ga-3' (sense), 5'-aag tgt ccc ctg ttt gga gg-3' (antisense); SOCS4: 5'-aga gaa ggt cct atg act gg-3' (sense), 5'-ggg cat cga tgc cat cat aa-3' (antisense); SOCS5: 5'-cag ttc tgc aat tcc aca ag-3 (sense), 5'-gac tgg ttc tgc ttc caa cc-3 (antisense). PCR products were electrophoresed on a 1% agarose gel and visualized by ethidium bromide staining.

Western blot analysis

U937 cells were treated with the indicated concentration (50 ng/ml) of PMA for 0~24 h. The treated cells and untreated control cells were washed with PBS (WelGENE, Daegu,

Korea) and immediately resuspended in ice-cold lysis buffer (M-PER® Mammalian protein extraction reagent, Pierce, Rockford, IL) for 20 min. Supernatant fractions were recovered by centrifugation at 12,000 rpm for 15 min. For immunoblotting, cell lysates containing equivalent amounts of total protein were mixed with 5X sample buffer and resolved on a SDS-polyacrylamide gel. Separated proteins were transferred to a Hybond-P (Amersham Biosciences, Burkes, UK) membrane. Membranes were incubated in 5% skim milk (BD Biosciences, San Jose, CA) in TBS-T (20 mM Tris-HCL, pH 7.5, 137 mM NaCl, 0.05% tween-20) for 1 h at room temperature, washed with TBS-T and incubated with primary antibody overnight at 4°C. The membranes were washed with TBS-T and incubated with secondary antibody for 2 h at room temperature. Protein bands were visualized by enhanced chemiluminescence (SuperSignal West Pico Chemiluminescent Substrate, Pierce, Rockford, IL) using a LAS 3000 imaging system (FUJIFILM Corporation, Tokyo, Japan).

Antibodies for western blotting

Antibodies against the following proteins were used in this study: STAT3 and phospho-STAT3 (Tyr705) (Cell Signaling, Danvers, MA); SOCS1, SOCS2, SOCS3, and actin (Santa Cruz, CA); NDRG2 (produced by our laboratory). Peroxidase-conjugated AffiniPure goat anti-rabbit IgG (Jackson Immuno-Research Laboratories, West Grove, PA), bovine anti-goat IgG HRP (Santa Cruz), and goat anti-mouse IgG (Sigma-Aldrich) were also used.

Cell proliferation using MTT assay and hemocytometer

Cells (2×10^4 cells/well) were cultured in 96-well culture plates for 24 h and treated with MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) for 2 h. Cells were harvested and centrifuged at 1,200 rpm for 5 min. The pellets were dissolved in 200 µl DMSO. Absorbance was measured at 540 nm using an ELISA reader (Victor 3™, PerkinElmer, Waltham, MA). For cell counting by microscope, cells (1×10^4 cells/well) in 6-well culture plates were cultured for 1 to 4 days and centrifuged at 1,200 rpm for 5 min. Viable cells, which are visible by light microscopy, were enumerated using a hemocytometer.

Cytokine detection by ELISA

IL-10 production was assessed after 24 h in culture by collecting the supernatants from U937 cells (3×10^5 cells/ml) that

had undergone PMA treatment, IL-10 levels were measured using the human IL-10 ELISA kit (Peprotech, Rocky Hill, NJ) according to the manufacturer's instructions. Each sample was tested in duplicate and quantified using the Multi Detection Microplate SpectraMax M5 (Molecular Devices, Sunnyvale, CA) at 405 nm and 650 nm absorbance.

Statistical analysis

Statistical analysis for the comparison between groups was performed using student's *t*-test and represented by the mean \pm SD.

RESULTS

NDRG2 inhibits IL-10 production in U937 cells

To examine IL-10 production in U937 cells, cells were stimulated with 50 ng/ml of PMA, and IL-10 mRNA was measured by RT-PCR. IL-10 expression increased in a time-dependent manner upon PMA stimulation in U937 cells (Fig. 1A). After 12 h of stimulation with PMA, IL-10 transcripts increased in a dose-dependent manner. Similarly, PMA treatment induced IL-10 protein production in U937 cells when measured by ELISA (Fig. 1B). We hypothesized that NDRG2 gene expression might inhibit IL-10 production. To determine the relationship between NDRG2 and IL-10 expression, U937-mock cells and U937 cells overexpressing NDRG2 (U937-NDRG2) were treated with PMA, and IL-10 mRNA levels were determined by RT-PCR. As shown in Fig. 2A, IL-10 mRNA in U937-NDRG2 cells was suppressed compared to wild type or mock-transfected cells. The measurement of IL-10 by ELISA showed that whereas IL-10 production was remarkably increased in U937-mock cells upon PMA stimulation, it was strongly reduced in U937-NDRG2 cells (Fig. 2B). Although overexpression of the gene itself did not affect cell proliferation (Fig. 2C), IL-10 production was corrected based on cell numbers because U937-NDRG2 cells showed growth rate faster than wild-type cells upon PMA stimulation (data not shown). These data indicate the negative effect of NDRG2 expression on both IL-10 protein production and IL-10 expression levels.

Expression of IL-10 and SOCS genes upon PMA stimulation

Based on our previous report (3), we hypothesized that NDRG2 might inhibit IL-10 production through SOCS and STAT signaling. SOCS genes (SOCS1-5) have emerged as key




Figure 1. PMA stimulation of U937 cells induces IL-10 production. (A) Dose- and time-dependent increase in IL-10 mRNA after PMA stimulation. U937 cells were stimulated with PMA for different time periods and harvested for mRNA analysis. Transcript levels of IL-10 and β -actin (loading control) were determined by RT-PCR. (B) U937 cells were stimulated with different concentrations of PMA for 24 h. Cells were harvested, and IL-10 protein levels in the supernatants were determined by ELISA. Data represent the mean \pm SD of duplicates.

physiological regulators of cytokine responses, including those that regulate the immune system (27). To test this hypothesis, U937-mock and U937-NDRG2 cells were treated with PMA (50 ng/ml) for 0, 15, 30, and 60 min and SOCS gene expression was detected by RT-PCR. Interestingly, SOCS3 expression was significantly upregulated upon PMA stimulation in U937 cells but was only slightly increased in U937-NDRG2 cells. In contrast, there was no significant difference in the levels of other SOCS family members (SOCS1, 2, 4, 5) between U937-mock and U937-NDRG2 cells (Fig. 3A). These expression patterns were maintained at 24 h after PMA treatment (Fig. 3B). Although PMA treatment appeared to




Figure 2. IL-10 production is reduced by NDRG2 expression. (A) U937, U937-mock and U937-NDRG2 cells were stimulated with PMA (50 ng/ml) for different time periods and harvested for mRNA analysis. Transcript levels of IL-10 and β -actin were determined by RT-PCR. (B) U937-mock and U937-NDRG2 cells (3×10^5 cells/ml) were stimulated with PMA (50 ng/ml) for 24 h. The levels of IL-10 in culture supernatants were determined by ELISA and corrected based on cell numbers after cell counting. Data represent the mean \pm SD of duplicates. (C) Cells (2×10^4 cells/well) were cultured in 96-well culture plates for 24 h and treated with MTT for 2 h. The cells were harvested and centrifuged at 1,200 rpm for 5 min. The pellets were dissolved in 200 μ l of DMSO, and absorbance was measured at 540 nm using an ELISA reader (upper). For cell counting by microscope, cells (5×10^5 cells/well) in 6-well culture plates were cultured for 1 to 4 days and centrifuged at 1,200 rpm for 5 min. Viable cells, which are visible by light microscopy, were enumerated using a hemocytometer. Data represent the mean \pm SD of triplicates.

slightly increase NDRG2 transcript levels as a result of the vector used for the transfection, its effect on protein level was negligible (see below). These results indicate that NDRG2 expression inhibits IL-10 induction in U937 cells when cells are simulated with PMA, and the expression pattern of SOCS3 mirrors that of IL-10.

Effect of NDRG2 expression on SOCS3 and STAT3 expression after PMA stimulation

IL-10 production has effects on many immune cells, affirming its crucial role as a feedback regulator of diverse immune responses (15). As mentioned above, p38-MAPK, NF- κ B and JAK/STAT pathways are known signaling pathways for innate

IL-10 production. In particular, STAT3 has an important role in IL-10 production and is associated with several cellular signals. In addition, it has been reported that targeted deletion of SOCS3 in macrophages results in markedly enhanced STAT3 activation (28). When we measured activation of STAT3, we found that phosphorylated STAT3 levels significantly decreased in U937-mock cells after PMA stimulation, whereas these levels were largely unchanged in U937-NDRG2 cells (Fig. 4A). Because we observed decreased levels of SOCS3 mRNA when NDRG2 was overexpressed, SOCS3 expression and STAT3 phosphorylation were examined together. After PMA stimulation, protein levels of SOCS1, 2, and 3 and phosphorylated STAT3 were assessed by western blot




Figure 3. IL-10 and SOCS3 mRNA levels after PMA stimulation are reduced by NDRG2 expression. (A) U937-mock and U937-NDRG2 cells were stimulated at various times with 50 ng/ml of PMA. Cells were harvested for analysis. Transcript levels of IL-10 and SOCS family members were determined by RT-PCR. (B) Cells were stimulated with PMA for 24 h and harvested for mRNA analysis. Transcript levels of IL-10 and SOCS genes were determined by RT-PCR.

at various time points. The data showed that phosphorylated STAT3 decreased in a time-dependent manner in U937-mock cells, but not in U937-NDRG2 cells (Fig. 4B). In contrast, SOCS3 levels were decreased in cells overexpressing NDRG2 but maintained at high levels in U937-mock cells. As SOCS1 and SOCS2 proteins remained at basal levels during the experimental times, it appeared that the inhibitory effect of




Figure 4. NDRG2 expression inhibits induction of SOCS3 but promotes STAT3 phosphorylation after PMA stimulation. (A) U937-mock and U937-NDRG2 cells were stimulated with 50 ng/ml of PMA for 15 or 30 min. Cell lysates were harvested and equal amounts of lysates were subjected to electrophoresis on SDS-PAGE. Western blotting was performed using STAT3-, pSTAT3- and NDRG2-specific antibodies. (B) Cells were stimulated at various times with 50 ng/ml of PMA. Equal amounts of whole-cell lysates were subjected to electrophoresis on SDS-PAGE, and then western blotting was performed using specific antibodies.

NDRG2 is SOCS3-specific. Therefore, NDRG2 inhibits SOCS3 expression and maintains STAT3 phosphorylation under conditions of PMA stimulation.

Increased IL-10 production upon STAT3 inhibition

To determine whether STAT3 inhibition affected IL-10 production induced by PMA, cells were pretreated with the STAT3 inhibitor, cucurbitacin I, for 2 h and further incubated with PMA for 48 h. As shown in Fig. 5A, IL-10 expression in U937-mock cells was enhanced upon PMA treatment, but it was markedly increased by pretreatment with cucurbitacin I, indicating a role for STAT3 in inhibiting IL-10 production. In line with these data, IL-10 expression in U937-NDRG2 cells after PMA stimulation was also significantly upregulated by pretreatment with cucurbitacin I or AG490 (Fig. 5B). Although the level of IL-10 transcripts was relatively low, increased lev-




Figure 5. Inhibition of STAT3 phosphorylation increases IL-10 production. (A) U937-mock cells were pretreated with cucurbitacin I for 2 h followed by PMA treatment for 48 h. Cells were harvested for mRNA analysis. Transcript levels of IL-10 were determined by RT-PCR. (B) U937-NDRG2 cells were pretreated with cucurbitacin I or AG490 for 2 h followed by PMA treatment for 48 h. Cells were harvested for mRNA analysis. Transcript levels of IL-10 were assessed by RT-PCR. (C) U937-NDRG2 cells were treated in the same manner as in (B). The levels of IL-10 in culture supernatants were determined by ELISA. Data represent the mean \pm SD of duplicates.

els of IL-10 protein in U937-NDRG2 cells was also observed after pretreatment with STAT3 inhibitors followed by PMA stimulation (Fig. 5C). Together, these data suggest that high levels of phosphorylated STAT3 in NDRG2-transfected U937 cells may play a role in the inhibition of IL-10 upon stimulation with PMA.

Effect of SOCS3- or NDRG2-specific siRNAs on U937 proliferation

Next, to test whether inhibition of SOCS3 affected IL-10 production, U937-mock and U937-NDRG2 cells were transfected with specific small interfering RNAs (siRNAs; siRNA control, siSOCS3, siNDRG2). We first examined the proliferation capacity of siRNA-transfected cells by plating the cells in 60 mm dishes and counting the cells daily for 4 days using a hemocytometer. The data showed that there were no differences in cell proliferation between the five cell types (Fig. 6).




Figure 6. Proliferation of siRNA-transfected U937 cells. U937-mock and U937-NDRG2 cells were transfected with siRNA control, SOCS3- or NDRG2-specific siRNAs (100 nM) for 48 h. After transfection, cells (1×10^4) were plated in 60 mm dishes and counted daily for 4 days.

IL-10 is down-regulated upon SOCS3 inhibition and recovered upon NDRG2 inhibition

To determine the effects of SOCS3 inhibition on IL-10 mRNA levels, siRNA-transfected U937-mock and U937-NDRG2 cells were stimulated with PMA, and gene expression was assessed by RT-PCR. First, we measured the level of SOCS3 and NDRG2 expression in siRNA-transfected cells to confirm the knockdown of the genes. As shown in Fig. 7A, SOCS3 expression was attenuated upon transfection with siSOCS3 in both U937-mock and U937-NDRG2 cells, and NDRG2 expression was attenuated upon transfection with siNDRG2 in U937-NDRG2 cells. Interestingly, reduced levels of SOCS3 elicited the down-regulation of IL-10 mRNA in U937-mock cells, and IL-10 mRNA levels were recovered in U937-NDRG2 cells upon knockdown of NDRG2. These results suggest that SOCS3 could be involved in IL-10 induction in U937 cells. Because the induction of IL-10 mRNA appeared to be associated with SOCS3 expression in U937-mock and U937-NDRG2 cells, we investigated whether IL-10 protein levels were also being decreased by SOCS3 inhibition. U937-mock and U937-NDRG2 cells were cultured for 24 h with PMA, and IL-10 production was measured by ELISA. As shown in Fig. 7B, IL-10 levels in U937-mock cells transfected with siSOCS3 were significantly reduced, whereas IL-10 levels in U937-NDRG2 cells remained unchanged in both control siRNA- and siSOCS3-transfected cells. The latter result indicates that SOCS3 expression was already down-regulated by NDRG2-overexpression in U937-NDRG2 cells. As with the mRNA ex-




Figure 7. Effect of SOCS3 siRNA and NDRG2 siRNA on IL-10 mRNA or protein expression. (A) U937-mock and U937-NDRG2 cells were transfected with siRNA control, SOCS3- or NDRG2-specific siRNAs (100 nM) for 48 h. Cells were then stimulated with PMA and harvested for mRNA analysis. Transcript levels of IL-10, SOCS3, and NDRG2 genes were determined by RT-PCR. (B) After siRNA transfection for 48 h, U937-mock and U937-NDRG2 cells (3×10^7 cells/ml) were stimulated with PMA for 24 h. The levels of IL-10 in culture supernatants were determined by ELISA. Data represent the mean \pm SD of duplicates.

pression data in Fig. 7A, IL-10 protein production was restored by knockdown of NDRG2. Thus, NDRG2 expression exerts a negative effect on IL-10 production through the modulation of SOCS3 and STAT3.

DISCUSSION

NDRG2, a member of the N-myc downstream-regulated gene family that includes NDRG1-4, is highly expressed in many normal cells and tissues but it is expressed at low levels or not at all in various tumors and tumor cell lines (29). NDRG2 is generally not expressed in monocytes or macrophages but is specifically expressed in dendritic cells (DCs) differentiated

from monocytes, CD34⁺ progenitor cells, and the myelomonocytic leukemic cell lines. In addition, it has been reported that NDRG2 expression in DCs modulates the T-cell stimulating capabilities of DCs via co-stimulatory factor expression (30). Therefore, NDRG2 may regulate immune responses via secretion of cytokine such as IL-10 or IL-12.

Phorbol 13-myristate 12-acetate (PMA) is a known activator of protein kinase C (PKC) and the Raf/MEK/Erk pathway (31,32). It has been reported that preincubation of cells with PMA inhibits the IL-6-induced STAT3 and STAT1 activation (33,34). This negative effect was found to be dependent on the activation of MAP kinases and transcription and protein synthesis. Because PMA leads to a rapid upregulation of SOCS3, a negative regulator of the JAK/STAT pathway, it is conceivable that the inhibitory effect of PMA on the JAK/STAT pathway is mediated by SOCS3 (35). In the present study, we chose PMA as a stimulator for IL-10 induction to investigate SOCS3 and STAT3 signaling.

We used the NDRG2-transfected human monocytic cell line, U937, to determine whether the overexpression of NDRG2 in U937 cells affected the ability of cells to produce IL-10. Our results showed that whereas IL-10 mRNA after PMA stimulation increased in U937-mock cells, it was significantly suppressed upon NDRG2 expression. As we were interested in SOCS3 signaling for IL-10 induction, mRNA levels of SOCS genes (SOCS1-5) were investigated. Interestingly, SOCS3 expression increased upon PMA stimulation in U937-mock cells but was not significantly changed in U937-NDRG2 cells, whereas the levels of other SOCS genes (SOCS1, 2, 4, 5) showed no significant difference between U937-mock and U937-NDRG2 cells. Based on these data, it is reasonable to assume that IL-10 production is affected by SOCS3 expression.

STAT3, one of signaling molecules that controls IL-10 production, is unphosphorylated and resides in the cytoplasm in normal resting cells. However, in response to various stimuli, STAT3 is activated through phosphorylation of tyrosine 705 or serine 727, which promotes its homodimerization and subsequent entry into the nucleus where the homodimers activate transcription (36). In particular, activation of STAT3 via phosphorylation of the tyrosine 705 residue induces the transcription of genes involved in inflammatory responses and cell proliferation and differentiation. STAT3 in immune cells can also induce genes involved in the inhibition of innate and adaptive immune responses that render resistance to tumors, and so it is expected that STAT3 negatively regulates immune

responses against tumors (37).

SOCS proteins are controlled in a STAT-dependent manner and inhibit the JAK/STAT pathway by a negative feedback loop. It has been reported that activation of ERK is essential for SOCS3 induction by either cyclic AMP or PKC (38). In recent years, the SOCS proteins, which were first identified as rapidly inducible negative regulators of cytokine signaling, have emerged as key modulators of the cellular response to numerous biological stimuli. The SOCS proteins regulate the duration of the cytokine signaling response by both specific and non-specific regulatory mechanisms (39). Overexpression of SOCS3 in cell lines has shown that SOCS proteins can inhibit many cytokine-induced responses (23). For example, it has been reported that SOCS3 overexpression in human periodontal ligament (PDL) cells reduces the phosphorylation in downstream signaling molecules including STAT (40). Furthermore, enhanced SOCS3 gene expression has been found to promote IL-10 production, suggesting that STAT3 and SOCS3 may play an important role in regulation of cytokine-induced anti-inflammatory responses (41).

We examined protein levels of STAT3 and SOCS3 by western blot after PMA stimulation to determine the relationship between IL-10 production and STAT3, SOCS3, and NDRG2 expression. We found that phosphorylated STAT3 was decreased in a time-dependent manner in U937-mock cells, but it was unchanged in U937-NDRG2 cells. By contrast, SOCS3 was suppressed in cells overexpressing NDRG2 but maintained at high levels in U937-mock cells. Increased SOCS3 expression after PMA stimulation might promote IL-10 production. The fact that SOCS3 levels were reduced by NDRG2 overexpression but STAT3 levels were maintained suggests that the negative feedback of SOCS3 was blocked. Blocking of STAT3 by specific inhibitors, such as cucurbitacin I, resulted in increased IL-10 production in both U937-mock and U937-NDRG2 cells, indicating a critical role for STAT3 in signaling for IL-10 production.

To investigate the role of SOCS3 in IL-10 production in the context of NDRG2 expression, U937-mock and U937-NDRG2 cells were transfected with specific siRNAs (siRNA control, siSOCS3, siNDRG2). Upon stimulation with PMA, IL-10 mRNA was decreased in U937-mock cells after siRNA-mediated knockdown of SOCS3. Although there is a great deal of evidence showing that SOCS3 can interfere with cytokine-induced phosphorylation of STAT3 in immune cells (42,43), it is presently unclear whether STAT3 recovery is directly responsible for the reduction of IL-10 in SOCS3-specific

siRNA-transfected cells. Nonetheless, together with enhanced IL-10 gene and protein expression upon treatment with STAT3 inhibitors, our data clearly indicate that PMA-induced SOCS3 expression negatively regulates STAT3, leading to increased production of IL-10. In this context, our data support the hypothesis that SOCS3 induced by IL-10 can participate in the inhibition of STAT3 that results in reduced expression of proinflammatory molecules.

In this study, we report that SOCS3 expression was reduced by NDRG2-overexpression in U937-NDRG2 cells, and the decreased IL-10 production was restored by NDRG2 inhibition. These results indicate that NDRG2 inhibits IL-10 production through the inhibition of SOCS3 upon PMA stimulation. Taken together, our data suggest that NDRG2 expression modulates SOCS3 and STAT3 activity, leading to the inhibition of IL-10 production. Further study is required to identify the detailed mechanism of NDRG2 effects associated with MAPK or other possible signaling pathways on IL-10 production.

ACKNOWLEDGEMENTS

This work was financially supported by the Research Center for Women's Diseases of the KOSEF (R11-2005-017-06001) and by the Korea Research Foundation Grant (KRF-2009-0074435) funded by the Korean Government (MEST).

CONFLICTS OF INTEREST

The authors have no financial conflict of interest.

REFERENCES

1. Kim YJ, Yoon SY, Kim JT, Choi SC, Lim JS, Kim JH, Song EY, Lee HG, Choi I, Kim JW: NDRG2 suppresses cell proliferation through down-regulation of AP-1 activity in human colon carcinoma cells. *Int J Cancer* 124;7-15, 2009
2. Choi SC, Kim KD, Kim JT, Kim JW, Yoon DY, Choe YK, Chang YS, Paik SG, Lim JS: Expression and regulation of NDRG2 (N-myc downstream regulated gene 2) during the differentiation of dendritic cells. *FEBS Lett* 553;413-418, 2003
3. Park Y, Shon SK, Kim A, Kim KI, Yang Y, Cho DH, Lee MS, Lim JS: SOCS1 induced by NDRG2 expression negatively regulates STAT3 activation in breast cancer cells. *Biochem Biophys Res Commun* 363;361-367, 2007
4. Liu N, Wang L, Li X, Yang Q, Liu X, Zhang J, Zhang J, Wu Y, Ji S, Zhang Y, Yang A, Han H, Yao L: N-Myc downstream-regulated gene 2 is involved in p53-mediated apoptosis. *Nucleic Acids Res* 36;5335-5349, 2008

5. O'Garra A, Barrat FJ, Castro AG, Vicari A, Hawrylowicz C: Strategies for use of IL-10 or its antagonists in human disease. *Immunol Rev* 223;114-131, 2008
6. Choi YK, Kim YJ, Park YH, Choi KS, Park JG: Effective inhibition of glomerulosclerosis by adenoviral vector expressing human IL-10. *Korean J Immunol* 22;187-195, 2000
7. Staples KJ, Smallie T, Williams LM, Foey A, Burke B, Foxwell BM, Ziegler-Heitbrock L: IL-10 induces IL-10 in primary human monocyte-derived macrophages via the transcription factor Stat3. *J Immunol* 178;4779-4785, 2007
8. Bettelli E, Das MP, Howard ED, Weiner HL, Sobel RA, Kuchroo VK: IL-10 is critical in the regulation of autoimmune encephalomyelitis as demonstrated by studies of IL-10- and IL-4-deficient and transgenic mice. *J Immunol* 161;3299-3306, 1998
9. Dai WJ, Köhler G, Brombacher F: Both innate and acquired immunity to *Listeria monocytogenes* infection are increased in IL-10-deficient mice. *J Immunol* 158;2259-2267, 1997
10. Weber-Nordt RM, Riley JK, Greenlund AC, Moore KW, Darnell JE, Schreiber RD: Stat3 recruitment by two distinct ligand-induced, tyrosine-phosphorylated docking sites in the interleukin-10 receptor intracellular domain. *J Biol Chem* 271;27954-27961, 1996
11. Wehinger J, Gouilleux F, Groner B, Finke J, Mertelmann R, Weber-Nordt RM: IL-10 induces DNA binding activity of three STAT proteins (Stat1, Stat3, and Stat5) and their distinct combinatorial assembly in the promoters of selected genes. *FEBS Lett* 394;365-370, 1996
12. Finbloom DS, Winestock KD: IL-10 induces the tyrosine phosphorylation of tyk2 and Jak1 and the differential assembly of STAT1 alpha and STAT3 complexes in human T cells and monocytes. *J Immunol* 155;1079-1090, 1995
13. O'Garra A, Vieira P: Regulatory T cells and mechanisms of immune system control. *Nat Med* 10;801-805, 2004
14. Akira S, Takeda K: Toll-like receptor signalling. *Nat Rev Immunol* 4;499-511, 2004
15. Saraiva M, O'Garra A: The regulation of IL-10 production by immune cells. *Nat Rev Immunol* 10;170-181, 2010
16. Williams L, Bradley L, Smith A, Foxwell B: Signal transducer and activator of transcription 3 is the dominant mediator of the anti-inflammatory effects of IL-10 in human macrophages. *J Immunol* 172;567-576, 2004
17. Stumhofer JS, Silver JS, Laurence A, Porrett PM, Harris TH, Turka LA, Ernst M, Saris CJ, O'Shea JJ, Hunter CA: Interleukins 27 and 6 induce STAT3-mediated T cell production of interleukin 10. *Nat Immunol* 8;1363-1371, 2007
18. Batten M, Kljavin NM, Li J, Walter MJ, de Sauvage FJ, Ghilardi N: Cutting edge: IL-27 is a potent inducer of IL-10 but not FoxP3 in murine T cells. *J Immunol* 180;2752-2756, 2008
19. Xu J, Yang Y, Qiu G, Lal G, Wu Z, Levy DE, Ochando JC, Bromberg JS, Ding Y: c-Maf regulates IL-10 expression during Th17 polarization. *J Immunol* 182;6226-6236, 2009
20. Kalliolias GD, Ivashkiv LB: IL-27 activates human monocytes via STAT1 and suppresses IL-10 production but the inflammatory functions of IL-27 are abrogated by TLRs and p38. *J Immunol* 180;6325-6333, 2008
21. Hebenstreit D, Wirnsberger G, Horejs-Hoeck J, Duschl A: Signaling mechanisms, interaction partners, and target genes of STAT6. *Cytokine Growth Factor Rev* 17;173-188, 2006
22. Qin H, Niyongere SA, Lee SJ, Baker BJ, Benveniste EN: Expression and functional significance of SOCS-1 and SOCS-3 in astrocytes. *J Immunol* 181;3167-3176, 2008
23. Alexander WS, Hilton DJ: The role of suppressors of cytokine signaling (SOCS) proteins in regulation of the immune response. *Annu Rev Immunol* 22;503-529, 2004
24. Yoshimura A, Nishinakamura H, Matsumura Y, Hanada T: Negative regulation of cytokine signaling and immune responses by SOCS proteins. *Arthritis Res Ther* 7;100-110, 2005
25. Ding Y, Chen D, Tarcsfalvi A, Su R, Qin L, Bromberg JS: Suppressor of cytokine signaling 1 inhibits IL-10-mediated immune responses. *J Immunol* 170;1383-1391, 2003
26. Yasukawa H, Ohishi M, Mori H, Murakami M, Chinen T, Aki D, Hanada T, Takeda K, Akira S, Hoshijima M, Hirano T, Chien KR, Yoshimura A: IL-6 induces an anti-inflammatory response in the absence of SOCS3 in macrophages. *Nat Immunol* 4;551-556, 2003
27. Alexander WS: Suppressors of cytokine signalling (SOCS) in the immune system. *Nat Rev Immunol* 2;410-416, 2002
28. Murray PJ: STAT3-mediated anti-inflammatory signalling. *Biochem Soc Trans* 34;1028-1031, 2006
29. Zhou RH, Kokame K, Tsukamoto Y, Yutani C, Kato H, Miyata T: Characterization of the human NDRG gene family: a newly identified member, NDRG4, is specifically expressed in brain and heart. *Genomics* 73;86-97, 2001
30. Choi SC, Kim KD, Kim JT, Kim JW, Lee HG, Kim JM, Jang YS, Yoon DY, Kim KI, Yang Y, Cho DH, Lim JS: Expression of human NDRG2 by myeloid dendritic cells inhibits down-regulation of activated leukocyte cell adhesion molecule (ALCAM) and contributes to maintenance of T cell stimulatory activity. *J Leukoc Biol* 83;89-98, 2008
31. Bradshaw CD, Ella KM, Qi C, Sansbury HM, Wisheart-Johnson AE, Meier KE: Effects of phorbol ester on phospholipase D and mitogen-activated protein kinase activities in T-lymphocyte cell lines. *Immunol Lett* 53;69-76, 1996
32. Schultz H, Engel K, Gaestel M: PMA-induced activation of the p42/44ERK- and p38RK-MAP kinase cascades in HL-60 cells is PKC dependent but not essential for differentiation to the macrophage-like phenotype. *J Cell Physiol* 173;310-318, 1997
33. Sengupta TK, Talbot ES, Scherle PA, Ivashkiv LB: Rapid inhibition of interleukin-6 signaling and Stat3 activation mediated by mitogen-activated protein kinases. *Proc Natl Acad Sci U S A* 95;11107-11112, 1998
34. Terstegen L, Gatsios P, Bode JG, Schaper F, Heinrich PC, Graeve L: The inhibition of interleukin-6-dependent STAT activation by mitogen-activated protein kinases depends on tyrosine 759 in the cytoplasmic tail of glycoprotein 130. *J Biol Chem* 275;18810-18817, 2000
35. Terstegen L, Maassen BG, Radtke S, Behrmann I, Schaper F, Heinrich PC, Graeve L, Gatsios P: Differential inhibition of IL-6-type cytokine-induced STAT activation by PMA. *FEBS Lett* 478;100-104, 2000
36. Lagunas L, Clipstone NA: Deregulated NFATc1 activity

- transforms murine fibroblasts via an autocrine growth factor-mediated Stat3-dependent pathway. *J Cell Biochem* 108;237-248, 2009
37. Wang T, Niu G, Kortylewski M, Burdelya L, Shain K, Zhang S, Bhattacharya R, Gabrilovich D, Heller R, Coppola D, Dalton W, Jove R, Pardoll D, Yu H: Regulation of the innate and adaptive immune responses by Stat-3 signaling in tumor cells. *Nat Med* 10;48-54, 2004
 38. Borland G, Bird RJ, Palmer TM, Yarwood SJ: Activation of protein kinase Calpha by EPAC1 is required for the ERK- and CCAAT/enhancer-binding protein beta-dependent induction of the SOCS-3 gene by cyclic AMP in COS1 cells. *J Biol Chem* 284;17391-17403, 2009
 39. Boyle K, Robb L: The role of SOCS3 in modulating leukaemia inhibitory factor signalling during murine placental development. *J Reprod Immunol* 77;1-6, 2008
 40. Fukushima A, Kajiya H, Izumi T, Shigeyama C, Okabe K, Anan H: Pro-inflammatory cytokines induce suppressor of cytokine signaling-3 in human periodontal ligament cells. *J Endod* 36;1004-1008, 2010
 41. Dong Q, Fan R, Zhao S, Wang Y: Over-expression of SOCS-3 gene promotes IL-10 production by JEG-3 trophoblast cells. *Placenta* 30;11-14, 2009
 42. Berlato C, Cassatella MA, Kinjyo I, Gatto L, Yoshimura A, Bazzoni F: Involvement of suppressor of cytokine signaling-3 as a mediator of the inhibitory effects of IL-10 on lipopolysaccharide-induced macrophage activation. *J Immunol* 168;6404-6411, 2002
 43. Niemand C, Nimmesgern A, Haan S, Fischer P, Schaper F, Rossaint R, Heinrich PC, Müller-Newen G: Activation of STAT3 by IL-6 and IL-10 in primary human macrophages is differentially modulated by suppressor of cytokine signaling 3. *J Immunol* 170;3263-3272, 2003