

Nucleoporin MOS7/Nup88 contributes to plant immunity and nuclear accumulation of defense regulators

Marcel Wiermer,^{1,†} Hugo Germain,^{1,‡} Yu Ti Cheng,^{1,2} Ana V. García,³ Jane E. Parker³ and Xin Li^{1,2,*}

¹Michael Smith Laboratories; and ²Department of Botany; University of British Columbia; Vancouver, BC Canada; ³Department of Plant-Microbe Interactions; Max-Planck Institute for Plant Breeding Research; Cologne, Germany

Current addresses: ¹Department of Plant Cell Biology; Albrecht-von-Haller Institute for Plant Sciences; Georg August University Göttingen; Göttingen, Germany; ²Natural Resources Canada; Canadian Forest Service; Laurentian Forestry Centre; Stn. Sainte-Foy, QC Canada

Key words: plant immunity, nucleoporins, nucleocytoplasmic trafficking, *snc1*, *mos*

Submitted: 02/26/10

Accepted: 03/30/10

Previously published online:
www.landesbioscience.com/journals/nucleus/article/12109

*Correspondence to: Xin Li;
Email: xinli@interchange.ubc.ca

Extra View to: Cheng YT, Germain H, Wiermer M, Bi D, García AV, Wirthmueller L, et al. Nuclear pore complex component MOS7/Nup88 is required for innate immunity and nuclear accumulation of defense regulators in Arabidopsis. *Plant Cell* 2009; 21:2503–16; PMID: 19700630; DOI: 10.1105/tpc.108.064519.

Controlled nucleocytoplasmic trafficking is an important feature for fine-tuning signaling pathways in eukaryotic organisms. Nuclear pore complexes (NPCs) composed of nucleoporin proteins (Nups) are essential for the exchange of macromolecules across the nuclear envelope. A recent genetic screen in our laboratory identified a partial loss-of-function mutation in Arabidopsis *MOS7/Nup88* that causes defects in basal immunity, Resistance (R) protein-mediated defense and systemic acquired resistance. In *Drosophila* and mammalian cells, exportin-mediated nuclear export of activated Rel/NFκB transcription factors is enhanced in *nup88* mutants resulting in immune response failure. Consistent with Nup88 promoting nuclear retention of NFκB, our functional analyses revealed that *MOS7/Nup88* is required for appropriate nuclear accumulation of the autoactivated R protein *snc1*, as well as the key immune regulators EDS1 and NPR1. These results suggest that controlling the nuclear concentrations of specific immune regulators is fundamental for defining defense outputs.

Eukaryotes have evolved elaborate immune systems allowing them to discriminate between self and non-self. In animals and plants, innate immune responses of individual cells constitute a major barrier to pathogen infection. There are two levels of innate immunity in the plant kingdom. The first, termed PAMP-triggered immunity (PTI),

is mediated by cell surface-resident Pattern Recognition Receptors (PRRs) that sense conserved pathogen associated molecular patterns (PAMPs). The second, pathogen-specific branch of the immune system, known as effector-triggered immunity (ETI), recognizes and responds to isolate-specific microbial effector molecules or their actions on host molecular targets.^{1,2} Pathogen effectors are often secreted into host cells during infection to increase virulence, in part by suppressing PTI. Recognition of specific effectors is conferred by R proteins. Most characterized R proteins are intracellular and have conserved Nucleotide-Binding and Leucine-Rich-Repeat (NB-LRR) domains that are also found in animal NOD (Nucleotide-Binding/Oligomerization Domain)-LRR immune receptors.

The role of the NPC and nucleocytoplasmic trafficking machinery in plant innate immunity was first revealed in our Arabidopsis genetic screen aimed at identifying components contributing to auto-immune responses mediated by the deregulated Toll-Interleukin-1 Receptor (TIR)-type NB-LRR *R* gene *snc1* (*suppressor of npr1-1, constitutive 1*).^{3,4} Auto-activation of *snc1* is caused by a point mutation resulting in an E₅₅₂K change in the linker region between the NB and LRR domains.³ As a consequence, *snc1* mutant plants are dwarf, accumulate high levels of the defense hormone salicylic acid (SA) and exhibit enhanced disease resistance to virulent pathogens.^{5,3} Intriguingly, certain mutations in the same region of human

immune NOD-LRR receptor NOD2 also result in constitutive activation and are associated with a chronic inflammatory disorder known as Crohn's disease.⁶

The *snc1* suppressor screen was designed to identify *modifier of snc1* (*mos*) mutants that resemble wild-type morphology and abolish constitutive pathogen resistance in *snc1*.⁷ We previously reported the isolation of *MOS3*,⁴ encoding the homolog of vertebrate Nup96 implicated in immunity-related mRNA export in mice and the importin α *MOS6*.^{8,9} *MOS3* and *MOS6* point to a requirement for mRNA export and nuclear localization signal (NLS)-mediated protein nuclear import pathways in plant immunity.

A defining feature of eukaryotic cells is the physical separation of the nucleoplasm from the cytoplasm by the double lipid bilayer of the nuclear envelope (NE). This separation necessitates a gateway and an elaborate trafficking machinery to facilitate the coordinated exchange of information between the two compartments. At the same time, this compartmentalization provides eukaryotic cells with a potent means of spatial and temporal control of signaling events. For example, restraining nuclear factor- κ B (NF κ B) transcription factors (TFs) outside the nucleus through association with inhibitory I τ B proteins in the cytoplasm represents a major regulatory step in animal innate immunity. In uninduced cells I κ B α /NF κ B complexes shuttle between the nucleus and the cytoplasm but steady-state localization appears to be almost exclusively cytosolic due to dominant NES-dependent nuclear export over nuclear localization signal (NLS)-dependent nuclear import.^{10,11} Inducing stimuli such as ligand recognition of certain PRRs, trigger proteasome-mediated I κ B degradation that alters the dynamic balance between I κ B α /NF κ B cytosolic and nuclear localization signals to favor nuclear accumulation of released NF κ B dimers and transcription of target genes.¹²

Recent data in *Drosophila* and mammalian cells suggest an additional mechanism controlling the nuclear accumulation of NF κ B and Rel-like TFs at the level of nucleocytoplasmic transport that is dependent on the dynamic inhibition

of NF κ B nuclear export rates by the nuclear pore complex protein Nup88 (Xylourgidis, et al. 2006). Nuclear pore complexes (NPCs) form numerous perforations in the NE and are composed of multiple copies of ~30 distinct Nups. Nups are modularly assembled in distinct subcomplexes of defined composition and arranged radially around a central channel that serves as the sole conduit and dynamic barrier for the selective bidirectional exchange of molecular cargoes to and from the nucleus. Translocation typically depends on the recognition of NLS and/or NES motifs on the cargo by nuclear transport receptors (NTRs) of the karyopherin family that facilitate nuclear import (importins) or export (exportins).

In *Drosophila*, the exportin CRM1 mediates nuclear export of Dorsal and Dif, members of the Rel protein family which includes NF κ B. Activation of the innate immune response in *Drosophila* is dependent on the nuclear activity of Dorsal and Dif that accumulate in the nucleus upon degradation of the I κ B homolog Cactus (Fig. 1A). Notably, *members only* (*mbo*) mutants, encoding the *Drosophila* homolog of vertebrate Nup88, fail to accumulate Dorsal and Dif in the nucleus or activate an effective immune response upon bacterial infection.¹³ Nup88 is localized on the cytoplasmic side of the NPC where it associates with Nup214 (Fig. 1A). Since the Nup88/Nup214 complex appears to sequester CRM1 at the NE, the cytoplasmic localization of Dorsal and Dif in *mbo* mutants likely results from excess CRM1 cargo export activity. This suggests that Nup88 acts as an attenuator of CRM1-mediated nuclear protein export to modulate the expression of NF κ B/Rel target genes, thereby controlling the relative strength and duration of innate immune responses at the level of the NPC (Fig. 1A).^{14,15} Consistent with Nup88 promoting nuclear retention of Rel proteins, depletion of Nup88 in mammalian cells prevents nuclear accumulation of NF κ B and inhibits NF κ B-dependent target gene expression whereas enhanced expression of Nup88 in malignant melanoma cells might contribute to constitutive NF κ B activation.¹⁶

We recently identified and functionally characterized *mos7-1*, a partial

loss-of-function mutation in the *Arabidopsis* homolog of vertebrate and *Drosophila* Nup88, which fully suppresses all known autoimmune phenotypes of *snc1*.¹⁷ *mos7-1* single mutant plants exhibit defects not only in basal defense against virulent pathogens and ETI conditioned by several NB-LRR R proteins of both the TIR- and Coiled-Coil (CC)-type, but also in systemic acquired resistance (SAR). SAR represents a long-lasting and broad spectrum disease resistance that is induced throughout the plant after a local immune response and accumulation of SA.¹⁸ SAR induction is mediated by the transcriptional coactivator Nonexpressor of Pathogenesis-Related genes 1 (NPR1). In uninduced tissues NPR1 is retained partially in the cytoplasm as a homooligomeric complex formed through intermolecular disulfide bonds (Fig. 1C). In response to pathogen attack, a change in the cellular redox potential leads to thioredoxin-mediated reduction of disulfide bonds allowing nuclear accumulation of NPR1 monomers, possibly due to the exposure of normally obscured NLSs (Mou, et al. 2003; Tada, et al.^{19,20} Inside the nucleus, NPR1 interacts with members of the TGA family of bZIP TFs to regulate downstream *Pathogenesis-Related* (*PR*) gene expression (Fig. 1C).^{21,22,23}

Since *mos7-1* mutant plants are impaired in SAR and the nuclear release of NPR1 is reminiscent of NF κ B signaling in animal immunity, we investigated the contribution of *MOS7* to NPR1 nuclear accumulation. We found that significantly lower amounts of a translational NPR1-GFP fusion expressed under control of the native *NPR1* promoter accumulated in nuclei of *mos7-1* than in wild type transgenic plants before and after SAR induction.¹⁷ As NPR1 nuclear accumulation is essential for SAR, NPR1 may not be able to attain sufficient threshold abundance in the nucleus for activation of SAR in *mos7-1*. Our cellular fractionation further suggests that a small portion of the cellular NPR1 pool is present in nuclei of uninduced tissues. Recent work by Spoel, et al. shows that, indeed, NPR1 monomer continuously translocates to the nucleus in uninduced tissues at a low rate. Here NPR1 amounts are kept low via proteasome-mediated degradation to prevent




Figure 1. Nup88 promotes nuclear retention of animal and plant immune regulators. (A) The *Drosophila* homolog of vertebrate Nup88, Members only (Mbo), is required for nuclear accumulation of the Rel-type transcription factors Dorsal and Dif. Toll receptor signaling upon microbial infection releases the NF- κ B homologs Dorsal/Dif from the inhibitory I κ B homolog Cactus, allowing Dorsal/Dif nuclear translocation. During this immune response, Mbo/Nup88 attenuates nuclear export of Dorsal/Dif by sequestering the Dorsal/Dif-loaded export receptor CRM1 at the nuclear rim. This attenuation in nuclear export results in nuclear accumulation of Dorsal/Dif and efficient defense gene expression. In contrast, *mbo* mutant animals show enhanced nuclear export of Dorsal/Dif and fail to activate an immune response (Roth et al. 2003; Uv et al. 2000; Xylourgidis et al. 2006).^{14,13,15} (B) In *Arabidopsis*, an E₅₅₂K change in SNC1 renders this TIR-NB-LRR immune receptor constitutively active without a pathogen stimulus.³ Autoactive *snc1* localizes to the cytoplasm and the nucleus and possibly shuttles between the two compartments via NLS/Importin-mediated nuclear import and NES/XPO1-mediated nuclear export. Autoimmunity of *snc1* requires functional MOS7 (Nup88) since a partial loss-of-function mutation in *mos7-1* results in XPO1-mediated nuclear leakage of *snc1* and suppression of all known *snc1* auto-immune phenotypes. This is consistent with MOS7 promoting nuclear retention and accumulation of autoactive *snc1* as a critical process in the constitutive activation of immune responses.¹⁷ (C) MOS7 (Nup88) is also required for proper nuclear accumulation of the plant-specific defense regulators NPR1 and EDS1. NPR1 translocates to nuclei of uninduced tissues at a low rate.²⁴ Induction by the plant defense hormone SA and subsequent changes in the cellular redox state promote NPR1 monomerization via thioredoxin-mediated reduction of intermolecular disulfide bonds and nuclear accumulation of NPR1 monomers, a process required for efficient expression of NPR1-regulated defense genes and induction of systemic immunity.¹⁹ EDS1 is a key component of immunity triggered by *snc1* and other intracellular TIR-NB-LRR immune receptors.^{25,5} Nucleocytoplasmic EDS1 forms several distinct defense regulatory complexes in the cytoplasm and the nucleus (not shown) and is capable of XPO1-mediated nuclear export.^{26,27} Altered nuclear translocation rates of NPR1 and EDS1 in *mos7-1* might affect NPR1 and EDS1 protein stability that becomes sensed and equilibrated across the NE over time.

target gene transcription in uninduced cells.²⁴

As *mos7-1* suppresses *snc1* fully whereas *npr1* does not, we reasoned that MOS7 must regulate nuclear traffic of further components in *snc1*-triggered resistance besides increasing nuclear NPR1 levels to induce SAR. To gain further mechanistic insight to the function of MOS7 in innate immunity, we investigated the effect of *mos7-1* on the subcellular distribution of another nucleocytoplasmic immune regulator, Enhanced Disease Susceptibility1 (EDS1), which is an indispensable component of *snc1* auto-immune responses and able to pass through NPCs via the *Arabidopsis* CRM1 homolog XPO1.^{26,27,28,5}

As with NPR1, EDS1 total protein was reduced in *mos7-1* mutant plants compared to wild type.¹⁷ Reduced EDS1 and NPR1 accumulation was not observed at the transcript level indicating a major effect of *mos7-1* on protein synthesis or stability. In *mos7-1*, the ratio of EDS1 distribution in the cytosol and nucleus was not strongly affected, but overall lower accumulation of EDS1 resulted in very low levels being detected in nuclei of *mos7-1*. This likely contributes to the ability of *mos7-1* to suppress *snc1* because interfering with nuclear accumulation of EDS1 impairs resistance mediated by another TIR-type NB-LRR receptor, RPS4.²⁷ Moreover, pathogen activation of RPS4 resistance triggers an

early increase in the nuclear EDS1 pool that directs EDS1-dependent changes in defense-related gene expression. However, such perturbations of nuclear EDS1 levels apparently become sensed and equilibrated with the EDS1 cytoplasmic pool which is also required for full resistance.²⁷ Therefore, reduced nuclear retention of EDS1 in *mos7-1* might perturb proper coordination of EDS1 pools across the NE and thereby alter EDS1 protein stability. This would lower the amount in the cytosolic pool available for nuclear import and result in the observed proportional depletion of EDS1 in both nuclear and cytoplasmic compartments in *mos7-1* (Fig. 1C).¹⁷

Two EDS1-dependent immune receptors, Arabidopsis RPS4 and tobacco N, have been shown to partially localize to and function inside the plant nucleus to trigger immune responses upon activation by their cognate pathogen effector.^{29,30} The specific effector detected by wild-type SNC1 in nature is unknown, but the protein contains a predicted NLS and two predicted NES motifs suggesting it might serve as karyopherin cargo substrate for bidirectional transport through NPCs (Fig. 1B).¹⁷ Since *mos7-1* was isolated as a genetic suppressor of *snc1* and *MOS7* is required for the expression of all known *snc1* phenotypes, we investigated whether the constitutively active immune receptor *snc1* shows a *MOS7*-dependent cellular distribution. A functional, *snc1*-GFP fusion protein expressed under the native *SNC1* promoter localizes to the cytoplasm and the nucleus of transgenic plants.¹⁷ While total amounts of *snc1*-GFP in *mos7-1* and wild type are similar, the ratio of cytoplasmic to nuclear localized *snc1*-GFP is significantly increased in *mos7-1*, suggesting that resistance conferred by this auto-activated R protein depends on attaining a sufficient concentration in the nucleus (Fig. 1B). In support of this hypothesis, enhancing nuclear export of *snc1*-GFP through translational fusion to an additional NES reduces its autoimmunity, a similar effect as observed in *mos7-1*.¹⁷ In summary, these results suggest a function of *MOS7* in promoting the nuclear retention and thus accumulation of auto-active *snc1* as an important step in defense activation.

We previously proposed a model in which nuclear activation of some nucleocytoplasmic NB-LRR receptors is a consequence of enhanced nuclear import and/or decreased nuclear export of the activated R protein, resulting in efficient defense gene expression after the nuclear R protein concentration reaches a certain threshold. Effector-mediated activation could either expose an obscured NLS via a conformational change or recruit additional interacting proteins to alter nuclear shuttling or retention. This model is based on our finding from the *mos* screen that defects in the NPC or nucleocytoplasmic transport machinery alter resistance responses, and upsetting the ratio of cytoplasmic-to

nuclear-localized R protein pools or their downstream regulators causes an immune deficiency in *mos7-1* (Cheng, et al. 2009; Palma, et al. 2008; Wiermer, et al. 2007).^{17,31,32} Based on previous work showing direct association between the CC-NB-LRR R protein MLA and plant-specific WRKY transcription factors, which suggests a direct link between certain R protein activation and transcriptional reprogramming,³³ we speculate that *snc1* may be able to target components of the transcriptional machinery inside the nucleus. Whether nuclear accumulation of wild type *SNC1* is required for defense activation and what kind of proteins *snc1*/*SNC1* interacts with inside nuclei awaits future investigation.

One question that remains is how specific *MOS7* functions are to plant immune responses, since perturbations in a conserved housekeeping machinery such as the NPC would be expected to negatively impact a number of signaling pathways. The role of *MOS7* in multiple branches of plant immunity, such as basal, systemic acquired and NB-LRR R protein-triggered resistance, suggests it may be more broadly required for modulating the transit of additional, yet unknown, factors during innate immunity. A role for *MOS7* as a rather general attenuator of XPO1/CRM1-mediated export is supported by the fact that a chimeric nucleocytoplasmic shuttle protein²⁸ shows enhanced NES-dependent nuclear export in *mos7-1* compared with wild type after transient transfection into Arabidopsis mesophyll protoplasts.¹⁷ Moreover, null mutations in *MOS7* are lethal, consistent with a lethality phenotype of null *mbol/nup88* mutations in *Drosophila*,¹³ indicating that wild type *MOS7*/*Mbo* is necessary for cargo-bound XPO1/CRM1-translocation of unknown proteins required for proper growth and development.

On the other hand, the phenotype of *mos7-1* mutants appears to be surprisingly selective. *mos7-1* plants do not show obvious pleiotropic defects in development, salt tolerance or plant hormone responses.¹⁷ Also, the subcellular localization and cellular abundance of several nuclear, nucleocytoplasmic or cytoplasmic proteins is unaffected in *mos7-1*, suggesting that the defects of *mos7-1* in nuclear

retention of *snc1*, NPR1 and EDS1 are rather specific.¹⁷ It is possible that the four amino acid deletion in *mos7-1* causes a slight change in protein conformation so that it affects the nuclear pore structure in a way that most acutely influences export of immunity-related nucleocytoplasmic regulators. One intriguing piece of data we recently obtained indicates that overexpression not only of wild type *MOS7*, but also *mos7-1*, can revert *snc1 mos7-1* back to *snc1* phenotypes (Cheng Y and Li X, unpublished). This suggests a more complex model in which *MOS7* may also contribute to the regulation of numbers of pores on the nuclear envelope. Thus, overexpression of *mos7-1* may help to revert the defects of *mos7-1* by increasing the frequency of pores on the envelope. At present this idea is purely speculative and future structural research on plant Nups with cell biology tools will enable us to test these hypotheses more directly.

It is conceivable that *MOS7* modulates the transit of certain XPO1/CRM1-cargo complexes upon activation of immune responses. This suggests that nuclear retention either depends on the conformation of a given cargo-NTR complex which determines its binding sites within and thus transport route through the NPC or that retention depends on additional pathway-specific signals that might alter the properties of *MOS7* to control NTR-cargo translocation. Such mechanisms dovetail with recent evidence that NTR conformation can be influenced by its specific cargo substrate³⁴ and that nuclear transport rates for a given cargo are regulated by altering Nup properties through posttranslational modifications.³⁵ Multiple signaling pathways through the NPC could thus be controlled by distinct Nups and functionally independent routes through the NPC might be directed by providing different binding sites for cargo-bound NTRs within the NPC.

In conclusion, the studies on Nup88 in *Drosophila*, mammals and Arabidopsis support a conserved function of Nup88 in modulating host immunity (Fig. 1). While the overall mechanisms of nucleocytoplasmic transport and the structural organization of the NPC are conserved among eukaryotes, some components of the NPC and transport mechanisms seem to be

unique to plants.^{36,37} Research on Nups in plant species is an emerging field. The genetic resources available in Arabidopsis provide huge potential for revealing the biological functions of plant Nups, many of which await characterization.

Acknowledgements

We thank Jacqueline Monaghan for careful reading of the manuscript. We are grateful for a postdoctoral fellowship from the Alexander von Humboldt Foundation to M.W., Le Fonds Québécois de la Recherche sur la Nature et les Technologies to H.G., a Natural Sciences and Engineering Research Council of Canada (NSERC) post-graduate scholarship to Y.T.C., and funds from NSERC, the Michael Smith Laboratories (MSL) and the Canadian Foundation for Innovation (CFI) to X.L. J.E.P. acknowledges support of DFG SFB 670 and an IMPRS fellowship to A.V.G.

References

- Jones JD, Dangl JL. The plant immune system. *Nature* 2006; 444:323-9.
- Lukasik E, Takken FL. STANDING strong, resistance proteins instigators of plant defence. *Curr Opin Plant Biol* 2009; 12:427-36.
- Zhang Y, Goritschnig S, Dong XN, Li X. A gain-of-function mutation in a plant disease resistance gene leads to constitutive activation of downstream signal transduction pathways in suppressor of *npr1-1*, constitutive 1. *Plant Cell* 2003; 15:2636-46.
- Zhang Y, Li X. A Putative Nucleoporin 96 Is Required for Both Basal Defense and Constitutive Resistance Responses Mediated by suppressor of *npr1-1*, constitutive 1. *Plant Cell* 2005; 17:1306-16.
- Li X, Clarke JD, Zhang YL, Dong XN. Activation of an EDS1-mediated R-gene pathway in the *sncl* mutant leads to constitutive, NPR1-independent pathogen resistance. *Mol Plant Microbe Interact* 2001; 14:1131-9.
- Yamamoto S, Ma X. Role of Nod2 in the development of Crohn's disease. *Microbes Infect* 2009; 11:912-8.
- Monaghan J, Germain H, Weihmann T, Li X. Dissecting plant defence signal transduction—modifiers of *sncl* in Arabidopsis. *Can J Plant Pathol* 2010; 32:35-42.
- Faria AM, Levay A, Wang Y, Kamphorst AO, Rosa ML, Nussenzweig DR, et al. The nucleoporin Nup96 is required for proper expression of interferon-regulated proteins and functions. *Immunity* 2006; 24:295-304.
- Palma K, Zhang Y, Li X. An Importin α Homolog, MOS6, Plays an Important Role in Plant Innate Immunity. *Curr Biol* 2005; 15:1129-35.
- Huang TT, Kudo N, Yoshida M, Miyamoto S. A nuclear export signal in the N-terminal regulatory domain of I κ B α controls cytoplasmic localization of inactive NF κ B/I κ B α complexes. *Proc Natl Acad Sci USA* 2000; 97:1014-9.
- Malek S, Chen Y, Huxford T, Ghosh G. I κ B β , but Not I κ B α , Functions as a Classical Cytoplasmic Inhibitor of NF κ B Dimers by Masking both NF κ B Nuclear Localization Sequences in Resting Cells. *J Biol Chem* 2001; 276:45225-35.
- Hayden MS and Ghosh S. Shared principles in NF κ B signaling. *Cell* 2008; 132:344-62.
- Uv AE, Roth P, Xylourgidis N, Wickberg A, Cantera R, Samakovlis C. members only encodes a Drosophila nucleoporin required for Rel protein import and immune response activation. *Genes Dev* 2000; 14:1945-57.
- Roth P, Xylourgidis N, Sabri N, Uv A, Fornerod M, Samakovlis C. The Drosophila nucleoporin DNup88 localizes DNup214 and CRM1 on the nuclear envelope and attenuates NES-mediated nuclear export. *J Cell Biol* 2003; 163:701-6.
- Xylourgidis N, Roth P, Sabri N, Tsarouhas V, Samakovlis C. The nucleoporin Nup214 sequesters CRM1 at the nuclear rim and modulates NF κ B activation in Drosophila. *J Cell Sci* 2006; 119:4409-19.
- Takahashi N, van Kilsdonk JW, Ostendorf B, Smeets R, Bruggeman SW, Alonso A, et al. Tumor marker nucleoporin 88 kDa regulates nucleocytoplasmic transport of NF κ B. *Biochem Biophys Res Commun* 2008; 374:424-30.
- Cheng Y, Germain H, Wiermer M, Bi D, García AV, Wirthmueller L, et al. Nuclear Pore Complex Component MOS7/Nup88 Is Required for Innate Immunity and Nuclear Accumulation of Defense Regulators in Arabidopsis. *Plant Cell* 2009; 21:2503-16.
- Durrant WE, Dong X. Systemic acquired resistance. *Annu Rev Phytopathol* 2004; 42:185-209.
- Mou Z, Fan WH, Dong XN. Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* 2003; 113:935-44.
- Tada Y, Spoel SH, Pajeroska-Mukhtar K, Mou Z, Song J, Wang C, et al. Plant Immunity Requires Conformational Changes of NPR1 via S-Nitrosylation and Thioredoxins. *Science* 2008; 321:952-6.
- Després C, DeLong C, Glaze S, Liu E, Fobert PR. The Arabidopsis NPR1/NIM1 protein enhances the DNA binding activity of a subgroup of the TGA family of bZIP transcription factors. *Plant Cell* 2000; 12:279-90.
- Rochon A, Boyle P, Wignes T, Fobert PR, Després C. The Coactivator Function of Arabidopsis NPR1 Requires the Core of Its BTB/POZ Domain and the Oxidation of C-Terminal Cysteines. *Plant Cell* 2006; 18:3670-85.
- Zhang Y, Fan W, Kinkema M, Li X, Dong X. Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the PR-1 gene. *Proc Natl Acad Sci USA* 1999; 96:6523-8.
- Spoel SH, Mou Z, Tada Y, Spivey NW, Genschik P, Dong X. Proteasome-Mediated Turnover of the Transcription Coactivator NPR1 Plays Dual Roles in Regulating Plant Immunity. *Cell* 2009; 137:860-72.
- Aarts N, Metz M, Holub E, Staskawicz BJ, Daniels MJ, Parker JE. Different requirements for EDS1 and NDR1 by disease resistance genes define at least two R gene-mediated signaling pathways in Arabidopsis. *Proc Natl Acad Sci USA* 1998; 95:10306-11.
- Feys BJ, Wiermer M, Bhat RA, Moisan LJ, Medina-Escobar N, Neu C, et al. Arabidopsis SENESCENCE-ASSOCIATED GENE101 stabilizes and signals within an ENHANCED DISEASE SUSCEPTIBILITY1 complex in plant innate immunity. *Plant Cell* 2005; 17:2601-13.
- García AV, Blanvillain-Baufumé S, Huibers RP, Wiermer M, Li G, Rietz S, et al. Balanced nuclear and cytoplasmic activities of EDS1 are required for a complete plant innate immune response. *Plos Pathogens* 2010; In press.
- Haasen D, Kohler C, Neuhaus G, Merkle T. Nuclear export of proteins in plants: AtXPO1 is the export receptor for leucine-rich nuclear export signals in *Arabidopsis thaliana*. *Plant Journal* 1999; 20:695-705.
- Burch-Smith TM, Schiff M, Caplan JL, Tsao J, Czymbek K, Dinesh-Kumar SP. A Novel Role for the TIR Domain in Association with Pathogen-Derived Elicitors. *PLoS Biol* 2007; 5:68.
- Wirthmueller L, Zhang Y, Jones JD, Parker JE. Nuclear Accumulation of the Arabidopsis Immune Receptor RPS4 Is Necessary for Triggering EDS1-Dependent Defense. *Curr Biol* 2007; 17:2023-9.
- Palma K, Wiermer M, Li X. Marshalling the troops: Intracellular dynamics in plant pathogen defense. *Annual Plant Reviews* 2008; 34:177-219.
- Wiermer M, Palma K, Zhang Y, Li X. Should I stay or should I go? Nucleocytoplasmic trafficking in plant innate immunity. *Cell Microbiol* 2007; 9:1880-90.
- Shen QH, Saijo Y, Mauch S, Biskup C, Bieri S, Keller B, et al. Nuclear activity of MLA immune receptors links isolate-specific and basal disease-resistance responses. *Science* 2007; 315:1098-103.
- Wohlwend D, Strasser A, Dickmanns A, Ficner R. Structural basis for RanGTP independent entry of spliceosomal U snRNPs into the nucleus. *J Mol Biol* 2007; 374:1129-38.
- Kosako H, Yamaguchi N, Aranami C, Ushiyama M, Kose S, Imamoto N, et al. Phosphoproteomics reveals new ERK MAP kinase targets and links ERK to nucleoporin-mediated nuclear transport. *Nat Struct Mol Biol* 2009; 16:1026-35.
- Lu Q, Tang X, Tian G, Wang F, Liu K, Nguyen V, et al. Arabidopsis homolog of the yeast TREX-2 mRNA export complex: components and anchoring nucleoporin. *Plant* 2010; 61:259-70.
- Zhao Q, Brkljacic J, Meier I. Two Distinct Interacting Classes of Nuclear Envelope-Associated Coiled-Coil Proteins Are Required for the Tissue-Specific Nuclear Envelope Targeting of Arabidopsis RanGAP. *Plant Cell* 2008; 20:1639-51.