

Interaction of Transcriptional Repressor ArgR with Transcriptional Regulator FarR at the *argB* Promoter Region in *Corynebacterium glutamicum*[∇]

Soo Youn Lee,^{1,2†} Jae-Min Park,^{3†} Jin Hyung Lee,² Suk-Tai Chang,⁴ Jin-Soo Park,⁵
Yang-Hoon Kim,^{3*} and Jiho Min^{1*}

Department of Bioprocess Engineering, Chonbuk National University, 664-14 Deokjin-dong, 1Ga Deokjin-Gu, Jeonju 561-756, South Korea¹; Bio-IT Convergence Center, Korea Institute of Ceramic Engineering & Technology, 103 Fashion Danji-gil, Geumcheon-gu, Seoul 153-801, South Korea²; Department of Microbiology, Chungbuk National University, 410 Sungbong-Ro, Heungduk-Gu, Cheongju 361-763, South Korea³; School of Chemical Engineering and Materials Science, Chung-Ang University, 221 Heukseok-dong, Dongjak-gu, Seoul 156-756, South Korea⁴; and Department of Environmental Engineering, College of Engineering, Sangmyung University, 300 Anseo-dong, Dongnam-gu, Cheonan 330-720, South Korea⁵

Received 7 July 2010/Accepted 17 November 2010

In *Corynebacterium glutamicum*, the ArgR protein, a transcriptional repressor, affects the expression level of the *argB* gene through binding to its promoter region. The *argB* promoter region (positions –77 to –25) has been found by *in vitro* electrophoretic mobility shift assay (EMSA) results and *in silico* analysis to be important for the DNA binding of ArgR. Proline supplementation prevented the DNA binding of ArgR to the *argB* promoter region and triggered an increase of the *argB* mRNA level. Additional mutational analyses of the *argB* promoter region found nucleotides critical for ArgR binding (G located at position –58, C at position –55, and A at position –41 of the *argB* promoter) in that region. Another transcriptional repressor, FarR, was also demonstrated to bind to the *argB* promoter region. This binding was delimited to positions –57 to –77 on the *argB* promoter. FarR has only one putative binding domain located at positions –57 to –77, but this region exactly overlapped with the binding region located from positions –55 to –77 for the binding of ArgR within the *argB* promoter; thus, if ArgR bound with the *argB* promoter first, the binding of FarR was not observed in this region. However, if FarR bound to the binding domain located at positions –57 to –77 first, ArgR could bind other binding sites located at positions –49 to –25 within the *argB* promoter. Finally, this study suggests that ArgR can affect FarR binding to the *argB* promoter region, as protein binding is dominated by the protein most able to do so.

The main regulator of a set of *arg* regulons (19), arginine repressor (ArgR), acts as a transcriptional repressor by binding to the hexameric structure of its target sequences, known as “ARG boxes”(7). Detailed studies of ArgR have been conducted with many bacteria, for example, *Escherichia coli* (19), *Pseudomonas aeruginosa* (22), and *Bacillus stearothermophilus* (6). In addition, the mechanism of ArgR is unusually well conserved across a wide range of divergent bacteria: both Gram-negative and Gram-positive bacteria (35).

Corynebacterium glutamicum is a Gram-positive soil bacterium widely used in the production of amino acids (18, 32). Bioinformatics tools have recently been used to detect the potential transcription regulators of winged helix-turn-helix (HTH) binding proteins, including ArgR, that were previously predicted from the genome sequence (1, 2, 11). Experimental

data and *in silico* analyses of a diverse range of bacteria show a surprising conservation of the arginine repressor proteins and their respective target sites. For instance, the identical 18.8-kDa polypeptides and the folded structure derived from the amino acid sequence of ArgR molecules from *C. glutamicum* closely match the winged-helix structures and N-terminal DNA-binding domains of several other species (14, 20).

Previous studies have shown that the biosynthesis of ornithine, an intermediate molecule of arginine’s biosynthesis, depends upon the DNA binding of ArgR to the operating regions of *arg* genes (14). In addition to its regulatory function, ArgR has a particular activity for the upstream region of the *argB* gene, encoding the *N*-acetylglutamate kinase enzyme, an enzyme relevant to ornithine biosynthesis in *C. glutamicum* (14, 16).

Fatty acyl-responsive regulator (FarR) (30), a previously uncharacterized transcription factor of the HTH GntR family (27) similar to HutC/FarR, seems to be involved in the regulation of amino acid biosynthesis in *C. glutamicum* (8). The transcription of the *argB* gene was influenced by FarR. Interestingly, both ArgR and FarR control ornithine and arginine levels by repressing the transcription of the *arg* genes (8, 14). However, the mechanism by which FarR stimulates *argB* transcription remains unclear.

This paper’s focus is the ArgR-mediated regulation of *argB*

* Corresponding author. Mailing address for Yang-Hoon Kim: Department of Microbiology, Chungbuk National University, 410 Sungbong-Ro, Heungduk-Gu, Cheongju 361-763, South Korea. Phone: 82-43-261-3575. Fax: 82-43-264-9600. E-mail: kyh@chungbuk.ac.kr. Mailing address for Jiho Min: Department of Bioprocess Engineering, Chonbuk National University, 664-14 Deokjin-dong, 1Ga Deokjin-Gu, Jeonju 561-756, South Korea. Phone: 82-63-270-2436. Fax: 82-63-270-2306. E-mail: jihomin@jbnu.ac.kr.

† S.Y.L. and J.-M.P. contributed equally to this work.

∇ Published ahead of print on 29 November 2010.

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Relevant characteristic(s)	Source or reference
Strains		
<i>C. glutamicum</i>	Wild type	ATCC 13032
<i>C. glutamicum</i> SJC 8074	Deletion of the <i>argR</i> , <i>argF</i> , and <i>proB</i> genes	10
<i>E. coli</i> BL21(DE3)	F ⁻ <i>ompT hsdS_B(r_B⁻ m_B⁻) gal dcm</i> (DE3)	Novagen, Madison, WI
Plasmids		
pET-21a	Ap ^r ; f1 origin; 6× histidyl fusion vector	Novagen, Madison, WI
pEMBTLSY0	pET-21a containing the <i>argR</i> structural gene	14
pEMBTLSY5	pET-21a containing the <i>farR</i> structural gene	13

expression in *C. glutamicum*. To establish how ArgR's operating site acts on the *argB* gene, the effects of ArgR on DNA-binding affinity were examined *in vitro*. The effects of the DNA-binding sites of FarR on the promoters of *arg* genes were also analyzed by *in vivo* chromatin immunoprecipitation (ChIP) assays of *C. glutamicum*. Furthermore, a new relationship of the two transcriptional regulators FarR and ArgR to the *argB* promoter is provided through the detailed analysis of the interaction of FarR with the promoter site of the *argB* gene.

MATERIALS AND METHODS

Bacterial strains, media, and growth conditions. Table 1 lists the bacterial strains and plasmids used in this study. Wild-type strain *C. glutamicum* ATCC 13032 (American Type Culture Collection, Manassas, VA) and mutant strain *C. glutamicum* SJC 8074 (provided by Sangji University, South Korea) (10) were grown at 30°C in Luria-Bertani medium (29) and, for the production of ornithine and arginine, in mineral medium containing yeast extract (MMY) [0.8 g KH₂PO₄, 10 g (NH₄)₂SO₄, 1 g MgSO₄ · 7H₂O, 1.2 g Na₂HPO₄, 20 mg MnSO₄ · H₂O, 20 mg FeSO₄ · 7H₂O, 10 mg ZnSO₄ · 7H₂O, 10 g yeast extract, 20 g CaCO₃, and 60 g glucose liter⁻¹] (17). In a 250-ml shake flask, a 1-ml preculture was inoculated into 50 ml of MMY medium. Cultivation was performed at 30°C at 150 rpm on a rotary shaker. *E. coli* BL21(DE3) cells (Novagen Chemicals, Inc., Germany) were grown at 37°C in Luria-Bertani medium (29). Selection for the presence of plasmids was carried out by using ampicillin (50 µg ml⁻¹ for *E. coli*). Shake flask culture growth was monitored by measuring the optical density at 600 nm (OD₆₀₀) using a UV/Vis spectrophotometer (Mecasys Co., Ltd., South Korea).

Genetic manipulations. Chromosomal DNA was prepared from *C. glutamicum* by using a Wizard SV genomic DNA kit (Promega, Madison, WI). Plasmid DNA was prepared from *E. coli* cells using an alkaline lysis technique with a QIAspin miniprep kit (Qiagen, Germany). DNA modification, analysis by agarose gel electrophoresis, and ligation were performed according to standard procedures (29). The PCR experiments were carried out by using a T Gradient thermocycler (Biometra, Germany), Ex *Taq* DNA polymerase (Takara Bio, Inc., Japan), and chromosomal DNA as the template. The PCR products were purified by using a QIAquick PCR purification kit (Qiagen, Germany). The oligonucleotides used for PCR amplification were purchased from AccuOligo (Bioneer Co., South Korea).

Purification of histidine-tagged ArgR and FarR. The construction of pEMBTLSY0 carrying the *argR* coding region and pEMBTLSY5 carrying the *farR* coding region were described in previous studies (Table 1) (13, 14). pEMBTLSY5 and pEMBTLSY0, were used to transform *E. coli* BL21(DE3) cells by electroporation. The syntheses of ArgR and FarR fused with a six-histidine tag were induced in recombinant *E. coli* BL21 cells (pEMBTLSY0 and pEMBTLSY5, respectively) by the addition of 0.5 mM isopropyl-β-D-thiogalactopyranoside (IPTG) after the culture had reached an OD₆₀₀ of 0.6. The cells were grown for 3 h, harvested, and then disrupted by using a Vibra Cell Sonic disruptor (Sonics & Materials, Inc.). Purification of the fusion proteins was carried out by Ni-nitrilotriacetic acid (NTA) affinity chromatography according to the instructions provided by Qiagen (Hilden, Germany). The purified fusion proteins were used directly for the production of polyclonal rabbit antibodies for

TABLE 2. Oligonucleotides used in this study

Primer and purpose	Sequence (5'-3')	Description (reference)
RT-PCR		
argB-F	ATATTGGTTTGGTCGGAGA	Amplification of <i>argB</i> cDNA (this study)
argB-R	TACAGTTCCCATCCTTGT	
16S rRNA-F	TCCTGGTGTAGCGGTGAAA	Amplification of 16S rRNA (this study)
16S rRNA-R	CCCACCTTCCTCCGAGTTA	
ChIP assay		
argC-F	TGCACTTCCAGGTGGT	PCR primers for ChIP detection of <i>arg</i> genes (14)
argC-R	AGTTACACCATACACG	
argJ-F	CTTAAGCGTTGGTTTTG	
argJ-R	CGGTAATGCCTTTTTCT	
argB-F	TCGAACCACTGACCTGA	
argB-R	CAGCGAGGACATTTGGG	
argF-F	TGGTGATCACCCGACGAA	
argF-R	AAACCTCTGCCTGCTCT	
argG-F	GCACCACTTAAAGCG	
argG-R	AGAACGATGCGGTTAG	
argH-F	CTCCAAGATCGCTAACA	
argH-R	TCCATGTGGTGTCTTC	

ChIP assays (Ab Frontier, Inc., South Korea) and electrophoretic mobility shift assay (EMSA) experiments.

EMSA. Short DNA probes that were approximately 30 bp long were generated by the annealing of complementary single oligonucleotides. For the labeling of the DNA and the setup of the reaction mixture containing purified His₆-ArgR or His₆-FarR for the EMSA, a digoxigenin (DIG) gel shift kit (Roche, Mannheim, Germany) was used according to the manufacturer's instructions. The oligonucleotides were blunt ended, since labeling were performed with terminal transferase and digoxigenin-11-dUTP (DIG-ddUTP). The labeled and unlabeled probes were incubated with His₆-ArgR or His₆-FarR. Binding reactions were performed with binding buffer [20 mM HEPES (pH 7.6), 1 mM EDTA, 10 mM (NH₄)₂SO₄, 1 mM dithiothreitol (DTT), 0.2% (wt/vol) Tween 20, 30 mM KCl, 1 µg poly(dI-dC), 0.2 µg poly-L-lysine] for 30 min at room temperature. Separation by gel electrophoresis was performed with native 6% polyacrylamide gels using 0.5× Tris-borate-EDTA (TBE) buffer as a running buffer. Subsequently, the labeled DNA was blotted onto a positively charged nylon membrane (Roche, Mannheim, Germany) by electroblotting. For the detection of the labeled DNA, X-ray films were used. Each sample was analyzed in triplicate.

DNA-protein cross-linking and ChIP assays. A previously described ChIP protocol (14) was adapted as follows. Individual strains were grown at 30°C for several generations in MMY medium. At an OD₆₀₀ of 1.9 to 2.0, samples (10 ml) from 100-ml cultures were transferred into new tubes and treated with formaldehyde to a final concentration of 1% and incubated for 10 min at 30°C with gentle agitation. The cells were harvested by centrifugation (10 min at 1,618 × g at 4°C) and washed twice in chilled phosphate-buffered saline. The cells were resuspended in 0.5 ml of a solution containing 1% sodium dodecyl sulfate, 10 mM EDTA, 50 mM Tris-HCl (pH 8.1), 1 mM phenylmethylsulfonyl fluoride, and 5 µg ml⁻¹ RNase A. The cells were then incubated at 30°C for 10 min and chilled on ice. The lysates were sonicated for 10 cycles, each lasting 20 s. The chromosomal DNA of the lysate was sheared to give a mean fragment size of 200 to 1,000 bp. The cell debris was used for the ChIP assay. The experiments were performed by using a ChIP assay kit (Upstate) according to the manufacturer's protocol, and the DNA-protein complexes in the supernatant were immunoprecipitated by using 3 µg ml⁻¹ affinity-purified polyclonal rabbit antibodies raised against the purified hexahistidine-tagged ArgR or FarR protein. Subsequently, PCR amplification of the immunoprecipitated DNA was carried out for 27 cycles using the primers designed to amplify the six *arg* genes (Table 2). Each sample was analyzed in triplicate.

RT-PCR. The levels of *argB* mRNA were quantified by real-time reverse transcription (RT)-PCR using SYBR green PCR master mix (ABI 7700; Applied Biosystems, CA). In brief, the total RNA from the same biomass of *C. glutamicum* was extracted by using TRIzol reagent (Gibco-BRL) according to the manufacturer's instructions. Reverse transcription was first performed to synthesize cDNA using total RNA (0.5 µg), random primers (16-mers) (Bioneer Co., Daejeon, South Korea), deoxynucleoside triphosphates (dNTPs) (1 mmol/liter), 4.5 units of avian myeloblastosis virus (AMV) reverse transcriptase (Promega Co.), and 20 units of RNase inhibitor (Promega Co.). cDNA corresponding to 50 ng of RNA was then added to SYBR green Ampli *Taq* master mix and 0.9 µmol/liter each specific primer in a total volume of 50 µl. Table 2 lists the primer pairs used for *argB* and the 16S



FIG. 1. Putative ARG box sequences. An alignment of the *argB* promoter region with the ARG box sequences in the *argJ*, *artP*, and *hisJ* promoter from *E. coli* (Ec) (3) using the ClustalX program (31) and a picture representing the conservation of bases at each position in the inferred consensus operator sequence (generated by WebLogo 2.8.2 software) (4) are shown. The conserved nucleotides in at least eight boxes are shaded and asterisked. Bases substituted in the *argB* promoter region used for EMSAs are indicated, and the new base is shown above. Cg, *C. glutamicum*.

rRNA gene (internal standard). PCR was carried out with a real-time PCR cycler (ABI 700; Applied Biosystems, CA). The thermal cycling conditions were 48°C for 30 min, 95°C for 10 min, and 40 cycles at 95°C for 15 s and 60°C for 1 min. At the end of each phase at 60°C, the fluorescence was measured and used for quantitation. Each sample was analyzed in triplicate.

Determination of amino acid concentration. High-performance liquid chromatography (HPLC) (Waters Alliance 2690 analytical HPLC system; Waters Co.) was used to determine amino acids. The system was equipped with a Nova-Pak C₁₄ column and a Waters 747 scanning fluorescence detector (Waters Co.). All the results represent the data from at least three independent experiments and include a mean value.

RESULTS

In vitro binding of ArgR to the *argB* promoter region. ArgR is a transcriptional regulator that controls ornithine biosynthesis through an interactive pathway of arginine biosynthesis in *C. glutamicum* (19). As previously reported, DNA sites for ArgR binding occur at four genes, located at the upstream region of *argC*, *argB*, *argF*, and *argG*. It was also shown previously that the DNA-binding affinity of ArgR for the upstream region of the *argB* gene is affected only by proline supplementation (14, 16). It was reported that ArgR not only recognizes but also binds weakly conserved operator sequences of the form 5'-TNTGAATwwwATTCANW-3' (where N is any base and W/w is A or T [with w in the linker region between two palindromic sequences of ArgR binding sites]) in *E. coli*, located in the three known arginine-inducible promoters (3, 19, 20). These promoters contain ARG boxes: the promoters of *artJ* (*P_{artJ}* Ec), *artPIQM* (*P_{artP}* Ec), and the *hisJQMP* operon (*P_{hisJ}* Ec) (3). An alignment of the three ARG boxes with the *argB* promoter region of *C. glutamicum* showed the 27-bp consensus sequence and highlighted the conserved nucleotides at each position (Fig. 1). To corroborate this finding further, EMSA was explored to see whether ArgR bound to the *argB* promoter region *in vitro* (Fig. 2). The two subfragments (subfragments B₁ and B₂) of the promoter regions of the *argB* gene were incubated with His₆-ArgR (Fig. 2A). Competition and supershift assays showed that ArgR did not bind to two other sites located at positions -102 to -72 and -27 to +3 under the promoter regions of *argB* (data not shown). These results defined the ArgR-responsive sequence as subfragments B₁ and B₂ located at positions -77 to -25 of the *argB* promoter. As shown in Fig. 2B (right), His₆-ArgR has clearly bound to the two subfragments. However, competitive assays showed that His₆-ArgR did not bind to the two subfragments. Therefore, this suggests that an important site for the DNA binding of

ArgR is between positions -77 and -25 of the *argB* promoter region.

The *argB* promoter region is regulated in response to proline. Previous studies showed that a reduced DNA-binding affinity of ArgR for the upstream region of the *argB* gene provokes an increase in levels of ornithine biosynthesis (14, 16). Thus, the DNA-binding activity of His₆-ArgR *in vitro* was analyzed by using the electrophoretic mobility shift assay (EMSA) technique, testing subfragments (subfragments B₁ and B₂) of the *argB* promoter region to verify whether the DNA binding of ArgR to the *argB* promoter is affected directly by proline (Fig. 3A). Consistent with previously reported results of *in vivo* experiments (14), the addition of proline had an effect on DNA binding by His₆-ArgR although dissolving the affinity of His₆-ArgR. Furthermore, to test whether proline influences the expression of the *argB* gene, the transcriptional activation of *argB* by proline was examined. Concurring with data from previous studies, it was clear that under the 10 mM proline-supplemented condition, the amount of *argB* mRNA increased (Fig. 3B). In conclusion, the results indicate that proline supplementation prevents the DNA binding of ArgR to subfragments B₁ and B₂ but triggers an increase in the level of *argB* transcription. Therefore, proline is a potent inducer of *argB* gene transcription regulated by ArgR.

Mutational analysis of ArgR-binding sites within the *argB* promoter region. Furthermore, to investigate which nucleotides within the consensus sequence were necessary for ArgR binding, single-nucleotide exchanges were made in the *argB* promoter region (Fig. 1). The most conserved bases were substituted to introduce transversion from AT to CG and from CG to AT: T₁→G, G₆→T, C₉→A, A₁₀→C, T₁₅→G, A₂₁→C, A₂₃→C, and A₂₇→C (Table 2). In addition, C₅, which was previously identified as the major position for ArgR binding by DNase I footprinting in *E. coli* (3), was also replaced with A. As shown in Fig. 4, His₆-ArgR clearly bound all DNA fragments except G₆T, C₉A, and A₂₃C. In particular, the fragments that contained C₉A and A₂₃C belonging to the highlighted and conserved nucleotides of Fig. 1 showed a very weak, or no, band shift compared with subfragments B₁ and B₂ as wild-type promoter regions of the *argB* gene (Fig. 2B). These results define the C₉ and A₂₃ nucleotides as major points of binding of ArgR to the *argB* promoter.

FarR is related to the biosynthesis of ornithine and arginine in *C. glutamicum*. Another transcriptional regulator, FarR, was found previously to be involved in the glutamate and arginine




FIG. 2. Binding of ArgR to the putative ARG box sequence in the *argB* promoter region. (A) Schematic representation of the upstream region of the *argB* gene. The numbers indicate the ends of the fragments relative to the proposed translational start site (position +1) in this study. Vertical arrows indicate the positions of base substitutions (Fig. 1). The box indicates the putative ARG box, the gray bar represents the putative recognition region of FarR binding, P_{argB} indicates the DNA fragment used for the *in vivo* DNA-binding affinities of ArgR (14) and FarR by ChIP assays, arrows indicate the bases substituted in the *argB* promoter region, and asterisks represent missing contact probing of ArgR. (B) EMSAs were performed with two subfragments (subfragments B₁ and B₂) of the promoter region. His₆-ArgR (2,400 nM) was incubated with the DIG-labeled probe, and the protein-DNA complexes were resolved by electrophoresis on native 6% polyacrylamide gels. Comp. denotes competitor assays (40-fold excess DIG-unlabeled oligonucleotides) of ArgR.

metabolic pathway in *C. glutamicum* (8). As shown in Fig. 5B, ChIP assays were used to measure the level of FarR binding to promoter regions of *arg* genes *in vivo*. For this purpose, putative promoter regions of six *arg* genes (*argC*, *argJ*, *argB*, *argF*, *argG*, and *argH*) were selected in accordance with previously

reported research (14). Compared with ChIP results and the growth curve of *C. glutamicum*, the DNA-binding affinity of FarR showed a gradual increase in trends at all target regions until an early stationary growth phase (Fig. 5). Interestingly, 12 to 14 h after inoculation, the productions of both ornithine and arginine reached a plateau (Fig. 5A). Additionally, the *argB* promoter region is more repressed by FarR than by other promoters during 12 h to 14 h due to the relative amount of DNA of FarR for each promoter region. In previous studies, another transcriptional repressor, ArgR, was shown to regulate ornithine biosynthesis by binding to the upstream regions of *arg* genes (15, 16). In particular, ornithine biosynthesis is highly regulated when ArgR acts on the *argB* promoter region under conditions of proline supplementation (14). This finding suggests that FarR also governs ornithine and arginine biosynthesis and the DNA binding of FarR to the *argB* promoter region, which is important for the physiological mechanisms.

***In vitro* binding of FarR to the *argB* promoter region.** According to data from previous studies, FarR binding to the *gdh* promoter region was confirmed by gel retardation experiments (positions -444 to -469 from the transcriptional start site) (8, 9). This study showed that FarR binds strongly to the *gdh* promoter region *in vivo* (data not shown). This site consists of a highly palindromic region with the sequence 5'-GCCAGGT TATATAACCAGTC-3' (8). Surprisingly, putative FarR-binding sites were located at positions -57 to -77 from the proposed translation start site (position +1) of the *argB* promoter (5'-ACGTGGAGATCAACTCCGCGT-3'), analyzed by multiple-sequence alignment with the FarR-binding site of the *gdh* promoter region and the consensus binding motif (Fig. 2A). HutC/FarR-type regulators of the GntR family bind by dimerization at the palindromic region of the DNA sequence in an




FIG. 3. Proline response of ArgR to the upstream regions of the *argB* gene. (A) DNA-binding activity of His₆-ArgR (2,400 nM) for subfragments B₁ and B₂ by EMSA in the presence of 0 mM, 1 mM, 5 mM, and 10 mM proline. (B) RT-PCR analysis of the *argB* gene in 10 mM proline-treated *C. glutamicum*. The mRNA expression level of the *argB* gene was calculated as a ratio of 16S rRNA gene expression. The results are reported as the means of data from three experiments.




FIG. 4. EMSAs were performed with substituted DNA fragments of subfragment B₁ and B₂ regions of the *argB* promoter. His₆-ArgR (2,400 nM) was incubated with the DIG-labeled probe. ArgR-DNA complexes are indicated by arrows; Comp. denotes competitor assays (40-fold excess mutated DIG-unlabeled oligonucleotides) of ArgR. Boxes represent alignments of subfragments B₁ and B₂ and their substituted DNA fragments.

inverted-repeated manner (27). Thus, a comparison of the sequence suggests that FarR may regulate the *argB* promoter region by its direct interaction with the DNA site for ornithine and arginine biosynthesis. Subsequently, to confirm the putative FarR-binding site, an EMSA was performed on purified His₆-FarR. Figure 6 shows that the presence of FarR, at a 1,169 nM concentration, led to a significant retardation of the DNA fragment (subfragment B₁) carrying the *argB* promoter region between positions -50 and -77, suggesting that FarR binds to the *argB* promoter region. The precise DNA binding of FarR was determined by competitive assays. As shown in Fig. 6, the binding of FarR to the *argB* promoter region was specific, as this binding was significantly inhibited in a dose-dependent manner by DIG-unlabeled subfragment B₁.

Relationship of binding of ArgR and FarR to the *argB* promoter region. From the mutational analysis of ArgR binding, the results indicate that G₆T, C₉A, and A₂₃C are the three critical sites for ArgR binding to the *argB* promoter region (Fig. 2A and 4). Among them, G₆T at sites at position -58 of the *argB* promoter is included in the putative FarR-binding site, located at positions -57 to -77 within the *argB* promoter region. Additionally, C₉A (at sites at position -55 of the *argB* promoter) was located in the vicinity of the putative FarR-binding site (Fig. 2A). This finding suggests that ArgR and FarR have some relation to regulate *argB* through the binding of the *argB* promoter region as a transcriptional regulator. Therefore, the binding of ArgR and FarR to the *argB* promoter region was further investigated by comparing *in vivo* ChIP assays with *in vitro* EMSAs. First, the DNA binding of FarR to the *arg* genes was examined by ChIP at 14 h after *C. glutamicum* SJC 8074 (*argR* mutant) inoculation, in order to elucidate a correlation between the binding of FarR and the binding of ArgR to the *argB* promoter region (Fig. 7A). The DNA binding of FarR to the *argB* promoter region was also

clearly observed in the absence of ArgR in *C. glutamicum*, and this phenomenon was observed for the promoter region of the *argB*, *argF*, *argG*, and *argH* genes. This means that FarR did directly bind to the *argB* promoter in *C. glutamicum* and not indirectly through ArgR.

In addition, the effect of the binding of purified ArgR and FarR to the *argB* promoter region was tested by EMSAs (Fig. 7B). Formations of protein-DNA complexes were observed for the individual EMSAs of ArgR and FarR, using DIG-labeled subfragment B₁ (positions -50 to -77). To further corroborate a relationship between these regulators, a different reaction order of ArgR and FarR for contact with subfragment B₁ was performed. In this assay, the ArgR-DNA complex was observed only when FarR was incubated with the ArgR-DNA complex after first binding ArgR with subfragment B₁, meaning that FarR did not bind to the ArgR-DNA complex. Whereas both ArgR and FarR formed a complex with the DNA fragment (ArgR-FarR), they formed protein-DNA complexes individually when ArgR was incubated with the FarR-DNA complex after first binding FarR with subfragment B₁. This suggests that FarR has only one putative binding domain located at positions -57 to -77, but this region exactly overlapped with subfragment B₁ for the binding of ArgR within the *argB* promoter; thus, if ArgR bound with the *argB* promoter first, the binding of FarR would be difficult. However, if FarR bound to the binding domain located at positions -57 to -77 first, ArgR could bind other binding sites located at positions -49 to -25 within the *argB* promoter.

DISCUSSION

Previous *in vivo* studies suggested that the upstream region of the *argB* gene on the *arg* operon plays an important role in interacting with ArgR under proline-supplemented conditions




FIG. 5. Time profiles of *in vivo* DNA binding of ArgR. (A) Cell growth and ornithine and arginine production during cultivation of *C. glutamicum*. Cell growth is indicated by circles, and ornithine and arginine concentrations are indicated by open and closed triangles, respectively. The results are reported as the means \pm standard deviations (SD) ($n = 3$). (B) *In vivo* binding of FarR to individual promoter regions of *arg* genes analyzed by ChIP assays. The *C. glutamicum* wild-type strain was treated with formaldehyde to cross-link FarR to promoters and lysed, and FarR complexes were immunoprecipitated for analysis.

in *C. glutamicum* (14, 16). Clarification of the specific effect of the transcriptional repressor ArgR on the action of another transcriptional regulator, FarR, on the *argB* promoter region in *C. glutamicum* is the aim of this work.

In bacteria, the *argB* gene encodes *N*-acetylglutamate kinase, a key enzyme for ornithine biosynthesis. The enzyme's importance comes from its feedback inhibition control: it regulates its own pathway (28, 34). The *C. glutamicum argB* gene can be transcribed from an internal promoter located in its upstream region (28). In the consensus *C. glutamicum* promoter, the prominent feature is a conserved extended -10 region, $\text{tgngnTA}(c/t)\text{aaTgg}$ (with the less-conserved nucleotides lowercase), while the -35 region is much less conserved (23). Caldara et al. previously described how liganded ArgR and RNA polymerase effectively compete *in vivo* by binding to partially overlapping sites (3). Sequence comparison allows the observation that the hypothesized -35 region of the *argB* promoter does not overlap the putative ARG box deduced by EMSAs (data not shown). However, the *argB* gene is consid-




FIG. 6. EMSAs were performed with a subfragment (subfragment B₁) of the promoter region in the *argB* promoter region. The His₆-FarR protein was incubated with a DIG-labeled probe containing subfragment B₁ and subjected to EMSA. (Left) Binding of His₆-FarR. The concentrations of FarR were 501 nM (lane 1), 835 nM (lane 2), 1,169 nM (lane 3), and 1,670 nM (lane 4). Lane 0, no protein. (Right) Competitor assay of His₆-FarR (1,670 nM) using excess DIG-unlabeled oligonucleotides.

ered a member of the *C. glutamicum* ArgR regulon, even though none of the core promoter elements overlap the ARG boxes (3, 24).

The protein building block proline has other important functions, including being a source of energy, carbon, and nitrogen and being an osmolyte (21, 25). The first and controlling step of the synthesis of proline from glutamate is catalyzed by glutamate-5-kinase (G5P), and it was reported previously that this enzyme is feedback inhibited by proline (25). A particularly interesting finding from previous studies, which relates to this work, was that proline also expedited ornithine biosynthesis, a




FIG. 7. Comparative analysis of *in vivo* and *in vitro* bindings of ArgR and FarR to the *argB* promoter region. (A) ChIP assay of FarR binding to promoter regions of the *argB* gene. *C. glutamicum* strain SJC 8074 (*argR* mutant) was treated with formaldehyde to cross-link FarR to promoters and lysed, and FarR complexes were immunoprecipitated for analysis. (B) EMSAs of His₆-ArgR and His₆-FarR. EMSA was done with subfragment B₁ as described in the legend of Fig. 2. The vertical arrows indicate the order of incubation of His₆-FarR (1,670 nM) and His₆-ArgR (2,400 nM) with subfragment B₁.

catalyzed synthesis with glutamate as a primary metabolite in *C. glutamicum* (14, 16). In this study, proline was observed to improve *argB* gene transcription due to its ability to decrease the efficacy of the binding of ArgR to the *argB* promoter region (Fig. 3). The combined results support the hypotheses that the enhancement of ornithine biosynthesis under conditions of proline supplementation might be mediated by two physiological effects: (i) metabolic flux from glutamate favors ornithine biosynthesis rather than proline biosynthesis, as G5P is feedback inhibited by proline supplementation (25), and (ii) proline acts as an antirepressor by binding to a specific region of the ArgR structure (14). The determination of the three-dimensional (3-D) structure of the ArgR-proline complex, essential for clarifying this issue, remains but a sought-after goal.

It has been known for some time that FarR is a fatty acid- and fatty acyl coenzyme A (acyl-CoA)-responsive DNA-binding protein. Its new function might be connected to amino acid biosynthesis and central carbon energy metabolism (8), since the *farR* gene in the tricarboxylic acid (TCA) cycle gene cluster of *E. coli* (*gltA-sdhCDAB-sucABCD-farR*) is autoregulated by the FarR protein (26). The ChIP results provide evidence that FarR binds to the upstream regions of *arg* genes (Fig. 6) as well as to the *gdh* promoter, which converts an intermediate of the TCA cycle to glutamate (15). Indeed, glutamate biosynthesis is closely related to fatty acid synthesis (5). The inactivation of DtsR (detergent sensitivity rescuer), which is assumed to be involved in fatty acid synthesis, triggers glutamate overproduction in coryneform bacteria (12, 36). Thus, it would be beneficial to clarify how FarR participates in fatty acid and amino acid synthesis in *C. glutamicum*.

The ArgR and FarR proteins have calculated molecular masses of 29.3 and 18.8 kDa, respectively (16, 33). In addition, the FarR protein is predicted to form dimers, in accordance with the behavior of other members of the GntR family of transcriptional regulators (27). Therefore, it may be presumed that ArgR and FarR in *C. glutamicum* have differing structures as hexamers and dimers, respectively. This is a likely cause of the distinct patterns of binding of ArgR and FarR to the *argB* promoter region (Fig. 7B). This study tries to provide insight into the complex regulation of *argB* expression. The regulator proteins ArgR and FarR likely bind to the *argB* upstream region in a manner similar to the previously reported upstream binding of FarR to *gdh* (8). In addition to FarR, the expression of *gdh* in *C. glutamicum* was also found to be regulated by the binding of the global transcriptional regulator protein, AmtR, to two separate sites in the *gdh* upstream region, at positions -184 to -209 and -334 to -359 from the site of the start of transcription, in a recent study of the stringent response (9). Among the latter positions is a putative ARG box sequence (14) as well as contact with the -10 to -35 region of the *gdh* promoter (9). In this study, FarR has only one putative binding domain located at positions -57 to -77, but this region exactly overlapped with subfragment B₁ for binding ArgR within the *argB* promoter; thus, if ArgR bound with the *argB* promoter first, the binding of FarR would be difficult. However, if FarR bound to the binding domain located at positions -57 to -77 first, ArgR could bind other binding sites located at positions -49 to -25 within the *argB* promoter. Therefore, both ArgR and FarR formed a complex with the DNA fragment (ArgR-FarR) and formed protein-DNA complexes individually when

ArgR was incubated with the FarR-DNA complex after first binding FarR with subfragment B₁.

In conclusion, these results demonstrate that *C. glutamicum* ArgR regulates *argB* gene transcription as a repressor and that this repression is regulated by the intracellular molecule proline. Moreover, this study extends our understanding of the molecular mechanisms involved in the transcription regulation of the *argB* gene by demonstrating the interaction of ArgR with another transcription factor, FarR. The findings have a number of hopeful implications for future studies of ArgR that may elucidate its regulatory mechanisms in greater detail.

ACKNOWLEDGMENT

This study was supported by the 21C Frontier Microbial Genomics and Applications Center Program, grant no. 11-2008-10-002-00, Ministry of Education, Science & Technology, Republic of Korea. We are grateful for their support.

REFERENCES

1. Brinkroff, K., I. Brune, and A. Tauch. 2007. The transcriptional regulatory network of the amino acid producer *Corynebacterium glutamicum*. *J. Biotechnol.* **129**:191–211.
2. Brune, I., K. Brinkroff, J. Kalinowski, A. Pühler, and A. Tauch. 2005. The individual and common repertoire of DNA-binding transcriptional regulators of *Corynebacterium glutamicum*, *Corynebacterium efficiens*, *Corynebacterium diphtheriae* and *Corynebacterium jeikeium* deduced from the complete genome sequence. *BMC Genomics* **6**:86.
3. Caldara, M., P. N. Minh, S. Bostoen, J. Massant, and D. Charlier. 2007. ArgR-dependent repression of arginine and histidine transport genes in *Escherichia coli* K-12. *J. Mol. Biol.* **373**:251–267.
4. Crooks, G. E., G. Hon, J. M. Chandonia, and S. E. Brenner. 2004. WebLogo: a sequence logo generator. *Genome Res.* **14**:1188–1190.
5. Eggeling, L., and M. Bott. 2005. Handbook of *Corynebacterium glutamicum*. CRC Press, Taylor & Francis Group, Boca Raton, FL.
6. Ghochikyan, A., et al. 2002. Arginine operator binding by heterologous and chimeric ArgR repressors from *Escherichia coli* and *Bacillus stearothermophilus*. *J. Bacteriol.* **184**:6602–6614.
7. Grandori, R., et al. 1995. The DNA-binding domain of the hexameric arginine repressor. *J. Mol. Biol.* **254**:150–162.
8. Hänfler, E., et al. 2007. FarR, a putative regulator of amino acid metabolism in *Corynebacterium glutamicum*. *Appl. Environ. Microbiol.* **76**:625–632.
9. Hänfler, E., et al. 2009. A game with many players: control of *gdh* transcription in *Corynebacterium glutamicum*. *J. Biotechnol.* **142**:114–122.
10. Hwang, J. H., G. H. Hwang, and J. Y. Cho. 2008. Effect of increased glutamate availability on L-ornithine production in *Corynebacterium glutamicum*. *J. Microbiol. Biotechnol.* **18**:704–710.
11. Ikeda, M., and S. Nakagawa. 2003. The *Corynebacterium glutamicum* genome: features and impacts on biotechnological processes. *Appl. Microbiol. Biotechnol.* **62**:99–109.
12. Kimura, E., C. Abe, Y. Kawahara, T. Nakamatsu, and H. Tokuda. 1997. A *dtsR* gene-disrupted mutant of *Brevibacterium lactofermentum* requires fatty acids for growth and efficiently produces L-glutamate in the presence of an excess of biotin. *Biochem. Biophys. Res. Commun.* **234**:157–161.
13. Lee, S. Y., et al. 2010. Utilization of phenol and naphthalene affects synthesis of differential amino acids in *Corynebacterium glutamicum*. *Curr. Microbiol.* **61**:596–600.
14. Lee, S. Y., H. S. Shin, J.-S. Park, Y. H. Kim, and J. Min. 2010. Proline reduces the binding of transcriptional regulator ArgR to upstream of *argB* in *Corynebacterium glutamicum*. *Appl. Microbiol. Biotechnol.* **86**:235–242.
15. Lee, S. Y., Y. H. Kim, and J. Min. 2010. Conversion of phenol to glutamate and proline in *Corynebacterium glutamicum* is regulated by transcriptional regulator ArgR. *Appl. Microbiol. Biotechnol.* **85**:713–720.
16. Lee, S. Y., Y. H. Kim, and J. Min. 2009. The effect of ArgR-DNA binding affinity on ornithine production in *Corynebacterium glutamicum*. *Curr. Microbiol.* **59**:483–488.
17. Lee, Y. J., and J. Y. Cho. 2006. Genetic manipulation of a primary metabolic pathway for L-ornithine production in *Escherichia coli*. *Biotechnol. Lett.* **28**:1849–1856.
18. Leuchtenberger, W., K. Huthmacher, and K. Drauz. 2005. Biotechnological production of amino acids and derivatives: current status and prospects. *Appl. Microbiol. Biotechnol.* **69**:1–8.
19. Maas, W. K. 1994. The arginine repressor of *Escherichia coli*. *Microbiol. Rev.* **58**:631–640.
20. Makarova, K. S., A. A. Mironov, and M. S. Gelfand. 2001. Conservation of the binding site for the arginine repressor in all bacterial lineages. *Genome Biol.* **2**:1–8.

21. Marco-Marín, C., et al. 2007. A novel two-domain architecture within the amino acid kinase enzyme family revealed by the crystal structure of *Escherichia coli* glutamate 5-kinase. *J. Mol. Biol.* **367**:1431–1446.
22. Park, S. M., C. D. Lu, and A. T. Abdelal. 1997. Cloning and characterization of *argR*, a gene that participates in regulation of arginine biosynthesis and catabolism in *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* **179**:5300–5308.
23. Pátek, M., J. Nešvera, A. Guyonvarch, O. Reyes, and G. Leblon. 2003. Promoters of *Corynebacterium glutamicum*. *J. Biotechnol.* **10**:311–323.
24. Paul, L., P. K. Mishra, R. M. Blumenthal, and R. G. Matthews. 2007. Integration of regulatory signals through involvement of multiple global regulators: control of the *Escherichia coli* *gltBDF* operon by Lrp, IHF, Crp, and ArgR. *BMC Microbiol.* **7**:1–17.
25. Pérez-Arellano, L., V. Rubio, and J. Cervera. 2006. Mapping active site residue in glutamate-5-kinase. The substrate glutamate and the feed-back inhibitor proline bind at overlapping sites. *FEBS Lett.* **580**:6247–6253.
26. Quail, M. A., C. E. Dempsey, and J. R. Guest. 1994. Identification of a fatty acyl responsive regulator (FarR) in *Escherichia coli*. *FEMS Lett.* **356**:183–187.
27. Rigali, S., A. Derouaux, F. Giannotta, and J. Dusart. 2002. Subdivision of the helix-turn-helix GntR family of bacterial regulators in the FadR, HutC, MocR, and YtrA subfamilies. *J. Biol. Chem.* **277**:12507–12515.
28. Sakanyan, V., et al. 1996. Genes and enzymes of the acetyl cycle of arginine biosynthesis in *Corynebacterium glutamicum*: enzyme evolution in the early steps of the arginine pathway. *Microbiology* **142**:99–108.
29. Sambrook, J., and D. W. Russell. 2001. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
30. Sampaio, M., et al. 2004. Phosphotransferase-mediated transport of the osmolyte 2-O- α -mannosyl-D-glycerate in *Escherichia coli* occurs by the product of the *mngA* (*hrsA*) gene and is regulated by the *mngR* (*farR*) gene product acting as repressor. *J. Biol. Chem.* **279**:5537–5548.
31. Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins. 1997. The Clustal X Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**:4876–4882.
32. Udaka, S. 1960. Screening method for microorganisms accumulating metabolites and its use in the isolation of *Micrococcus glutamicus*. *J. Bacteriol.* **79**:754–755.
33. van Duyn, G. D., G. Ghosh, W. K. Maas, and P. B. Sigler. 1994. Structure of the oligomerization and L-arginine binding domain of the arginine repressor of *Escherichia coli*. *J. Mol. Biol.* **256**:377–391.
34. Xu, Y., B. Labedan, and N. Glansdorff. 2007. Surprising arginine biosynthesis: a reappraisal of the enzymology and evolution of the pathway in microorganisms. *Microbiol. Mol. Biol. Rev.* **71**:36–47.
35. Xu, Y., et al. 2003. Regulation of arginine biosynthesis in the psychrophilic bacterium *Moritella profunda*: in vivo repressibility and in vitro repressor-operator contact probing. *J. Mol. Biol.* **326**:353–369.
36. Yao, W., et al. 2009. Double deletion of *dstR1* and *pyc* induce efficient L-glutamate overproduction in *Corynebacterium glutamicum*. *J. Ind. Microbiol. Biotechnol.* **36**:911–921.