

Fatty-Acid Binding Proteins Modulate Sleep and Enhance Long-Term Memory Consolidation in *Drosophila*

Jason R. Gerstner^{1*†}, William M. Vanderheyden², Paul J. Shaw², Charles F. Landry³, Jerry C. P. Yin^{1,4,5*}

1 Department of Genetics, University of Wisconsin-Madison, Madison, Wisconsin, United States of America, **2** Anatomy and Neurobiology, Washington University School of Medicine, St. Louis, Missouri, United States of America, **3** Scarab Genomics, LLC, Madison, Wisconsin, United States of America, **4** Department of Neurology, University of Wisconsin-Madison, Madison, Wisconsin, United States of America, **5** Waisman Center, University of Wisconsin-Madison, Madison, Wisconsin, United States of America

Abstract

Sleep is thought to be important for memory consolidation, since sleep deprivation has been shown to interfere with memory processing. However, the effects of augmenting sleep on memory formation are not well known, and testing the role of sleep in memory enhancement has been limited to pharmacological and behavioral approaches. Here we test the effect of overexpressing the brain-type fatty acid binding protein (Fabp7) on sleep and long-term memory (LTM) formation in *Drosophila melanogaster*. Transgenic flies carrying the murine Fabp7 or the *Drosophila* homologue dFabp had reduced baseline sleep but normal LTM, while Fabp induction produced increases in both net sleep and LTM. We also define a post-training consolidation “window” that is sufficient for the observed Fabp-mediated memory enhancement. Since Fabp overexpression increases consolidated daytime sleep bouts, these data support a role for longer naps in improving memory and provide a novel role for lipid-binding proteins in regulating memory consolidation concurrently with changes in behavioral state.

Citation: Gerstner JR, Vanderheyden WM, Shaw PJ, Landry CF, Yin JCP (2011) Fatty-Acid Binding Proteins Modulate Sleep and Enhance Long-Term Memory Consolidation in *Drosophila*. PLoS ONE 6(1): e15890. doi:10.1371/journal.pone.0015890

Editor: Wim Crusio, Université de Bordeaux and Centre National de la Recherche Scientifique, France

Received: September 4, 2010; **Accepted:** November 25, 2010; **Published:** January 27, 2011

Copyright: © 2011 Gerstner et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was funded by the National Institutes of Health (MH60774 to JC), its Training Grant Program (GM07507), and start up funds from the Waisman Center (to JCY). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: gerstner@upenn.edu (JRG); jcyin@wisc.edu (JCPY)

† Current address: Center for Sleep and Circadian Neurobiology, Translational Research Laboratories, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania, United States of America

Introduction

Although sleep is an essential behavioral process conserved across phyla from fruit flies to humans, its functions remain elusive. There are many theories about the function of sleep, including roles in metabolic balance [1], excitotoxic repair [2], and memory consolidation [3]. The process of memory is also widely conserved in the animal kingdom, and the relationship between sleep and memory formation continues to be controversial [4].

Studies in both rodents [5] and Aplysia [6] show a clear time-of-day effect on memory formation. In flies, short-term memory has been shown to be regulated by processes involving both circadian rhythms [7] and sleep [8]. In addition, a temporal window for the effects of sleep deprivation on memory has been shown in both invertebrate and vertebrate species. Four hours of sleep deprivation (SD) immediately following courtship training abolishes memory retention in flies [9], while SD later during the sleeping period has no effect. In rodents, REM SD either just before [10], or immediately after behavioral training [11], has been shown to abolish contextual (hippocampal dependent) memory formation. It has also been suggested that REM sleep is necessary during later consolidation periods of memory in rodents [12]. While these results imply a role for sleep in the consolidation of memory, they describe the consequences of reducing sleep (or specific components of sleep) on the formation of memory. An alternate method for understanding the influence of sleep on memory formation would be to test the effects of augmenting sleep using genetic

approaches. While molecular targets involving circadian- and sleep-dependent memory formation are beginning to be identified [13,14], specific molecular targets responsible for augmenting sleep and memory concomitantly are not known.

A number of laboratories have used microarrays to identify sleep (or deprivation)-responsive genes [15–19], and some of these include immediate-early genes (IEGs), many of which have been shown to be involved in learning and memory consolidation [20–22]. Levels of some IEGs, such as *Zif268*, are elevated in response to enriched environments or long-term potentiation (LTP) during subsequent REM sleep episodes in rodents [23,24]. The regulation of synaptic plasticity-related molecules in sleep/wake behavior suggests that the sleep/wake system may re-deploy some of the molecular components underlying learning and memory [25]. Previously we used microarray analysis to show that brain-type fatty-acid binding protein (Fabp7) shows circadian oscillation in its expression [26], and that its cycling pattern is widespread throughout mammalian brain [27]. Here we utilize the *Drosophila* model to investigate the functional role of fatty-acid binding proteins in sleep and memory consolidation, and describe a novel molecular player for improving memory through increases in daytime sleep.

Results

Characterization of *Drosophila* Fabp

Fatty-acid binding proteins (Fabps) comprise a group of soluble proteins that bind small hydrophobic lipids and act as transporters.

The mammalian Fabp family consists of ~9 separate genes, which are expressed in various cell and tissue types. In the adult mammalian central nervous system (CNS), only three of these appear to be expressed, and include heart-, epidermal-, and brain-type fatty-acid binding proteins (Fabp3, 5, 7, respectively). In contrast, there is a single Fabp locus (CG6783) in the *Drosophila* genome, which we have termed dFabp, that encodes at least 3 alternatively spliced isoforms of *Drosophila* Fabp (dFabp) (Fig. 1C). The second dFabp isoform (CG6783-B) shares the greatest homology to mouse Fabp7 (54% Identical, 68% Positive). An alignment of the amino-acid sequences for Fabps from various species, and a derived consensus sequence, are shown in Fig. 1A.

Since the mouse Fabp7 transcript and protein have been shown to follow diurnal changes in expression, we asked whether dFabp also oscillates across the day/night cycle. Analysis of dFabp mRNA from fly head extracts show changes based on time-of-day, which reflect circadian changes previously observed [28]. We identified an elevation in mRNA levels during sleep and reduced levels during the wake period (Fig. 1B). The increase in dFabp mRNA preceded an accumulation of dFabp protein, which peaked early in the waking phase (Fig. 1B; Figure S1A). These results mirror oscillations in Fabp7 expression observed in mouse

brain [27], suggesting a conserved function over the day/night cycle. Immunocytochemical (ICC) analysis using antibodies raised against dFabp-B revealed a broad expression pattern throughout the fly brain (Fig. 1D). ICC staining against dFabp appeared to be much stronger in nuclei (identified using the nuclear marker Topro-3) than in the cytoplasm (Fig. 1D), however the signal cannot be verified to be solely due to endogenous expression.

Fabp expression regulates sleep and memory

Genes that are involved in regulating behavioral states (wake, sleep) might be predicted to show coordinated, brain-wide changes in their expression levels across the day/night cycle. Since both flies (Fig. 1B) and mammals [27] exhibit a diurnal fluctuation of lipid-binding protein expression in the CNS, we examined the effects of altering Fabp expression on sleep behavior. We generated heat-shock inducible transgenic flies carrying the mouse Fabp7 open reading frame (ORF) in the background strain *w(isoCG1)*, an isogenized wild-type strain. Western blots probed with species-specific antibodies showed that the transgenic flies overexpress protein, even at lower temperatures (20°C; Figure S1B). Surprisingly, this overexpression of Fabp7 resulted in a dramatic reduction in baseline sleep relative to the *w(isoCG1)*




Figure 1. Characterization and diurnal regulation of the *Drosophila* fatty-acid binding protein, dFabp. (A) Amino acid sequence homology of dFabp is compared to brain-type fatty-acid binding protein (Fabp7) of mouse, rat, human, zebra fish, and worm (lbp-7). A consensus sequence of all species is also shown. (B) Densitometric analysis of Northern and Western blots reveals a diurnal regulation of dFabp mRNA (blue solid line) and protein (red dashed line) in fly head. dFabp mRNA levels are normalized to Gapdh, with elevated expression during the sleep period (mean \pm s.e.m., N=2/timepoint, 50–100 heads per timepoint, asterisk $P<0.05$, t-test ZT18 vs ZT6). dFabp protein levels are normalized to laminin, and are elevated toward wake onset (mean \pm s.e.m., N=4/timepoint, 50–100 heads per timepoint, $t P=0.051$, t-test ZT3 vs ZT15). Open bar = lights on, filled bar = lights off. (C) Schematic representation of the dFabp locus. Open bar = non-coding, filled bar = coding region. (D) Projection slice of brain stained with an antibody against dFabp (red), and the nuclear stain Topro-3 (blue), as viewed by confocal microscopy. ML = midline, OL = optic lobe, AL = antennal lobe, scale bar = 120 μ m. doi:10.1371/journal.pone.0015890.g001

background flies (housed at $\sim 22^{\circ}\text{C}$), principally from decreased daytime sleep (Fig. 2B, D, E). These effects on baseline sleep also occur in male flies (data not shown) and two additional lines carrying the Fabp7 ORF (Figure S2). We observed similar reductions in total and daytime sleep in transgenic flies that carry the *Drosophila* dFabp-B ORF (Fig. 2C, D, E). The average daytime bout length and the average maximum daytime bout length are both significantly reduced in Fabp7 and dFabp flies (Fig. 2E, F). The bout length is considered to be a measurement of consolidated sleep. Two additional dFabp overexpressing fly lines also showed similar significant results in each of these sleep parameters (Figure S3). The reduction in daytime sleep was not due to hyperactivity, since daytime locomotor activity, as measured by counts per waking minute, was in fact reduced in Fabp7 and dFabp flies (Fig. 2G; Figures S2G; S3G). The total number of sleep bouts over a 24-hour period for the transgenic flies was not different from controls (data not shown).

Given the reduction in daytime sleep in Fabp7 and dFabp flies, and the growing appreciation for the importance of sleep in memory formation, we tested the effect of altering post-training Fabp protein levels on long-term memory formation. Wild type and transgenic flies were trained with 10 cycles of spaced training, and then stored at various temperatures to modulate the post-acquisition levels of Fabp protein expression. Despite the reduced baseline sleep phenotype observed in Fabp flies at lower temperatures (Fig. 2), trained flies maintained at 20°C did not show any changes in performance as compared to the *w(isoCJ1)* background strain, (Fig. 3A). However, inducing Fabp7 overexpression in flies stored at 30°C during the maintenance period following training showed a significant enhancement of 7-day LTM (Fig. 3A). There was a strong correlation between temperature and memory when the ratio of performance between

Fabp7: *w(isoCJ1)* flies was plotted, and this paralleled the increase in Fabp7 protein expression (Fig. 3B). Learning scores for Fabp7 flies were comparable to the background strain *w(isoCJ1)* (Figure S4A), further supporting the conclusion that the temperature-induced enhancement of memory was not due to an effect on learning. This result suggests that some aspect of Fabp overexpression during the 7-days following training enhances memory formation.

Since the transgenic flies showed a temperature-dependent enhancement of memory formation, we examined whether there are changes in sleep as a function of temperature. A significant increase in total sleep occurred in Fabp7 flies when the transgene was induced, compared to *w(isoCJ1)* flies (Fig. 4B, D). This increase in total sleep resulted primarily from increases in daytime sleep (Fig. 4E), since night-time sleep was similarly reduced in all flies (Fig. 4A, B, C). A more detailed analysis showed that Fabp7 and dFabp transgene induction increased the maximum daytime bout length (Fig. 4G), without significantly affecting the average daytime bout length (Fig. 4F) or bout number (Fig. 4H), compared to *w(isoCJ1)* controls. Therefore, increasing the temperature from 20°C to 30°C increased Fabp protein expression and correlated with a net increase in longer daytime sleep bouts.

Temporal window for Fabp-mediated memory enhancement

In order to further determine when, after behavioral training, Fabp7 and dFabp act to enhance memory formation, we tested various temporal “windows” of overexpression. It has previously been shown that aPKM transgenic flies enhance memory formation when the transgene is induced shortly after the end of behavioral training [29]. However, increasing Fabp7 immediately after the end of LTM training did not enhance 24-hour memory (data not shown).




Figure 2. Reduced sleep in Fabp7 and dFabp overexpressing flies. Hourly diurnal sleep profile of the (A) *w(isoCJ1)* background strain, (B) Fabp7, and (C) dFabp flies. Open bars = lights on, filled bars = lights off. Analysis of various sleep parameters: (D) Total sleep over 24 hours, (E) Total daytime sleep, (F) Average daytime sleep bout length, (G) Maximum daytime sleep bout length, (H) Daytime locomotor activity per waking minute. Open bars = *w(isoCJ1)*, light grey bars = Fabp7, dark grey bars = dFabp. Results are mean \pm s.e.m.; $n = 23-24$, representative of 3-5 experiments. Single asterisk, $P < 0.05$; double asterisk, $P < 0.01$; triple asterisk, $P < 0.001$, t -test. doi:10.1371/journal.pone.0015890.g002




Figure 3. *Fabp7* overexpression enhances 7-day long-term memory (LTM). (A) Post-training induction of the heat shock (HS) promoter-driven *Fabp7* flies produces a temperature-dependent enhancement of memory after 10x spaced training if flies are maintained at higher temperatures (30°C) for 7 days prior to testing. Grey bars, *Fabp7*, open bars, *w(isoCJ1)* background strain. Results are mean \pm s.e.m.; $n=8-10$ groups. Asterisk $P<0.05$, t -test. (B) Ratio of Performance Index (PI) scores (A) for *Fabp7/w(isoCJ1)* (diamonds; y-axis left) shows a correlation of increasing performance with temperature ($R^2=0.95$), similar to the correlation of increase in protein expression (black circles; y-axis right) with temperature ($R^2=0.89$). Inset: Western blot of *Fabp7* protein expression in *Fabp7* staged at 20°, 25°, and 30°C for 7 days, and the immunoreactive background of *w(isoCJ1)* (*w*) at room temperature (RT), in nuclear fraction (lysate sample: $\sim 50-100$ fly heads per lane). doi:10.1371/journal.pone.0015890.g003

Next we tested if *Fabp7* overexpression during the first four days after training is sufficient to enhance seven day LTM formation. Flies were trained at 25°C and immediately placed at 30°C for four days, after which they were shifted back to 20°C for the remaining time. This incubation schedule preserved memory enhancement (Fig. 5A), suggesting that the effects of *Fabp7* on LTM require sustained periods of overexpression. Shifting the overexpression window to a later consolidation period (between post-training days 2 and 7) also preserved enhancement of LTM in *Fabp7* flies (Fig. 5B). Further, this effect replicated when we analyzed *dFabp* flies using this same regimen (Fig. 5C). While

Fabp7 flies showed elevated olfactory avoidance responses following five days at 30°C (data not shown), we still observed an increase in LTM when *Fabp7* flies were shifted back to 20°C for two days prior to testing (Fig. 5A). This result suggested that elevated *Fabp7* expression for the time period just prior to recall (Fig. 5B, C) is not responsible for the increase in LTM observed. However, in order to rule out the possible effects of *Fabp7* on sensory-systems during retrieval, we examined the effects of increasing *Fabp7* on memory recall. We found that an acute induction of *Fabp7* two hours prior to testing for LTM actually inhibited performance (Figure S4C). This suggests that acute *Fabp7* expression at the time of retrieval is able to impede memory recall processes. Together, these data provide evidence that the *Fabp7*-mediated enhancement effects on LTM occur during the consolidation period, and are not due to effects on learning, acquisition or recall.

We have shown that at low temperatures where the transgene induction is mild, the *Fabp7* or *dFabp* proteins reduce the steady-state amount of daytime sleep. However, when the transgenes are induced to higher levels, the net gain in daytime sleep that flies obtained (relative to their isogenic control line) was substantial. Memory formation was enhanced when flies were stored at inducing temperatures. Is the total amount of sleep gained correlated with the memory enhancement? To test if these two phenotypes are correlated, we measured the cumulative change in sleep of *Fabp7* and *dFabp* flies maintained at 30°C for four consecutive days following two days baseline at 20°C. This is the temperature regimen that produces enhancement of memory formation during the consolidation phase. There was a significant increase in minutes and percentage of sleep gained in *Fabp7* and *dFabp* flies, as compared to *w(isoCJ1)* flies, on the second, third, and fourth days at 30°C (Fig. 6A). In addition, the cumulative increase in total sleep (Fig. 6B), and percentage of total sleep gained (Fig. 6C) was significantly higher in both *Fabp7* and *dFabp* flies compared to *w(isoCJ1)* flies for the four days at 30°C. Taken together, these results show that at elevated temperatures following spaced training, *Fabp7* and *dFabp* flies undergo an increase in cumulative sleep during the consolidation period, and this increase correlates with an enhancement in LTM performance.

Discussion

Previously we have shown that the mRNA levels for *Fabp7* are diurnally regulated and are highest during the sleeping period of mice [26,27]. Here we characterized the *Fabp7* *Drosophila* homologue *dFabp*, and show that *dFabp* mRNA expression also cycles across the day, with elevated levels during the sleeping period. Further, in order to determine whether increasing *Fabp7* expression influences complex behavior, we show that induction of *Fabp7* or *dFabp* in transgenic flies is sufficient to produce sleep and long-term memory enhancement phenotypes. These data suggest a common theme for cycling molecules in the formation of long-term memory, and suggest that like other known circadian genes, *Fabp7*-mediated signaling is likely implicated in both sleep/wake and memory-related processes.

Regulation of *Fabp7* expression

What are the upstream molecular signaling cascades currently known to regulate *Fabp7* expression? While the characterization of specific molecular pathways involved in modulating *Fabp7* levels across the day/night cycle has not yet been reported, *cis*-acting elements regulating *Fabp7* expression during CNS development have begun to be identified. Pax6, a transcription factor known to promote developmental neuronal differentiation, was shown to regulate *Fabp7* expression, in a manner believed to be critical for




Figure 4. Increased net sleep in *Fabp7* and *dFabp* overexpressing flies. Hourly change in sleep profile of the (A) *w(isoCJ1)* background strain, (B) *Fabp7*, and (C) *dFabp* flies shifted from 20°–30°C. Open bars = lights on, filled bars = lights off. Analysis of change in various sleep parameters of flies shifted from 20°–30°C: (D) Total sleep over 24 hours, (E) Total daytime sleep, (F) Average daytime sleep bout length, (G) Maximum daytime sleep bout length, (H) Number of daytime bouts. Open bars = *w(isoCJ1)*, light grey bars = *Fabp7*, dark grey bars = *dFabp*. Results are mean \pm s.e.m.; $n = 20$ –30. Single asterisk, $P < 0.05$; double asterisk, $P < 0.01$; triple asterisk, $P < 0.001$, t -test. doi:10.1371/journal.pone.0015890.g004

the proper maintenance of neuroepithelial cells [30]. *Fabp7* has also been shown to be a downstream target of Notch signaling pathways during CNS development [31]. Since Notch signaling has also been implicated in LTM formation in both rodents [32] and flies [33,34], the Notch cascade could represent a conserved molecular pathway mediating *Fabp*-induced memory-related processes in the adult CNS across animal phylogeny. Since Notch-effector binding sites are observed in both mammalian and fly promoters for *Fabp7* [31] and *dFabp* (J. Gerstner, unpublished observations), respectively, future studies aimed at the identification of these and other phylogenetically conserved *cis*-acting elements will provide novel targets for sleep and cognitive therapeutics.

Fabp7 knock-out (KO) mice have previously been shown to have altered behavioral responses, including an increase in anxiety-like behavior [35], and deficits in prepulse inhibition, a hallmark of schizophrenia [36]. *Fabp7* mRNA expression is known to be enriched in the dendritic layers of the hippocampus [37], where it has also been shown to be induced following kainate injection [38], a procedure that initiates epileptiform activity. Further, following the induction of a constitutively active form of CREB, *Fabp7* mRNA shows an elevation in expression within the hippocampus that mirrors the temporal profile of CREB-induced BDNF mRNA expression [39]. Taken together, these data suggest a role for *Fabp7* in plasticity-related mechanisms in mammals, and implicate *Fabp7* in pathways downstream of well established memory-related molecular signaling cascades.

Genes, sleep and memory

There is an emerging notion that sleep, like the circadian system, is under genetic control [40,41]. In addition, mounting




Figure 5. *Fabp7*- and *dFabp*-mediated LTM enhancement is preserved in a late consolidation period. (A) *Fabp7* (grey fill) flies show a significant enhancement in 7 day LTM performance after 10X Spaced training, when shifted from 30°–20°C on day 4, compared to *w(isoCJ1)* background flies (open fill bar). Results are mean \pm s.e.m.; $n = 6$ groups. Asterisk $P < 0.05$, t -test. (B) *Fabp7* (light grey fill) and (C) *dFabp* (dark grey fill) flies show a significant enhancement in 7 day LTM performance after 10X Spaced training, when shifted from 20°–30°C on day 2, as compared to *w(isoCJ1)* background flies (open fill bars, B,C). Results are mean \pm s.e.m.; $n = 7$ –8 groups. Asterisk $P < 0.05$, t -test. d1 = day 1, d2 = day 2, d4 = day 4, d7 = day 7. doi:10.1371/journal.pone.0015890.g005




Figure 6. Cumulative increased sleep in Fabp7 and dFabp flies correlates with LTM enhancement. (A) Fabp7 (light grey square, solid line) and dFabp (dark grey triangle, solid line) flies have an increase in sleep above the *w(isoCJ1)* background strain (open circle, dashed line) following a shift from 20°–30°C. Results are mean \pm s.e.m.; $n=20-30$; double asterisk $P<0.01$, (Fabp7 vs. *w(isoCJ1)*), *t*-test. (B) Cumulative increases in minutes of total sleep gained and (C) percentage of sleep increased over four days when maintained at 30°C are higher in Fabp7 and dFabp flies, as compared to the *w(isoCJ1)* background strain. Open bars = *w(isoCJ1)*, light grey bars = Fabp7, dark grey bars = dFabp. Results are mean \pm s.e.m.; $n=20-30$; double asterisk $P<0.01$, triple asterisk $P<0.001$, *t*-test. doi:10.1371/journal.pone.0015890.g006

evidence suggests that genes involved in the circadian system are able to influence memory processes [42,43]. For example, disruption of core clock components (such as CLOCK and BMAL1) has been shown to inhibit long-term memory in mice [44]. In *Drosophila*, a null mutation of the circadian gene *period* blocked memory, while overexpression of the *period* gene protein product PER was able to enhance memory formation [45]. Recently it has been shown that *period* gene expression was also able to regulate experience-dependent increases in sleep in *Drosophila* [13]. These data are interesting because they suggest that transcription of specific genes that are known regulators of circadian rhythms are able to regulate other complex neurological phenomena, such as learning and memory formation.

Interestingly, while Fabp7 KO mice have not yet been examined for changes in circadian rhythms or behavioral state, KO mice for another Fabp expressed in the mammalian brain, Fabp3, have recently been shown to have increased locomotor activity over the light-dark cycle [46]. Further, Fabp3 has been shown to interact with the dopamine receptor (D2R), and is highly expressed in acetylcholine interneurons and at glutamatergic terminals [46], implicating lipid-transport pathways in the

regulation of dopaminergic signaling within the CNS. Together, these data suggest that lipid-mediated pathways could modulate diurnal changes in locomotor activity, perhaps through a Fabp-regulated mechanism involving signaling cascades downstream of D2R activation.

Influence of lipid metabolites on sleep and memory

Prostaglandin (PG) D2 is lipid-derived eicosanoid that has been shown to promote sleep following the activation of PGD2 receptors (DP1R) and subsequent release of adenosine [47]. In rats, cerebrospinal fluid levels of PGD2 have been shown to exhibit a circadian rhythm [48], and also increase following sleep deprivation [49], suggesting that PGD2 regulates normal sleep in mammals. Infusion of PGD2 into the third ventricle of rats during their wake-period resulted in a dose-dependent increase in both REM and NREM sleep [50]. Dose-dependent increases in sleep were also observed when PGD2 was infused into the lateral ventricle of wild-type (WT) mice, but not in DP1R KO mice [51]. Pharmacological inhibitors of the enzyme which produces PGD2 out of arachidonic acid derivatives, PGD synthase (PGDS), generate a reversible and dose-dependent inhibition of sleep in rats [52]. Interestingly, PGDS is a part of a superfamily of lipid-binding proteins called lipocalins that also include Fabps. Together, these results suggest that lipid signaling pathways regulate normal sleep, and that lipid-binding proteins likely mediate both synthesis and transport of lipid-derived signaling cascades involved in sleep behavior.

Diurnal fluctuations in other lipid species have also been identified in the CNS. Time-of-day cycling has been observed broadly in mammalian brain for the major endocannabinoids anandamide and 2-arachidonoyl-glycerol (2-AG) [53]. Within the rat hippocampus, higher levels of anandamide have been observed during the dark period [53,54], while higher levels of 2-AG were observed during the light period [53]. Further, fatty acid amide hydrolase (FAAH), the enzyme which catabolizes anandamide, has levels that also vary based on time-of-day within various regions of the brain [53,55], suggesting that lipid-metabolism likely also varies over the light/dark cycle. Inhibition of FAAH has recently been shown to promote memory processing [56], implicating anandamide-metabolism in the regulation of memory formation. Oleylethanolamide (OEA), a monounsaturated analogue of anandamide and an endogenous peroxisome proliferator activated receptor (PPAR)- α agonist, failed to show any diurnal fluctuations within the hippocampus [54]. However, post-training administration of OEA, or exogenous PPAR- α agonists, have recently been shown to enhance memory for inhibitory avoidance and Morris water maze tasks [57]. These data suggest that an increase in specific lipid-signaling pathways are able to enhance memory formation during the consolidation period, and together indicate that increased levels of anandamide or its derivatives are able to enhance memory formation.

Interestingly, Fabp7 has recently been shown to be an intracellular carrier of anandamide [58], raising the possibility that subcellular transport of certain lipid molecules, such as anandamide or OEA, via Fabps may activate PPAR-mediated transcription in the CNS. Indeed, various Fabps have already been shown to traffic specific ligands from the cytoplasm to nuclear receptors, such as PPARs, in order to induce transcriptional activation [59-62]. Since PPARs are involved in the circadian clockwork in various tissues and cell types [63-66], these data provide additional evidence for Fabp signaling in gating temporal and memory-enhancing processes.

Growing evidence suggests that sleep is critical for memory consolidation processes [67]. Previously, post-training temporal

windows of sleep were found to be necessary for rodent memory [11,68]. Improved cognitive performance has been found to correlate with sleep and “naps”, while deterioration in memory formation is observed when post-training sleep is disrupted [69–74]. Sleep has previously been shown to recover performance deficits observed in various cognitive tasks in humans [75,76]. More recently sleep (and not merely the passage of time) has been suggested to enhance hippocampus-dependent memory in mice [77] and auditory discrimination performance in European starlings [78]. However, there are few examples of improved performance when post-training sleep is increased or otherwise enhanced. Two such studies involve pharmacological stimulation of REM activity and the “rescue” of sleep deprivation’s disruptive effects on performance [79,80]. While all of these studies point toward a role of sleep in memory consolidation, direct evidence for the effects of augmented sleep on memory formation has been lacking. Furthermore, evidence for genetic control of sleep with concomitant beneficial effects on cognitive ability has not been demonstrated. Here we show that Fabp enhances LTM concurrently with an increase in sleep during post-training consolidation periods in *Drosophila*. Since Fabp overexpression increases consolidated daytime sleep bouts, these data support a role for longer naps in the enhancement of memory. Further, these data suggest a late temporal “consolidation window” which occurs after the initial post-training ~24 hour period usually associated with “synaptic consolidation”. It is conceivable that chronic Fabp-mediated increases in sleep could cause performance improvements during recall, providing an alternative to a pure consolidation effect. This question remains open for future study. Our experimental design also excludes memory improvement resulting from indirect effects on arousal, motivation, or stress during training and retrieval. Since our data is based on a simple associative behavioral task in an invertebrate model, the requirement for sleep in memory consolidation may be surprisingly broad. Finally, this work points toward a novel molecular link between sleep, memory formation, and fatty acid signaling, and may lead to new treatments relating lipid metabolism with sleep, learning, and cognitive disorders.

Methods

Transgenic Flies

The background strain *w(isoC71)* is an isogenic line previously described [29]. Murine Fabp7 open-reading frame (ORF) cDNA was generated by PCR using a forward primer containing the *EcoRI* restriction sequence and a Kozak sequence, and a reverse primer containing a *BglII* sequence. The resulting fragment was subcloned into the p(CaSpeR-hs) vector. An identical strategy was used for the *Drosophila* dFabp-B ORF. All plasmids were sequenced and injected into *w(isoC71)* embryos (BestGene, Inc.).

Fabp7 forward primer: 5'-ACCACCGAATTC AACATGGTAGATGCTTTCTGCGCAACCTG -3'; Fabp7 reverse primer 5'- ACCACAGATCTCTATGCCTTTTCATAAACCGCAA-CAGC-3'. dFabp forward primer: 5'- ACCACCGAATTC CAACATGTCTTTTCGTTGGCAAGAAGTACAAG -3'; dFabp reverse primer 5'- ACCACAGATCTTTAGACGGCCTTGTA-GACGCGC -3'.

Northern Blotting

Fly brain RNA was extracted using Trizol (Invitrogen) and Northern blot analysis done as described [27]. Briefly, brains were dissected and total RNA was isolated using TRIZOL (Invitrogen, Carlsbad, CA), according to the manufacturer’s specifications, and stored at -80°C . Prior to loading, each sample was incubated with

sample buffer (7.5%Formaldehyde, 43%Formamide, 12% 10XMOPS, 0.12%Ethidium Bromide) for 5 min at 56°C , and then cooled on ice for 5 min. Northern Blots consisted of $1\ \mu\text{g}$ of total RNA per lane and were electrophoresed on a 1.2% agarose gel. RNA was transferred overnight onto a sheet of GeneScreen Plus nitrocellulose (NEN Life Science Products, Boston, MA). DNA templates generated by PCR were labeled with ^{32}P using the Megaprime DNA labeling system (Amersham Biosciences, England). Antisense dFabp cDNA probes were PCR amplified from above products. *Drosophila* Gapdh forward primer: 5' - TTTCTCAGCCATCACAGTCG - 3'; reverse primer: 5' - CGACCTCCTCATCGGTGTAT -3'. Probe was mixed with hybridization buffer (Hybrisol, and incubated overnight at 42°C . Blots were washed three times in $2\times\text{SSC}$ ($1\times\text{SSC} = 150\ \text{mM NaCl}$, $15\ \text{mM Na citrate}$, pH, 7.0), 1% SDS at room temperature, then two times for 30 min in $2\times\text{SSC}$, 1% SDS at 50°C , and finally one time for 30 min in $0.5\times\text{SSC}$, 1% SDS at 60°C . Following washes, blots were exposed to a phosphor screen for 3 days and image analysis was performed using the Storm 860 and ImageQuant 5.2 software (Molecular Dynamics, Sunnyvale, CA).

Immunocytochemistry

Fly brains were fixed and stained as described [81]. Antibodies against murine Fabp7 or dFabp-B were generated by purifying MAT-FLAG-tag (Sigma-Aldrich) fused ORFs expressed in *E.coli* using a metal affinity column, and injected into rabbits (Cocalico Biologicals, PA). Anti-Fabp7 or dFabp-B was diluted 1:100, Topro-blue 1:500. All images were captured with the Bio-Rad MRC-1024 Confocal System (Keck Imaging Center, UW-Madison).

Western Blotting

Fly head extracts were lysed and Western blot analysis done as described [27]. Protein samples prepared were subjected to SDS-polyacrylamide gel electrophoresis by separation on a 12–15% gel, transferred to $0.2\ \mu\text{m}$ Protran nitrocellulose membranes (PerkinElmer, Boston, MA), blocked in 5% dried milk powder in 50 mM Tris-HCl, pH 7.5, 15 mM NaCl, 0.5% Tween-20 (TBST), washed briefly in TBST and incubated in primary antibody in TBST overnight at 4°C . Fabp7 or dFabp antibodies were used at ~1:1500, laminin antibody (Hybridoma bank, U Iowa) ~1:1000. Blots were then washed 3 times in TBST, incubated for 1 hour in anti-rabbit horse radish peroxidase-conjugated secondary antibody (1:7500 in TBST), washed 3 times in TBST and incubated in ECL plus Western blotting detection reagent for 5 minutes based on the manufacturer’s instructions (Thermo Scientific, Rockford, IL). Visualization was performed by chemoluminescent exposure to X-Ray film (Eastman Kodak, Rochester, NY) and quantitation was done using ImageJ (NIH).

Behavioral Testing

Flies were maintained on standard molasses-yeast-cornmeal food at 22°C , and entrained to a 12-h:12-h light:dark cycle for 2-3 d before being assayed for sleep or olfactory avoidance conditioning. For analysis of sleep behavior, ~3–4 day old female flies were assayed using the *Drosophila* Activity Monitoring System (Trikinetics), carried out as previously described [82]. Olfactory avoidance conditioning was done as described [29]. Briefly, the performance index (PI) was calculated by subtracting the number of flies making the incorrect choice from those making the correct one, dividing by the total number of flies, and multiplying by 100. To avoid odor-avoidance biases, we calculate the PI of each single n by taking an average performance of two groups of flies, one group trained with the CS+ being OCT, the

other with the CS+ being MCH. Methylcyclohexanol was diluted 1:100, Octanol was diluted 1:40 in mineral oil. *w(isoC71)* and Fabp transgenics were combined for training and testing in the same environments at the same time for most sleep and memory experiments.

Supporting Information

Figure S1 Western blots of dFabp and Fabp7 protein in fly heads (A) Western blots of fly head lysates showing the diurnal regulation of the dFABP band at 15kD, as compared to a loading control (laminin). 50-100 head equivalents of protein were loaded in each lane. (B) Western blots of *w(isoC71)* (*w*) background strain and Fabp7 overexpressing fly head lysates showing the presence and overexpression of Fabp7 using antibodies raised against Fabp7 at 20°C and 30°C. 50-100 head equivalents of protein per lane. ZT = zeitgeber time; RT = room temperature. (EPS)

Figure S2 Reduced sleep in multiple Fabp7 overexpressing fly strains. Hourly sleep profile of the (A) *w(isoC71)* background strain, (B) a second Fabp7 (103-5) strain, and (C) a third Fabp7 (103-8) strain. Open bars = lights on, filled bars = lights off. Analysis of various sleep parameters: (D) Total sleep over 24 hours, (E) Total daytime sleep, (F) Average daytime sleep bout length, (G) Maximum daytime sleep bout length, (H) Daytime locomotor activity per waking minute. Open bars = *w(isoC71)*, light grey bars = 103-5, dark grey bars = 103-8. Results are mean \pm s.e.m.; $n = 11-14$, $t, P \leq 0.0502$; single asterisk, $P < 0.05$; double asterisk, $P < 0.01$; triple asterisk, $P < 0.001$, t -test. (EPS)

Figure S3 Reduced sleep in multiple dFabp overexpressing fly strains. Hourly diurnal sleep profile of the (A) *w(isoC71)* background strain, (B) a second dFabp (101-2) strain, and (C) a third dFabp (101-6) strain. Open bars = lights on, filled bars = lights off. Analysis of various sleep parameters: (D) Total

sleep over 24 hours, (E) Total daytime sleep, (F) Average daytime sleep bout length, (G) Maximum daytime sleep bout length, (H) Daytime locomotor activity per waking minute. Open bars = *w(isoC71)*, light grey bars = 101-2, dark grey bars = 101-6. Results are mean \pm s.e.m.; $n = 13$ *w(isoC71)*, $n = 22-24$; single asterisk, $P < 0.05$; double asterisk, $P < 0.01$; triple asterisk, $P < 0.001$, t -test. (EPS)

Figure S4 Acquisition and retrieval in Fabp flies. (A) Fabp7 (grey fill) and (B) dFabp (dark grey fill) flies exhibit normal learning after a single training trial (1X). Results are mean \pm s.e.m.; $n = 8$ groups. (C) An acute 30 minute 34°C heat shock (+) of Fabp7 flies 2 hours prior to testing inhibits LTM retrieval following 10X Spaced (10XSP) training of flies maintained at 20°C for 6 days, compared to non-heat shock (-) controls. Results are mean \pm s.e.m.; $n = 8$ groups. Asterisk $P < 0.05$, t -test. (D) An acute 30 minute 34°C heat shock (+) of the *w(isoC71)* background strain flies 2 h prior to testing has no effect on LTM retrieval following 10X Spaced (10XSP) training of flies maintained at 20°C for 6 days, compared to non-heat shock (-) controls. Results are mean \pm s.e.m.; $n = 8$ groups. Grey bars, Fabp7, open bars, *w(isoC71)* background strain. d1 = day 1, d6 = day 6. (EPS)

Acknowledgments

We thank H. Zhou and C. Worley for technical assistance on Western blot and Immunocytochemical assays. J. Elliott for advice on Immunocytochemistry, T. Tubon for advice on antibody generation, E. Drier for advice on behavioral assays.

Author Contributions

Conceived and designed the experiments: JRG JCPY PJS CFL. Performed the experiments: JRG WMV. Analyzed the data: JRG WMV. Contributed reagents/materials/analysis tools: PJS. Wrote the paper: JRG.

References

- Knutson KL, Spiegel K, Penev P, Van Cauter E (2007) The metabolic consequences of sleep deprivation. *Sleep Med Rev* 11(3): 163–178.
- Schulze G (2004) Sleep protects excitatory cortical circuits against oxidative damage. *Med Hypotheses* 63(2): 203–207.
- Stickgold R (2005) Sleep-dependent memory consolidation. *Nature* 437(7063): 1272–1278.
- Vertes RP, Siegel JM (2005) Time for the sleep community to take a critical look at the purported role of sleep in memory processing. *Sleep* 28(10): 1228–9; discussion 1230-3.
- Chaudhury D, Colwell CS (2002) Circadian modulation of learning and memory in fear-conditioned mice. *Behav Brain Res* 133(1): 95–108.
- Lyons LC, Rawashdeh O, Eskin A (2006) Non-ocular circadian oscillators and photoreceptors modulate long term memory formation in aplysia. *J Biol Rhythms* 21(4): 245–255.
- Lyons LC, Roman G (2008) Circadian modulation of short-term memory in drosophila. *Learn Mem* 16(1): 19–27.
- Li X, Yu F, Guo A (2009) Sleep deprivation specifically impairs short-term olfactory memory in drosophila. *Sleep* 32(11): 1417–1424.
- Ganguly-Fitzgerald I, Donlea J, Shaw PJ (2006) Waking experience affects sleep need in drosophila. *Science* 313(5794): 1775–1781.
- McDermott CM, LaHoste GJ, Chen C, Musto A, Bazan NG, et al. (2003) Sleep deprivation causes behavioral, synaptic, and membrane excitability alterations in hippocampal neurons. *J Neurosci* 23(29): 9687–9695.
- Graves LA, Heller EA, Pack AI, Abel T (2003) Sleep deprivation selectively impairs memory consolidation for contextual fear conditioning. *Learn Mem* 10(3): 168–176.
- Smith C, Kelly G (1988) Paradoxical sleep deprivation applied two days after end of training retards learning. *Physiol Behav* 43(2): 213–216.
- Donlea JM, Ramanan N, Shaw PJ (2009) Use-dependent plasticity in clock neurons regulates sleep need in drosophila. *Science* 324(5923): 105–108.
- Gerstner JR, Lyons LC, Wright KP, Jr., Loh DH, Rawashdeh O, et al. (2009) Cycling behavior and memory formation. *J Neurosci* 29(41): 12824–12830.
- Tononi G, Cirelli C (2001) Modulation of brain gene expression during sleep and wakefulness: A review of recent findings. *Neuropsychopharmacology* 25(5 Suppl): S28–35.
- Cirelli C, Faraguna U, Tononi G (2006) Changes in brain gene expression after long-term sleep deprivation. *J Neurochem* 98(5): 1632–1645.
- Terao A, Greco MA, Davis RW, Heller HC, Kilduff TS (2003) Region-specific changes in immediate early gene expression in response to sleep deprivation and recovery sleep in the mouse brain. *Neuroscience* 120(4): 1115–1124.
- Terao A, Wisor JP, Peyron C, Apte-Deshpande A, Wurts SW, et al. (2005) Gene expression in the rat brain during sleep deprivation and recovery sleep: An affymetrix GeneChip(R) study. *Neuroscience*.
- Zimmerman JE, Rizzo W, Shockley KR, Raizen DM, Naidoo N, et al. (2006) Multiple mechanisms limit the duration of wakefulness in drosophila brain. *Physiol Genomics* 27(3): 337–350.
- Bozon B, Davis S, Laroche S (2003) A requirement for the immediate early gene zif268 in reconsolidation of recognition memory after retrieval. *Neuron* 40(4): 695–701.
- Jones MW, Errington ML, French PJ, Fine A, Bliss TV, et al. (2001) A requirement for the immediate early gene Zif268 in the expression of late LTP and long-term memories. *Nat Neurosci* 4(3): 289–296.
- Hall J, Thomas KL, Everitt BJ (2001) Cellular imaging of zif268 expression in the hippocampus and amygdala during contextual and cued fear memory retrieval: Selective activation of hippocampal CA1 neurons during the recall of contextual memories. *J Neurosci* 21(6): 2186–2193.
- Ribeiro S, Goyal V, Mello CV, Pavlides C (1999) Brain gene expression during REM sleep depends on prior waking experience. *Learn Mem* 6(5): 500–508.
- Ribeiro S, Mello CV, Velho T, Gardner TJ, Jarvis ED, et al. (2002) Induction of hippocampal long-term potentiation during waking leads to increased extrahippocampal zif-268 expression during ensuing rapid-eye-movement sleep. *J Neurosci* 22(24): 10914–10923.
- Gilestro GF, Tononi G, Cirelli C (2009) Widespread changes in synaptic markers as a function of sleep and wakefulness in drosophila. *Science* 324(5923): 109–112.

26. Gerstner JR, Vander Heyden WM, Lavaute TM, Landry CF (2006) Profiles of novel diurnally regulated genes in mouse hypothalamus: Expression analysis of the cysteine and histidine-rich domain-containing, zinc-binding protein 1, the fatty acid-binding protein 7 and the GTPase, ras-like family member 11b. *Neuroscience* 139(4): 1435–1448.
27. Gerstner JR, Bremer QZ, Vander Heyden WM, Lavaute TM, Yin JC, et al. (2008) Brain fatty acid binding protein (Fabp7) is diurnally regulated in astrocytes and hippocampal granule cell precursors in adult rodent brain. *PLoS ONE* 3(2): e1631.
28. Ceriani MF, Hogenesch JB, Yanovsky M, Panda S, Straume M, et al. (2002) Genome-wide expression analysis in drosophila reveals genes controlling circadian behavior. *J Neurosci* 22(21): 9305–9319.
29. Drier EA, Tello MK, Cowan M, Wu P, Blace N, et al. (2002) Memory enhancement and formation by atypical PKM activity in drosophila melanogaster. *Nat Neurosci* 5(4): 316–324.
30. Arai Y, Funatsu N, Numayama-Tsuruta K, Nomura T, Nakamura S, et al. (2005) Role of Fabp7, a downstream gene of Pax6, in the maintenance of neuroepithelial cells during early embryonic development of the rat cortex. *J Neurosci* 25(42): 9752–9761.
31. Anthony TE, Mason HA, Gridley T, Fishell G, Heintz N (2005) Brain lipid-binding protein is a direct target of notch signaling in radial glial cells. *Genes Dev* 19(9): 1028–1033.
32. Costa RM, Honjo T, Silva AJ (2003) Learning and memory deficits in notch mutant mice. *Curr Biol* 13(15): 1348–1354.
33. Ge X, Hannan F, Xie Z, Feng C, Tully T, et al. (2004) Notch signaling in drosophila long-term memory formation. *Proc Natl Acad Sci U S A* 101(27): 10172–10176.
34. Presente A, Boyles RS, Serway CN, de Belle JS, Andres AJ (2004) Notch is required for long-term memory in drosophila. *Proc Natl Acad Sci U S A* 101(6): 1764–1768.
35. Owada Y, Abdelwahab SA, Kitanaka N, Sakagami H, Takano H, et al. (2006) Altered emotional behavioral responses in mice lacking brain-type fatty acid-binding protein gene. *Eur J Neurosci* 24(1): 175–187.
36. Watanabe A, Toyota T, Owada Y, Hayashi T, Iwayama Y, et al. (2007) Fabp7 maps to a quantitative trait locus for a schizophrenia endophenotype. *PLoS Biol* 5(11): e297.
37. Zhong J, Zhang T, Bloch LM (2006) Dendritic mRNAs encode diversified functionalities in hippocampal pyramidal neurons. *BMC Neurosci* 7: 17.
38. Owada Y, Yoshimoto T, Kondo H (1996) Increased expression of the mRNA for brain- and skin-type but not heart-type fatty acid binding proteins following kainic acid systemic administration in the hippocampal glia of adult rats. *Brain Res Mol Brain Res* 42(1): 156–160.
39. Barco A, Patterson S, Alarcon JM, Gromova P, Mata-Roig M, et al. (2005) Gene expression profiling of facilitated L-LTP in VP16-CREB mice reveals that BDNF is critical for the maintenance of LTP and its synaptic capture. *Neuron* 48(1): 123–137.
40. Tafti M, Franken P (2002) Invited review: Genetic dissection of sleep. *J Appl Physiol* 92(3): 1339–1347.
41. Cirelli C (2009) The genetic and molecular regulation of sleep: From fruit flies to humans. *Nat Rev Neurosci* 10(8): 549–560.
42. Gerstner JR (2010) The aging clock: To 'BMAL'icious toward learning and memory. *Aging (Albany NY)* 2(5): 251–254.
43. Gerstner JR, Yin JC (2010) Circadian rhythms and memory formation. *Nat Rev Neurosci* 11(8): 577–588.
44. Kondratova AA, Dubrovsky YV, Antoch MP, Kondratov RV (2010) Circadian clock proteins control adaptation to novel environment and memory formation. *Aging (Albany NY)* 2(5): 285–297.
45. Sakai T, Tamura T, Kitamoto T, Kidokoro Y (2004) A clock gene, period, plays a key role in long-term memory formation in drosophila. *Proc Natl Acad Sci U S A* 101(45): 16058–16063.
46. Shioda N, Yamamoto Y, Watanabe M, Binas B, Owada Y, et al. (2010) Heart-type fatty acid binding protein regulates dopamine D2 receptor function in mouse brain. *J Neurosci* 30(8): 3146–3155.
47. Huang ZL, Urade Y, Hayaishi O (2007) Prostaglandins and adenosine in the regulation of sleep and wakefulness. *Curr Opin Pharmacol* 7(1): 33–38.
48. Pandey HP, Ram A, Matsumura H, Hayaishi O (1995) Concentration of prostaglandin D2 in cerebrospinal fluid exhibits a circadian alteration in conscious rats. *Biochem Mol Biol Int* 37(3): 431–437.
49. Ram A, Pandey HP, Matsumura H, Kasahara-Orita K, Nakajima T, et al. (1997) CSF levels of prostaglandins, especially the level of prostaglandin D2, are correlated with increasing propensity towards sleep in rats. *Brain Res* 751(1): 81–89.
50. Inoue S, Honda K, Komoda Y, Uchizono K, Ueno R, et al. (1984) Differential sleep-promoting effects of five sleep substances nocturnally infused in unrestrained rats. *Proc Natl Acad Sci U S A* 81(19): 6240–6244.
51. Mizoguchi A, Eguchi N, Kimura K, Kiyohara Y, Qu WM, et al. (2001) Dominant localization of prostaglandin D receptors on arachnoid trabecular cells in mouse basal forebrain and their involvement in the regulation of non-rapid eye movement sleep. *Proc Natl Acad Sci U S A* 98(20): 11674–11679.
52. Matsumura H, Takahata R, Hayaishi O (1991) Inhibition of sleep in rats by inorganic selenium compounds, inhibitors of prostaglandin D synthase. *Proc Natl Acad Sci U S A* 88(20): 9046–9050.
53. Valenti M, Vigano D, Casico MG, Rubino T, Steardo L, et al. (2004) Differential diurnal variations of anandamide and 2-arachidonoyl-glycerol levels in rat brain. *Cell Mol Life Sci* 61(7-8): 945–950.
54. Murillo-Rodriguez E, Desarnaud F, Prospero-Garcia O (2006) Diurnal variation of arachidonylethanolamine, palmitoylethanolamide and oleoylethanolamide in the brain of the rat. *Life Sci* 79(1): 30–37.
55. Glaser ST, Kaczocha M (2009) Temporal changes in mouse brain fatty acid amide hydrolase activity. *Neuroscience* 163(2): 594–600.
56. Mazzola C, Medalie J, Scherma M, Panlilio LV, Solinas M, et al. (2009) Fatty acid amide hydrolase (FAAH) inhibition enhances memory acquisition through activation of PPAR-alpha nuclear receptors. *Learn Mem* 16(5): 332–337.
57. Campolongo P, Roozendaal B, Trezza V, Cuomo V, Astarita G, et al. (2009) Fat-induced satiety factor oleoylethanolamide enhances memory consolidation. *Proc Natl Acad Sci U S A* 106(19): 8027–8031.
58. Kaczocha M, Glaser ST, Deutsch DG (2009) Identification of intracellular carriers for the endocannabinoid anandamide. *Proc Natl Acad Sci U S A* 106(15): 6375–6380.
59. Tan NS, Shaw NS, Vinckenbosch N, Liu P, Yasmin R, et al. (2002) Selective cooperation between fatty acid binding proteins and peroxisome proliferator-activated receptors in regulating transcription. *Mol Cell Biol* 22(14): 5114–5127.
60. Helledie T, Antonius M, Sorensen RV, Hertzog AV, Bernlohr DA, et al. (2000) Lipid-binding proteins modulate ligand-dependent trans-activation by peroxisome proliferator-activated receptors and localize to the nucleus as well as the cytoplasm. *J Lipid Res* 41(11): 1740–1751.
61. Wolfrum C, Borrmann CM, Borchers T, Spener F (2001) Fatty acids and hypolipidemic drugs regulate peroxisome proliferator-activated receptors alpha- and gamma-mediated gene expression via liver fatty acid binding protein: A signaling path to the nucleus. *Proc Natl Acad Sci U S A* 98(5): 2323–2328.
62. Schroeder F, Petrescu AD, Huang H, Atshaves BP, McIntosh AL, et al. (2008) Role of fatty acid binding proteins and long chain fatty acids in modulating nuclear receptors and gene transcription. *Lipids* 43(1): 1–17.
63. Oishi K, Shirai H, Ishida N (2005) CLOCK is involved in the circadian transactivation of peroxisome-proliferator-activated receptor alpha (PPARalpha) in mice. *Biochem J* 386(Pt 3): 575–581.
64. Canaple L, Rambaud J, Dkhissi-Benyahya O, Rayet B, Tan NS, et al. (2006) Reciprocal regulation of brain and muscle arnt-like protein 1 and peroxisome proliferator-activated receptor alpha defines a novel positive feedback loop in the rodent liver circadian clock. *Mol Endocrinol* 20(8): 1715–1727.
65. Kohsaka A, Bass J (2007) A sense of time: How molecular clocks organize metabolism. *Trends Endocrinol Metab* 18(1): 4–11.
66. Schmutz I, Ripperger JA, Baeiriswyl-Aebischer S, Albrecht U (2010) The mammalian clock component PERIOD2 coordinates circadian output by interaction with nuclear receptors. *Genes Dev* 24(4): 345–357.
67. Diekelmann S, Born J (2010) The memory function of sleep. *Nat Rev Neurosci* 11(2): 114–126.
68. Smith C, Rose GM (1996) Evidence for a paradoxical sleep window for place learning in the morris water maze. *Physiol Behav* 59(1): 93–97.
69. Mednick SC, Nakayama K, Cantero JL, Atienza M, Levin AA, et al. (2002) The restorative effect of naps on perceptual deterioration. *Nat Neurosci* 5(7): 677–681.
70. Mednick S, Nakayama K, Stickgold R (2003) Sleep-dependent learning: A nap is as good as a night. *Nat Neurosci* 6(7): 697–698.
71. Gais S, Plihal W, Wagner U, Born J (2000) Early sleep triggers memory for early visual discrimination skills. *Nat Neurosci* 3(12): 1335–1339.
72. Stickgold R, James L, Hobson JA (2000) Visual discrimination learning requires sleep after training. *Nat Neurosci* 3(12): 1237–1238.
73. Wamsley EJ, Tucker MA, Payne JD, Stickgold R (2010) A brief nap is beneficial for human route-learning: The role of navigation experience and EEG spectral power. *Learn Mem* 17(7): 332–336.
74. Tucker MA, Fishbein W (2008) Enhancement of declarative memory performance following a daytime nap is contingent on strength of initial task acquisition. *Sleep* 31(2): 197–203.
75. Fenn KM, Nusbaum HC, Margoliash D (2003) Consolidation during sleep of perceptual learning of spoken language. *Nature* 425(6958): 614–616.
76. Brawn TP, Fenn KM, Nusbaum HC, Margoliash D (2008) Consolidation of sensorimotor learning during sleep. *Learn Mem* 15(11): 815–819.
77. Cai DJ, Shuman T, Gorman MR, Sage JR, Anagnostaras SG (2009) Sleep selectively enhances hippocampus-dependent memory in mice. *Behav Neurosci* 123(4): 713–719.
78. Brawn TP, Nusbaum HC, Margoliash D (2010) Sleep-dependent consolidation of auditory discrimination learning in adult starlings. *J Neurosci* 30(2): 609–613.
79. Datta S, Li G, Auerbach S (2008) Activation of phasic pontine-wave generator in the rat: A mechanism for expression of plasticity-related genes and proteins in the dorsal hippocampus and amygdala. *Eur J Neurosci* 27(7): 1876–1892.
80. Wetzel W, Wagner T, Balschun D (2003) REM sleep enhancement induced by different procedures improves memory retention in rats. *Eur J Neurosci* 18(9): 2611–2617.
81. Thum AS, Jenett A, Ito K, Heisenberg M, Tanimoto H (2007) Multiple memory traces for olfactory reward learning in drosophila. *J Neurosci* 27(41): 11132–11138.
82. Andretic R, Shaw PJ (2005) Essentials of sleep recordings in drosophila: Moving beyond sleep time. *Methods Enzymol* 393: 759–772.