

# Violation of the Leggett–Garg inequality with weak measurements of photons

M. E. Goggin<sup>a,b,1</sup>, M. P. Almeida<sup>a</sup>, M. Barbieri<sup>a,c</sup>, B. P. Lanyon<sup>a</sup>, J. L. O’Brien<sup>d</sup>, A. G. White<sup>a</sup>, and G. J. Pryde<sup>e</sup>

<sup>a</sup>Department of Physics and Centre for Quantum Computer Technology, University of Queensland, Brisbane 4072, Australia; <sup>b</sup>Physics Department, Truman State University, Kirksville, MO 63501; <sup>c</sup>Groupe d’Optique Quantique, Laboratoire Charles Fabry, Institut d’Optique—Graduate School, F91127 Palaiseau Cedex, France; <sup>d</sup>Centre for Quantum Photonics, H. H. Wills Physics Laboratory, and Department of Electrical and Electronic Engineering, University of Bristol, Merchant Venturers Building, Woodland Road, Bristol, BS8 1UB, United Kingdom; and <sup>e</sup>Centre for Quantum Dynamics and Centre for Quantum Computer Technology, Griffith University, Brisbane, 4111, Australia

Edited by Yakir Aharonov, Tel Aviv University, Tel Aviv, Israel, and approved December 15, 2010 (received for review April 28, 2010)

**By weakly measuring the polarization of a photon between two strong polarization measurements, we experimentally investigate the correlation between the appearance of anomalous values in quantum weak measurements and the violation of realism and nonintrusiveness of measurements. A quantitative formulation of the latter concept is expressed in terms of a Leggett–Garg inequality for the outcomes of subsequent measurements of an individual quantum system. We experimentally violate the Leggett–Garg inequality for several measurement strengths. Furthermore, we experimentally demonstrate that there is a one-to-one correlation between achieving strange weak values and violating the Leggett–Garg inequality.**

entanglement | quantum communication | quantum nonlocality | quantum measurement

There has been much debate in quantum physics over the question of whether measurable quantities have definite values prior to their measurement. Key ideas addressing this question include the Bell inequality, which considers correlations between measurements on components of a composite system that are space-like separated (1, 2) and contextuality tests, which examine whether identical experiments produce results in different “classically equivalent” contexts (3, 4). A conceptually elegant extension to these ideas is the Leggett–Garg inequality (LGI) (5), which is an inequality constructed from the correlation functions of a series of three consecutive measurements on a single system. Leggett and Garg derive limits based on the joint assumptions of (i) macroscopic realism: An observable for a system will have a definite value at all times; and (ii) noninvasive measurement: It is possible to determine this value with arbitrarily small disturbance on the subsequent evolution of the system. The limits on the value of the inequality derived from these assumptions differ from the predictions of quantum mechanics. Thus the LGI tests the limits of measurement and macroscopic realism.

Here we present an experimental test of a generalized LGI using weak measurements (6–9) of the polarization of single photons and measure violations by up to 14 standard deviations. Additionally, we experimentally demonstrate a one-to-one relation (10, 11) between LGI violations and strange weak-valued measurements (6–8), which also arise from the inability to assign values to physical quantities between an earlier and a later measurement.

Testing the LGI requires monitoring the system without projecting it into a specific state. For a quantum system a quantum nondemolition (QND) experiment (12–14) would be one way to do this. But QND measurements are not the only way to perform a noninvasive measurement. A generalization of the QND measurement is the so-called weak measurement (6). A weak measurement is one for which it is possible to adjust the strength of the measurement and, in principle, to reduce the back action on the system to an arbitrarily small amount. In other words, a weak measurement is one for which the level of “invasiveness” can be controlled.

Shortly after Leggett and Garg introduced their inequality, Aharonov et al. (7) suggested that observing the result of a weak measurement conditioned on a specific result of a separate projective measurement leads to unusual results. One unusual property, dubbed strange weak values, is that the value assigned in this way may lie outside the eigenspectrum of the observable being measured (7). Because such strange weak values have been observed (15–19), the idea that the measured value lies outside the operator’s eigenspectrum raises questions about macroscopic realistic descriptions of the system’s state and of the measurement process (20, 21). In this sense, strange weak values explore the same concepts (or raise the same problems) as the LGI. A formal connection between a generalized LGI and weak values has been recently proposed by Williams and Jordan (10, 11). Specifically, they propose that the LGI is violated if and only if the experiment yields a strange weak value.

We want to be clear about what we mean by the term macroscopic realism and how it applies to systems of an arbitrary size. We use the same definition of macroscopic realism as Leggett and Garg: A system is described by a probability distribution for definite values of observables prior to measurement (5). The origin of the term macroscopic realism relates to the fact that one expects an ensemble of very large systems to be described by a classical distribution of definite values. However, this expectation in no way constrains the description to this class of objects. The term macroscopic realism, therefore, is not fundamentally about the size of the investigated system (which can be any size, in accordance with the generality of the Leggett–Garg treatment) but rather about the assignment of definite values to measurable quantities, either definitely (e.g., speed of a tennis ball) or probabilistically (e.g., speeds of particles in an ideal gas). This has been pointed out previously in a theoretical description connecting Leggett–Garg inequalities for Rydberg atoms (22) to hidden variables. The previous discussion holds true for the generalization proposed in ref. 8, where the probability distribution is also parametrized by a hidden-variable.

Recently, Jordan and coworkers (8, 10, 11) proposed a correlation function that generalizes the original Leggett–Garg correlation function to include nonprojective weak measurements. With this generalization, the ability to test the LGI at differing measurement strengths becomes possible, allowing for an experimental investigation of the measurement process. The generalized Leggett–Garg correlation function (8, 10, 11) is

$$B = \langle \mathcal{M}_a \mathcal{M}_b \rangle + \langle \mathcal{M}_b \mathcal{M}_c \rangle - \langle \mathcal{M}_a \mathcal{M}_c \rangle, \quad [1]$$

Author contributions: M.E.G., M.P.A., M.B., B.P.L., J.L.O., A.G.W., and G.J.P. designed research; M.E.G., M.P.A., and M.B. performed research; M.E.G., M.P.A., M.B., B.P.L., and G.J.P. analyzed data; and M.E.G., M.P.A., M.B., B.P.L., J.L.O., A.G.W., and G.J.P. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. E-mail: mgoggin@truman.edu.


where  $\{\mathcal{M}_a, \mathcal{M}_b, \mathcal{M}_c\}$  is a set of three consecutive weak measurements on the photon polarization. We use letters to denote the sequence of measurements in order to distinguish the ordering of the measurements from the physical quantities we measured, the Stokes' parameters, which are denoted with numerical subscripts according to conventional notation. For a two-level system with the properties of both macroscopic realism and noninvasive detection,  $B$  must satisfy  $-3 \leq B \leq 1$ , which is the LGI. On the other hand, quantum measurements produce values of  $B$  satisfying (8):  $-3 \leq B \leq 1.5$ . Thus, there is a violation of the LGI if  $1 < B \leq 1.5$ . Although the original formulation of the generalized LGI, Eq. 1, is in terms of three consecutive weak measurements, the detailed analysis presented in refs. 8, 10, and 11 shows that, according to quantum mechanics, the violation of the LGI does not depend on the measurement strength of either  $\mathcal{M}_a$  or  $\mathcal{M}_c$ . Thus, we can relax the requirement that all three measurements are weak and perform strong projective measurements for  $\mathcal{M}_a$  or  $\mathcal{M}_c$  thereby simplifying the experiment.

We feel it necessary to point out that the LGI as we have tested it differs from sequential contextuality tests (23). Contextuality tests using sequential measurements require the measured observables be compatible. The LGI has no such requirement; it requires noninvasive measurement of incompatible observables. Therefore some of the criticisms levelled at contextuality tests (see, e.g., ref. 24 and references therein) do not apply to the LGI. It may be that there is a deeper connection between contextuality and noninvasive measurement, but we leave that for future work.

A conceptual scheme of the experimental setup is shown in Fig. 1. We input a "signal" photon in the state  $|\sigma_{in}\rangle = \cos\frac{\theta}{2}|H\rangle + \sin\frac{\theta}{2}|V\rangle$  where  $|H\rangle$  and  $|V\rangle$  represent horizontal and vertical polarization states. We consider  $|\sigma_{in}\rangle$  as the state prepared by the first measurement  $\mathcal{M}_a$ ; in this way, we can deterministically assign the value 1 to  $\mathcal{M}_a$ . The weak measurement,  $\mathcal{M}_b$ , measures the  $\mathcal{S}_1$  Stokes' parameter of the signal photon (14), corresponding to the degree of polarization in the horizontal-vertical basis. The final measurement,  $\mathcal{M}_c$ , is the Stokes' parameter  $\mathcal{S}_2$  of the signal photon that quantifies the degree of polarization in the diagonal-antidiagonal basis  $|D\rangle = \frac{1}{\sqrt{2}}(|H\rangle + |V\rangle)$ ,  $|A\rangle = \frac{1}{\sqrt{2}}(|H\rangle - |V\rangle)$ . Thus, in our case we can rewrite Eq. 1 as (10, 11)

$$B = \langle \mathcal{S}_1 \rangle + \langle \mathcal{S}_1 \mathcal{S}_2 \rangle - \langle \mathcal{S}_2 \rangle. \quad [2]$$

Note that  $\mathcal{S}_1$  is a weak measurement in the sense of Jordan et al. (8), and  $\mathcal{S}_2$  is a projective measurement.



**Fig. 1.** Conceptual representation of the experiment. A "signal" photon is prepared with an arbitrary linearly polarized state using a half waveplate ( $\lambda/2$ ),  $\mathcal{M}_a$ . A weak measurement of the polarization is then made ( $\mathcal{M}_b$ ) by interacting the signal photon with a "meter" photon via a C-Sign gate, which operates via a measurement-induced nonlinearity (14, 25–29). Conditional on detecting a single photon in each of the two output modes (a coincidence measurement) the gate ideally performs the operation  $|H\rangle\langle H| \otimes \sigma_y + |V\rangle\langle V| \otimes \sigma_x$  on the two-qubit polarization Hilbert space  $\mathcal{H}_{\text{signal}} \otimes \mathcal{H}_{\text{meter}}$ , where  $\sigma_y$  ( $\sigma_x$ ) is the 1-qubit identity (Pauli  $\mathcal{X}$ ) operator and  $|H\rangle = |0\rangle$ ,  $|V\rangle = |1\rangle$ . See the main text for a full description of  $\mathcal{M}_a$ ,  $\mathcal{M}_b$ , and  $\mathcal{M}_c$  and *Materials and Methods* for more experimental details.

The weak measurement is implemented using a nondeterministic linear optical controlled-sign gate (25–29). The polarization of the signal qubit  $|\sigma_{in}\rangle$  in the gate's control mode becomes entangled with that of a second, "meter," photon in the target mode. A subsequent projective measurement of the meter photon realizes the weak measurement of the signal. The strength of the measurement can be set by changing the meter input state  $|\mu_{in}\rangle = \gamma|D\rangle + \tilde{\gamma}|A\rangle$  (with  $\gamma^2 + \tilde{\gamma}^2 = 1$ , and  $\gamma, \tilde{\gamma} \in \mathbb{R}$  without loss of generality). The strength of the measurement is quantified by the knowledge (14),  $K = 2\gamma^2 - 1$ . When  $K = 1$  ( $\gamma = 1$ ), full information is extracted about the polarization of the signal photon, while when  $K = 0$  ( $\gamma = 1/\sqrt{2}$ ), the photon is left completely undisturbed and unmeasured. Between those two extremes, the photon polarization undergoes partial disturbance, and one can extract the average value of  $\mathcal{S}_1$  as (14):

$$\langle \mathcal{S}_1 \rangle = \frac{P(D) - P(A)}{K}, \quad [3]$$

where  $P(D)$  and  $P(A)$  are the probabilities of measuring the values  $D$  and  $A$ , respectively, for the meter photon. The weak character is reflected in the fact that an increasing number of copies are necessary to estimate  $\langle \mathcal{S}_1 \rangle$  to a given precision as  $K$  decreases. The weak value (6, 7) is calculated by postselecting on a measurement of  $\mathcal{M}_c$  that results in the value  $D$  for the signal photon. In our implementation, we considered the weak value:

$${}_D\langle \mathcal{S}_1 \rangle = \frac{P(D|D) - P(A|D)}{K}, \quad [4]$$

where  $P(A|D)$  is the probability of measuring the value  $A$  for the meter photon, conditioned on the postselection of the signal photon registering a value of  $D$  and similarly for  $P(D|D)$ . We expect strange weak values ( ${}_D\langle \mathcal{S}_1 \rangle > 1$ ) to emerge when  $|\sigma_{in}\rangle$  approaches the antidiagonal state  $|A\rangle$ . We refer to weak values as the postselected results of any weak measurement. Notice that these coincide with the original formulation of Aharonov et al. in the limit  $K \rightarrow 0$  (7). Even for finite strengths, strange weak values may appear (13, 30). Fig. 2A shows the measured values for the generalized LG correlation function,  $B$ , defined in Eq. 2 and the weak values,  $WV$ , for the choice  $K = 0.5445 \pm 0.0083$  as  $\theta$  is varied in the range  $[0, 2\pi]$ . The maximum value we observe is  $B_{\text{max}} = 1.312 \pm 0.022$ , which is 14 standard deviations above the classical limit of  $B = 1$ . The results show that there is a clear correlation between the violation of the LGI and the appearance of strange positive weak values. There are discrepancies between the theoretical curves and the data points for values of  $\theta$  where we do not expect to see a violation of the LGI. We discuss these in more detail in *Materials and Methods*. We also see strange negative weak values corresponding to values of  $B$  that do not violate the LGI of Eq. 2. This does not contradict the prediction of Williams and Jordan (10, 11) because they predict that there are strange weak values when a generalized LGI is violated. There are four distinct generalized LGIs corresponding to different assignments of the  $\pm 1$  eigenvalues for the operators in Eq. 2. It is also possible to define four different weak values, by eigenvalue reassignment and varying the final postselection state of the signal. See *Materials and Methods* for more details.

The condition—which applies across the whole parameter range—is that whenever a strange weak value is obtained one of these four LGIs will also be violated. As shown in Fig. 2B, if we assign our eigenvalues such that  $\mathcal{M}_b = -\mathcal{S}_1$  rather than  $\mathcal{M}_b = \mathcal{S}_1$  as in Fig. 2A, a violation of the LGI,  $B > 1$ , corresponds to the strange weak values, in the range where previously there was no violation.

To understand the effect of the measurement strength on violation of the LGI, we decreased the knowledge to  $K = 0.1598 \pm 0.0091$ , Fig. 3. This corresponds to a weaker measure-





macroscopic realism becomes the only condition left to violate. Yet there is little difference between our result with a finite knowledge ( $K = 0.1598 \pm 0.0091$ ) and the theoretical limit of zero knowledge. In fact, we see a significant violation with a knowledge of 0.5445. So with a better than 50/50 shot of knowing  $\mathcal{S}_1$ , we get a violation of the LGI.

Bounded weak values also require noninvasive measurement to be observed. (10). Yet the weak values become less bounded as the measurement gets weaker. This leaves open questions about weak values as markers for a violation of local realism.

More significantly, both the LGI violation and strange weak values disappear when the measurement is strong enough. Presumably the system did not become macroscopically real as the measurement became stronger; the measurement forced a definite value on the system. These results put into the spotlight questions that are as old as quantum theory. Is measurement the only necessary condition for reality? Is macroscopic realism a superfluous condition? To answer this last question we need to devise an experimentally meaningful test of macroscopic realism that is independent of conditions on measurement strength.

## Materials and Methods

**Generalized Leggett-Garg Inequalities and Corresponding Weak Value Measurements.** Here we give more details on the different LGIs and weak value expressions that can be defined in our experiment. The two measurement operators in Eq. 2 of the main text have eigenvalues of  $\pm 1$ , corresponding to two possible outcomes. Experimentally, two detectors are required for each measurement: one at the transmission output port of a polarizing beamsplitter and the other at the reflection output port. We must choose how to assign clicks at each detector to the eigenvalues. There are two choices for each measurement, yielding four possible configurations in total and subsequently four distinct LGIs:

$$B_1 = +\langle \mathcal{S}_1 \rangle + \langle \mathcal{S}_1 \mathcal{S}_2 \rangle - \langle \mathcal{S}_2 \rangle, \quad [5]$$

$$B_2 = -\langle \mathcal{S}_1 \rangle - \langle \mathcal{S}_1 \mathcal{S}_2 \rangle - \langle \mathcal{S}_2 \rangle, \quad [6]$$

$$B_3 = +\langle \mathcal{S}_1 \rangle - \langle \mathcal{S}_1 \mathcal{S}_2 \rangle + \langle \mathcal{S}_2 \rangle, \quad [7]$$

$$B_4 = -\langle \mathcal{S}_1 \rangle + \langle \mathcal{S}_1 \mathcal{S}_2 \rangle + \langle \mathcal{S}_2 \rangle. \quad [8]$$

A violation of the assumptions of Leggett and Garg occurs if any of the above expressions exceeds 1. This is shown in (10). Briefly, if we write  $\mathcal{M}_q$ , with  $q = a, b, c$  as

$$\mathcal{M}_q = \mathcal{S}_j + \epsilon_j, \quad q = a, b, c \quad \text{and} \quad j = 1, 2$$

where  $\epsilon$  is "noise" from the experiment. The assumptions of macroscopic realism and noninvasive measurement imply

$$\langle \epsilon_j \rangle = 0$$

and

$$\langle \mathcal{S}_j \epsilon_k \rangle = 0, j, k = 1, 2.$$

Therefore

$$-1 \leq \langle \mathcal{M}_q \mathcal{M}_{q'} \rangle \leq 1,$$

giving the limits of  $\langle B \rangle$ .

We can also define four weak values for the signal photon, given the two choices of eigenvalue assignment and two choices of the postselected signal state:

$$WV_1 = {}_D \langle \mathcal{S}_1 \rangle = \frac{P(D|D) - P(A|D)}{K}, \quad [9]$$

$$WV_2 = {}_D \langle -\mathcal{S}_1 \rangle = \frac{P(A|D) - P(D|D)}{K}, \quad [10]$$

$$WV_3 = {}_A \langle \mathcal{S}_1 \rangle = \frac{P(D|A) - P(A|A)}{K}, \quad [11]$$


$$WV_4 = {}_A \langle -\mathcal{S}_1 \rangle = \frac{P(A|A) - P(D|A)}{K}, \quad [12]$$

where  $P(A|D)$  is the probability of measuring the value  $A$  for the meter photon, conditioned on the postselection of the signal photon in state  $D$  and similarly for  $P(D|D)$ . Thus we have a weak value for each LG inequality. The condition that holds for all input signal states (parameterized by  $\theta$ ) is that whenever a weak value is observed one of the LGIs is violated and vice versa. Showing all eight of the above relations would produce plots that would be very difficult to read. The interested reader can see the missing relations by imagining Figs. 2 and 3 horizontally reflected about a line through  $180^\circ$ . We also note that these LGIs and weak values would produce the mirror image (horizontally again) of Fig. 4 of the main text with the horizontal range changed from  $[180^\circ, 360^\circ]$  to  $[0^\circ, 180^\circ]$ . Finally we note that future experiments and investigation could consider more general measurements with more than two outcomes.


**Experimental Details.** Here we give a simple description of our apparatus, illustrated in Fig. 7. Single photons are produced in pairs by spontaneous parametric down conversion (SPDC) in a bismuth borate (BiBO) nonlinear crystal. The pump beam is obtained by frequency doubling a 100-fs pulsed Ti:Sa laser lasing at 820 nm, with a repetition rate 82 MHz and bandwidth 8nm. We used a pump power of approximately 50 mW in order to reduce double pair emission to a negligible level. Single modes are selected by narrowband interference filters (IF) and fiber launchers (FL), coupling them to single mode fibers. The rate of pair production is approximately  $50 \text{ s}^{-1}$  at the output of the gate.

Their original  $H$  polarization state is recovered by means of polarization controllers in the optical fibers between the SPDC source and the free-space output couplers and then prepared by means of a glan-taylor polarizing beamsplitter (PBS) and half-wave plate ( $\lambda/2$ ). This state preparation coincides with the first measurement  $\mathcal{M}_a$ , which therefore assumes the value  $+1$  deterministically.

The photons then arrive on our C-Sign gate. This is based on a single partially polarizing beam splitter (PPBS), whose transmittivities for the horizontal,  $H$ , and vertical,  $V$ , polarizations are ideally  $\eta_H = 1$  and  $\eta_V = 1/\sqrt{3}$  (25–27). Due to quantum interference, the event  $|H_s H_m\rangle$  acquires a  $\pi$  phase shift, while the others are left unaltered—here  $s$  and  $m$  label the signal and the meter. Polarization-dependent loss induced by the gate can be either compensated directly in the state preparation (as we did for the meter arm) or by



**Fig. 5.** The three correlation functions that comprise the generalized Leggett-Garg correlation function for a range of input states  $|\sigma_{in}\rangle$  parameterized by  $\theta$ . The measurement strength is  $K = 0.5445 \pm 0.0083$ . Error bars show the standard deviation and arise from poissonian counting statistics.




**Fig. 6.** The three correlation functions that comprise the generalized Leggett-Garg correlation function for a range of input states  $|\sigma_{in}\rangle$  parametrized by  $\theta$ . The measurement strength is  $K = 0.1598 \pm 0.0091$ . Error bars show the standard deviation and arise from poissonian counting statistics.

adding extra loss (the second PPBS in the signal arm). The gate is nondeterministic, but its correct functioning is heralded by a coincidence count between two distinct output arms.

The basic principle for the nondestructive measurement  $\mathcal{M}_b$  derives from the action of the gate in the rotated basis  $|D\rangle = (|H\rangle + |V\rangle)/\sqrt{2}$  and  $|A\rangle = (|H\rangle - |V\rangle)/\sqrt{2}$ . If  $|V\rangle_s|D\rangle_m$  are injected, they emerge in the same state at the output. Conversely, when  $|H\rangle_s|D\rangle_m$  is injected, there occurs the phase shift flipping the polarization of the meter from  $|D\rangle_m$  to  $|A\rangle_m$ . Therefore, one can assess the polarization of the signal by a measurement on the meter arm distinguishing between the two rotated polarizations: A measurement of  $|D\rangle_m$  flags the state  $|H\rangle_s$ , the same for the orthogonal complement. This is easily obtained by means of a half-wave plate and a glan-taylor polarizing beam splitter (PBS). Notice that the injection of  $|H\rangle_m$  in the meter leads no information about the state of the signal, because this remains unaffected in any case.

An intermediate situation occurs if the polarization  $|\mu\rangle_m = \gamma|D\rangle_m + \bar{\gamma}|A\rangle_m$  is used. This is unaltered upon injection of  $|V\rangle$  and rotated to  $|\bar{\mu}\rangle = \bar{\gamma}|D\rangle_m + \gamma|A\rangle_m$  in the other case. Nevertheless  $\|\langle \bar{\mu} | \mu \rangle\| < 1$ , in the general case, and the two events can not be perfectly distinguished. We notice that a measure in the  $\{|D\rangle_m, |A\rangle_m\}$  basis is the optimal discrimination strategy (32). Consequently, only partial information can be retrieved about the state of the signal. The measurement is then nondestructive and its strength can be tuned at will by properly setting the polarization  $|\mu\rangle_m$ . Because the signal photon is not destroyed, it can be subsequently measured in the standard way by means of a half-wave plate and PBS, so to perform the final step  $\mathcal{M}_c$ . We note that the measurement results are independent of the detection efficien-



**Fig. 7.** Schematic of experimental setup. See *Experimental Details* for a complete description of the experiment. The calcite beam displacers are in the setup for other experiments. They are not integral to the current experiment and are included here for completeness. The  $\phi$  waveplate is used to adjust the relative phase between the beam displacer interferometers and is an artifact of having the beam displacers.

cies. The operation of the gate is such that we make all three measurements on the same system when we have a coincidence event.

**Error Analysis.** Here we examine the systematic error in the experiment. As was noted in the main text there are values of  $\theta$  for which the experimental results diverge from the theoretical results by more than the standard deviation. This error is due to imperfect mode-matching at the PPBS. We checked the outcomes of the measurement results for each of the measurements independently, as shown in Figs. 5 and 6. We can compare the measured results with the expected results for that measurement, which is known theoretically, which helps us to identify that the error is in  $\delta_1$  but only for certain values of  $\theta$ . This is consistent with our experimental observations that suggest that mode matching is the biggest error factor in the experiment. The mode matching is input-dependent because of a steering effect from rotating the input waveplate. In this case, the mode matching is optimized for the regime of interest,  $180 \leq \theta \leq 360^\circ$  as demonstrated by the fact that the  $\delta_1$  measurement gives the correct result, to within statistical error, in that range. Therefore, we claim violation for the LGIs involving  $B_1$  and  $B_2$ , while systematic error might play a role for  $B_3$  and  $B_4$ . We also note that  $\langle \delta_1 \delta_2 \rangle = 0$  over the range  $180 \leq \theta \leq 360^\circ$  indicating that  $\mathcal{M}_b$  is indeed a quantum nondemolition measurement over this range.

**ACKNOWLEDGMENTS.** We thank Howard Wiseman for helpful discussions. We acknowledge financial support from the Australian Research Council and the Army Research Office Contract W911NF-0397 funded by the Intelligence Advanced Research Projects Activity (IARPA). M.E.G. would like to thank the University of Queensland for support and hospitality during his sabbatical visit there.

1. Bell JS (1964) On the Einstein Podolski Rosen paradox. *Physics* 1:195–200.
2. Bell JS (1966) On the problem of hidden variables in quantum mechanics. *Rev Mod Phys* 38:447–452.
3. Kochen S, Specker EP (1967) The problem of hidden variables in quantum mechanics. *J Math Mech* 17:59–87.
4. Kirchmair G, et al. (2009) State-independent experimental test of quantum contextuality. *Nature* 460:494–497.
5. Leggett AJ, Garg A (1985) Quantum mechanics versus macroscopic realism: Is the flux there when nobody looks? *Phys Rev Lett* 54:857–860.
6. Aharonov Y, Vaidman L (1990) Properties of a quantum system during the time interval between two measurements. *Phys Rev A* 41:11–20.
7. Aharonov Y, Albert DZ, Vaidman L (1988) How the result of the measurement of a component of the spin of a spin-1/2 particle can turn out to be 100. *Phys Rev Lett* 60:1351–1354.
8. Jordan AN, Korotkov AN, Büttiker M (2006) Leggett-Garg inequality with a kicked quantum pump. *Phys Rev Lett* 97:026805.
9. Palacios-Laloy A, et al. (2010) Experimental violation of a Bells inequality in time with weak measurement. *Nature Phys* 6 442–447.
10. Williams NS, Jordan AN (2008) Weak values and the Leggett-Garg inequality in solid-state qubits. *Phys Rev Lett* 100:026804.

11. Williams NS, Jordan AN (2009) Erratum: Weak values and the Leggett-Garg inequality in solid-state qubits. *Phys Rev Lett* 103:089902.
12. Grangier P, Levenson JA, Poizat J-P (1998) Quantum non-demolition measurements in optics. *Nature* 396:537–542.
13. Ralph TC, Bartlett SD, O'Brien JL, Pryde GJ, Wiseman HM (2006) Quantum nondemolition measurements for quantum information. *Phys Rev A* 73:012113.
14. Pryde GJ, O'Brien JL, White AG, Bartlett SD, Ralph TC (2004) Measuring a photonic qubit without destroying it. *Phys Rev Lett* 92:190402.
15. Ritchie NWM, Story JG, Hulet RG (1991) Realization of a measurement of a “weak value”. *Phys Rev Lett* 66:1107–1110.
16. Resch KJ, Lundeen JS, Steinberg AM (2004) Experimental realization of the quantum box problem. *Phys Lett A* 324:125–131.
17. Pryde GJ, O'Brien JL, White AG, Ralph TC, Wiseman HM (2005) Measurement of quantum weak values of photon polarization. *Phys Rev Lett* 94:220405.
18. Hosten O, Kwiat PG (2008) Observation of the spin Hall effect of light via weak measurements. *Science* 319:787–790.
19. Dixon PB, Starling DJ, Jordan AN, Howell JC (2009) Ultrasensitive beam deflection measurement via interferometric weak value amplification. *Phys Rev Lett* 102:173601.
20. Leggett AJ (1989) Comment on “How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100”. *Phys Rev Lett* 62:2325–2325.

21. Peres A (1989) Quantum measurements with postselection. *Phys Rev Lett* 62:2326–2326.
22. Huelga SF, Marshall TW, Santos E (1995) Proposed test for realist theories using Rydberg atoms coupled to a high-Q resonator. *Phys Rev A* 52:R2497–R2500.
23. Gühne O, et al. (2010) Compatibility and noncontextuality for sequential measurements. *Phys Rev A* 81:022121.
24. Barrett J, Kent A (2004) Non-contextuality, finite precision measurement and the Kochen–Specker theorem. *Stud Hist Phil Mod Phys* 35:151–176.
25. Langford NK, et al. (2005) Demonstration of a simple entangling optical gate and its use in bell-state analysis. *Phys Rev Lett* 95:210504.
26. Kiesel N, Schmid C, Weber U, Ursin R, Weinfurter H (2005) Linear optics controlled-phase gate made simple. *Phys Rev Lett* 95:210505.
27. Okamoto R, Hofmann HF, Takeuchi S, Sasaki K (2005) Demonstration of an optical quantum controlled-not gate without path interference. *Phys Rev Lett* 95:210506.
28. O'Brien JL, Pryde GJ, White AG, Ralph TC, Branning D (2003) Demonstration of an all-optical quantum controlled-NOT gate. *Nature* 426:264–267.
29. O'Brien JL, et al. (2004) Quantum process tomography of a controlled-NOT gate. *Phys Rev Lett* 93:080502.
30. Wiseman HM (2002) Weak values, quantum trajectories, and the cavity-QED experiment on wave-particle correlation. *Phys Rev A* 65:032111.
31. Fuchs CA (1998) Information gain vs. state disturbance in quantum theory. *Fortschr Phys* 46:535–565.
32. Helstrom CW (1976) Quantum Detection and Estimation Theory. *Mathematics in Science and Engineering*, (Academic, New York), 123.