

Complete genome sequence of *Ferrimonas balearica* type strain (PAT^T)

Matt Nolan¹, Johannes Sikorski², Karen Davenport^{1,3}, Susan Lucas¹, Tijana Glavina Del Rio¹, Hope Tice¹, Jan-Fang Cheng¹, Lynne Goodwin^{1,3}, Sam Pitluck¹, Konstantinos Liolios¹, Natalia Ivanova¹, Konstantinos Mavromatis¹, Galina Ovchinnikova¹, Amrita Pati¹, Amy Chen⁴, Krishna Palaniappan⁴, Miriam Land^{1,5}, Loren Hauser^{1,5}, Yun-Juan Chang^{1,5}, Cynthia D. Jeffries^{1,5}, Roxanne Tapia^{1,3}, Thomas Brettin^{1,3}, John C. Detter^{1,3}, Cliff Han^{1,3}, Montri Yasawong⁶, Manfred Rohde⁶, Brian J Tindall², Markus Göker², Tanja Woyke¹, James Bristow¹, Jonathan A. Eisen^{1,7}, Victor Markowitz⁴, Philip Hugenholtz¹, Nikos C. Kyrpides¹, Hans-Peter Klenk^{2*}, and Alla Lapidus¹

¹ DOE Joint Genome Institute, Walnut Creek, California, USA

² DSMZ - German Collection of Microorganisms and Cell Cultures GmbH, Braunschweig, Germany

³ Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico, USA

⁴ Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, California, USA

⁵ Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁶ HZI – Helmholtz Centre for Infection Research, Braunschweig, Germany

⁷ University of California Davis Genome Center, Davis, California, USA

*Corresponding author: Hans-Peter Klenk

Keywords: chemoorganotroph, iron(III)-reducing bacterium, facultatively anaerobic, *Ferrimonadaceae*, *Gammaproteobacteria*, GEBA

Ferrimonas balearica Rossello-Mora *et al.* 1996 is the type species of the genus *Ferrimonas*, which belongs to the family *Ferrimonadaceae* within the *Gammaproteobacteria*. The species is a Gram-negative, motile, facultatively anaerobic, non spore-forming bacterium, which is of special interest because it is a chemoorganotroph and has a strictly respiratory metabolism with oxygen, nitrate, Fe(III)-oxyhydroxide, Fe(III)-citrate, MnO₂, selenate, selenite and thiosulfate as electron acceptors. This is the first completed genome sequence of a member of the genus *Ferrimonas* and also the first sequence from a member of the family *Ferrimonadaceae*. The 4,279,159 bp long genome with its 3,803 protein-coding and 144 RNA genes is a part of the *Genomic Encyclopedia of Bacteria and Archaea* project.

Introduction

Strain PAT^T (= DSM 9799 = CCM 4581) is the type strain of the species *Ferrimonas balearica*, which is the type species of its genus *Ferrimonas* [1,2]. Currently, there are five species in the genus *Ferrimonas* [3]. The generic name derives from the Latin word '*ferrum*' meaning 'iron' and the Greek word '*monas*' meaning 'unit', referring to an iron(III)-reducing cell. The species epithet is also derived from the Latin word '*balearica*' meaning 'of the Balearic Islands', referring to the place where the strain was isolated [1]. *Ferrimonas* is the type genus of the family *Ferrimonadaceae* and one of two genera in the family *Ferrimonadaceae*

[4]. Strain PAT^T was described in 1995 by Rossello-Mora *et al.* [1] who isolated the strain from the upper few centimeters of marine sediment of the Palma de Mallorca harbor, Spain [1,5]. Here we present a summary classification and a set of features for *F. balearica* PAT^T, together with the description of the complete genomic sequencing and annotation.

Classification and features

The 16S rRNA gene sequence of PAT^T is 99% identical to four culturable strains, which are reported

in GenBank [6]. Two strains, A2A-18 (AB193752) and A3B-47-3 (AB193753), were isolated from marine sand [7]. The culturable strain S8-05 (EU620413) was isolated from Palk Bay sediment in Thondi, India and another strain with accession number AY158002 was isolated from Ala Wai Canal sediment in Honolulu, USA. The 16S rRNA gene of strain PAT^T shares 93.5-97.4% sequence identity with the sequences of the type strains from the other members of the family *Ferrimonadaceae* [8]. The environmental samples database (env_nt) contains the marine metagenome clone 1096626783183 (96% sequence identity, AACY020355234). The genomic survey sequences database (gss) contains the uncultured bacterium

clone BYUP987.b1 (92%, EF996742), isolated from a fecal sample of adult woman who gave birth after 11 months [9]. Altogether, strains belonging to the species *F. balearica* or the genus *Ferrimonas* are rather rare in the habitats screened so far (status September 2010).

Figure 1 shows the phylogenetic neighborhood of *F. balearica* PAT^T in a 16S rRNA based tree. The sequences of the seven 16S rRNA gene copies in the genome differ from each other by up to five nucleotides, and differ by up to four nucleotides from the previously published sequence (X93021), which contains two ambiguous base calls.




Figure 1. Phylogenetic tree highlighting the position of *F. balearica* PAT^T relative to the type strains of the other species within the family *Ferrimonadaceae* and to the type of the neighboring family *Psychromonadaceae*. The trees were inferred from 1,449 aligned characters [10,11] of the 16S rRNA gene sequence under the maximum likelihood criterion [12] and rooted with the type strain of the order *Alteromonadaceae*. The branches are scaled in terms of the expected number of substitutions per site. Numbers above branches are support values from 650 bootstrap replicates [13] if larger than 60%. Lineages with type strain genome sequencing projects registered in GOLD [14] are shown in blue, published genomes in bold (CP000510) [15].

Strain PAT^T is a Gram-negative, nonspore-forming, facultatively anaerobic bacterium [1]. The cells are straight rods (0.3-0.5 × 1.2-1.5 μm) with rounded ends (Figure 2, Table 1) [1,5] and appear singly, occasionally in pairs or short chains and usually not encapsulated [1,5]. Strain PAT^T is motile by means of monotrichous flagella (not visible in Figure 2, but 10% of the cells in the original liquid culture were highly motile) [1]. Colonies produce a black iron precipitate when the cells are grown on TSI agar [1]. Although initially isolated using

TSI based media this strain grows better on Marine Broth. Colonies are often brown and mucous when the cells are grown under aerobic conditions [5]. Fresh isolates of this species may not form colonies on PYG agar medium, but the colonies are formed after several subcultivations in enrichment medium [1,5]. Resting stages of strain PAT^T are not known [5]. Cells of the strain undergo autolysis within five days under aerobic conditions [1,5]. Strain PAT^T does not contain polyhydroxybutyrate (PHB) or other intracellular inclu-

sions [2]. The strain is chemoorganotrophic. Under anaerobic conditions, the reduction of Fe(III)-oxyhydroxide is coupled to the utilization of lactate as the electron donor, which yields magnetite [1,5]. Strain PAT^T uses oxygen, nitrate, Fe(III)-oxyhydroxide, Fe(III)-citrate, MnO₂, selenate, selenite and thiosulfate as electron acceptors [1,5,25]. Strain PAT^T requires a minimum of 0.5% NaCl for growth, with a range of NaCl tolerance of 0.5%-7.5% [1]. It does not grow at 5°C or 44°C but does grow at 42°C [1]. The pH range for growth is 6-9 [1]. Enzymatic reactions are positive for catalase, oxidase, phenylalanine deaminase, DNase

and lipase (Tween 20 and Tween 80), but negative for amylase, arginine dihydrolase, gelatinase, lysine decarboxylase, Simmons citrate and urease [1,5]. The strain does not hydrolyze starch [1]. The genus *Ferrimonas* can be distinguished from other strictly respiratory Gram-negative genera of the *Gammaproteobacteria* based on its ability to reduce Fe(III), denitrification, growth at 42°C, presence of phenylalanine deaminase activity, inability to grow in NaCl-free media, lack of gelatinase, urease and a negative reaction of Simmons citrate test [5].




Figure 2. Scanning electron micrograph of *F. balearica* PAT^T

Chemotaxonomy

The quinone profiles of strain PAT^T are MK-7 (62.9%), Q-8 (20.4%) and Q-7 (16%) [7]. The presence of both menaquinones and ubiquinones being indicative of the ability of this organism to grow aerobically (with ubiquinones) and anaerobically (with menaquinones). The presence of menaquinones and ubiquinones with different distributions of isoprenoid side chains is a feature also shared by members of the genus *Shewanella* [26-28] and *Paraferrimonas* [29]. The major cellular fatty acids of strain PAT^T, when grown on PYG medium, given in the original species description are C_{17:1ω8c} (27.5%), iso-C_{15:0} (14.5%), C_{17:0} (7.8%), iso-C_{13:0} (5.8%), C_{16:1ω7c} (4.7%), C_{15:0} (4.5%), C_{13:0}

(4.5%), C_{14:0} (4.2%), C_{18:1ω9c} (4.0%) and C_{12:0} 3-OH (1.8%), C_{17:1ω6c} (1.6%) and C_{18:1ω7c} (1.2%) [1]. More recent data show a somewhat different fatty acid pattern [7], with the fatty acids comprising iso-C_{15:0} (9.8%), C_{15:0} (1.8%) iso-C_{16:1ω9c} (10.4%) iso-C_{16:1ω7c} (5.2%), C_{16:0} (13.4%) iso-C_{17:0} (2.1%) C_{17:1ω8c} (12.6%) C_{17:0} (7.9%) C_{18:1ω9c} (17.6%) C_{18:1ω7c} (4.9%) and C_{18:0} (3.9%). Hydroxylated fatty acids were not reported. Interestingly the fatty acids reported in a subsequent paper [25] that are based on the work of Kasuta *et al.* [7] omit the iso-C_{16:1} fatty acids. The fatty acids reported in the original publication [1] show a number of features also found in members of the genera *Shewanella*

and *Paraferrimonas* [29,30]. Data generated in the DSMZ during the course of this work indicates that the fatty acids comprise, iso-C_{13:0} (3.7%), C_{13:0} (2.7%), C_{12:0} 3OH (2.2%), iso-C_{14:0} (1.1%), C_{14:0} (1.0%), iso-C_{13:0} 3OH (3.7%), C_{13:0} 3OH (1.9%), iso-C_{15:0} (16.1%), C_{15:1 w8c} (2.1%), C_{15:0} (4.5%), C_{14:0} 3-OH (2.9%), C_{16:1 w9c} (8.1%), C_{16:1w7c} (4.9%), C_{16:0} (8.4%), iso-C_{15:0} 3OH, (0.9%), iso-C_{17:0} (1.4%), C_{17:1 w8c} (14.7%), C_{17:0} (5.6%), C_{18:1 w9c} (7.8%) and C_{18:1 w7c} (1.4%). These results are more consistent with those published in the original description [1], but there are differences that cannot be attributed to

differences in the growth conditions. The complete absence of hydroxylated fatty acids in the work of Kasuta *et al.* [7] suggests that no attempt was made to detect them. The presence of at least two positional isomers in unsaturated fatty acids with the same chain length is indicative of the presence of at least two enzymatic pathways for introducing the double bonds. A fairly simple polar lipid pattern has been reported for *Ferrimonas futtsuensis*, comprising, phosphatidylglycerol, phosphatidylethanolamine and an unidentified aminophospholipid [29].

Table 1. Classification and general features of *F. balearica* PAT^T according to the MIGS recommendations [16].

MIGS ID	Property	Term	Evidence code
		Domain <i>Bacteria</i>	TAS [17]
		Phylum <i>Proteobacteria</i>	TAS [18-20]
		Class <i>Gammaproteobacteria</i>	TAS [18,21]
	Current classification	Order <i>Alteromonadales</i>	TAS [18,22]
		Family <i>Ferrimonadaceae</i>	TAS [4]
		Genus <i>Ferrimonas</i>	TAS [1,2]
		Species <i>Ferrimonas balearica</i>	TAS [1,2]
		Type strain PAT	TAS [1,2]
	Gram stain	negative	TAS [1]
	Cell shape	straight rods with rounded ends	TAS [1,5]
	Motility	yes	TAS [1]
	Sporulation	nonspore-forming	TAS [1]
	Temperature range	37°C-42°C	TAS [1,5]
	Optimum temperature	37°C	NAS
	Salinity	0.5%-7.5% (w/v) NaCl	TAS [1,5]
MIGS-22	Oxygen requirement	facultatively anaerobic	TAS [1]
	Carbon source	lactate	TAS [1]
	Energy source	chemoorganotroph	TAS [1,5]
MIGS-6	Habitat	marine sediment	TAS [1]
MIGS-15	Biotic relationship	free-living	NAS
MIGS-14	Pathogenicity	none	NAS
	Biosafety level	1	TAS [23]
	Isolation	marine sediment	TAS [1]
MIGS-4	Geographic location	Palma de Mallorca harbor, Spain	TAS [1]
MIGS-5	Sample collection time	1995 or before	TAS [1]
MIGS-4.1	Latitude	39.57	NAS
MIGS-4.2	Longitude	2.63	NAS
MIGS-4.3	Depth	not report	
MIGS-4.4	Altitude	below the sea level	TAS [1,5]

Evidence codes - IDA: Inferred from Direct Assay (first time in publication); TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from of the Gene Ontology project [24]. If the evidence code is IDA, then the property was directly observed by one of the authors or an expert mentioned in the acknowledgements.

Genome sequencing and annotation

Genome project history

This organism was selected for sequencing on the basis of its phylogenetic position [31], and is part of the *Genomic Encyclopedia of Bacteria and Archaea* project [32]. The genome project is deposited in the Genome OnLine Database [14] and the

complete genome sequence is deposited in GenBank. Sequencing, finishing and annotation were performed by the DOE Joint Genome Institute (JGI). A summary of the project information is shown in Table 2.

Table 2. Genome sequencing project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	Finished
MIGS-28	Libraries used	Two genomic Sanger libraries: 8 kb pMCL200 library, fosmid (40 kb) library
MIGS-29	Sequencing platforms	ABI3730
MIGS-31.2	Sequencing coverage	9.8 × Sanger
MIGS-30	Assemblers	Phrap
MIGS-32	Gene calling method	Prodigal 1.4, GenePRIMP
	INSDC ID	CP002209
	Genbank Date of Release	October 1, 2010
	GOLD ID	Gc01378
	NCBI project ID	30799
	Database: IMG-GEBA	2502082106
MIGS-13	Source material identifier	DSM 9799
	Project relevance	Tree of Life, GEBA

Growth conditions and DNA isolation

F. balearica PAT^T, DSM 9799, was grown in DSMZ medium 514 (Bacto Marine Broth) [33] at 28°C. DNA was isolated from 0.5-1 g of cell paste using Qiagen Genomic 500 DNA Kit (Qiagen, Hilden, Germany) following the standard protocol as recommended by the manufacturer, with modification st/L for cell lysis as described in Wu *et al.* [32].

Genome sequencing and assembly

The genome was sequenced using the Sanger sequencing platform (6 and 40 kb DNA libraries). All general aspects of library construction and sequencing performed at the JGI can be found at <http://www.jgi.doe.gov/>. The [Phred/Phrap/Consed](#) software package was used for sequence assembly and quality assessment. After the shotgun stage, reads were assembled with parallel phrap (High Performance Software, LLC). Possible mis-assemblies were corrected with Dupfinisher or transposon bombing of bridging clones (Epicentre Biotechnologies, Madison, WI) [34]. Gaps between contigs were closed by editing in Consed, custom primer walk or PCR amplification. A total of 404 additional custom primer reactions were necessary to close gaps and to raise the quality of the finished sequence. The completed genome sequence contains 48,554 reads, achieving an average of 9.8-fold

sequence coverage with an error rate less than 1 in 100,000.

Genome annotation

Genes were identified using [Prodigal](#) [35] as part of the Oak Ridge National Laboratory genome annotation pipeline, followed by a round of manual curation using the JGI [GenePRIMP](#) pipeline [36]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) non-redundant database, UniProt, TIGRFam, Pfam, PRIAM, KEGG, COG, and InterPro databases. Additional gene prediction analysis and functional annotation was performed within the Integrated Microbial Genomes - Expert Review (IMG-ER) platform [37].

Genome properties

The genome consists of a 4,279,159 bp long chromosome with a 60.2% GC content (Table 3 and Figure 3). Of the 3,947 genes predicted, 3,803 were protein-coding genes, and 144 RNAs; twenty one pseudogenes were also identified. The majority of the protein-coding genes (72.5%) were assigned a putative function while the remaining ones were annotated as hypothetical proteins. The distribution of genes into COGs functional categories is presented in Table 4.

Table 3. Genome Statistics

Attribute	Value	% of Total
Genome size (bp)	4,279,159	100.00%
DNA coding region (bp)	3,842,563	89.80%
DNA G+C content (bp)	2,576,887	60.22%
Number of replicons	1	
Extrachromosomal elements	0	
Total genes	3,947	100.00%
RNA genes	144	3.65%
rRNA operons	7	
Protein-coding genes	3,803	96.35%
Pseudo genes	21	0.53%
Genes with function prediction	2,860	72.46%
Genes in paralog clusters	462	11.71%
Genes assigned to COGs	2,929	74.21%
Genes assigned Pfam domains	3,089	78.26%
Genes with signal peptides	1,154	29.24%
Genes with transmembrane helices	981	24.85%
CRISPR repeats	0	




Figure 3. Graphical circular map of the genome. From outside to the center: Genes on forward strand (color by COG categories), Genes on reverse strand (color by COG categories), RNA genes (tRNAs green, rRNAs red, other RNAs black), GC content, GC skew.

Table 4. Number of genes associated with the general COG functional categories

Code	value	%age	Description
J	189	5.8	Translation, ribosomal structure and biogenesis
A	1	0.0	RNA processing and modification
K	213	6.5	Transcription
L	138	4.2	Replication, recombination and repair
B	1	0.0	Chromatin structure and dynamics
D	35	1.1	Cell cycle control, cell division, chromosome partitioning
Y	0	0.0	Nuclear structure
V	61	1.9	Defense mechanisms
T	178	5.5	Signal transduction mechanisms
M	219	6.7	Cell wall/membrane/envelope biogenesis
N	133	4.1	Cell motility
Z	0	0.0	Cytoskeleton
W	0	0.0	Extracellular structures
U	128	3.9	Intracellular trafficking and secretion, and vesicular transport
O	155	4.8	Posttranslational modification, protein turnover, chaperones
C	238	7.3	Energy production and conversion
G	105	3.2	Carbohydrate transport and metabolism
E	248	7.6	Amino acid transport and metabolism
F	85	2.6	Nucleotide transport and metabolism
H	167	5.1	Coenzyme transport and metabolism
I	99	3.0	Lipid transport and metabolism
P	184	6.7	Inorganic ion transport and metabolism
Q	53	1.6	Secondary metabolites biosynthesis, transport and catabolism
R	338	10.4	General function prediction only
S	287	8.8	Function unknown
-	1,018	25.8	Not in COGs

Acknowledgements

We would like to gratefully acknowledge the help of Regine Fähnrich for growing *F. balearica* cultures and Susanne Schneider for DNA extraction and quality analysis (both at DSMZ). This work was performed under the auspices of the US Department of Energy Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under contract No. DE-

AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, and Los Alamos National Laboratory under contract No. DE-AC02-06NA25396, UT-Battelle and Oak Ridge National Laboratory under contract DE-AC05-00OR22725, as well as German Research Foundation (DFG) INST 599/1-2 and Thailand Research Fund Royal Golden Jubilee Ph.D. Program No. PHD/0019/2548 for MY.

References

- Rossello-Mora R, Ludwig W, Kämpfer P, Amann R, Schleifer KH. *Ferrimonas balearica* gen. nov., spec. nov., a new marine facultative Fe(III)-reducing bacterium. *Syst Appl Microbiol* 1995; **18**:196-202.
- Validation List No. 56. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Bacteriol* 1996; **46**:362-363. [doi:10.1099/00207713-46-1-362](https://doi.org/10.1099/00207713-46-1-362)
- Garrity G. NamesforLife. BrowserTool takes expertise out of the database and puts it right in the browser. *Microbiol Today* 2010; **7**:1.
- Ivanova EP, Flavier S, Christen R. Phylogenetic relationships among marine *Alteromonas*-like proteobacteria: emended description of the family *Alteromonadaceae* and proposal of *Pseudoalteromonadaceae* fam. nov., *Colwelliaceae* fam. nov., *Shewanellaceae* fam. nov., *Moritellaceae*

- fam. nov., *Ferrimonadaceae* fam. nov., *Idiomarinaceae* fam. nov. and *Psychromonadaceae* fam. nov. *Int J Syst Evol Microbiol* 2004; **54**:1773-1788. [PubMed](#) [doi:10.1099/ijs.0.02997-0](https://doi.org/10.1099/ijs.0.02997-0)
5. Rossello-Mora R, Kämpfer P, Ludwig W, Amann R, Schleifer KH. 2005. Genus IV. *Ferrimonas*, p. 454-456. In D. J. Brenner, N. R. Krieg, J. T. Staley, and G. M. Garrity (ed.), *Bergey's Manual of Systematic Bacteriology*, 2nd ed, vol. 2. Springer, New York.
 6. Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank. *Nucleic Acids Res* 2009; **37**:D26-D31. [PubMed](#) [doi:10.1093/nar/gkn723](https://doi.org/10.1093/nar/gkn723)
 7. Katsuta A, Adachi K, Matsuda S, Shizuri Y, Kasai H. *Ferrimonas marina* sp. nov. *Int J Syst Evol Microbiol* 2005; **55**:1851-1855. [PubMed](#) [doi:10.1099/ijs.0.63689-0](https://doi.org/10.1099/ijs.0.63689-0)
 8. Chun J, Lee JH, Jung Y, Kim M, Kim S, Kim BK, Lim YW. EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. *Int J Syst Evol Microbiol* 2007; **57**:2259-2261. [PubMed](#) [doi:10.1099/ijs.0.64915-0](https://doi.org/10.1099/ijs.0.64915-0)
 9. Vaishampayan PA, Kuehl JV, Froula JL, Morgan JL, Ochman H, Francino MP. Comparative metagenomics and population dynamics of the gut microbiota in mother and infant. *Genome Biol Evol* 2010; **2**:53-66. [PubMed](#) [doi:10.1093/gbe/evp057](https://doi.org/10.1093/gbe/evp057)
 10. Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; **17**:540-552. [PubMed](#)
 11. Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics* 2002; **18**:452-464. [PubMed](#) [doi:10.1093/bioinformatics/18.3.452](https://doi.org/10.1093/bioinformatics/18.3.452)
 12. Stamatakis A, Hoover P, Rougemont J. A rapid bootstrap algorithm for the RAxML Web servers. *Syst Biol* 2008; **57**:758-771. [PubMed](#) [doi:10.1080/10635150802429642](https://doi.org/10.1080/10635150802429642)
 13. Pattengale ND, Alipour M, Bininda-Emonds ORP, Moret BME, Stamatakis A. How many bootstrap replicates are necessary? *Lect Notes Comput Sci* 2009; **5541**:184-200. [doi:10.1007/978-3-642-02008-7_13](https://doi.org/10.1007/978-3-642-02008-7_13)
 14. Liolios K, Chen IM, Mavromatis K, Tavernarakis N, Hugenholtz P, Markowitz VM, Kyrpides NC. The Genomes On Line Database (GOLD) in 2009: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res* 2009; **38**:D346-D354. [PubMed](#) [doi:10.1093/nar/gkp848](https://doi.org/10.1093/nar/gkp848)
 15. Auman AJ, Breezee JL, Gosink JJ, Kämpfer P, Staley JT. *Psychromonas ingrahamii* sp. nov., a novel gas vacuolate, psychrophilic bacterium isolated from Arctic polar sea ice. *Int J Syst Evol Microbiol* 2006; **56**:1001-1007. [PubMed](#) [doi:10.1099/ijs.0.64068-0](https://doi.org/10.1099/ijs.0.64068-0)
 16. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](#) [doi:10.1038/nbt1360](https://doi.org/10.1038/nbt1360)
 17. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](#) [doi:10.1073/pnas.87.12.4576](https://doi.org/10.1073/pnas.87.12.4576)
 18. Validation List No. 106. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Evol Microbiol* 2005; **55**:2235-2238. [doi:10.1099/ijs.0.64108-0](https://doi.org/10.1099/ijs.0.64108-0)
 19. Garrity GM, Bell JA, Lilburn T. 2005. Phylum XIV. *Proteobacteria* phyl. nov. In DJ Brenner, NR Krieg, JT Staley, and GM Garrity (eds), *Bergey's Manual of Systematic Bacteriology*, 2nd ed, vol. 2. Springer, New York.
 20. Garrity GM, Holt JG. The Road Map to the Manual. In: Garrity GM, Boone DR, Castenholz RW (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 1, Springer, New York, 2001, p. 119-169.
 21. Garrity GM, Bell JA, Lilburn T. 2005. Class III. *Gammaproteobacteria* class. nov., p. 1. In DJ Brenner, NR Krieg, JT Staley, and GM Garrity (eds), *Bergey's Manual of Systematic Bacteriology*, 2nd ed, vol. 2. Springer, New York.
 22. Bowman JP, McMeekin TA. 2005. Order X. *Alteromonadales* ord. nov., p. 443. In D J Brenner, NR Krieg, JT Staley, and GM Garrity (eds), *Bergey's Manual of Systematic Bacteriology*, 2nd ed, vol. 2. Springer, New York.
 23. Classification of bacteria and archaea in risk groups. TRBA 466.
 24. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene Ontology: tool for the unification of biology. *Nat Genet* 2000; **25**:25-29. [PubMed](#) [doi:10.1038/75556](https://doi.org/10.1038/75556)

25. Nakagawa T, Iino T, Suzuki K, Harayama S. *Ferrimonas futtsuensis* sp. nov. and *Ferrimonas kyonanensis* sp. nov., selenate-reducing bacteria belonging to the *Gammaproteobacteria* isolated from Tokyo Bay. *Int J Syst Evol Microbiol* 2006; **56**:2639-2645. [PubMed](#) doi:10.1099/ijs.0.64399-0
26. Moule AL, Wilkinson SG. Polar lipids, fatty acids, and isoprenoid quinones of *Alteromonas putrefaciens* (*Shewanella putrefaciens*). *Syst Appl Microbiol* 1987; **9**:192-198.
27. Akagawa-Matsushita M, Itoh T, Katayama Y, Kuraishi H, Yamasato K. Isoprenoid quinone composition of some marine *Alteromonas*, *Marinomonas*, *Deleya*, *Pseudomonas* and *Shewanella* species. *J Gen Microbiol* 1992; **138**:2275-2281.
28. Frolova GM, Pavel KG, Shparteeva AA, Nedashkovskaya OI, Gorshkova NM, Ivanova EP, Mikhailov VV. Lipid composition of novel *Shewanella* species isolated from Far Eastern seas. *Microbiology* 2005; **74**:664-669. doi:10.1007/s11021-005-0121-9
29. Khan ST, Harayama S. *Paraferrimonas sedimenticola* gen. nov., sp. nov., a marine bacterium of the family *Ferrimonadaceae*. *Int J Syst Evol Microbiol* 2007; **57**:1493-1498. [PubMed](#) doi:10.1099/ijs.0.64529-0
30. Leonardo MR, Moser DP, Brabieri E, Brantner CA, MacGregor BJ, Paster BJ, Stackebrandt E, Nea1son KH. *Shewanella pealeana* sp. nov., a member of the microbial community associated with the accessory nidamental gland of the squid *Loligo pealei*. *Int J Syst Bacteriol* 1999; **49**:1341-1351. [PubMed](#) doi:10.1099/00207713-49-4-1341
31. Klenk HP, Göker M. En route to a genome-based classification of *Archaea* and *Bacteria*? *Syst Appl Microbiol* 2010; **33**:175-182. [PubMed](#) doi:10.1016/j.syapm.2010.03.003
32. Wu D, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova NN, Kunin V, Goodwin L, Wu M, Tindall BJ, et al. A phylogeny-driven genomic encyclopaedia of *Bacteria* and *Archaea*. *Nature* 2009; **462**:1056-1060. [PubMed](#) doi:10.1038/nature08656
33. List of growth media used at DSMZ: http://www.dsmz.de/microorganisms/media_list.php.
34. Sims D, Brettin T, Detter JC, Han C, Lapidus A, Copeland A, Glavina Del Rio T, Nolan M, Chen F, Lucas S, et al. Complete genome sequence of *Kytococcus sedentarius* type strain (541^T). *Stand Genomic Sci* 2009; **1**:12-20. doi:10.4056/sigs.761
35. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 2010; **11**:119. [PubMed](#) doi:10.1186/1471-2105-11-119
36. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. Gene-PRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 2010; **7**:455-457. [PubMed](#) doi:10.1038/nmeth.1457
37. Markowitz VM, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 2009; **25**:2271-2278. [PubMed](#) doi:10.1093/bioinformatics/btp393