

Published in final edited form as:

Metabolism. 2011 July ; 60(7): 965–968. doi:10.1016/j.metabol.2010.09.005.

Low Salt Diet Increases Insulin Resistance in Healthy Subjects

Rajesh Garg¹, Gordon H. Williams¹, Shelley Hurwitz^{1,2}, Nancy J. Brown³, Paul N. Hopkins⁴, and Gail K. Adler¹

¹Division of Endocrinology, Diabetes, and Hypertension, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

²Department of Biostatistics, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

³Division of Clinical Pharmacology, Vanderbilt University School of Medicine, Nashville, TN

⁴Cardiovascular Genetics, Cardiology Division, University of Utah, Salt Lake City, UT

Abstract

Objective—Low salt (LS) diet activates the renin-angiotensin-aldosterone and sympathetic nervous systems, both of which can increase insulin resistance (IR). We investigated the hypothesis that LS diet is associated with an increase in IR in healthy subjects.

Methods—Healthy individuals were studied after 7 days of LS diet (urine sodium <20 mmol/day) and 7 days of high salt (HS) diet (urine sodium >150 mmol/day) in a random order. IR was measured after each diet and compared statistically, unadjusted and adjusted for important covariates.

Results—One hundred fifty two healthy men and women, age 39.1 ± 12.5 years (range 18–65), body mass index (BMI) 25.3 ± 4.0 kg/m², were included in this study. Mean (SD) homeostasis model assessment (HOMA) was significantly higher on LS compared with HS diet (2.8 ± 1.6 Vs 2.4 ± 1.7 ; $P < 0.01$). Serum aldosterone (21.0 ± 14.3 Vs 3.4 ± 1.5 ng/dl; $p < 0.001$), 24 hour urine aldosterone (63.0 ± 34.0 Vs 9.5 ± 6.5 µg/day; $p < 0.001$) and 24 hour urine norepinephrine excretion (78.0 ± 36.7 Vs 67.9 ± 39.8 µg/day; $p < 0.05$) were higher on LS diet compared with HS diet. LS diet was significantly associated with higher HOMA independent of age, gender, blood pressure, BMI, serum sodium and potassium, serum angiotensin II, plasma renin activity, serum and urine aldosterone, and urine epinephrine and norepinephrine.

Conclusions—Low salt diet is associated with an increase in IR. The impact of our findings on the pathogenesis of diabetes and cardiovascular disease needs further investigation.

© 2010 Elsevier Inc. All rights reserved.

Address for Correspondence: Rajesh Garg, M.D., Division of Endocrinology, Diabetes, and Hypertension, Brigham and Women's Hospital, 221 Longwood Ave., Boston, MA 02115, U.S.A., Telephone: 617-732-7672, Fax: 617-277-1568, rgarg@partners.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure Statement: The authors have no disclosures.

Disclosure statement: Authors have no conflicts of interest

Author contributions: RG conceived the idea, analyzed and interpreted data and wrote the manuscript; GHW helped in interpretation of data and reviewed the manuscript; SH helped with statistical analysis; NB recruited patients and reviewed the manuscript; PNH recruited patients and reviewed the manuscript and GKA helped in interpretation of data and revised the manuscript.

Keywords

HOMA; Renin - angiotensin - Aldosterone; Salt intake

Introduction

Low dietary salt intake is recommended as one of the public health measures to decrease the risk of cardiovascular disease [1]. However, low salt intake stimulates aldosterone production through activation of the renin-angiotensin-aldosterone system (RAAS) [2]. We recently demonstrated an association between aldosterone and insulin resistance (IR) in healthy subjects [3]. Moreover, sympathetic nervous system is activated by low salt diet as shown by an increase in urine norepinephrine levels [4,5]. Activation of the sympathetic nervous system may also increase IR [6]. Therefore, low dietary salt intake may be associated with an increase in IR. Previous studies on the effect of salt intake on IR in healthy subjects have shown contradictory results. Some studies showed an increase in IR [7,8], while others found no such effect or a decrease in IR [9–11]. To investigate this further, we analyzed data from a large cohort of healthy subjects studied under carefully controlled conditions. Further, we investigated whether the effect of salt intake on IR could be explained by known risk factors for IR.

Methods

This is an analysis of data collected during the studies conducted as part of the International HyperPath (Hypertensive Pathotype) cohort [12] and includes subjects from three sites--- Boston, Salt Lake City and Nashville. Study subjects included men and women, aged 18–65 years who were generally free of any significant medical or psychiatric problems. The studies were approved by the Human Research Committee at each site and informed consent was obtained. All participants were studied after seven days of an isocaloric, high-salt (HS) diet and low salt (LS) diet in a random order. Compliance with salt intake was monitored by measuring 24 hour urine sodium. Those with urine sodium <150 mmol/day on HS diet or >20 mmol/day on LS diet were excluded.

After seven days of respective diet, participants collected 24-hour urine for sodium, creatinine, aldosterone, cortisol, epinephrine and norepinephrine. They were then kept fasting and supine overnight for 8–10 hours. Next day, baseline blood pressure was measured. Mean arterial blood pressure (MAP) was calculated as: $MAP = \text{Diastolic Blood Pressure} + (\text{Systolic} - \text{Diastolic})/3$. Fasting supine blood samples were obtained for plasma renin activity (PRA), aldosterone, angiotensin II, cortisol, sodium, potassium, glucose and insulin. Angiotensin II (Bachem Inc., Weil am Rhein, Germany) was infused intravenously at 3 ng/kg/min for 45 minutes following which serum aldosterone was measured again. Blood pressure and heart rate were monitored every 2 minutes during the angiotensin II infusion.

All laboratory assays were performed at a central laboratory as previously described [13]. Urine and serum aldosterone were measured by solid phase RIA by the Coat-A-Count method (Diagnostic Products Corporation, Los Angeles, CA). Urine and serum cortisol was measured by Access Cortisol assay (Beckman Coulter, Chaska, MN). Urine epinephrine and norepinephrine were measured by radioimmunoassay by 1st converting them enzymatically to N-acetylmetanephrine and N-acetylnormetanephrine (LDN 2 CAT RIA, Immuno Biological Laboratories, Inc, Minneapolis, MN, 55432). Urine and serum sodium and potassium were assayed by flame photometry (Nova Biomedical, Waltham, MA). Urine creatinine was measured with the ACE Creatinine Reagent (Alfa Wasserman, West Caldwell, NJ). Insulin

was assayed by chemiluminescence immunoassay using the Access Immunoassay System (Beckman Coulter, Chaska, MN).

The homeostasis model assessment (HOMA) was used as the measure of insulin resistance and was calculated as $HOMA = (\text{plasma glucose (mmol/L)} \times \text{plasma insulin } (\mu\text{U/mL})) / 22.5$ [14]. The data were summarized using means and standard deviations. Measurements obtained on LS diet were compared with measurements obtained on HS diet by paired T test. Mixed model analyses were performed to assess the impact of diet on the natural log of HOMA while controlling for single and multiple covariates (SAS version 9.1).

Results

We identified 227 subjects studied on both LS and HS diets. Seventy five subjects were excluded from analysis due to urine sodium >20 mmol/day on LS diet or <150 mmol/day on HS diet. One hundred fifty two subjects were included in this analysis. The subjects were 39.1 ± 12.5 years old and included 55% women. Eighty percent of subjects were Caucasian. Subject characteristics and laboratory values obtained on LS and HS diets are shown in table 1. BMI and blood pressure were lower and measures of RAAS were higher on LS compared to HS diet. Urine norepinephrine levels were also higher on LS diet, while the urine cortisol levels were low. HOMA was significantly higher on LS compared with HS diet (2.82 ± 1.6 Vs 2.45 ± 1.69 , $p < 0.01$). Change in HOMA from LS to HS diet (Δ HOMA) did not correlate with changes in serum aldosterone, PRA, angiotensin II, stimulated serum aldosterone, urine aldosterone and urine norepinephrine (all $p = \text{NS}$). On mixed model analyses, the effect of diet on HOMA remained significant after controlling individually for each of the covariates in table 1 (all $p < 0.05$). A multiple variable model was developed to assess the impact of diet on HOMA while controlling for multiple potentially important variables simultaneously. Potential predictors besides diet were age, sex, BMI, serum potassium, serum sodium, MAP, serum aldosterone, serum angiotensin II, serum and urine cortisol, urine aldosterone and urine norepinephrine. The final parsimonious model contained only diet, age, and BMI (all $p < 0.001$). HOMA remained significantly higher on LS diet in the multiple variable model that controlled for age and BMI.

Discussion

Our study shows that low salt intake is associated with higher insulin resistance. The effect of dietary salt intake on insulin sensitivity has been controversial. Some previous studies have shown an increase in IR with low salt diet [7,8] while others found no such effect or even a decrease in IR [9–11]. These discrepancies may be due to differences in study populations, levels of salt restriction, study methods or to the relatively small sample size in previous studies of healthy individuals. Two studies that showed a decrease or no change in IR on LS diet included only 7 – 8 subjects and excluded women [9,10]. Another study that showed no change in IR included 34 men, but the subjects were not provided a standardized diet and were studied after a moderate salt restriction (urine sodium 70 ± 45.1 mmol/day on LS diet and 175 ± 72.1 mmol/day on HS diet) [11]. Our results are consistent with those of Townsend et al who studied 20 healthy subjects, both men and women, and compared IR after standardized diets containing 20 mmol/day versus 200 mmol/day sodium for 6 days [8]. Our study is the largest study in healthy subjects, includes both men and women, was performed under carefully controlled conditions, and utilized a sodium content for the HS diet that is comparable to the average sodium intake in the US population [15].

We found an increase in aldosterone and norepinephrine, both of which may contribute to an increase in IR on LS diet. Angiotensin II levels were also higher on LS diet and angiotensin II has been shown to interfere with insulin signaling pathways [16]. However, none of these

factors could individually explain the increase in IR; salt intake remained an independent predictor of HOMA after including them individually in the analysis. Moreover, change in HOMA on LS versus HS diet did not correlate with corresponding changes in components of the RAAS. This lack of correlation may indicate a different mechanism for LS diet induced IR or may be due to the large variability in RAAS components with dietary salt restriction. Consistent with previous studies, urine cortisol was lower on LS diet [17,18], thus, ruling out the possibility of high cortisol production as a mediator of IR. The possibility of other unidentified factors involved in LS diet induced IR cannot be ruled out.

High salt intake is considered a public health problem in USA. High dietary salt intake is associated with higher blood pressure, a known cardiovascular disease risk factor. In a recent study, using computerized models, it was estimated that reduction in salt intake could reduce the number of deaths by 44,000 to 92,000 [19]. However, this study and other similar studies were based on the estimated cardiovascular disease risk reduction associated with a fall in blood pressure. Activation of RAAS and changes in IR were not included in these models. Available data do not support universal recommendation for any particular level of dietary salt intake [20].

In conclusion, in healthy subjects LS diet is associated with an increase in IR. The impact of our findings on the pathogenesis of diabetes and cardiovascular disease warrants further investigation.

Abbreviations

RAAS	Renin-angiotensin-aldosterone system
IR	Insulin resistance
HS	High salt
LS	Low salt
MAP	Mean arterial blood pressure
PRA	Plasma renin activity
HOMA	Homeostasis model assessment

Acknowledgments

Funding: Supported by NIH grant HL 087060

References

1. Most Americans Should Consume Less Sodium. Available from: <http://www.cdc.gov/salt/>.
2. Hollenberg NK, Chenitz WR, Adams DF, et al. Reciprocal influence of salt intake on adrenal glomerulosa and renal vascular responses to angiotensin II in normal man. *J Clin Invest.* 1974; 54(1):34–42. [PubMed: 4365595]
3. Garg R, Hurwitz S, Williams GH, et al. Aldosterone production and insulin resistance in healthy adults. *J Clin Endocrinol Metab.* 2010; 95(4):1986–1990. [PubMed: 20130077]
4. Lilavivathana U, Campbell RG. The influence of sodium restriction on orthostatic sympathetic nervous activity. *Arch Intern Med.* 1980; 140(11):1485–1489. [PubMed: 7436645]
5. Warren SE, Vieweg WV, O'Connor DT. Sympathetic nervous system activity during sodium restriction in essential hypertension. *Clin Cardiol.* 1980; 3(5):348–351. [PubMed: 7002404]
6. Jamerson KA, Julius S, Gudbrandsson T, et al. Reflex sympathetic activation induces acute insulin resistance in the human forearm. *Hypertension.* 1993; 21(5):618–623. [PubMed: 8491496]

7. Perry CG, Palmer T, Cleland SJ, et al. Decreased insulin sensitivity during dietary sodium restriction is not mediated by effects of angiotensin II on insulin action. *Clin Sci (Lond)*. 2003; 105(2):187–194. [PubMed: 12691602]
8. Townsend RR, Kapoor S, McFadden CB. Salt intake and insulin sensitivity in healthy human volunteers. *Clin Sci (Lond)*. 2007; 113(3):141–148. [PubMed: 17425514]
9. Donovan DS, Solomon CG, Seely EW, et al. Effect of sodium intake on insulin sensitivity. *Am J Physiol*. 1993; 264(5 Pt 1):E730–E734. [PubMed: 8498495]
10. Fliser D, Fode P, Arnold U, et al. The effect of dietary salt on insulin sensitivity. *Eur J Clin Invest*. 1995; 25(1):39–43. [PubMed: 7705385]
11. Grey A, Braatvedt G, Holdaway I. Moderate dietary salt restriction does not alter insulin resistance or serum lipids in normal men. *Am J Hypertens*. 1996; 9(4 Pt 1):317–322. [PubMed: 8722434]
12. Kosachunhanun N, Hunt SC, Hopkins PN, et al. Genetic determinants of nonmodulating hypertension. *Hypertension*. 2003; 42(5):901–908. [PubMed: 14530292]
13. Hopkins PN, Lifton RP, Hollenberg NK, et al. Blunted renal vascular response to angiotensin II is associated with a common variant of the angiotensinogen gene and obesity. *J Hypertens*. 1996; 14(2):199–207. [PubMed: 8728297]
14. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985; 28:412–419. [PubMed: 3899825]
15. Brown IJ, Tzoulaki I, Candeias V, et al. Salt intakes around the world: implications for public health. *Int J Epidemiol*. 2009; 38(3):791–813. [PubMed: 19351697]
16. Diamond-Stanic MK, Henriksen EJ. Direct inhibition by angiotensin II of insulin-dependent glucose transport activity in mammalian skeletal muscle involves a ROS-dependent mechanism. *Arch Physiol Biochem*. 2010; 116(2):88–95. [PubMed: 20384568]
17. Wambach G, Bleienheuft C, Bonner G. Sodium loading raises urinary cortisol in man. *J Endocrinol Invest*. 1986; 9(3):257–259. [PubMed: 3020116]
18. Lewicka S, Nowicki M, Vecsei P. Effect of sodium restriction on urinary excretion of cortisol and its metabolites in humans. *Steroids*. 1998; 63(7–8):401–405. [PubMed: 9654646]
19. Bibbins-Domingo K, Chertow GM, Coxson PG, et al. Projected effect of dietary salt reductions on future cardiovascular disease. *N Engl J Med*. 2010; 362(7):590–599. [PubMed: 20089957]
20. Alderman MH. Evidence relating dietary sodium to cardiovascular disease. *J Am Coll Nutr*. 2006; 25(3 Suppl):256S–261S. [PubMed: 16772637]

Table 1

Comparison of subject characteristics and laboratory parameters on high salt versus low salt diet.

	Low Salt Diet	High Salt Diet	P
BMI (kg/m ²)	24.6 ± 4.1	25.3 ± 4.1	<0.0001
Mean arterial blood pressure (mmHg)	78.1 ± 8.2	82.6 ± 9.2	<0.0001
Serum Na (mmol/L)	139.0 ± 5.4	140.0 ± 5.3	NS
Serum K (mmol/L)	4.1 ± 0.4	4.1 ± 0.4	NS
Serum Cortisol (µg/dL)	12.3 ± 4.4	11.9 ± 4.1	NS
Serum aldosterone (ng/dL)	21.0 ± 13.5	3.4 ± 1.4	<0.0001
PRA (ng/ml/h)	3.1 ± 2.1	0.39 ± 0.40	<0.0001
Angiotensin II (pg/mL)	43.9 ± 22.6	28.1 ± 15.9	<0.0001
Serum aldosterone after Angiotensin II infusion (ng/dL)	45.2 ± 21.8	11.2 ± 6.6	<0.0001
Urine Na (mmol/day)	7.5 ± 4.9	237.8 ± 70.9	<0.0001
Urine aldosterone (µg/day)	63.0 ± 34.0	9.6 ± 6.5	<0.0001
Urine cortisol (µg/day)	37.4 ± 24.4	52.7 ± 25.9	<0.0001
Urine epinephrine (µg/day)	11.5 ± 6.3	12.2 ± 6.5	NS
Urine norepinephrine (µg/day)	78.0 ± 36.7	67.9 ± 39.8	<0.05
Urine creatinine (mg/day)	1367 ± 408	1401 ± 397	NS
HOMA	2.82 ± 1.6	2.45 ± 1.69	0.002