

Inhibition of Prostaglandin Synthesis after Metabolism of Menadione by Cultured Porcine Endothelial Cells

Aaron Barchowsky, Katayoun Tabrizi, Richard S. Kent,* and A. Richard Whorton

Divisions of Pulmonary Medicine and Cardiology,* Departments of Pharmacology and Medicine, Duke University Medical Center, Durham, North Carolina 27710

Abstract

We have examined the effects of menadione on porcine aortic endothelial cell prostaglandin synthesis. Addition of 1–20 μM menadione caused a dose- and time-dependent inhibition of stimulated prostaglandin synthesis with an IC_{50} of 5 μM at 15 min. Concentrations greater than 100 μM menadione were necessary to increase ^{51}Cr release from prelabeled cells. Recovery of enzyme inactivated by menadione required a 6-h incubation in 1% serum. In a microsomal preparation, menadione was shown to have no direct effect on conversion of arachidonic acid to prostaglandins. In intact cells menadione caused only a 40% inhibition of the conversion of PGH_2 to prostacyclin. Enzymes involved in the incorporation and the release of arachidonic acid were not affected by menadione (20 μM , 15 min). Menadione undergoes oxidation/reduction reactions in intact cells leading to partial reduction of oxygen-forming, reactive oxygen species. In our cells menadione was found to increase KCN-resistant oxygen consumption. Further, an increased accumulation of H_2O_2 was observed with a time course consistent with menadione-induced inhibition of prostaglandin synthesis. We conclude that menadione at sublethal doses caused inhibition of prostaglandin synthesis. The mechanism involves inactivation of PGH_2 synthase by a reactive species resulting from metabolism of menadione by endothelial cells.

Introduction

Numerous studies have shown that damage to vascular endothelium is a primary event in oxidative injury caused by hyperoxia (1–3), hypoxia (4), activated neutrophils (1, 5), endotoxin (6, 7), or certain chemotherapeutic agents (8, 9). In these studies, cytotoxicity was demonstrated by measuring gross morphologic changes (4, 7–9), by demonstrating loss of endothelial-dependent organ function (8), or by assessing loss of cell viability (1, 5, 8). Comparatively few studies have addressed the effects of sublethal concentrations of oxidants or oxidant-generating species on biological functions of the endothelial cell. One important endothelial function which has recently been demonstrated to be inhibited after exposure to sublethal

concentrations of H_2O_2 and superoxide anion (O_2^-) is the capacity of these cells to produce prostacyclin (PGI_2 ; 10, 11). It is well known that PGI_2 synthesized by the endothelium plays a role in maintaining local blood flow (12, 13), in protecting against vascular injury by limiting the activation and adherence of neutrophils and platelets (12–15), and perhaps in modulating release of catecholamines (13). Thus, loss of the ability to produce PGI_2 after oxidative stress may have profound implications on the role of the endothelium in maintaining vascular homeostasis.

The conversion of arachidonic acid to prostaglandins is known to be tightly regulated by oxidants, especially at the level of PGH_2 synthase. In fact, the activity of this enzyme becomes rate limiting when the enzyme is fully reduced (a lack of oxidants) or when the enzyme metabolizes a sufficient concentration of hydroperoxide to cause autoinactivation (16, 17). In isolated systems, low levels of exogenous hydroperoxides have been shown to activate PGH_2 synthase by reducing the lag time for arachidonic acid metabolism (16, 17). The regulatory mechanisms in intact cells are not completely understood. However, it has been shown that, in intact vessels (18) and in cultured endothelial cells, PGH_2 synthase is irreversibly inactivated through autoinhibition (19) or after metabolism of exogenous hydroperoxides (10, 11, 20). Recovery of inactivated prostaglandin synthetic capacity in these cells requires new enzyme synthesis (11).

Damaging concentrations of reactive oxygen species arise after metabolism of certain compounds by endothelial cells. Quinones represent an important group of such compounds, whose therapeutic effectiveness is limited by their ability to cause oxidative injury (8, 21, 22). Quinone-induced toxicity has been observed in many organs and cell types and is thought to result from univalent reduction of the quinone to form the semiquinone radical (21–24). This reduction is catalyzed by flavin-containing, NAD(P)H-dependent enzymes in the microsomal and mitochondrial fractions of the cell (21–24). Oxidation of the semiquinone in the presence of molecular oxygen regenerates the quinone and causes the formation of O_2^- . O_2^- quickly dismutates to yield H_2O_2 (21–24) which may affect cell function. High doses of menadione (vitamin K_3) have been shown to cause the production of O_2^- and other reactive oxygen species in endothelial cells (25) and to alter cell membrane integrity. The effects of lower concentrations of quinones, which would presumably cause a lower level of intracellular oxidant production, on endothelial cell integrity and function have not been characterized.

Therefore, we chose to examine the effects of sublethal concentrations of menadione on prostaglandin production in endothelial cells. The concentration- and time-dependent effects of menadione on synthesis of PGI_2 and prostaglandin E_2 (PGE_2) were investigated and the activities of the enzymes in the synthetic cascade were determined to establish the site of

Portions of this work were presented in abstract form (1988. *FASEB [Fed. Am. Soc. Exp. Biol.] J.* 2:A1188).

Address reprint requests to Dr. Barchowsky at his present address: Division of Clinical Pharmacology, Thomas Jefferson University, 1100 Walnut Street, Philadelphia, PA 19107.

Received for publication 5 July 1988 and in revised form 27 September 1988.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/89/04/1153/07 \$2.00

Volume 83, April 1989, 1153–1159

action of menadione. In addition, the time course and dose dependence of the cytotoxic effects of menadione were evaluated and the requirements for recovery from menadione-induced effects were examined. The data presented demonstrate that low concentrations of menadione irreversibly inhibit endothelial prostaglandin synthesis. This inhibition appears to result from increased intracellular production of H_2O_2 .

Methods

Materials. Menadione, arachidonic acid, bradykinin triacetate, catalase, superoxide dismutase, ibuprofen, and cytodex beads were obtained from Sigma Chemical Co. (St. Louis, MO). The calcium ionophore, A23187, was obtained as the free acid from Calbiochem-Behring Corp. (San Diego, CA). Dulbecco's modified Eagle's media (DME), Hanks' balanced salt solution (HBSS), medium 199, and other tissue culture reagents were obtained from Gibco (Grand Island, NY). Fetal calf serum (FCS) was obtained from HyClone Laboratories (Logan, UT). Deuterated prostaglandins, [3,3,4,4,- 2H]6-keto-prostaglandin $F_{1\alpha}$ and [3,3,4,4,- 2H]prostaglandin E_2 were purchased from Merck, Sharp & Dohme (Montreal, Quebec). [5,6,8,9,11,12,14,15- 3H]arachidonic acid (60 Ci/mmol) and [^{51}Cr]Na $_2$ CrO $_4$ (200 Ci/g) were purchased from NEN Research Products (Boston, MA). All solvents were HPLC grade or of the highest purity available.

Cell culture. Endothelial cells were enzymatically isolated from fresh porcine aortas as previously described (10, 11) and cultured according to established methods (10, 11). The cells were grown in 25-cm 2 flasks (NUNC, USA Plastics, Ocala, FL) containing DME plus 10% FCS and were maintained at 37°C under an atmosphere of 10% CO $_2$ /90% air. The cells were subcultured using 0.1% trypsin-EDTA and confluent monolayers of second to fourth-passage cells were used for experimentation. The endothelial cells were characterized as previously reported (11, 26) with more than 90–95% of the cells in random cultures containing factor VIII antigen.

Culture of cells on cytodex 1 beads was essentially as described above. After isolation, cells were plated in 25-cm 2 flasks and allowed to grow to confluence. After demonstrating the purity of the culture, an aliquot of beads was added and the cells were cultured for an additional 2 d. The beads were removed by gentle agitation and added to a stirred flask containing complete media plus 3 mg/ml fresh beads. Half of the media was exchanged every 3 d and cell numbers were increased by adding more media and beads.

Exposure to menadione. Endothelial cells were washed twice with HEPES-buffered HBSS (HHBS), pH 7.4, to remove the culture medium which contained serum. Menadione, dissolved in DMSO and diluted in HHBS, was added, incubated with the cells for the desired time period, and removed by another two rinses. The final concentration of DMSO was < 0.5% and an equal amount was added to controls. The capacity of the endothelial monolayers to produce prostaglandins was evaluated by addition of the sodium salt of arachidonic acid (20 μ M) dissolved in HHBS and incubated at 37°C for 3 min. The buffer was collected, deuterated internal standards were added, and the samples were stored at -20°C until analysis.

Prostaglandin analysis. Prostaglandin levels were determined according to previously published procedures by selective ion monitoring using a gas chromatograph-mass spectrometer equipped with an electron impact source (10, 11, 18). Briefly, the samples were acidified with formic acid (pH 3.0) and extracted twice with 2 vol of ethyl acetate. After forming the methyl ester-methoxime-trimethylsilyl ether derivatives, the samples were injected on to a 6-ft \times 4-mm glass column packed with 3% OV-1 on 80–100 Supelcoport (Supelco Inc., Bellefonte, PA) at a carrier gas flow of 25 ml/min and an oven temperature of 245°C. These column conditions allowed complete baseline resolu-

tion of the derivatives of 6-keto-prostaglandin $F_{1\alpha}$ (the stable nonenzymatic metabolite of PGI $_2$) and of PGE $_2$.

Assay for cytotoxicity. The cytotoxic effect of menadione was examined by following ^{51}Cr release from cells prelabeled for 5 h with [^{51}Cr]Na $_2$ CrO $_4$ (10 μ Ci/flask). The cells were washed twice with HHBS containing 1.0% bovine serum albumin (BSA) and incubated for 10 min before two additional rinses. Menadione or H $_2$ O $_2$ was then added in HHBS plus 1.0% BSA and aliquots of the incubation buffer were sampled for radioactivity at 30, 60, and 90 min. Data are expressed as the percent of total releasable ^{51}Cr .

Assay of microsomal PGH $_2$ synthase. Microsomes were prepared from ram seminal vesicles (Wayne State University, Detroit, MI) and resuspended (10 mg/ml) in 0.1 M potassium phosphate buffer, pH 7.4, containing 1 mM EDTA. The microsomes were preincubated for 5 min at 37°C in buffer containing hemoglobin (0.1 mg/ml), phenol (2 mM), and tryptophan (1 mM) before addition of saturating levels of tritiated arachidonic acid (100 μ g/ml arachidonic acid plus 1.0 μ Ci [3H]arachidonic acid) for 2 min. The reaction was stopped by addition of methanol-formic acid (5:2 vol/vol). The incubation mixtures were extracted twice with ethyl acetate (2 vol) and the organic layers were combined, evaporated, and redissolved in 5 ml of hexane-diethyl ether (9:1). The samples were applied to a 0.75-g silicic acid column, and unreacted arachidonic acid was eluted with 20 ml of the same solvent. The sample vial was rinsed with and hydroxylated products were eluted with 20 ml of hexane-ether (8:2). The sample vial was finally rinsed with and prostaglandins were eluted with ethyl acetate-methanol (9:1). Greater than 80% of the initial 3H added was recovered, with 10–15% being lost to formation of [3H]H $_2$ O. Labeled prostaglandins, 15-hydroxyicosatetraenoic acid, and arachidonic acid were used to standardize the chromatography.

Release of arachidonic acid. To examine the effects of menadione on the release of arachidonic acid from endothelial cells, monolayers were incubated overnight in DME containing 1.0% FCS and 2 μ Ci [3H] arachidonic acid. The culture medium was removed by two washes with HHBS containing 0.25% BSA and the cells were exposed to menadione for 15 min. The flasks were rinsed with three buffer washes and release of label was followed by sampling an aliquot of the buffer every 5 min. After establishing basal release, either bradykinin (1.0 μ g/ml) or A23187 (1.0 μ M) was added to stimulate arachidonic acid release. Radioactivity was quantified by liquid scintillation counting.

Oxygen uptake assay. Oxygen consumption by endothelial cells grown on cytodex beads was measured at 37°C using a Clark electrode fitted into a stirred chamber. The change in electrode potential was monitored with an oxygen monitor (Yellow Springs Instruments, Yellow Springs, OH) with output to a strip chart recorder. The electrode was calibrated using air equilibrated HHBS. An aliquot of beads was added to the chamber in the same buffer and data obtained from different experiments were normalized for protein content of the aliquot.

Measurement of H $_2$ O $_2$. To measure formation of H $_2$ O $_2$ after addition of menadione, monolayers of cells were rinsed and incubated for 15 min with 2 ml of HHBS containing menadione. Separate groups of flasks were treated similarly with buffer containing 1 mM NaN $_3$. After the exposure period, the buffer was removed and assayed for H $_2$ O $_2$ content using the colorimetric assay described by Thurman et al. (27). NaN $_3$ did not alter the assay results. Specificity of the procedure was determined by quantifying H $_2$ O $_2$ in aliquots of buffer treated with 10 U of catalase. All values were compared to production in nontreated cells.

Enzyme assays. Catalase activity was assayed by determining the rate of H $_2$ O $_2$ reduction (decrease in absorbance at 280 nm) as described by Bergmeyer (28). Superoxide dismutase activity was determined by the ability to prevent reduction of acetylated cytochrome c as described by McCord and Fridovich (29).

Statistics. To test for significant differences, data were analyzed by analysis of variance following Bartlett's test for normal distribution. When necessary, log transforms were made to normalize the variance. The means of groups were compared using Student-Newman-Keuls or

1. *Abbreviations used in this paper:* HHBS, HEPES-buffered Hanks' balanced salt solution.

Dunnett's tests. Data are presented as mean \pm standard deviation unless designated otherwise.

Results

We have previously demonstrated that addition of 20 μ M arachidonic acid to confluent monolayers of porcine aortic endothelial cells for 3 min results in maximal production of prostaglandins (11). The primary prostaglandin produced by these cells is PGI₂, with a minor amount of PGE₂ also being formed (10, 11, 26). Pretreatment of the cells for 10, 15, or 30 min with varying concentrations of menadione caused a time- and concentration-dependent inhibition of prostaglandin synthesis (Fig. 1) with an IC₅₀ = 5 μ M at 15 min. Although the data presented represent only the production of PGI₂, the synthesis of PGE₂ decreased in a parallel fashion (76.9 \pm 13.1% inhibition after 15-min treatment with 20 μ M menadione, *n* = 9). Exposure of monolayers to higher concentrations of menadione also inhibited basal prostaglandin production as indicated by a > 100% inhibition of the capacity to convert exogenous arachidonic acid.

Since menadione was found to affect the function of endothelial cells, we chose to examine the cytotoxicity of the compound. Cytotoxicity was measured by determining the rate of ⁵¹Cr release from prelabeled cells exposed to increasing concentrations of menadione. There was no increase in release of ⁵¹Cr above control values at concentrations of menadione that maximally inhibit prostaglandin synthesis. However, when cells were exposed to much larger doses of menadione (1 mM), there was a significant increase in ⁵¹Cr release 60 min after addition of the drug (Fig. 2). This effect on membrane integrity occurred at a dose that is 200-fold greater than that required for inhibition of PGI₂ synthesis. There was no obvious detachment of cells during the exposure period.

Although these cytotoxic concentrations of menadione have been shown to cause generation of reactive oxygen metabolites in endothelial cells (25), it is not known whether




Figure 1. Menadione-induced inhibition of PGI₂ synthesis. Endothelial cell monolayers were exposed to increasing concentrations of menadione for 10 (● --- ●), 15 (▲ ····· ▲), or 30 (■ — ■) min, before rinsing. Prostaglandin synthetic capacity was tested by incubating the cells with 20 μ M arachidonic acid for 3 min and measuring the amount of PGI₂ released into the buffer as described in Methods. This production in non-menadione-treated cells after addition of arachidonate was 122 \pm 20 ng/flask above the basal production of 22.5 \pm 5.4 ng/flask. Data are presented as the percentage of the total capacity of nontreated cells to produce prostaglandins. Each point represents the mean \pm SD, *n* = 3.




Figure 2. Time course of menadione-induced ⁵¹Cr release. Cells were prelabeled with ⁵¹Cr and rinsed as described in Methods. Release of label was followed with time from nontreated cells (○ ····· ○) or cells treated with 2 mM H₂O₂ (● --- ●), 0.1 mM menadione (□ - · - · □), or 1 mM menadione (■ — ■) at *t* = 0. Data are presented as the percentage of label relative to the total releasable ⁵¹Cr within the monolayer and represent mean \pm SD of groups of four flasks of cells. At 60 min, 1 mM menadione-treated flasks released significantly more ⁵¹Cr than controls (*P* < 0.01); no other differences seen. At 90 min both 1 mM menadione and 2 mM H₂O₂-treated flasks released significantly more ⁵¹Cr than controls (*P* < 0.01) or 0.1 mM menadione-treated flasks which were not different from controls.

much lower, sublethal doses have the same effect. We examined this possibility by measuring oxygen consumption by monolayers of cells grown on beads. As shown in Fig. 3, the basal rate of O₂ consumption (3.43 \pm 1.71 nmol/min per mg protein) was increased 10-fold after addition of 20 μ M menadione (34.8 \pm 5.4 nmol/min per mg protein). Addition of 130 U of catalase resulted in a release of O₂ into the system followed by a decrease in the rate of consumption to half the menadione-induced rate (14.6 \pm 7.5 nmol/min per mg protein). This effect of catalase could be explained by liberation of O₂ from H₂O₂ accumulated in the buffer and equilibration to a rate governed by the stoichiometry of the reaction: 2H₂O₂ \rightarrow 2H₂O + O₂ (25, 31). Menadione-driven O₂ consumption was not inhibited by 1 mM KCN or by 2 μ M rotenone, demonstrating that the increase was not dependent on mitochondrial respira-




Figure 3. Oxygen consumption by endothelial cells grown on cyto-dex beads. Cells were cultured as described in Methods and loaded into a 2.8-ml stirred chamber, which was maintained at 37°C. Oxygen consumption was measured with a Clark electrode fitted into the chamber. At the designated times, menadione (20 μ M), catalase (130 U), KCN (1 mM), or rotenone (2 μ M) was added. The protein content was 0.327 mg. The data are representative of four separate experiments.

tion (21, 22, 30, 31). The accelerated rate after addition of KCN probably reflects inhibition of endogenous catalase (30, 31).

Since menadione-induced oxygen consumption was slowed by addition of catalase, we decided to quantify the amount of H₂O₂ diffusing from monolayers of endothelial cells grown in 25-cm² flasks. The basal rate of H₂O₂ accumulation was below the detection limits of our assay conditions (< 0.5 nmol/15 min, Table I). Menadione caused a dose-dependent increase in H₂O₂ formation. Addition of NaN₃ increased this drug-induced accumulation of H₂O₂ twofold. These data indicate that metabolism of menadione by endothelial cells leads to measurable H₂O₂ formation, although the accumulation of H₂O₂ in the buffer probably underestimates the total amount of H₂O₂ formed.

It has been suggested that menadione can induce the formation of reactive oxygen species in the extracellular media as the fully reduced menadione diffuses from the cell and reoxidizes to menadione in the media (21, 25). We have previously demonstrated that extracellular generation of O₂⁻ or addition of H₂O₂ causes inhibition of endothelial cell prostaglandin synthesis (10). This inhibition can be completely prevented by addition of catalase to the media (10). To test whether menadione inhibited prostaglandin synthesis via extracellular generation of oxidants, we preincubated monolayers for 15 min with increasing concentrations of menadione in the presence and absence of superoxide dismutase and catalase. We observed no effect of these enzymes on menadione-induced inhibition of prostaglandin synthesis (*P* < 0.05). This indicates that extracellular generation of H₂O₂ is not the mechanism for menadione inhibition of PGI₂ synthesis. Therefore, the actions of menadione must be dependent on intracellular generation of inhibitory species or the drug has a direct effect on the synthetic enzymes of the prostaglandin cascade.

To examine the direct effects of menadione on PGH₂ synthase, we chose to use a preparation of ram seminal vesicle microsomes, which are enriched with the enzyme. Ibuprofen, a known inhibitor of PGH₂ synthase, was used as a positive control in these experiments. Nontreated microsomes converted 14.0±1.9% and 31.2±2.3% of [³H]arachidonate to hydroxylated products and prostaglandins, respectively. Ibupro-

fen prevented this conversion of arachidonate as indicated by a decrease in ³H products and an increase in radioactivity eluting as unconverted arachidonic acid compared to control (94.5±0.5% vs. 55.2±2.1%). In contrast, preincubation of microsomes for 5 min with 20 μM menadione did not alter the elution profile indicating that menadione has no direct effect on conversion of arachidonic acid. Thus menadione inhibits PGI₂ synthesis indirectly suggesting that the drug is metabolized by the endothelial cell to create an inhibitory species.

In addition to examining the effects of quinones on the activity of PGH₂ synthase, it is also important to investigate effects on PGI₂ synthase activity as well. This is the second enzyme in the prostaglandin synthetic cascade which is known to be sensitive to oxidants (18, 32, 33). We exposed monolayers to menadione for 15 min before addition of PGH₂, the substrate for PGI₂ synthase. As shown in Fig. 4, 5 μM menadione caused a 40% decrease in conversion of PGH₂ to PGI₂. There was no further inhibition observed in monolayers treated with 20 μM of the drug. However, this effect on PGI₂ synthase can only partially account for the inhibition of PGI₂ synthesis from arachidonic acid.

We observed inhibition of arachidonic acid conversion at concentrations of menadione (Fig. 1) which have previously been shown to increase Ca²⁺ ionophore-stimulated release of PGI₂ (34). It is possible that menadione leads to loss of PGH₂ synthase activity by increasing the availability of free arachidonate. Metabolism of this arachidonate to PGI₂ during the exposure period might cause autoinhibition of the enzyme. Therefore, we examined the effects of menadione on stimulated arachidonic acid release from endothelial cells prelabeled with [³H] arachidonic acid for 18 h. As shown in Fig. 5 a, pretreatment with menadione (20 μM) for 15 min did not affect either basal or A23187-stimulated release of label. We also examined the effect of menadione on receptor-mediated release. In these experiments, there was also no alteration of basal release, nor was there an effect on release stimulated by the agonist, bradykinin (Fig. 5 b). Finally, to examine the effect of menadione on incorporation of arachidonic acid, we pretreated monolayers for 15 min with 20 μM of the drug before adding 1.26 μCi of [³H]arachidonic acid. 200 μl of buffer were sampled at 5, 30, 45, and 60 min to determine amount of unincorporated radioactivity. There was no difference in the rate of incorporation of label into nontreated and treated cells

Table I. Accumulation of H₂O₂ after Addition of Menadione

Treatment	H ₂ O ₂ nmol/flask
Control	ND
1 mM NaN ₃	ND
5 μM menadione	2.75±0.19
5 μM menadione + NaN ₃	5.11±0.56
20 μM menadione	4.32±0.22
20 μM menadione + NaN ₃	8.41±0.81

Cells, in 25-cm² flasks, were rinsed and exposed for 15 min to menadione diluted in 2 ml of buffer. The buffer was collected and assayed for H₂O₂ content. Data represent mean±SD, *n* = 3. ND designates values which were below the detection limits of the assay (0.5 nmol of H₂O₂). Treatment with 20 μM menadione was different from treatment with 5 μM menadione (*P* < 0.01). NaN₃ significantly enhanced the effect of either treatment (*P* < 0.01).




Figure 4. Menadione-induced inhibition of endothelial PGI₂ synthase. Monolayers were exposed to menadione for 15 min. After rinsing, buffer containing 5 μg/ml PGH₂ was added and the cells were incubated for an additional 5 min. The buffer was then removed and assayed for PGI₂. Data are mean±SD, *n* = 3. All flasks incubated with PGH₂ produced significantly more PGI₂ than controls (*P* < 0.01). Menadione at both doses inhibited conversion (*P* < 0.05).




Figure 5. Effect of menadione on (a) A23187- or (b) bradykinin (BK)-stimulated arachidonate release. Cells, prelabeled for 18 h with [³H]arachidonate, were exposed to menadione (20 μ M, 15 min). After rinsing, release of label was followed by sampling buffer every 5 min. At 15 or 20 min, BK (100 ng/ml) or A23187 (1 μ M), respectively, was added. Data are mean \pm SD, $n = 4$. Menadione did not alter release of label from either control, A23187-, or BK-stimulated cells. Both A23187- and BK-treated cells released significantly more label than controls at all time points after addition of the agonist ($P < 0.01$).

(0.734 ± 0.008 vs. 0.736 ± 0.010 μ Ci/flask per h respectively, $n = 4$).

We have previously demonstrated that sublethal concentrations of H₂O₂ cause irreversible inactivation of PGH₂ synthase, and that new enzyme must be synthesized in order to regain prostaglandin synthetic capacity in the endothelial cell (11). If menadione-induced inhibition of prostaglandin synthesis is also the result of irreversible enzyme inactivation and if, as shown in Fig. 2, this inhibition occurs at sublethal concentrations of menadione, then the monolayers should be able to recover after exposure. To test this hypothesis, we exposed monolayers to 20 μ M menadione for 15 min. The cells were then washed and incubated for varying periods of time in medium 199 plus 1% FCS before testing the capacity to produce prostaglandins. As shown above, this dose of menadione inhibited the conversion of exogenous arachidonic acid to PGI₂. A full 6 h was required before prostaglandin synthetic capacity returned to normal (Fig. 6). This recovery time course is identical to that we have reported for the recovery of PGH₂ synthase following inhibition by H₂O₂ (11).

Discussion

The results of this investigation demonstrate that menadione causes irreversible inhibition of endothelial prostaglandin synthesis. This inhibition is both concentration- and time-related,




Figure 6. Recovery of endothelial prostaglandin synthesis inhibited by menadione. Cells were exposed to menadione (20 μ M) for 15 min, rinsed, and incubated for varying times with medium 199 containing 1% FCS. The culture media was removed and the prostaglandin synthetic capacity was tested. Bars (\square) represent production in nontreated cells, while closed circles (\bullet) represent production in menadione-treated cells. Data represent mean \pm SEM, $n = 6$.

yet occurs at lower concentrations and with a shorter time course than menadione-induced release of ⁵¹Cr. Thus menadione-induced loss of endothelial cell function greatly precedes loss of cell integrity.

The primary mechanism for menadione cytotoxicity is formation of reactive oxygen species. Rosen and Freeman (25) demonstrated that menadione, in high concentrations, induced formation of O₂⁻ in endothelial cells. These studies demonstrated that 100 μ M menadione increased both intracellular and extracellular production of O₂⁻ and subsequent formation of H₂O₂. Our present data demonstrate that menadione increases endothelial oxygen consumption by 10-fold at a dose five times lower than that used previously. That this consumption represents partial reduction of molecular oxygen leading to the formation of H₂O₂ is supported by several findings: (a) accumulation of H₂O₂ is dependent on menadione concentration; (b) addition of extracellular catalase slows the rate of oxygen consumption; (c) H₂O₂ accumulation is augmented by addition of NaN₃. The most likely mechanism for the effect of NaN₃ is inhibition of intracellular catalase (30, 31); although NaN₃ may act to inhibit mitochondrial respiration, thus providing more reducing equivalents to drive menadione redox cycling. However, since endothelial cells contain few mitochondria (35), this contribution of reducing equivalents would be minimal.

The exact mechanism for menadione-dependent H₂O₂ formation in endothelial cells has not been investigated, but it is probably qualitatively similar to mechanisms observed in other cell types. In the rat hepatocyte, menadione has been shown to be reduced by enzymes in the mitochondria and endoplasmic reticulum to form the semiquinone (21–24). This radical is a strong nucleophile and can react with a variety of intracellular constituents including membrane proteins, DNA, GSH, and O₂ (23, 24). The semiquinone is reoxidized to menadione, causing generation of O₂⁻ (21–24). A continued recycling of the semiquinone to the quinone and detoxification of the oxygen metabolites produced leads to depletion of reducing equivalents within the cell and cytotoxicity ensues (23, 36). Although this pathway appears universal, cells differ in their sensitivities to menadione (23). These differences are quite pronounced and form the basis for use of menadione as an antineoplastic agent (37, 38) or to sensitize tumor cells to var-

ious forms of oxidative anticancer therapies (26, 27). It is interesting that we observe inhibition of endothelial cell function within the range of menadione concentrations found to be effective in killing a variety of tumor cells in vitro (37, 38), but loss of endothelial cell viability occurred at a 200-fold higher concentration.

Differential sensitivities of various cell types to the effects of menadione may be due to differences in number of mitochondria (23), in the relative contribution of extramitochondrial metabolism of the drug (22), in the intracellular content of NADPH (23), or in the cellular content of GSH (23, 36). Jones et al. (23) proposed that renal epithelial cells were more sensitive to menadione cytotoxicity than hepatocytes, because menadione cytotoxicity is dependent on mitochondrial metabolism of the drug and the renal cells contain more mitochondria. This would predict that endothelial cells would be less sensitive than hepatocytes to menadione effects, since they contain few mitochondria (35). However, our data indicate that the dose relationship for menadione-induced cytotoxicity in the endothelial cell is similar to those observed in hepatocytes (21, 39). For menadione to have a similar potency in both cell types, endothelial cells must have a great capacity for microsomal metabolism of the drug. This is supported by the fact that endothelial cells are predominately glycolytic and are capable of increasing their level of NADPH many fold upon demand (35). Such an increase in NADPH could augment deleterious univalent reduction of menadione by the microsomal electron transport systems.

The time course and dose dependence of menadione-induced H_2O_2 formation are consistent with menadione-induced inhibition of PGI_2 synthesis, supporting our hypothesis that it is H_2O_2 which mediates this inhibition. We have previously demonstrated that sublethal doses of H_2O_2 inhibit endothelial cell prostaglandin synthesis by irreversibly inactivating the PGH_2 synthase (10, 11). In the present study, menadione was shown to cause a similar inhibition although this drug has a minor effect on the PGI_2 synthase as well. PGI_2 synthase is known to be sensitive to lipid hydroperoxides (18, 32, 33), but menadione-induced formation of these species is unlikely, since menadione is known to prevent propagation of lipid peroxides (40, 41). Since we have previously demonstrated that endothelial cell PGI_2 synthase is not affected by H_2O_2 , it is likely that menadione, or the semiquinone, has a direct effect on this enzyme.

Nolan and Eling (34) demonstrated that menadione had a biphasic effect on A23187-stimulated PGI_2 synthesis in endothelial cells, with concentrations of $> 1 \mu M$ augmenting production. The mechanism for this stimulation was not described, although an analogue of menadione, vitamin K_1 , was shown to attenuate PGI_2 synthesis by preventing release of endogenous arachidonate. Since the stimulatory effect of menadione was not characterized, it is difficult to explain the difference between the findings of this previous study and our present data. However, since we did not find an effect of menadione on either bradykinin- or A23187-stimulated release, we conclude that the inhibitory effects of menadione on endothelial prostaglandin production are limited to the synthetic enzymes of the cascade.

Consistent with our previous findings (11), we demonstrated that, although PGH_2 synthase is irreversibly inactivated after treatment with menadione, the endothelial cell can recover lost capacity to produce prostaglandins. In H_2O_2 -treated

endothelial cells, this recovery was shown to be dependent on synthesis of both new mRNA and protein (11). In a recent study, we found that this RNA synthesis occurs with a half-life of 3 h, while the half-life for new protein synthesis was only 1.3 h (42). These rates are consistent with the time required for full recovery of lost prostaglandin synthetic capacity.

In conclusion, these data demonstrate that sublethal concentrations of menadione cause a rapid, irreversible inactivation of endothelial cell prostaglandin production. This inhibition appears to be dependent on formation of reactive oxygen species after metabolism of the drug by the endothelial cell. Consistent with our previous findings (10, 11), this oxidant-mediated inhibition is limited to the synthetic enzymes of the prostaglandin cascade and the activity of these enzymes can be restored by synthesizing new enzyme. These data indicate that the effects of menadione on vascular cell function can greatly proceed the cytotoxic effects of the drug.

Acknowledgments

A. Barchowsky was supported by NIEHS training grant T32-E507031. The authors wish to thank Lori Tripp for her excellent help in preparing this manuscript.

This work was supported by grants AG-02868 and HL-31992 from the National Institutes of Health and by a grant from the Burroughs Wellcome Fund.

References

1. Suttorp, N., and L. M. Simon. 1982. Lung cell oxidant injury. Enhancement of polymorphonuclear leukocyte-mediated cytotoxicity in lung cells exposed to sustained in vitro hyperoxia. *J. Clin. Invest.* 70:342-350.
2. Tate, R. M., and J. E. Repine. 1983. Neutrophils and adult respiratory distress syndrome. *Am. Rev. Respir. Dis.* 128:552-559.
3. Autor, A. P., A. C. Bonham, and R. L. Theis. 1984. Toxicity of oxygen radicals in cultured pulmonary endothelial cells. *J. Toxicol. Environ. Health.* 13:387-395.
4. Sjostrom, K., and J. D. Crapo. 1983. Structural and biochemical adaptive changes in rat lungs after exposure to hypoxia. *Lab. Invest.* 48:68-79.
5. Orr, F. W., and D. J. A. Warner. 1987. Effects of neutrophil-mediated pulmonary injury on the localization of circulating Walker carcinoma cells. *Invasion Metastasis.* 7:183-196.
6. Reidy, M. A., and S. M. Schwartz. 1983. Endothelial injury and regeneration. IV. Endotoxin: a non-denuding injury to aortic endothelium. *Lab. Invest.* 48:25-34.
7. Brigham, K. L., and B. Meyrick. 1986. Endotoxin and lung injury. *Am. Rev. Respir. Dis.* 133:913-927.
8. Lazo, J. S. 1986. Endothelial injury caused by antineoplastic agents. *Biochem. Pharmacol.* 35:1919-1923.
9. Adamson, I. Y. R. 1984. Drug-induced pulmonary fibrosis. *Environ. Health. Perspect.* 55:25-36.
10. Whorton, A. R., M. E. Montgomery, and R. S. Kent. 1985. Effect of hydrogen peroxide on prostaglandin production and cellular integrity in cultured porcine aortic endothelial cells. *J. Clin. Invest.* 76:295-302.
11. Barchowsky, A., R. S. Kent, and A. R. Whorton. 1987. Recovery of porcine aortic endothelial cell prostaglandin synthesis following inhibition by sublethal concentrations of hydrogen peroxide. *Biochim. Biophys. Acta.* 927:372-381.
12. Gryglewski, R. J., and S. Moncada. 1987. Secretory function of vascular endothelium. *Adv. Prostaglandin Thromboxane Leukotriene Res.* 17:397-404.
13. Simpson, P. J., and B. R. Lucchesi. 1987. Free radicals and

- myocardial ischemia and reperfusion injury. *J. Lab. Clin. Med.* 110:13-30.
14. Czervionke, R. C., J. B. Smith, G. L. Fry, J. C. Hoak, and D. C. Haycraft. 1979. Inhibition of prostacyclin by treatment of endothelium with aspirin: correlation with platelet adherence. *J. Clin. Invest.* 63:1089-1092.
 15. Fantone, J. C., and D. A. Kinnes. 1983. Prostaglandin E₁ and prostaglandin I₂ modulation of superoxide production by neutrophils. *Biochem. Biophys. Res. Commun.* 113:506-512.
 16. Kulmacz, R. J. 1986. Prostaglandin H synthase and hydroperoxides: peroxidase reaction and inactivation. *Arch. Biochem. Biophys.* 249:273-285.
 17. Marshall, P. J., R. J. Kulmacz, and W. E. M. Lands. 1987. Constraints on prostaglandin biosynthesis in tissues. *J. Biol. Chem.* 262:3510-3517.
 18. Kent, R. S., S. L. Deidrich, and A. R. Whorton. 1983. Regulation of vascular prostaglandin synthesis by metabolites of arachidonic acid in perfused rabbit aorta. *J. Clin. Invest.* 72:455-465.
 19. Brotherton, A. F., and J. C. Hoak. 1983. Prostaglandin biosynthesis in cultured vascular endothelium is limited by deactivation of cyclooxygenase. *J. Clin. Invest.* 72:1255-1261.
 20. Madden, M. C., T. E. Eling, and M. Friedman. 1987. Ozone inhibits endothelial cell cyclooxygenase activity through formation of hydrogen peroxide. *Prostaglandins.* 34:445-463.
 21. Thor, H., M. T. Smith, P. Hartzell, G. Bellomo, S. A. Jewell, and S. Orrenius. 1982. The metabolism of menadione (2-methyl-1,4-naphthoquinone) by isolated hepatocytes. *J. Biol. Chem.* 257:12419-12425.
 22. de Groot, H., T. Noll, and H. Sies. 1985. Oxygen dependence and subcellular partitioning of hepatic menadione-mediated oxygen uptake. Studies with isolated hepatocytes, mitochondria and microsomes from rat liver in an oxytstat system. *Arch. Biochem. Biophys.* 243:556-562.
 23. Jones, T. W., H. Thor, and S. Orrenius. 1986. Cellular defense mechanisms against toxic substances. *Arch. Toxicol. Suppl.* 9:259-271.
 24. Frei, B., K. H. Winterhalter, and C. Richter. 1986. Menadione-(2-methyl-1,4-naphthoquinone)-dependent enzymatic redox cycling and calcium release by mitochondria. *Biochemistry.* 25:4438-4443.
 25. Rosen, G. M., and B. A. Freeman. 1984. Detection of superoxide generated by endothelial cells. *Proc. Natl. Acad. Sci. USA.* 81:7269-7273.
 26. Whorton, A. R., S. L. Young, J. L. Data, A. Barchowsky, and R. S. Kent. 1982. Mechanism of bradykinin-stimulated prostacyclin synthesis in porcine aortic endothelial cells. *Biochim. Biophys. Acta.* 712:79-87.
 27. Thurman, R. G., H. G. Ley, and R. Scholz. 1972. Hepatic microsomal ethanol oxidation. Hydrogen peroxide and the role of catalase. *Eur. J. Biochem.* 25:420-430.
 28. Bergmeyer, H. O. 1955. Zur messung von Katalase-aktivitäten. *Biochem. Z.* 327:255-258.
 29. McCord, J. M., and I. Fridovich. 1965. Superoxide dismutase. An enzymatic function for erythrocyte hemocuprein (hemocuprein). *J. Biol. Chem.* 244:6049-6055.
 30. Thayer, W. S. 1977. Adriamycin stimulated superoxide formation in submitochondrial particles. *Chem. Biol. Interact.* 19:265-278.
 31. Hassan, H. M., and I. Fridovich. 1979. Intracellular production of superoxide radical and of hydrogen peroxide by redox activity compounds. *Arch. Biochem. Biophys.* 196:385-395.
 32. Demopoulos, H. B., E. S. Flam, and D. D. Pietromyro. 1980. The free radical pathology and the microcirculation in the major central nervous system disorders. *Acta Physiol. Scand.* 492:91-119.
 33. Salmon, J. A., D. R. Smith, R. J. Flower, S. Moncada, and J. R. Vane. 1978. Further studies on the enzymatic conversion of prostaglandin endoperoxide into prostacyclin by porcine aorta microsomes. *Biochim. Biophys. Acta.* 523:250-262.
 34. Nolan, R. D., and T. E. Eling. 1986. Inhibition of prostacyclin synthesis in cultured bovine aortic cells by vitamin K₁. *Biochem. Pharmacol.* 35:4273-4281.
 35. Dobrina, A., and F. Rossi. 1983. Metabolic properties of freshly isolated bovine endothelial cells. *Biochim. Biophys. Acta.* 762:295-301.
 36. Bellomo, G., F. Mirabelli, D. DiMonte, P. Richelmi, H. Thor, C. Orrenius, and S. Orrenius. 1987. Formation and reduction of glutathione-protein mixed disulfides during oxidative stress. A study with isolated hepatocytes and menadione (2-methyl-1,4-naphthoquinone). *Biochem. Pharmacol.* 36:1313-1320.
 37. Chlebowski, R. T., S. A. Akman, and J. B. Block. 1985. Vitamin K in the treatment of cancer. *Cancer Treat. Rev.* 12:49-63.
 38. Su, Y-Z., T. E. Duarte, P. L. Dill, and L. M. Weisenthal. 1987. Selective enhancement by menadiol of in vivo drug activity in human lymphatic neoplasms. *Cancer Treat. Rep.* 71:619-625.
 39. DiMonte, D., G. Bellomo, H. Thor, P. Nicoteva, and S. Orrenius. 1984. Menadione-induced cytotoxicity is associated with protein thiol oxidation and alteration in intracellular Ca²⁺ homeostasis. *Arch. Biochem. Biophys.* 235:343-350.
 40. Talcott, R. E., M. T. Smith, and D. D. Giannini. 1985. Inhibition of microsomal lipid peroxidation by naphthoquinones: structure-activity relationships and possible mechanisms of action. *Arch. Biochem. Biophys.* 241:88-94.
 41. Younes, M., S. Cornelius, and C. P. Siegers. 1985. Fe²⁺-supported in vivo lipid peroxidation induced by compounds undergoing redox cycling. *Chem. Biol. Interact.* 54:97-103.
 42. Whorton, A. R., Z. Mirza, A. Barchowsky, and B. J. Buckley. 1988. Protein kinase C regulation of arachidonate release and metabolism in aortic endothelial cells. *FASEB (Fed. Am. Soc. Exp. Biol.) J.* 2:A1188. (Abstr.)