



Published in final edited form as:

Biochem Biophys Res Commun. 2011 January 7; 404(1): 424–427. doi:10.1016/j.bbrc.2010.11.137.

Visualization and quantification of endoplasmic reticulum Ca²⁺ in renal cells using confocal microscopy and Fluo5F

Andre C. Eaddy¹ and Rick G. Schnellmann^{1,2}

¹ Center for Cell Death, Injury, and Regeneration, Department of Pharmaceutical and Biomedical Sciences, Medical University of South Carolina, Charleston, SC 29425

² Ralph H. Johnson VA Medical Center Charleston, SC 29401

Abstract

Sarcoplasmic/endoplasmic reticulum (ER) Ca²⁺ is the most abundant store of intracellular Ca²⁺, and its release is an important trigger of physiological and cell death pathways. Previous work in our laboratory revealed the importance of ER Ca²⁺ in toxicant-induced renal proximal tubular cell (RPTC) death. The purpose of this study was to evaluate the use of confocal microscopy and Fluo5F, a low affinity Ca²⁺ indicator, to directly monitor changes in RPTC ER Ca²⁺. Fluo5F staining reflected ER Ca²⁺, resolved ER structure, and showed no colocalization with tetramethyl rhodamine methyl ester (TMRM), a marker of mitochondrial membrane potential. Thapsigargin, an ER Ca²⁺ pump inhibitor, decreased ER fluorescence by 30% and 55% at 5 and 15 min, respectively, whereas A23187, a Ca²⁺ ionophore caused more rapid ER Ca²⁺ release (55% and 75% decrease in fluorescence at 5 and 15 min).

Carbonylcyanide-p-trifluoromethoxyphenylhydrazone (FCCP), a mitochondrial uncoupler, added at the end of the experiment, further decreased ER fluorescence after thapsigargin treatment, revealing that thapsigargin did not release all ER Ca²⁺. In contrast, FCCP did not decrease ER fluorescence after A23187 treatment, suggesting complete ER Ca²⁺ release. ER Ca²⁺ release in response to A23187 or thapsigargin resulted in a modest but significant decrease in mitochondrial membrane potential. These data provide evidence that confocal microscopy and Fluo5F are useful and effective tools for directly monitoring ER Ca²⁺ in intact cells.

Keywords

calcium; endoplasmic reticulum; confocal microscopy; kidney; Fluo5F

INTRODUCTION

Ca²⁺ acts as a universal second messenger and regulates numerous cellular functions including metabolism, motility, and transport[1]. Loss of Ca²⁺ homeostasis is critical to

Author to whom correspondence should be addressed: Rick G. Schnellmann, Department of Pharmaceutical and Biomedical Sciences, Medical University of South Carolina, 280 Calhoun St, MSC 140, Charleston, SC 29425–1400, Phone: 843–792–3754, Fax: 843–792–2620, schnell@musc.edu.

Disclaimer. The contents of this manuscript do not represent the views of the Department of Veteran Affairs or the United States Government.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

many disease processes and is a major component of cell death pathways including necrosis, apoptosis, and autophagy [2,3,4,5,6,7,8]. Sarcoplasmic/endoplasmic reticulum (ER) Ca^{2+} is the most abundant store of intracellular Ca^{2+} and its disruption often initiates the deleterious cascade of events leading to cell death and dysfunction [5,6,7,8,9,10,11].

Traditional methods of measuring of ER Ca^{2+} are indirect or require difficult probe loading techniques (i.e. membrane permeabilization, microinjection, or fused cell hybrids). For example, human embryonic kidney cells require 1 h of dye loading (2 μM Fluo-3 at room temperature) and extracellular Ca^{2+} chelation using BAPTA or EGTA, just to monitor increases in cytosolic Ca^{2+} as an indirect measure of ER Ca^{2+} stores [12]. To monitor ER Ca^{2+} directly, Montero and Robert *et al.* engineered an ER-targeted aequorin chimera, a Ca^{2+} -sensitive photoprotein with a lower affinity for Ca^{2+} . However, in HeLa and skeletal muscle cells, the aequorin chimera was rapidly saturated by the high Ca^{2+} concentrations within the ER and required a non-physiological reduction of ER Ca^{2+} to detect changes in Ca^{2+} stores [13,14].

Primary cultures of renal proximal tubular cells (RPTC) are highly aerobic, gluconeogenic, and exhibit robust transport capacity, making them ideal for the study of kidney tubular function and injury [15,16]. Previous work in our laboratory revealed the importance of ER Ca^{2+} in toxicant-induced kidney injury; although, the mechanism by which Ca^{2+} plays such a pivotal role is not completely understood [5,7,9,10,17,18].

Confocal microscopy is a useful method for the visualization and quantification of fluorophores at the subcellular level in living cells, and is compatible with most Ca^{2+} indicators. Chemical fluorescent (UV and visible-wavelength excitation fluorescent indicators) and bioluminescent calcium indicators (Ca^{2+} binding photoproteins and GFP-based Ca^{2+} indicators) differ in their characteristics (loading requirements, excitation/emission spectrum, permeability, compartmentalization, relative brightness, and Ca^{2+} affinity) and have inherent drawbacks (i.e., dye leakage, cytotoxicity, bleaching, autofluorescence, intracellular buffering, and lack of ion specificity)[19]. Fluo5F is a chemical fluorescent Ca^{2+} indicator with a lower affinity for Ca^{2+} ($K_d \sim 2.3 \mu\text{M}$), limited cytotoxicity and bleaching, and high Ca^{2+} specificity, making it ideal for studying ER Ca^{2+} . The purpose of this study was to evaluate the use of confocal microscopy and Fluo5F for directly monitoring changes in ER Ca^{2+} in RPTC.

MATERIALS AND METHODS

Materials

Female New Zealand White rabbits (1.5–2.0 kg) were purchased from Myrtle's Rabbitry (Thompson Station, TN). Tetramethyl rhodamine methyl ester (TMRM) and Fluo-5F-AM were purchased from Molecular Probes, Invitrogen (Carlsbad, CA). All other chemical and materials were obtained from Sigma Chemical (St. Louis, MO).

Isolation of Rabbit RPTC and Culture Conditions

Rabbit RPTC were isolated using the iron oxide perfusion method and grown to confluence in 35-mm tissue culture dishes under improved conditions as previously described [15,16]. The culture medium was a 1:1 mixture of Dulbecco's modified Eagle's medium/Ham's F-12 medium (without glucose, phenol red, or sodium pyruvate) supplemented with 15 mM HEPES buffer, 2.5 mM L-glutamine, 1 μM pyridoxine HCl, 15 mM sodium bicarbonate, and 6 mM lactate. Hydrocortisone (50 nM), selenium (5 ng/ml), human transferrin (5 $\mu\text{g}/\text{ml}$), bovine insulin (10 nM), and L-ascorbic acid-2-phosphate (50 μM) were added daily to fresh culture medium.

RPTC Loading

RPTC were loaded with 2 μM Fluo-5F-AM (Fluo5F) and 100 nM TMRM for 20 min at 37° C, washed twice with 37° C phosphate buffered saline, and media was replaced. Then, 1 μM Fluo5F and 40 nM TMRM were added to the media to maintain dye equilibrium and incubated for 30 min at 37° C prior to imaging. The TMRM loading protocol was modified from Lemasters and Ramshesh [20].

Microscopic Imaging and Analysis

RPTC were imaged on a Leica Microsystems, TCS SP2 AOBS laser scanning confocal microscope (LSCM) using standardized pinhole, gain, and black level settings. A 63 \times 0.9 NA water-immersion objective was used in an upright microscope configuration. All images were acquired at 8-bit resolution and at 1024 \times 1024 with a line averaging of two.

RPTC were monitored for 10 min prior to treatment to establish baseline. Then, RPTC were treated and monitored an additional 15 min prior to Carbonylcyanide-p-trifluoromethoxyphenylhydrazone (FCCP, 1 μM) addition to depolarize mitochondria. Images were acquired in 1-min intervals during the course of experiments. To quantify data, images were analyzed using Adobe Photoshop. Mean intensities of images in the green and red channel after background subtraction were interpreted as a quantitative measure of ER Ca^{2+} and mitochondrial membrane potential, respectively. Values are graphed as the percent mean intensity of each subsequent image vs. mean intensity of the image taken at time zero. Images comprise of 7–10 cells.

Statistical Analysis

RPTC isolated from one animal represents one experiment (N=1). Data are presented as means \pm SE, and multiple means were compared using Student-Newman-Keuls test at each time point. Means with values $p < 0.05$ were considered statistically significant.

RESULTS AND DISCUSSION

To investigate the use of confocal microscopy to monitor ER Ca^{2+} dynamics in RPTC, Fluo5F fluorescence was monitored as described above (Fig. 1a). Fluo5F fluorescence revealed a reticular pattern around nuclei void of staining. The peripheral granularity of staining and scarcity of staining at the cell margins provided evidence that the probe is not fluorescing in the cytosol. To confirm that Fluo5F was not in the mitochondria, RPTC were loaded with tetramethyl rhodamine methyl ester (TMRM), a marker of polarized mitochondria. TMRM staining (Fig. 1b) had a punctuate/tubular pattern unlike Fluo5F staining with no co-localization (Fig. 1c), confirming that Fluo5F does not reflect mitochondrial Ca^{2+} , the second most significant storage/buffering compartment for intracellular Ca^{2+} [4].

We investigated the use of Fluo5F to monitor real-time changes in RPTC ER Ca^{2+} . TMRM was used to measure mitochondrial membrane potential. In control cells, both Fluo5F and TMRM fluorescence intensity, morphology, and resolution were maintained over time (Fig. 2). Carbonylcyanide-p-trifluoromethoxyphenylhydrazone (FCCP), a mitochondrial uncoupler, was added at the end of each experiment to cause mitochondrial depolarization and indirectly ER Ca^{2+} release, due to ATP depletion and cessation of sarcoplasmic/ER Ca^{2+} ATPase (SERCA) activity.

We then explored the effects of two pharmacological agents known to alter ER Ca^{2+} . RPTC were treated with the SERCA inhibitor, thapsigargin, and the Ca^{2+} ionophore, A23187. Thapsigargin decreased ER Ca^{2+} over several min (Fig. 2A), which was more indicative of

ER Ca²⁺ leakage than receptor-dependent ER Ca²⁺ release that is characterized by DAG, ryanodine, and IP₃ receptor sensitive stores [1,21,22]. In contrast, A23187 caused a more rapid release of ER Ca²⁺, an anticipated finding due to its ionophore activity.

Temporal and incremental changes in ER Ca²⁺ release and mitochondrial membrane potential were quantified (Fig. 3). A23187 decreased fluorescence by 50% and 70% at 5 and 15 min after treatment, respectively, and FCCP addition did not decrease fluorescence further. Thapsigargin decreased fluorescence by 25% and 55% at 5 and 15 min after treatment, respectively. In contrast to A23187, FCCP addition further decreased fluorescence after thapsigargin, providing evidence that thapsigargin did not completely release all ER Ca²⁺ stores under these conditions.

The advantage of this approach is that ER Ca²⁺ stores are measured directly, allowing observation of spatiotemporal differences between A23187- and thapsigargin-induced ER Ca²⁺ release that would otherwise be masked by measuring cytosolic free Ca²⁺. Interestingly, Jiang *et al.* reported a higher rate and magnitude of cytosolic Ca²⁺ increase in response to A23187 over thapsigargin using Fluo-3, but this approach required chelation of extracellular Ca²⁺ prior to treatment [12].

The effect of ER Ca²⁺ release on mitochondrial polarization was monitored using TMRM (Fig. 4). A23187- and thapsigargin-induced ER Ca²⁺ release decreased TMRM fluorescence by approximately 10%. In contrast, FCCP, the mitochondrial uncoupler added at the end of each experiment, decreased TMRM fluorescence by 60%, reflecting complete mitochondrial depolarization. Together, these data reveal that ER Ca²⁺ release produced by A23187 or thapsigargin partially decreases mitochondrial polarization and is consistent with previous reports [23].

In conclusion, confocal microscopy and Fluo5F are valuable tools for visualization of spatiotemporal dynamics of ER Ca²⁺ release in complex cellular systems in real time and will aid in our understanding of cellular dynamics and death.

RESEARCH HIGHLIGHTS

- A new method of monitoring ER Ca²⁺ directly in intact cells
- Confocal microscopy and low affinity Ca²⁺ indicator, Fluo5F, reveals ER Ca²⁺ morphology
- Fluo5F fluorescence shows no co-localization with mitochondria or cytosol
- Stimulation of ER Ca²⁺ release reveal spatiotemporal differences in ER Ca²⁺ release dynamics

Acknowledgments

This work was supported by an NIH fellowship (F30 ES015964) to AE and NIH grant (DK-062028) to RGS, and by the Biomedical Laboratory Research and Development Program of the Department of Veterans Affairs. Animal facilities were funded by NIH grant C06 RR-015455 and Cell and Molecular Imaging (CMI) facility was supported, in part, by grant P30 CA138313 to the Hollings Cancer Center, Medical University of South Carolina. The authors also thank Dr. Venkat Ramshesh and Dr. John Lemasters (Medical University of South Carolina) for their advice and constructive discussions during the execution of these experiments.

Abbreviations

ER endoplasmic reticulum

EGTA	ethylene glycol tetraacetic acid
RPTC	renal proximal tubular cell
TMRM	tetramethyl rhodamine methyl ester
Fluo5F	Fluo-5F-AM
LSCM	laser scanning confocal microscopy
FCCP	Carbonylcyanide-p-trifluoromethoxyphenylhydrazone
SERCA	sarcoplasmic/endoplasmic reticulum ATPase
DAG	diacylglycerol
IP3	inositol triphosphate
DMSO	dimethyl sulfoxide

References

1. Berridge MJ. Inositol trisphosphate and calcium signaling. *Ann N Y Acad Sci* 1995;766:31–43. [PubMed: 7486679]
2. Cummings BS, Kinsey GR, Bolchoz LJ, Schnellmann RG. Identification of caspase-independent apoptosis in epithelial and cancer cells. *J Pharmacol Exp Ther* 2004;310:126–134. [PubMed: 15028782]
3. Lemasters JJ, Nieminen AL, Qian T, Trost LC, Elmore SP, Nishimura Y, Crowe RA, Cascio WE, Bradham CA, Brenner DA, Herman B. The mitochondrial permeability transition in cell death: a common mechanism in necrosis, apoptosis and autophagy. *Biochim Biophys Acta* 1998;1366:177–196. [PubMed: 9714796]
4. Lemasters JJ, Qian T, Elmore SP, Trost LC, Nishimura Y, Herman B, Bradham CA, Brenner DA, Nieminen AL. Confocal microscopy of the mitochondrial permeability transition in necrotic cell killing, apoptosis and autophagy. *Biofactors* 1998;8:283–285. [PubMed: 9914830]
5. Schnellmann RG. Intracellular calcium chelators and oxidant-induced renal proximal tubule cell death. *J Biochem Toxicol* 1991;6:299–303. [PubMed: 1774771]
6. Waters SL, Sarang SS, Wang KK, Schnellmann RG. Calpains mediate calcium and chloride influx during the late phase of cell injury. *J Pharmacol Exp Ther* 1997;283:1177–1184. [PubMed: 9399991]
7. Waters SL, Wong JK, Schnellmann RG. Depletion of endoplasmic reticulum calcium stores protects against hypoxia- and mitochondrial inhibitor-induced cellular injury and death. *Biochem Biophys Res Commun* 1997;240:57–60. [PubMed: 9367881]
8. Pozzan T, Rizzuto R, Volpe P, Meldolesi J. Molecular and cellular physiology of intracellular calcium stores. *Physiol Rev* 1994;74:595–636. [PubMed: 8036248]
9. Liu X, Van Vleet T, Schnellmann RG. The role of calpain in oncotic cell death. *Annu Rev Pharmacol Toxicol* 2004;44:349–370. [PubMed: 14744250]
10. Harriman JF, Liu XL, Aleo MD, Machaca K, Schnellmann RG. Endoplasmic reticulum Ca(2+) signaling and calpains mediate renal cell death. *Cell Death Differ* 2002;9:734–741. [PubMed: 12058278]
11. Muruganandan S, Cribb AE. Calpain-induced endoplasmic reticulum stress and cell death following cytotoxic damage to renal cells. *Toxicol Sci* 2006;94:118–128. [PubMed: 16920763]
12. Jiang Y, Cypess AM, Muse ED, Wu CR, Unson CG, Merrifield RB, Sakmar TP. Glucagon receptor activates extracellular signal-regulated protein kinase 1/2 via cAMP-dependent protein kinase. *Proc Natl Acad Sci U S A* 2001;98:10102–10107. [PubMed: 11517300]
13. Robert V, De Giorgi F, Massimino ML, Cantini M, Pozzan T. Direct monitoring of the calcium concentration in the sarcoplasmic and endoplasmic reticulum of skeletal muscle myotubes. *J Biol Chem* 1998;273:30372–30378. [PubMed: 9804801]

14. Montero M, Brini M, Marsault R, Alvarez J, Sitia R, Pozzan T, Rizzuto R. Monitoring dynamic changes in free Ca²⁺ concentration in the endoplasmic reticulum of intact cells. *EMBO J* 1995;14:5467–5475. [PubMed: 8521803]
15. Nowak G, Schnellmann RG. Improved culture conditions stimulate gluconeogenesis in primary cultures of renal proximal tubule cells. *Am J Physiol* 1995;268:C1053–1061. [PubMed: 7733227]
16. Nowak G, Schnellmann RG. L-ascorbic acid regulates growth and metabolism of renal cells: improvements in cell culture. *Am J Physiol* 1996;271:C2072–2080. [PubMed: 8997210]
17. Cummings BS, McHowat J, Schnellmann RG. Role of an endoplasmic reticulum Ca²⁺-independent phospholipase A2 in cisplatin-induced renal cell apoptosis. *J Pharmacol Exp Ther* 2004;308:921–928. [PubMed: 14634037]
18. Cummings BS, McHowat J, Schnellmann RG. Role of an endoplasmic reticulum Ca(2+)-independent phospholipase A(2) in oxidant-induced renal cell death. *Am J Physiol Renal Physiol* 2002;283:F492–498. [PubMed: 12167600]
19. Takahashi A, Camacho P, Lechleiter JD, Herman B. Measurement of intracellular calcium. *Physiol Rev* 1999;79:1089–1125. [PubMed: 10508230]
20. Lemasters JJ, Ramshesh VK. Imaging of mitochondrial polarization and depolarization with cationic fluorophores. *Methods Cell Biol* 2007;80:283–295. [PubMed: 17445700]
21. Bian JH, Ghosh TK, Wang JC, Gill DL. Identification of intracellular calcium pools. Selective modification by thapsigargin. *J Biol Chem* 1991;266:8801–8806. [PubMed: 1827436]
22. Hirose K, Kadowaki S, Tanabe M, Takeshima H, Iino M. Spatiotemporal dynamics of inositol 1,4,5-trisphosphate that underlies complex Ca²⁺ mobilization patterns. *Science* 1999;284:1527–1530. [PubMed: 10348740]
23. Lemasters JJ, Theruvath TP, Zhong Z, Nieminen AL. Mitochondrial calcium and the permeability transition in cell death. *Biochim Biophys Acta* 2009;1787:1395–1401. [PubMed: 19576166]




Figure 1. Morphology of Renal Proximal Tubule Cells (RPTC) co-loaded with Fluo5F and TMRM using confocal microscopy

RPTC were co-loaded with Fluo5F (2 μM) and TMRM (100 nM) as described in the materials and methods. Green fluorescence of Fluo5F and red fluorescence of TMRM were imaged by laser scanning confocal microscopy. The white bar represents 10 μm . Fluo5F fluorescence (A) represents endoplasmic reticulum (ER) Ca^{2+} , TMRM fluorescence (B) represents polarized mitochondria, and there was no co-localization (C).




Figure 2. ER Ca²⁺ release RPTC exposed to A23187 and thapsigargin
 RPTC were loaded with Fluo5F and TMRM (see in Fig. 1), monitored for 10 min to establish baseline, exposed to diluent (DMSO, Control), A23187 (10 μ M), or thapsigargin (10 μ M) for 15 min, and then exposed to FCCP (1 μ M), a mitochondrial uncoupler. Open arrows represent time of A23187 or thapsigargin treatment and closed arrows represent the addition of FCCP. Green fluorescence (A) and red fluorescence (B) represents ER Ca²⁺ and mitochondrial polarization, respectively. Note the more rapid ER Ca²⁺ release (A) in response to A23187 exposure compared to thapsigargin and the rapid loss of mitochondrial polarization (B) in response to FCCP.




Figure 3. Quantification of A23187- and thapsigargin- induced ER Ca²⁺ release in RPTC
RPTC were co-loaded with Fluo5F and TMRM, treated as described in fig. 2, and images acquired as described in Materials and Methods. Values are mean intensities of images in the green channel \pm SE and loss of intensity represents ER Ca²⁺ release. The open arrow represents exposure to A23187, thapsigargin, or diluent control and the closed arrow represents the addition of FCCP. Note the more rapid ER Ca²⁺ release after A23187 exposure compared to thapsigargin and an additional loss of fluorescence with FCCP following thapsigargin exposure.




Figure 4. Quantification of A23187- and thapsigargin-induced mitochondrial depolarization in RPTC

See above for experimental details. Values are mean intensities of images in the red channel \pm SE and loss of intensity represents mitochondrial depolarization. The open arrow represents exposure to A23187, thapsigargin, or diluent control and the closed arrow represents the addition of FCCP. Note the decrease in mitochondrial polarization following A23187 or thapsigargin exposure compared with control and the rapid loss of mitochondrial polarization following FCCP exposure in all groups.