



Published in final edited form as:

*J Am Chem Soc.* 2011 February 23; 133(7): 2088–2091. doi:10.1021/ja110215b.

## Enantioselective Total Syntheses of (–)-Taiwaniaquinone H and (–)-Taiwaniaquinol B by Iridium-Catalyzed Borylation and Palladium-Catalyzed Asymmetric $\alpha$ -Arylation

Xuebin Liao, Levi M. Stanley, and John F. Hartwig\*

Department of Chemistry, University of Illinois, 600 South Mathews Avenue, Urbana, Illinois 61801-3602

### Abstract

We report a concise, enantioselective total synthesis of (–)-taiwaniaquinone H and the first enantioselective total synthesis of (–)-taiwaniaquinol B by a route that includes enantioselective, palladium-catalyzed  $\alpha$ -arylation of a ketone with an aryl bromide that was generated by sterically controlled halogenation via iridium-catalyzed C–H borylation. This asymmetric  $\alpha$ -arylation creates the benzylic, quaternary stereogenic center present in the taiwaniaquinoids. The synthesis was completed efficiently by developing a Lewis acid-promoted cascade to construct the [6,5,6] tricyclic core of an intermediate common to the synthesis of a number of taiwaniaquinoids. Through the preparation of these compounds, we demonstrate the utility of constructing benzylic, quaternary, stereogenic centers, even those lacking a carbonyl group in the  $\alpha$ -position, by asymmetric  $\alpha$ -arylation.

The taiwaniaquinoids are a family of unusual diterpenoids possessing a [6,5,6]-*abeo*-abietane skeleton that was unknown in nature prior to 1995.<sup>1–7</sup> Each member of this family contains a benzylic quaternary stereogenic center (Figure 1). Studies of the biological activities of taiwaniaquinoids are ongoing, but preliminary data have indicated that some members of this family exhibit antitumor activities.<sup>8–11</sup> The structural features and the biological activities of these compounds have led many groups to report total and formal syntheses of the taiwaniaquinoids.<sup>11–25</sup> However, the majority of these syntheses have targeted racemic versions of this class of natural products.

Asymmetric syntheses of the taiwaniaquinoids are challenging due to the limited number of enantioselective reactions that establish quaternary stereogenic centers.<sup>26–29</sup> One approach that eliminates this challenge is to prepare the material by a ring-contractive derivatization of enantioenriched terpenes. Alvarez-Manzaneda and co-workers prepared (–)-taiwaniaquinone G (**1**),<sup>22</sup> (–)-taiwaniaquinone H (**2**) and (–)-dichroanone (**3**) in 15 or 16 steps from (+)-sclareolide,<sup>23</sup> and Gademann and coworkers prepared (–)-taiwaniaquinone H (**2**) in 13 steps from methyl dehydroabiate.<sup>25</sup> Recently, Alvarez-Manzaneda and co-workers reported the syntheses of (–)-taiwaniaquinone A (**4**) in 15 steps and (–)-taiwaniaquinone F (**5**) in 16 steps from (+)-podocarpa-8(14)-en-13-one.<sup>30</sup> To date, only two enantioselective total syntheses of taiwaniaquinoids from achiral or racemic starting materials have been reported. McFadden and Stoltz reported an 11-step synthesis of (+)-dichroanone (*ent*-**3**) in which the quaternary stereogenic center was set by an asymmetric Tsuji allylation.<sup>17</sup>

jhartwig@illinois.edu.

**Supporting Information Available:** Experimental details, characterization data, and <sup>1</sup>H and <sup>13</sup>C NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Recently, Node and coworkers employed an enantioselective, intramolecular Heck reaction for the total syntheses of (–)-dichroanal **6** in 12 steps, (–)-dichroanone (**3**) in 14 steps, and (–)-taiwaniaquinone H (**2**) in 15 steps beginning with a tetrasubstituted arene.<sup>24</sup>


Previously, our group reported the asymmetric  $\alpha$ -arylation of ketones to form products containing quaternary benzylic stereogenic centers.<sup>31</sup> Although the taiwaniaquinoids do not contain a carbonyl group  $\alpha$  to the benzylic stereogenic center, we envisioned that asymmetric  $\alpha$ -arylation could be used to construct this center and that the ketone function could be integral to creation of the [6,5,6] tricyclic core. Further, we envisioned that the aryl electrophile for the proposed  $\alpha$ -arylation process could be generated from a simple arene by a sterically controlled iridium-catalyzed borylation and halogenation sequence developed by our group.<sup>32,33</sup>

Herein, we report concise, enantioselective total syntheses of (–)-taiwaniaquinone H (**2**) and (–)-taiwaniaquinol B (**7**) from simple achiral and racemic reactants through this halogenation process and an asymmetric  $\alpha$ -arylation for the first time in the synthesis of a natural product. A Lewis acid-promoted cascade then establishes the unusual [6,5,6] tricyclic core.

Scheme 1 illustrates the strategy we envisioned to achieve the enantioselective total syntheses of (–)-taiwaniaquinone H (**2**) and (–)-taiwaniaquinol B (**7**). Both molecules would be accessible from the same intermediate, tetrahydrofluorene **8**, by oxidation and demethylation procedures. We considered that the tricyclic core of tetrahydrofluorene **8** could be formed from epoxide **9** by a formal intramolecular addition of the electron-rich arene ring to the epoxide unit. Epoxide **9** would be derived from  $\alpha$ -aryl ketone **10**, which would be generated by our palladium-catalyzed asymmetric  $\alpha$ -arylation of ketones.

In the forward direction, the first step of the synthesis required aryl bromide **12**. Under traditional conditions for electrophilic aromatic substitution, the known resorcinol derivative **13**<sup>34</sup> undergoes bromination at the position located ortho to the two methoxy groups.<sup>16</sup> To overcome this electronic bias, we used a one-pot, two-step sequence to prepare aryl bromide **12** from **13** by the combination of a sterically controlled, iridium-catalyzed borylation and a subsequent bromination of the aryl boronate ester with copper(II) bromide (Scheme 2).<sup>32</sup> This method provided aryl bromide **12** in 75% yield on a multi-gram scale.


Access to aryl bromide **12** enabled us to explore the key enantioselective  $\alpha$ -arylation. The reaction of aryl bromide **12** with 2,2,6-trimethylcyclohexanone did not generate the  $\alpha$ -arylation product in the presence of a catalyst generated from Pd(dba)<sub>2</sub> and (*R*)-Segphos or (*R*)-Difluorophos. However, the reaction of aryl bromide **12** with the 2-methylcyclohexanone derivative **11**<sup>35,36</sup> in the presence of 10 mol % of the combination of Pd(dba)<sub>2</sub> and (*R*)-Difluorophos as catalyst formed the  $\alpha$ -aryl ketone **14** in 80% yield with 94% ee (eq 1).<sup>37</sup>



(1)

The synthesis of the key tetrahydrofluorene intermediate **8** was accomplished in five steps from enantioenriched ketone **14** (Scheme 3). Acid-promoted hydrolysis of ketone **14**, followed by deformylation of the resulting  $\beta$ -ketoaldehyde in the presence of aqueous sodium hydroxide, provided the deprotected ketone **15** in 91% yield over two steps. Dimethylation of **15** with methyl iodide and sodium bis(trimethylsilyl)amide (NaHMDS) as the base formed the tetra-substituted cyclohexanone **10** in 86% yield.

The absolute configuration of the stereogenic center in these products was established by iodination of **15**. The reaction of **15** with iodine and silver sulfate formed diiodide **16** (eq 2). Single-crystal X-ray diffraction of the major diastereomer of **16** enabled the assignment of the absolute configuration as (2*R*,6*S*)-**16** (see Figure S1 in the Supporting Information). Thus, the enantioselective  $\alpha$ -arylation of ketone **11** conducted with (*R*)-Difluorophos and Pd(dba)<sub>2</sub> generates (*R*)-**14** and creates the appropriate enantiomer to prepare (–)-taiwaniaquinone H (**2**) and (–)-taiwaniaquinol B (**7**).



(2)

Attempts to convert ketone **10** to epoxide **9** by Corey-Chaykovsky epoxidation with the common sulfur ylide, dimethylsulfoxonium methylide,<sup>38,39</sup> led to recovery of **10**, even under forcing conditions (80 °C). We presumed this lack of reactivity resulted from the steric hindrance surrounding the ketone functionality in **10**. Thus, we attempted a Corey-Chaykovsky epoxidation of ketone **10** with the smaller and more reactive sulfur ylide, dimethylsulfonium methylide.<sup>39,40</sup> This reaction formed epoxide **9** in 95% yield with >15:1 diastereoselectivity.

Some Lewis acids are known to promote the rearrangement of epoxides to carbonyl compounds.<sup>41–45</sup> We tested the ability of such Lewis acids to initiate the cascade shown in Scheme 3 in which epoxide **9** would form tetrahydrofluorene **8** via rearrangement of the epoxide to the aldehyde. This aldehyde would then be attacked by the electron rich arene in the presence of a Lewis acid to form the corresponding alcohol, which could undergo elimination to form the target indene **8**. After testing a series of Lewis-acid catalysts for this sequence (see the Table S1 in the Supporting Information) we found that treatment of

epoxide **9** with  $\text{BF}_3 \cdot \text{OEt}_2$  or a Bi(III) salt generated tetrahydrofluorene **8** in good yield. The cascade promoted by  $\text{BF}_3 \cdot \text{OEt}_2$  occurred to form tetrahydrofluorene **8** in up to 75% yield; however, the yields of this reaction varied from 50–75%. Slightly lower yields were observed for the formation of tetrahydrofluorene **8** from epoxide **9** in the presence of Bi(III) salts as the Lewis acid, but this reaction was less sensitive to variations in temperature. Ultimately, we conducted this cascade promoted by 20 mol %  $\text{BiCl}_3$  to generate tetrahydrofluorene **8** in 64% yield.

The synthesis of (–)-taiwaniaquinone H (**2**) was completed according to methods developed by Trauner and coworkers for the synthesis of (±)-taiwaniaquinone H.<sup>16</sup> Selective monodemethylation of tetrahydrofluorene **8** with  $\text{BBr}_3 \cdot \text{SMe}_2$ , followed by oxidation of the resulting tetrahydrofluoren-1-ol in the presence of salcomine formed (–)-taiwaniaquinone H (**2**) in 51% yield over two steps (Scheme 4).

Tetrahydrofluorene **8** is also a suitable intermediate for the synthesis of (–)-taiwaniaquinol B (**7**). The enantioselective total synthesis of (–)-taiwaniaquinol B was completed in four steps from tetrahydrofluorene **8** (Scheme 5). Hydroboration of **8**, followed by oxidation of the resulting alkylborane in the presence of basic hydrogen peroxide formed a diastereomeric mixture of hexahydrofluoren-9-ols. Oxidation of these alcohols with 2-iodoxybenzoic acid (IBX) gave tetrahydrofluoren-9-one **17** as a single diastereomer in 80% yield. Tetrahydrofluoren-9-one **17** was converted to (–)-taiwaniaquinol B according to procedures reported by Fillion<sup>13</sup> and Trauner<sup>16</sup> for syntheses of racemic taiwaniaquinol B. Selective monodemethylation of **17** with  $\text{BCl}_3$ , oxidation of the resulting 8-hydroxyfluoren-9-one intermediate with ceric ammonium nitrate (CAN), and reductive workup with sodium dithionite gave (–)-taiwaniaquinol B in 52% yield over two steps.

In summary, an enantioselective total synthesis of (–)-taiwaniaquinone H was accomplished in 14.6% overall yield in just 10 steps, and the first enantioselective total synthesis of (–)-taiwaniaquinol B was completed in 11.9% overall yield in just 12 steps. Two catalytic protocols reported by our group, a one-pot meta bromination of 2-isopropyl-1,3-dimethoxybenzene by iridium-catalyzed borylation followed by copper-mediated bromination of the resulting arylboronate ester, and a palladium-catalyzed, enantioselective  $\alpha$ -arylation, were used in early steps on multi-mmol scales. The synthesis was completed efficiently by developing a Lewis acid-promoted cascade to construct the [6,5,6] tricyclic core of an intermediate common to the synthesis of a number of taiwaniaquinoids. Finally, we wish to emphasize that the quaternary stereogenic center in the taiwaniaquinoids bears no carbonyl group in the  $\alpha$ -position, but is readily set by the asymmetric  $\alpha$ -arylation, and the ketone functionality serves as a key point of reaction to complete the syntheses. Thus, the enantioselective  $\alpha$ -arylation of ketones should be considered as a strategic method to establish quaternary benzylic stereogenic centers for complex-molecule synthesis. Further studies to increase the scope of this enantioselective process are ongoing.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

We thank the NIH for financial support of this work (GM-58108 to J.F.H. and GM84584 to L.M.S.), Zhijian Liu for contributions to early steps in the synthesis, Johnson-Matthey for gifts of iridium and palladium salts, and Allychem and BASF for gifts of bis(pinacolato)diboron.


## References

1. Majetich G, Shimkus JM. *J. Nat. Prod* 2010;73:284. [PubMed: 20102168]
2. Lin WH, Fang JM, Cheng YS. *Phytochemistry* 1995;40:871.
3. Lin WH, Fang JM, Cheng YS. *Phytochemistry* 1996;42:1657.
4. Kawazoe K, Yamamoto M, Takaishi Y, Honda G, Fujita T, Sezik E, Yesilada E. *Phytochemistry* 1999;50:493.
5. Ohtsu H, Iwamoto M, Ohishi H, Matsunaga S, Tanaka R. *Tetrahedron Lett* 1999;40:6419.
6. Chang CI, Chien SC, Lee SM, Kuo YH. *Chem. Pharm. Bull* 2003;51:1420. [PubMed: 14646321]
7. Chang CI, Chang JY, Kuo CC, Pan WY, Kuo YH. *Planta Med* 2005;71:72. [PubMed: 15678377]
8. Iwamoto M, Ohtsu H, Tokuda H, Nishino H, Matsunaga S, Tanaka R. *Bioorg. Med. Chem* 2001;9:1911. [PubMed: 11425594]
9. Minami T, Iwamoto M, Ohtsu H, Ohishi H, Tanaka R, Yoshitake A. *Planta Med* 2002;68:742. [PubMed: 12221600]
10. Hanson JR. *Nat. Prod. Rep* 2004;21:312. [PubMed: 15042151]
11. Katoh T, Akagi T, Noguchi C, Kajimoto T, Node M, Tanaka R, Nishizawa M, Ohtsu H, Suzuki N, Saito K. *Bioorg. Med. Chem* 2007;15:2736. [PubMed: 17287126]
12. Banerjee M, Mukhopadhyay R, Achari B, Banerjee AK. *Org. Lett* 2003;5:3931. [PubMed: 14535746]
13. Fillion E, Fishlock D. *J. Am. Chem. Soc* 2005;127:13144. [PubMed: 16173732]
14. Banerjee M, Mukhopadhyay R, Achari B, Banerjee AK. *J. Org. Chem* 2006;71:2787. [PubMed: 16555833]
15. Planas L, Mogi M, Takita H, Kajimoto T, Node M. *J. Org. Chem* 2006;71:2896. [PubMed: 16555851]
16. Liang GX, Xu Y, Seiple IB, Trauner D. *J. Am. Chem. Soc* 2006;128:11022. [PubMed: 16925409]
17. McFadden RM, Stoltz BM. *J. Am. Chem. Soc* 2006;128:7738. [PubMed: 16771478]
18. Li SL, Chiu P. *Tetrahedron Lett* 2008;49:1741.
19. Tang SC, Xu YF, He JM, He YP, Zheng JY, Pan XF, She XG. *Org. Lett* 2008;10:1855. [PubMed: 18393520]
20. Majetich G, Shimkus JM. *Tetrahedron Lett* 2009;50:3311.
21. Alvarez-Manzaneda E, Chahboun R, Cabrera E, Alvarez E, Alvarez-Manzaneda R, Meneses R, Es-Samti H, Fernandez A. *J. Org. Chem* 2009;74:3384. [PubMed: 19348490]
22. Alvarez-Manzaneda E, Chahboun R, Cabrera E, Alvarez E, Haidour A, Ramos JM, Alvarez-Manzaneda R, Hmamouchi M, Es-Samti H. *Chem. Commun* 2009;592
23. Alvarez-Manzaneda E, Chahboun R, Cabrera E, Alvarez E, Haidour A, Ramos JM, Alvarez-Manzaneda R, Charrah Y, Es-Samti H. *Org. Biomol. Chem* 2009;7:5146. [PubMed: 20024110]
24. Node M, Ozeki M, Planas L, Nakano M, Takita H, Mori D, Tamatani S, Kajimoto T. *J. Org. Chem* 2010;75:190. [PubMed: 19968244]
25. Jana CK, Scopelliti R, Gademann K. *Chem. Eur. J* 2010;16:7692.
26. *Quaternary Stereocenters. Challenges and Solutions in Organic Synthesis.* Wiley-VCH: Weinheim; 2005.
27. Trost BM, Jiang CH. *Synthesis* 2006;369
28. Christoffers J, Baro A. *Adv. Synth. Catal* 2005;347:1473.
29. Douglas CJ, Overman LE. *Proc. Natl. Acad. Sci. U.S.A* 2004;101:5363. [PubMed: 14724294]
30. Alvarez-Manzaneda ECR, Alvarez E, Tapia R, Alvarez-Manzaneda R. *Chem. Commun* 2010:9244.
31. Liao XB, Weng ZQ, Hartwig JF. *J. Am. Chem. Soc* 2008;130:195. [PubMed: 18076166]
32. Murphy JM, Liao X, Hartwig JF. *J. Am. Chem. Soc* 2007;129:15434. [PubMed: 18027947]
33. Mkhaliid IAI, Barnard JH, Marder TB, Murphy JM, Hartwig JF. *Chem. Rev* 2010;890. [PubMed: 20028025]

34. Engler TA, Sampath U, Naganathan S, Velde DV, Takusagawa F, Yohannes D. *J. Org. Chem* 1989;54:5712.
35. Chieffi A, Kamikawa K, Ahman J, Fox JM, Buchwald SL. *Org. Lett* 2001;3:1897. [PubMed: 11405739]
36. Woodward RB, Sondheimer F, Taub D, Heusler K, McLamore WM. *J. Am. Chem. Soc* 1952;74:4223.
37. .
38. Corey EJ, Chaykovsky M. *Journal of the American Chemical Society* 1962;84:867.
39. Corey EJ, Chaykovsky M. *Journal of the American Chemical Society* 1965;87:1353.
40. Corey EJ, Chaykovsky M. *Journal of the American Chemical Society* 1962;84:3782.
41. Rickborn, B. *Comprehensive Organic Synthesis*. Trost, BM., editor. Vol. Vol. 3. Pergamon: Oxford; 1991. p. 733
42. Ranu BC, Jana U. *J. Org. Chem* 1998;63:8212.
43. Sudha R, Narasimhan KM, Saraswathy VG, Sankararaman S. *J. Org. Chem* 1996;61:1877. [PubMed: 11667069]
44. Maruoka K, Murase N, Bureau R, Ooi T, Yamamoto H. *Tetrahedron* 1994;50:3663.
45. House HO. *J. Am. Chem. Soc* 1955;77:3070.




**Figure 1.**  
Representative taiwaniaquinoids.




**Scheme 1.**  
Retrosynthetic Analysis for the Syntheses of (-)-Taiwaniaquinone H (**2**) and (-)-Taiwaniaquinol B (**7**)





**Scheme 2.****One-pot Synthesis of Aryl Bromide 12a**

<sup>a</sup> B<sub>2</sub>pin<sub>2</sub> = bis(pinacolato)diboron; dtpbpy = 4,4'-di-*tert*-butylbipyridine.



**Scheme 3.**  
Synthesis of Tetrahydrofluorene **8**

**Scheme 4.**Synthesis of (-)-Taiwaniaquinone H (**2**)<sup>a</sup><sup>a</sup> Salcomine = *N,N'*-bis(salicylidene)ethylenediaminocobalt (II).



**Scheme 5.**  
Synthesis of (-)-Taiwaniaquinol B (7)