



Published in final edited form as:

Addict Behav. 2010 July ; 35(7): 721–725. doi:10.1016/j.addbeh.2010.03.009.

Family History of Alcohol Abuse Associated With Problematic Drinking Among College Students

Joseph W. LaBrie, Ph.D.,

Loyola Marymount University; Associate Professor, Department of Psychology, Director, Heads UP, 1 LMU Drive, Suite 4700, Los Angeles, CA 90045. (310) 338-5238, jlabrie@lmu.edu.

Savannah Migliuri, B.A.,

Loyola Marymount University; Research Coordinator, Heads UP, Department of Psychology, 1 LMU Drive, Suite 4700, Los Angeles, CA 90045. (310) 338-3753, smigliur@lmu.edu.

Shannon R. Kenney, Ph.D., and

Loyola Marymount University; Postdoctoral Research Fellow, Heads UP, Department of Psychology, 1 LMU Drive, Suite 4700, Los Angeles, CA 90045. (310) 338-7770, Shannon.kenney@lmu.edu.

Andrew Lac, M.A., M.P.A.

Loyola Marymount University; Statistical Consultant, Heads UP, Department of Psychology, 1 LMU Drive, Suite 4700, Los Angeles, CA 90045. andrew.lac@cgu.edu.

Abstract

Studies examining family history of alcohol abuse among college students are not only conflicting, but have suffered various limitations. The current report investigates family history of alcohol abuse (FH+) and its relationship with alcohol expectancies, consumption, and consequences. In the current study, 3753 student participants (35% FH+), completed online assessments. Compared to FH–same-sex peers, FH+ males and FH+ females endorsed greater overall positive expectancies, consumed more drinks per week, and experienced more alcohol-related negative consequences. Further, FH+ females evaluated the negative effects of alcohol to be substantially worse than FH– females. An ANCOVA, controlling for age, GPA, race, and alcohol expectancies, resulted in family history main effects on both drinking and consequences. An interaction also emerged between gender and family history, such that FH+ males were especially vulnerable to high levels of alcohol consumption. Results reveal the scope of FH+ individuals in the college environment and the increased risk for these students, particularly male FH+ students, suggesting a need for researchers and college health personnel to focus attention and resources on this issue.

Keywords

family history; alcohol use; college students; gender differences; alcohol expectancies; consequences

© 2010 Elsevier Ltd. All rights reserved.

†Correspondence concerning this article should be sent to Joseph LaBrie via e-mail at: jlabrie@lmu.edu or to Loyola Marymount University, 1 LMU Drive, Suite 4700, Los Angeles, CA 90045-0041, tel: 310-338-5238, fax: 310-338-7726.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

I. Introduction

Risky drinking among college students is of particular concern for university administrators and health professionals. Researchers have attempted to isolate correlates of risky drinking. A family history of alcohol abuse (FH+) is a well-documented risk factor for heavy alcohol use and alcohol-related problems (Chalder, Elgar, & Bennett, 2006; Cotton, 1979; Hussong, Curran, & Chassin, 1998; Kuntsche, 2004; Pullen, 1994; Turnbull, 1994; Warner, White, & Johnson., 2007). About 20% of college students are FH+ (Perkins, 2002) and the college environment may be more harmful for those students predisposed to alcohol problems. A few studies have revealed considerably higher rates of alcohol use (Kushner & Sher, 1993; LaBrie et al., 2009; Pullen, 1994) and alcohol-related problems (Leeman, Fenton, & Volpicelli, 2004) among FH+ compared with FH- college students. In contrast, other studies have found no relationship between family history and problematic alcohol use among college students (Engs, 1990; MacDonald, Fleming, & Barry, 1991; Harrell, Slane, & Klump, 2009). Further, there have been conflicting results on the role gender plays among FH+ college students. Some have found FH+ males to be more susceptible to risky drinking and consequences than FH+ females (e.g. Andersson et al., 2007; Jackson et al., 2001; Sher, et al., 1991), while Hartford, Parker and Grant (1992) found no such gender difference. Inconsistencies in existing research highlight the need to explicate how family history status may impact drinking behaviors and problems in collegiate populations.

Alcohol expectancies, the specific beliefs about the behavioral, emotional, and cognitive effects of alcohol (Leigh, 1987), are a potential psychosocial motivator of risky drinking. Stronger positive alcohol expectancies are associated with problem drinking (e.g. Anderson et al., 2005; Brown, Goldman, & Christiansen, 1985). Alcohol-outcome expectancies result from both personal experience with alcohol and from mirroring drinking behavior of individuals (Lundahl et al, 1997), and have thus been shown to differ by family history status in that FH+ individuals have endorsed stronger alcohol-related expectancies, particularly overall positive expectancies (Morean et al, 2009; Pastor & Evans, 2003). Further, FH+ individuals with stronger overall positive expectancies are most likely to experience alcohol-related problems (Conway, Swendsen, & Merikangas, 2003; VanVoorst & Quirk, 2003).

Much of the previous research on family history of alcohol abuse has focused on COAs (children of alcoholics) during adolescence (Barnow et al., 2002; Brown, Creamer, & Stetson, 1987; Chalder et al., 2006; Nash, McQueen, & Bray, 2005; Sher, 1991) and middle-late adulthood (Beaudoin et al., 1997; Cloninger, Sigvardsson, & Bohman, 1996; Curran, et al., 1999). Moreover, family history studies involving college students have suffered from various limitations, such as a relatively small sample size (e.g., Leeman, et al., 2004; Pullen, 1994), single-sex samples (e.g. LaBrie et al., 2009; Harrell, et al., 2009), or first-year student samples (e.g. Andersson et al., 2007, Gotham, Sher, & Wood, 2003; Jackson et al., 2001). The present study broadens previous research by offering unique insight into family history of alcohol abuse, alcohol-related behaviors and problems, and further examines the moderating effect of gender in family history status on alcohol consumption, alcohol expectancies, and alcohol-related consequences among a large, multi-site, ethnically diverse sample of male and female college students.

2. Methods

2.1 Participants

Participants were recruited from two west-coast universities, a large, public institution with 30,000 undergraduates and a mid-sized private institution with approximately 5,500

undergraduates. Of a randomly selected pool of 7,000 students, 3,753 (53.6%) consented to participate. Representative of the makeup of the corresponding institutions, participants' mean age was 19.88 ($SD = 1.36$) and the majority of the participants were female (61%). The sample consisted of 18.9% first-year students, 24.5% sophomores, 27.4% juniors, and 29.2% seniors. Racial representation was as follows: 57.4% Caucasian, 18.7% Asian, 10.7% Multiracial, 3.2% African American, and 10.0% reported other racial/ethnic groups. On average, participants consumed 6.04 ($SD = 8.58$) drinks over 1.59 ($SD = 1.53$) drinking days per week. Among the 67.5% of students who drank, they consumed an average of 8.94 ($SD = 9.11$) drinks per week and averaged 2.36 ($SD = 1.30$) drinking days.

2.2. Design and Procedure

At the start of the fall semester, 7,000 students (3,500 from each campus), received letters inviting them to participate in a study about alcohol use and perceptions of college-student drinking. The students were directed to a link for an online survey. After students clicked on the link and entered their individual pin, they were presented with a local IRB-approved consent form. Participants then completed a 20 minute survey, for which they received a \$20 compensation.

2.3 Measures

Demographics—Participants indicated their gender, age, most recent GPA, and race.

Family History—Participants indicated whether they had a biological relative that “has or has had a significant drinking problem—one that should or did lead to treatment.” This measure was previously developed and successfully used by Miller & Marlatt (1984).

Alcohol Consumption—The Daily Drinking Questionnaire (DDQ; Collins, Parks, & Marlatt, 1985; Kivlahan, Marlatt, Fromme, & Coppel, 1990) asked students to report, from the past 30 days, the typical number of drinks they consumed each day of the week. Responses were summed to form a total drinks per week variable used in this analysis.

Negative Consequences—The 25-item Rutgers Alcohol Problem Index (RAPI, White & Labouvie, 1989) ($\alpha = .925$) assessed alcohol-related consequences. Using a 0 (*never*) to 4 (*more than 10 times*) scale, participants indicated how many times in the past three months they had experienced each stated circumstance (e.g., “Caused shame or embarrassment to someone,” “Passed out or fainted suddenly,” or “Felt that you had a problem with school.”).

Alcohol Expectancies and Evaluations—The Comprehensive Effects of Alcohol (CEOA; Fromme, Stroot, & Kaplan, 1993) is a two-part questionnaire consisting of 76 items. In Part 1, representing items tapping expectancies, participants indicated expectations concerning how he or she may act or feel under the influence of alcohol (e.g., “I would enjoy sex more,” “I would act sociable”; 1 = “disagree” 4 = “agree”). In Part 2, representing evaluations, participants subjectively evaluated the effects of alcohol with the same 38 items as Part 1 of the questionnaire (e.g., “Enjoying sex more,” “Feeling sociable”; 1 = “bad” 3 = “neutral” 5 = “good”). Each of the expectancies and evaluations components may be further divided into positive factors (sociability, tension reduction, liquid courage, and sexuality) and negative factors (cognitive behavioral impairment, risk and aggression, and self-perception).

3. Results

A family history of alcohol abuse was reported by 35.0% of the total sample, and FH+ participants were more likely to have drunk in the past year than their FH- peers (81% vs.

74%; $\chi^2 = 9.63, p < .001$). Independent samples t-tests, separately conducted for males and females, revealed several systematic differences between FH+ and FH- respondents (Table 1). Among males, FH+ respondents averaged significantly higher than their FH- counterparts on drinks per week, negative consequences, overall positive expectancies, positive expectancies concerned with tension reduction and liquid courage, as well as positive evaluations concerned with tension reduction. Among females, FH+ respondents reported significantly higher drinks per week, negative consequences, overall positive expectancies, as well as positive expectancies concerned with sociability, tension reduction, and sexuality in comparison to the FH- participants. Typically, FH+ females reported negative evaluations (risk and aggression, and self-perception) to be worse than did FH- females.

Additional analyses show that, among males, FH+ participants drank 45.7% more drinks per week and experienced 43.6% more negative consequences than those classified as FH-. Among females, however, FH+ individuals consumed 14.4% more drinks and experienced 23.6% more negative consequences than their FH- counterparts. Such results, taken together, suggest that a family history of alcohol abuse may adversely impact males more than females in the college environment.

An ANCOVA model, controlling for age, GPA, race, overall positive and negative expectancies, and overall positive and negative evaluations, was performed to predict drinks per week. Family history status (FH+ or FH-) and respondent gender (male or female) served as the independent factors. After ruling out the statistical contribution of the covariates, main effects were found for both family history and gender, and their interaction also emerged (Table 2). This statistical interaction, presented in Figure 1, revealed that the difference between FH+ and FH- on drinking was more pronounced in males than females, and that FH+ males were especially vulnerable to higher levels of alcohol consumption.

A second ANCOVA model was conducted to predict alcohol negative consequences. Age, GPA, race, and drinks per week were entered as covariates, and family history and gender served as the independent variables. After the variance attributed to the covariates were accounted for in the model, family history remained statistically significant, but no gender main effect or interaction was discovered (Table 2).

4. Discussion

The present investigation uses a large multisite sample and corroborates extant literature by identifying family history as a significant risk factor for alcohol misuse and related consequences among male and female college students (Kushner & Sher, 1993; Leeman, et al., 2004; LaBrie et al., 2009; Pullen, 1994). More specifically, this study extends previous research by finding that, whether attributable to genetics or environmental upbringing, familial ties to alcoholism were considerably more hazardous for males than females in regard to excessive alcohol consumption. Compared to FH- same-sex peers, FH+ males drank 41% more drinks per week and FH+ females drank 14% more drinks per week. Notably, results covaried out other important predictors of drinking (e.g. age, GPA, race) to better assess how FH status and gender may be related to drinking in college, over and above such variables. By highlighting family history positive college students' heightened susceptibility to risky drinking and consequences, and male FH+ students' enhanced risk for alcohol misuse, the current results may help college personnel identify and target prevention efforts to at-risk students. Preventative interventions taking place early in college with FH+ students might help them better understand their heightened alcohol-related vulnerabilities and provide them with tools and motivation to reduce potential harm.

In addition, findings both confirm and extend relevant research examining the role that alcohol expectancies play in FH+ college students' alcohol behaviors and outcomes. Not only did students reporting familial alcohol abuse endorse significantly greater overall positive expectancies than same-sex FH- counterparts, but FH+ female respondents evaluated the negative effects of alcohol to be substantially "more bad" than FH- females. This paradoxical finding, in which women exposed to familial alcohol abuse judged alcohol's negative evaluations to be worse, yet were more likely to agree that drinking personally yielded positive effects (i.e., expectancies) may suggest that FH+ women may not equate their own drinking with that of alcoholic family members and thus may feel immune to the negative evaluations they themselves associate with alcohol. More concerning, however, is the possibility that these findings may be indicative of cognitive dissonance, whereby highly endorsed positive expectancies contribute to continued drinking, often heavy drinking, in students even though they have been exposed to, and thus recognize, the negative aspects of drinking. By rationalizing alcohol misuse through heightened expectancies FH+ college females may be able to reduce dissonance and fulfill strong, possibly genetically predisposed desires to imbibe. Regardless, FH+ students' apparent awareness of the negative effects of alcohol use through their own familial experience may be a promising avenue for intervening. Intervention with these students should allow them to reflect on and be mindful of their experiences with these negative effects thereby building motivation to avoid these same consequences while challenging positive alcohol expectancies.

The present findings are limited in that they do not account for environmental risk factors known to co-occur with FH+ status (e.g., histories of physical or sexual abuse or attraction to high-risk student groups) and that may confound the relationship between FH+ status and both alcohol expectancies and misuse. Future studies assessing such risk factors may be warranted. Another limitation of the current study is the use of one, nonspecific classification of FH+ status. Future research may benefit from distinguishing first, second, and third degree affiliation to alcohol abuse (e.g. parent vs. grandparent or aunt/uncle), gender of the relative with alcohol problems (e.g. mother vs. father), or familial history density (i.e., whether an individual has more than one family member with an alcohol problem). A more defined classification of FH status may also reveal environmental risk factors; for instance, the extent to which residential exposure to alcoholism may heighten risk.

The current study reveals that FH+ students make up a substantial percentage (35%) of the college population and that these students are at increased risk for problematic drinking and consequences as compared to their FH- peers. Despite this and previous research in concert with the current findings, preventative interventions targeting FH+ students are lacking. Both researchers and college health personnel may wish to invest resources in targeting these individuals.

Acknowledgments

Support for this research was provided by National Institute of Alcohol Abuse and Alcoholism grant R01-AA012547

References

- Ackerman RJ, Gondolf EW. Adult children of alcoholics: The effects of background and treatment on ACOA symptoms. *International Journal of the Addictions* 1991;26:1159–1172. [PubMed: 1743816]
- Anderson KG, Schweinsburg A, Paulus M, Brown SA, Tapert S. Examining personality and alcohol expectancies using Functional Magnetic Resonance Imaging (fMRI) with adolescents. *Journal of Studies on Alcohol* 2005;66:323–331. [PubMed: 16047521]

- Andersson C, Johnsson K, Berglund M, Öjehagen A. Alcohol involvement in Swedish university freshmen related to gender, age, serious relationship and family history of alcohol problems. *Alcohol and Alcoholism* 2007;42(5):448–455. [PubMed: 17360719]
- Barnow S, Schuckit M, Lucht M, John U, Freyberger HJ. The importance of positive family history of alcoholism, parental rejection and emotional warmth, behavioral problems and peer substance use for alcohol problems in teenagers: A path analysis. *Journal of Studies on Alcohol* 2002;63:305–315. [PubMed: 12086131]
- Beaudoin CM, Murray RP, Bond J JR, Barnes GE. Personality characteristics of depressed or alcoholic adult children of alcoholics. *Personality and Individual Differences* 1997;23:559–567.
- Brown SA, Creamer VA, Stetson BA. Adolescent alcohol expectancies in relation to personal and parental drinking patterns. *Journal of Abnormal Psychology* 1987;96:117–121. [PubMed: 3584659]
- Brown S, Goldman M, Christiansen B. Do alcohol expectancies mediate drinking patterns of adults? *Journal of Consulting and Clinical Psychology* 1985;53(4):512–519. [PubMed: 4031207]
- Chalder M, Elgar FJ, Bennett P. Drinking and motivations to drink among adolescent children of parents with alcohol problems. *Alcohol & Alcoholism* 2006;41(1):107–113. [PubMed: 16239352]
- Cloninger CR, Sigvardsson S, Bohman M. Type I and Type II Alcoholism: An update. *Alcohol Health and Research World* 1996;20:18–23.
- Collins RL, Parks GA, Marlett A. Social determinants of alcohol consumption: The effects of social interaction and model status on self-administration of alcohol. *Journal of Consulting and Clinical Psychology* 1985;53(2):189–200. [PubMed: 3998247]
- Conway KP, Swendsen JD, Merikangas KR. Alcohol expectancies, alcohol consumption, and problem drinking. The moderating role of family history. *Addictive Behaviors* 2003;28:823–836. [PubMed: 12788259]
- Cotton NS. The familial incidence of alcoholism: A review. *Journal of Studies on Alcohol* 1979;40(1):89–116. [PubMed: 376949]
- Curran GM, Stoltenberg SF, Hill EM, Mudd SA, Blow FC, Zucker RA. Gender differences in the relationship among SES, family history of alcohol disorders and alcohol dependence. *Journal of Studies on Alcohol* 1999;60:825–832. [PubMed: 10606495]
- Engs R. Family background of alcohol abuse and its relationship to alcohol consumption among college students: An unexpected finding. *Journal of Studies on Alcohol* 1990;51(6):542–547. [PubMed: 2270063]
- Fromme K, Stroot E, Kaplan D. Comprehensive effects of alcohol: Development and psychometric assessment of a new expectancy questionnaire. *Psychological Assessment* 1993;5(1):19–26.
- Gotham HJ, Sher KJ, Wood PK. Alcohol involvement and development task completion during young adulthood. *Journal of Studies on Alcohol* 2003;64:32–42. [PubMed: 12608481]
- Harford T, Parker D, Grant B. Family history, alcohol use and dependence symptoms among young adults in the United States. *Alcoholism: Clinical and Experimental Research* 1992;16(6):1042–1046.
- Harrell Z, Slane J, Klump K. Predictors of alcohol problems in college women: The role of depressive symptoms, disordered eating, and family history of alcoholism. *Addictive Behaviors* 2009;34(3):252–257. [PubMed: 19027241]
- Hussong AM, Curran PJ, Chassin L. Pathways of risk for accelerated heavy alcohol use among adolescent children of alcohol parents. *Journal of Abnormal Child Psychology* 1998;26(6):453. [PubMed: 9915652]
- Kivlahan DR, Marlatt GA, Fromme K, Coppel DB. Secondary prevention with college drinkers: Evaluation of an alcohol skills training program. *Journal of Consulting and Clinical Psychology* 1990;58:805–810. [PubMed: 2292630]
- Kuntsche E, Rehm J, Gmel G. Characteristics of binge drinkers in Europe. *Social Science & Medicine* 59:113–127.
- Kushner MG, Sher KJ. Comorbidity of alcohol and anxiety disorders among college students: Effects of gender and family history of alcoholism. *Addictive Behaviors* 1993;18:543–552. [PubMed: 8310873]
- Jackson KM, Sher KJ, Gotham HJ, Wood PK. Transition into and out of large-effect drinking in young adulthood. *Journal of Abnormal Psychology* 2001;110(3):378–391. [PubMed: 11502081]

- LaBrie JW, Kenney SR, Lac A, Migliuri SM. Differential drinking patterns of family history positive and family history negative first semester college females. *Addictive Behaviors* 2009;34:190–196. [PubMed: 18992994]
- Leeman RF, Miriam F, Volpicelli JR. Impaired control and undergraduate problem drinking. *Alcohol & Alcoholism* 2007;42(1):42–48. [PubMed: 17142826]
- Leigh BC. Evaluations of alcohol expectancies: Do they add to prediction of drinking patterns? *Psychology of Addictive Behaviors* 1987;1(3):135–139.
- Lundahl L, Davis T, Adesso V, Lukas S. Alcohol expectancies: Effects of gender, age, and family history of alcoholism. *Addictive Behaviors* 1997;22(1):115–125. [PubMed: 9022877]
- MacDonald R, Fleming M, Barry K. Risk factors associated with alcohol abuse in college students. *American Journal of Drug and Alcohol Abuse* 1991;17(4):439–449. [PubMed: 1746506]
- Miller, WR.; Marlatt, GA. Brief drinking profile. Odessa, FL: Psychological Assessment Resources; 1984.
- Morean ME, Corbin WR, Sinha R, O'Malley SS. Parental history of anxiety and alcohol-use disorders and alcohol expectancies as predictors of alcohol-related problems. *Journal of Studies on Alcohol and Drugs* 2009;70:227–236. [PubMed: 19261234]
- Nash SG, McQueen A, Bray JH. Pathways to adolescent alcohol use: Family environment, peer influence, and parental expectations. *Journal of Adolescent Health* 2005;37:19–28. [PubMed: 15963903]
- Pastor AD, Evans SM. Alcohol outcome expectancies and risk for alcohol use problems in women with and without a family history of alcoholism. *Drug and Alcohol Dependence* 2003;70:201–214. [PubMed: 12732414]
- Perkins HW. Surveying the damage: A review of research on consequences of alcohol misuse in college populations. *Journal of Studies on Alcohol* 2002;14:91–100.
- Pullen LM. The relationship among alcohol abuse in college students and selected psychological/demographic variables. *Journal of Alcohol and Drug Education* 1994;40(1):36–50.
- Sher KJ, Walitzer KS, Wood PK, Brent EE. Characteristics of children of alcoholics: Putative risk factors, substance use and abuse, and psychopathology. *Journal of Abnormal Psychology* 1991;100:427–448. [PubMed: 1757657]
- Turnbull JE. Early background variables as predictors of adult alcohol problems in women. *International Journal of the Addictions* 1994;29:707–728. [PubMed: 8034381]
- VanVoorst WA, Quirk SW. Are relations between parental history of alcohol problems and changes in drinking moderated by positive expectancies? *Alcoholism: Clinical and Experimental Research* 2003;26:25–30.
- White H, Labouvie E. Towards the assessment of adolescent problem drinking. *Journal of Studies on Alcohol* 1989;50(1):30–37. [PubMed: 2927120]
- Warner LA, White HR, Johnson V. Alcohol initiation experiences and family history of alcoholism as predictors of problem-drinking trajectories. *Journal of Studies on Alcohol and Drugs* 2007;68:56–65. [PubMed: 17149518]




Figure 1.
Family history status \times gender interaction on drinks per week.

Table 1
 Mean Differences on Drinking Variables by Family History, for Males and Females

Measure	Males				Females				t-test
	FH+		FH-		FH+		FH-		
	M	(SD)	M	(SD)	M	(SD)	M	(SD)	
Drinks Per Week	10.75	(12.83)	7.38	(10.43)	4.94	(6.04)	4.32	(5.68)	2.48*
Negative Consequences	6.72	(9.10)	4.68	(8.77)	4.55	(6.64)	3.68	(6.03)	3.11**
Overall Positive Expectancies	2.55	(0.55)	2.47	(0.58)	2.42	(0.58)	2.35	(0.59)	2.73**
Sociability	2.94	(0.68)	2.88	(0.70)	2.96	(0.73)	2.87	(0.73)	2.74**
Tension Reduction	2.63	(0.71)	2.53	(0.74)	2.32	(0.72)	2.24	(0.72)	2.53*
Liquid Courage	2.48	(0.69)	2.39	(0.72)	2.29	(0.71)	2.25	(0.73)	1.18
Sexuality	2.13	(0.73)	2.09	(0.73)	2.12	(0.76)	2.05	(0.75)	2.11*
Overall Negative Expectancies	2.28	(0.52)	2.28	(0.59)	2.20	(0.56)	2.20	(0.58)	0.21
Cognitive Behavioral Imp.	2.64	(0.60)	2.63	(0.65)	2.64	(0.65)	2.62	(0.68)	0.73
Risk and Aggression	2.31	(0.71)	2.26	(0.73)	2.10	(0.74)	2.11	(0.74)	-0.40
Self-Perception	1.89	(0.64)	1.95	(0.72)	1.87	(0.69)	1.87	(0.69)	0.25
Overall Positive Evaluations	3.48	(0.84)	3.40	(0.83)	3.18	(0.85)	3.19	(0.86)	-0.39
Sociability	3.84	(0.93)	3.80	(0.92)	3.66	(0.94)	3.65	(0.95)	0.27
Tension Reduction	3.82	(1.00)	3.67	(0.97)	3.41	(0.99)	3.40	(1.00)	0.26
Liquid Courage	2.95	(0.91)	2.90	(0.90)	2.72	(0.90)	2.79	(0.91)	-1.79
Sexuality	3.32	(1.07)	3.21	(1.02)	2.92	(1.08)	2.92	(1.05)	-0.20
Overall Negative Evaluations	1.89	(0.57)	1.92	(0.62)	1.68	(0.50)	1.76	(0.55)	-3.45***
Cognitive Behavioral Imp.	1.82	(0.64)	1.82	(0.68)	1.62	(0.56)	1.63	(0.58)	-0.19
Risk and Aggression	2.13	(0.83)	2.15	(0.84)	1.92	(0.76)	2.07	(0.81)	-4.25***
Self-Perception	1.71	(0.64)	1.79	(0.71)	1.49	(0.55)	1.58	(0.61)	-3.49***

* $p < .05$.** $p < .01$.

 $p < .001$

NIH-PA Author Manuscript

NIH-PA Author Manuscript

NIH-PA Author Manuscript

Table 2

ANCOVA Models Predicting Drinks Per Week and Negative Consequences

Variable	<i>df</i>	<i>MS</i>	<i>F test</i>
DV: Drinks Per Week			
Covariates			
Age	1	137.51	2.29
GPA	1	209.84	3.49
Race	1	5207.62	86.57***
Overall Positive Expectancies	1	10846.35	108.31***
Overall Negative Expectancies	1	6536.75	108.67***
Overall Positive Evaluations	1	6.15	0.10
Overall Negative Evaluations	1	2833.10	47.10***
Family History	1	1032.70	17.17***
Gender	1	8485.34	141.06***
Family History × Gender	1	1025.75	17.05***
DV: Negative Consequences			
Covariates			
Age	1	95.33	2.34
GPA	1	452.91	11.09***
Race	1	423.66	10.38**
Drinks Per Week	1	43439.92	1063.97***
Family History	1	460.67	11.28***
Gender	1	130.98	3.21
Family History × Gender	1	2.71	0.07

Note. Race (1 = Caucasian, 0 = non-Caucasian)

* $p < .05$.

** $p < .01$.

*** $p < .001$.