

20. **Thomas CD**, Feik SA, Clement JG. Regional variation of intracortical porosity in the midshaft of the human femur: age and sex differences. *J Anat* 2005;**206**:115–25.
21. **Goldman HM**, Thomas CD, Clement JG, *et al*. Relationships among microstructural properties of bone at the human midshaft femur. *J Anat* 2005;**206**:127–39.
22. **Drapeau MS**, Streeter MA. Modeling and remodeling responses to normal loading in the human lower limb. *Am J Phys Anthropol* 2006;**129**:403–9.
23. **Skedros JG**, Mason MW, Bloebaum RD. Differences in osteonal micromorphology between tensile and compressive cortices of a bending skeletal system: indications of potential strain-specific differences in bone microstructure. *Anat Rec* 1994;**239**:405–13.
24. **Nonaka K**, Fukuda S, Aoki K, *et al*. Regional distinctions in cortical bone mineral density measured by pQCT can predict alterations in material property at the tibial diaphysis of the cynomolgus monkey. *Bone* 2006;**38**:265–72.
25. **Lai YM**, Qin L, Hung VW, *et al*. Regional differences in cortical bone mineral density in the weight-bearing long bone shaft – a pQCT study. *Bone* 2005;**36**:465–71.
26. **Roldan EJ**, Capiglioni R, Cointy CR, *et al*. Postmenopausal changes in the distribution of the volumetric BMD of cortical bone. A pQCT study of the human leg. *J Musculoskelet Neuronal Interact* 2001;**2**:157–62.
27. **Mackelvie KJ**, McKay HA, Khan KM, *et al*. Lifestyle risk factors for osteoporosis in Asian and Caucasian girls. *Med Sci Sports Exerc* 2001;**33**:1818–24.
28. **Kontulainen SA**, Macdonald HM, Khan KM, *et al*. Examining bone surfaces across puberty: a 20-month pQCT trial. *J Bone Miner Res* 2005;**20**:1202–7.
29. **Crocker PR**, Bailey DA, Faulkner RA, *et al*. Measuring general levels of physical activity: preliminary evidence for the physical activity questionnaire for older children. *Med Sci Sports Exerc* 1997;**29**:1344–9.
30. **Kowalski KC**, Crocker PR, Faulkner RA. Validation of the physical activity questionnaire for older children. *Pediatr Exerc Sci* 1997;**9**:174–86.
31. **Baxter-Jones AD**, Mirwald RL, McKay HA, *et al*. A longitudinal analysis of sex differences in bone mineral accrual in healthy 8-19-year-old boys and girls. *Ann Hum Biol* 2003;**30**:160–75.
32. **Peterman MM**, Hamel AJ, Cavanagh PR, *et al*. In vitro modeling of human tibial strains during exercise in micro-gravity. *J Biomech* 2001;**34**:693–8.
33. **Sonoda N**, Chosa E, Totoribe K, *et al*. Biomechanical analysis for stress fractures of the anterior middle third of the tibia in athletes: nonlinear analysis using a three-dimensional finite element method. *J Orthop Sci* 2003;**8**:505–13.
34. **Binkley TL**, Specker BL, Wittig TA. Centile curves for bone densitometry measurements in healthy males and females ages 5–22 yr. *J Clin Densitom* 2002;**5**:343–53.
35. **Lorentzon M**, Mellstrom D, Ohlsson C. Age of attainment of peak bone mass is site specific in Swedish men – the GOOD study. *J Bone Miner Res* 2005;**20**:1223–7.
36. **Wang Q**, Nicholson PH, Suuriniemi M, *et al*. Relationship of sex hormones to bone geometric properties and mineral density in early pubertal girls. *J Clin Endocrinol Metab* 2004;**89**:1698–703.
37. **Moyer-Mileur L**, Xie B, Ball S, *et al*. Predictors of bone mass by peripheral quantitative computed tomography in early adolescent girls. *J Clin Densitom* 2001;**4**:313–23.
38. **Wang Q**, Alen M, Nicholson P, *et al*. Growth patterns at distal radius and tibial shaft in pubertal girls: a 2-year longitudinal study. *J Bone Miner Res* 2005;**20**:954–61.
39. **Lorentzon M**, Mellstrom D, Ohlsson C. Association of amount of physical activity with cortical bone size and trabecular volumetric BMD in young adult men: the GOOD study. *J Bone Miner Res* 2005;**20**:1936–43.
40. **Nikander R**, Sievanen H, Uusi-Rasi K, *et al*. Loading modalities and bone structures at nonweight-bearing upper extremity and weight-bearing lower extremity: a pQCT study of adult female athletes. *Bone* 2006;**39**:886–94.
41. **Heinonen A**, Sievanen H, Kyrolainen H, *et al*. Mineral mass, size, and estimated mechanical strength of triple jumpers' lower limb. *Bone* 2001;**29**:279–85.
42. **Heinonen A**, Sievanen H, Kannus P, *et al*. Site-specific skeletal response to long-term weight training seems to be attributable to principal loading modality: a pQCT study of female weightlifters. *Calcif Tissue Int* 2002;**70**:469–74.
43. **Haapasalo H**, Kontulainen S, Sievanen H, *et al*. Exercise-induced bone gain is due to enlargement in bone size without a change in volumetric bone density: a peripheral quantitative computed tomography study of the upper arms of male tennis players. *Bone* 2000;**27**:351–7.
44. **Kontulainen S**, Sievanen H, Kannus P, *et al*. Effect of long-term impact-loading on mass, size, and estimated strength of humerus and radius of female racquet-sports players: a peripheral quantitative computed tomography study between young and old starters and controls. *J Bone Miner Res* 2002;**17**:2281–9.
45. **Specker B**, Binkley T, Fahrenwald N. Rural versus nonrural differences in BMC, volumetric BMD, and bone size: a population-based cross-sectional study. *Bone* 2004;**35**:1389–98.
46. **Hangartner TN**. Thresholding technique for accurate analysis of density and geometry in QCT, pQCT and microCT images. *J Musculoskelet Neuronal Interact* 2007;**7**:9–16.
47. **Ward KA**, Adams JE, Hangartner TN. Recommendations for thresholds for cortical bone geometry and density measurement by peripheral quantitative computed tomography. *Calcif Tissue Int* 2005;**77**:275–80.
48. **Bousson V**, Bergot C, Meunier A, *et al*. CT of the middiaphyseal femur: cortical bone mineral density and relation to porosity. *Radiology* 2000;**217**:179–87.

CORRECTION

doi: 10.1136/bjsm.2007.037945corr1

There was an error in the article by Guerrero *et al* published in the July issue of the journal (Guerrero M, Guiu-Comadevall M, Cadefau JA, *et al*. Fast and slow myosins as markers of muscle injury. *Br J Sports Med* 2008;**42**:581–4). Table 1 was omitted from the article. The table is reproduced online at <http://bjsm.bmj.com/supplemental/>.



CORRECTION

Br J Sports Med 2008 42: 695
doi: 10.1136/bjsem.2007.037945corr1

Updated information and services can be found at:
<http://bjsm.bmj.com/content/42/8/695.full.html>

	<i>These include:</i>
Data Supplement	<i>"web only table"</i> http://bjsm.bmj.com/content/suppl/2008/07/21/42.8.695.DC1.html
Email alerting service	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>