

Published in final edited form as:

*Cancer Epidemiol Biomarkers Prev.* 2010 May ; 19(5): 1301–1310. doi:10.1158/1055-9965.EPI-10-0002.

## Meat mutagens and breast cancer in postmenopausal women -a cohort analysis

Kana Wu<sup>1</sup>, Rashmi Sinha<sup>4</sup>, Michelle D. Holmes<sup>2</sup>, Edward Giovannucci<sup>1,2,3</sup>, Walter Willett<sup>1,2,3</sup>, and Eunyong Cho<sup>1,2</sup>

<sup>1</sup> Department of Nutrition, Harvard School of Public Health, Boston, MA

<sup>2</sup> Channing Laboratory, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

<sup>3</sup> Department of Epidemiology, Harvard School of Public Health, Boston, MA

<sup>4</sup> Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, Rockville, MD

### Abstract

**Background**—Mutagenic compounds produced when meats are cooked at high temperatures have been hypothesized to increase risk of breast cancer.

**Methods**—We examined the association between intakes of the heterocyclic amines (HCAs) MeIQx (2-amino-3,8-dimethylimidazo (4,5,-f) quinoxaline), PhIP (2-amino-1-methyl-6-phenylimidazo (4,5-b) pyridine), DiMeIQx (2-amino-3,4,8-trimethylimidazo (4,5,-f)) and meat-derived mutagenic activity (MDM) and risk of breast cancer using a cooking method questionnaire administered in 1996 in the Nurses Health Study. Between 1996 and 2006, 2,317 breast cancer cases were diagnosed during 533,618 person years.

**Results**—Higher intake of HCAs or MDM was not associated with elevated risk of breast cancer (multivariate relative risk (RR) and 95% confidence interval (95% CI) for the highest vs. lowest quintile: MeIQx: 0.90 (0.79–1.03); PhIP: 0.92 (0.80–1.05); DiMeIQx: 0.92 (0.80–1.05) and MDM: 0.98 (0.85–1.12)). HCA or MDM were not associated with estrogen receptor positive/progesterone positive breast cancer risk either. There was some suggestion of a decreased risk of estrogen receptor negative/progesterone receptor negative breast cancer with higher intakes of MeIQx, DiMeIQx and PhIP, but none of the associations were statistically significant. There was little evidence for an interaction between intake of cruciferous vegetables and HCA or MDM intake and risk of breast cancer.

**Conclusion**—Higher consumption of mutagens from meats cooked at higher temperature and longer duration was not associated with increased risk of postmenopausal breast cancer.

**Impact**—Overall prospective data including results from our study do not provide support for a substantial increase in risk of breast cancer with higher intake of HCAs.

### Keywords

Cooking methods; heterocyclic amines; meat; breast cancer

## Introduction

Cooking meats at high temperatures and for long duration can result in the production of mutagenic compounds, such as heterocyclic amines (HCAs) (1). Findings from animal studies have long suggested a role of HCAs in breast carcinogenesis (2,3) and there is some evidence from epidemiological studies that higher intakes of well-done meats may raise breast cancer risk (4–6). However, only a limited number of epidemiological studies, including three recent prospective studies, have examined the associations between HCAs and breast cancer risk (5,7–13). In two prospective studies no evidence for an association between HCA intake and risk of postmenopausal breast cancer was observed (10,11) whereas another recent prospective study (13) reported a significant positive association between higher intake of MeIQx and a marginally significant positive association between higher intake DiMeIQx and risk of postmenopausal breast cancer. Considering the paucity of prospective data on the association between HCAs and breast cancer, we examined the association between intakes of major HCAs including MeIQx (2-amino-3,8-dimethylimidazo [4,5-f] quinoxaline), PhIP (2-amino-1-methyl-6-phenylimidazo [4,5-b] pyridine), DiMeIQx (2-amino-3,4,8-trimethylimidazo [4,5-f]) and meat-derived mutagenic activity (MDM) and risk of breast cancer using a cooking method questionnaire administered in 1996 in the Nurses Health Study (NHS), a large female cohort. The large number of cases allowed us to examine associations between HCAs and breast cancer risk by hormone receptor status as some recent evidence suggests that certain risk factors for breast cancer including red meat intake may differ by hormone receptor status (14,15). Only a limited number of epidemiological studies have examined associations between HCA intake and risk of breast cancer separately by hormone receptor status (5,11,13).

## Materials and Methods

### Study Population

More detailed information on the NHS cohort is provided elsewhere (16). In brief, the NHS cohort was founded in 1976 when 121,700 U.S. female nurses aged between 30–55 years were sent a questionnaire to obtain information on their lifestyle factors and medical histories. Follow-up questionnaires were sent every two years, and in 1980, 1984, 1986 and every four years thereafter semiquantitative food frequency questionnaires (FFQs) were added to the follow-up questionnaires. In 1996, a cooking method questionnaire containing questions on cooking methods and doneness of several meats was added to the questionnaire. The study was approved by the Human Subjects Committee of the Brigham and Women's Hospital.

### Cooking method questionnaire

All questions included in the 1996 cooking method questionnaire were based on results from a pilot study (17) designed to ascertain a set of questions that would best predict the intake of HCAs in the NHS. On this cooking method questionnaire participants were asked about frequency of intake (i.e. never, < 1/month, 1/month, 2–3/month, 1/week, 2–3/week, and 4+/week) and doneness (i.e. depending on type of meat: lightly browned, medium browned, well browned and blackened/charred) of cooked meats and fish (i.e. pan-fried, broiled and grilled chicken, broiled fish, roast beef, pan-fried steak, grilled or barbecued steak, and homemade beef gravy). In addition, participants were also asked whether they cooked and/or ate chicken with or without skin (“when you eat chicken, how often is it cooked with the skin on: always, most of the time, sometimes and never”, and “how often do you eat the skin: always, most of the time, sometimes and never”, for more information also refer to [www.charred.cancer.gov/nhs.html](http://www.charred.cancer.gov/nhs.html)). We estimated intake of pan-fried hamburgers by multiplying the frequency of intake of hamburger from the 1994 FFQ (also see below) with

the proportion of participants from the NHS in the pilot study that had reported eating hamburgers pan-fried (i.e. 0.33) (17), assuming the outside appearance to be medium brown (based on the median value for doneness obtained from the pilot study). Since only limited variability with regard to doneness levels of pan-fried bacon was observed in our pilot study, we estimated intake of pan-fried bacon using bacon intake from the 1994 FFQ and assumed bacon to be prepared at high degree of doneness (i.e. well-done) (17).

### **Assessment of Diet, Meat and HCA Intake**

The validity and reproducibility of the FFQs administered in the NHS cohort have been reported elsewhere (18,19). Since we did not request information on frequency of total meat intake on the 1996 cooking questionnaires (i.e. total meat intake that does not take into account specific cooking methods), we estimated total red, white and processed meat intake using information from the 1994–2002 FFQs and calculated a cumulative average intake of total red, white and processed meats. Cumulative average intake of meats represents the average intake of meats from all available FFQs up to the start of each follow-up interval (20).

For participants who had reported frequency of cooked meat intake on the cooking method questionnaire but did not report on the outside appearance of meat we imputed the median value for doneness, i.e. lightly brown for broiled fish, medium brown for the other cooked meat items. Participants who reported eating cooked chicken but who left the “cook/eat chicken with skin” sections blank were assigned to the “chicken not cooked with skin” category.

HCAs values were calculated using an online database the “Charred Database” (21,22) and (23) created by Sinha and colleagues from the National Cancer Institute that provides users with data on HCA and MDM from measured meat sample extracts. Briefly, the mutagenic activity of meat samples was assessed by the Ames/Salmonella test (24,25) and MeIQx, DiMeIQx and PhIP were determined using a method previously reported by Gross and Gruter (26). Intake of MeIQx, PhIP, DiMeIQx and MDM were calculated by multiplying the HCA or MDM values from the Charred Database (ng/gram meat for HCA and revertant colonies/gram meat for MDM) with standard portion sizes from the Charred Database and frequency of intake of the relevant cooked meat item obtained from the 1996 cooking questionnaire. HCA values for roast beef were 0 for all doneness categories and HCA values for broiled fish were not available from the “Charred Database” (21) and thus did not contribute to our calculations for HCAs and MDM intake. For more detailed information regarding HCA and MDM values used in this study could be found in the website for “Charred Database”. (21,22).

### **Ascertainment of Breast Cancer Cases**

Whenever a participant reported a breast cancer diagnosis during the past two years on a biennial follow-up questionnaire, we contacted them and requested their consent to obtain and review their medical records pertaining to this specific diagnosis. Investigators reviewed those medical records to a) confirm a breast cancer diagnosis and to b) extract information relevant to breast cancer including histology of breast cancer. In addition, we were able to extract information on hormone receptor status (estrogen receptor and progesterone receptor) from the pathology reports. After exclusion of ineligible participants (for exclusion criteria see below) a total of 2,317 breast cancer cases (diagnosed between 1996 and 2006) were included in our analysis. Of the 1,779 cases with information on hormone receptor status, 1,174 were estrogen receptor positive and progesterone receptor positive (ER+/PR+) cases, 295 were estrogen receptor negative and progesterone negative cases (ER-/PR-), 279

cases were ER+/PR- and 31 cases were ER-/PR+. Breast cancer cases with a carcinoma in situ histology were excluded from our analyses.

### Exclusion Criteria

Participants were ineligible for this study if they a) had reported a history of any cancer (except for non-melanoma skin cancer) prior to 1996, b) were premenopausal in 1996 because we had few premenopausal cases, c) had calculated energy intakes of <600 or >3,500 kcal per day or had left more than 70 food items on the FFQs blank and d) had left the entire cooking method section on the 1996 questionnaire blank. In addition, we also excluded participants for whom HCA/MDM could not be calculated e.g. due to missing information on bacon or hamburger intake on the 1994 FFQ or had reported information on doneness of meat but not on frequency of cooked meat intake. Thus our study population consisted of 54,440 women who were followed between 1996 and 2006 and contributed a total of 533,618 person years of follow up.

### Statistical Analysis

For every participant we calculated person-years of follow up from the date of return of the 1996 follow-up questionnaire to the end of our follow-up period (June 2006), death, date of breast cancer or other cancer diagnosis (except for non-melanoma skin cancer), whichever occurred first. Intake of HCA, MDM as well as red meat, white meat and processed meat intake were divided into quintiles. Frequency of intakes and doneness of cooked meat items were divided into relevant categories (Intake: pan-fried chicken: never, <1/mo, 1/mo, 2-3/mo, >=1/wk, broiled chicken: never, <1/mo, 1/mo, 2-3/mo, 1/wk, >=2-3/wk, grilled/barbequed chicken: never, <1/mo, 1/mo, 2-3/mo, >=1/wk, broiled fish: never, <1/mo, 1/mo, 2-3/mo, 1/wk, >=2-3/wk, roast beef: never, 1/mo, 2-3/mo, >=1/wk, pan-fried steak: never, <1/mo, 1/mo, >=2-3/mo, grilled/barbequed chicken: never, <1/mo, 1/mo, 2-3/mo, >=1/wk, homemade gravy: never, <1/mo, 1/mo, 2-3/mo, >=1/wk). To assess associations between quintiles of meat mutagen intake, quintiles/categories of total and cooked meat intake and risk of breast cancer we used a Cox Proportional Hazards Model. Because results obtained from the age-adjusted models were similar to those from the multivariate models, only multivariate RRs are presented. The multivariate models included known and suspected risk factors for breast cancer i.e., age in months (continuous), smoking status (never, past, current 1-14 cigarettes/day, current 15-24 cigarettes/day, current >=25 cigarettes/day), BMI (kg/m<sup>2</sup>: <25, 25-<30, >=30), height (inches <63, 63-<64, 64-<66, >=66), physical activity (hours per week: <1, 1-<2, 2-<4, 4-<7, >=7), age at menarche (years <=12, 13, >=14), family history of breast cancer (yes, no), history of benign breast disease (yes, no), parity and age at first birth (nulliparous, 1-2 children and age at first birth <25 years, 1-2 children and age at first birth 25-<30 years, 1-2 children and age at first birth >=30 years, 3-4 children and age at first birth <25 years, 3-4 children and age at first birth 25-<30 years, 3-4 children and age at first birth >=30 years, 5-8 children and age at first birth <25 years, 5-8 children and age at first birth 25-<30 years), postmenopausal hormone use (never, current, past), weight change 1996-2006 (<-2 kg, -2 to +2 kg, +2.1 kg to +5 kg, +5.1 kg to +10 kg, +10.1 to +20 kg, +20.1 kg to +25 kg, >=+25.1 kg), total energy and alcohol intake (continuous).

Trend test were calculated by adding the median of each quintile or category of the exposure variable as a continuous variable to the model. Associations were also examined separately by hormone receptor status i.e., ER+/PR+ and ER-/PR- breast cancers and after stratification by age (<65, >=65 years), body mass index (<25, >=25 kg/m<sup>2</sup>), smoking status (never, past, current) and postmenopausal hormone use (never, current, past). As there is some evidence that the mutagenic effects of HCAs may be modified by certain compounds found in fruits and vegetables such as chlorophyllin and isothiocyanates (27-29)

associations were also examined after stratification by intakes of cruciferous vegetables and total fruits and vegetables (<5.25 servings/day, ≥5.25 servings/day). To obtain P-values for interaction we added a cross term product consisting of the meat mutagen (medians of quintiles) and the exposure variable (as a binary variable) to the multivariate model and performed a Wald test. All p-values were two sided.

## Results

Baseline characteristics pertaining to intake of meat mutagens (lowest and highest quintile) are shown in Table 1. Participants with higher intake of meat mutagens were more likely to have a higher BMI as well as higher intake of calories, fat, alcohol, white meat, processed meat and red meat. Furthermore, women with higher MeIQx intake appeared to be less active than those with lower MeIQx intake. Participants with high and low meat mutagen intake did not appear to differ with regard to most reproductive factors, except that a slightly higher percentage of women with higher meat mutagen intake had had at least 3 children.

The amount of HCAs or MDM consumed by each participant varied according to the specific type of meat as well as the cooking method and the outside appearance of the cooked meat (Table 2). The top three contributors to PhIP intake were grilled chicken, grilled steak and broiled chicken. Pan-fried hamburgers, pan-fried bacon and pan-fried steak contributed most to MeIQx intake and grilled chicken, pan-fried hamburgers and pan-fried bacon contributed most to DiMeIQx intake. On the other hand, broiled and grilled chicken as well as grilled steak contributed most to MDM intake. Correlations between each individual HCAs were  $r=0.60$  (MeIQx vs. PhIP),  $r=0.83$  (DiMeIQx vs. MeIQx) and  $r=0.68$  (DiMeIQx vs. PhIP).

Higher intakes of red, white and processed meats, HCAs and MDM were not associated with increased risk of total breast cancer (Table 3) (multivariate relative risk (RR) and 95% confidence interval (95% CI) for the highest vs. lowest quintile: MeIQx: 0.90 (0.79–1.03); PhIP: 0.92 (0.80–1.05); DiMeIQx: 0.92 (0.80–1.05) and MDM: 0.98 (0.85–1.12)). When we examined associations between meat mutagens and breast cancer separately by hormone receptor status (Table 4), intakes of HCAs, MDM and PMH were not associated with risk of ER+/PR+ breast cancer. There was some suggestion of a lower risk of estrogen receptor negative and progesterone receptor negative (ER-/PR-) breast cancer with higher intakes of MeIQx, DiMeIQx and PhIP, but none of the associations were statistically significant (RR and 95% CI for the highest vs. lowest quintile: MeIQx: 0.79 (0.54–1.15), p value, test for trend (p) =0.23; PhIP: 0.73 (0.50–1.08), p=0.12; DiMeIQx: RR=0.74 (0.50–1.10), p=0.06). MDM, intake was not associated with risk of ER-/PR- breast cancer. Intakes of total red, white, and processed meat were not significantly associated with risk of breast cancer regardless of hormone receptor status.

We also investigated whether intakes of cooked meats that contributed to intake of HCAs (from the 1996 cooking method questionnaire) were associated with total breast cancer risk. In general, intakes of cooked meats and gravy (pan-fried, grilled chicken, broiled fish, roast beef, pan-fried steak, grilled or barbecued steak, and homemade beef gravy) did not appear to be associated with total breast cancer (data not shown). However, there was some evidence for a slightly and marginally increased risk of total breast cancer with higher intake of broiled chicken (highest vs. lowest category: RR=1.13 (95% CI=0.97–1.31); ptrend=0.05) and some evidence for a non-significant inverse association between higher intake of barbecued/grilled steak and breast cancer (highest vs. lowest category: RR=0.84 (95% CI=0.69–1.02; ptrend=0.12).

As there is some evidence that the mutagenic effects of HCAs may be modified by certain compounds found in fruits and vegetables such as chlorophyllin and isothiocyanates (27–29) we examined associations after stratification by intake of cruciferous vegetables. However, we found little evidence for interaction between intake of cruciferous vegetables and HCA or MDM intake and risk of total breast cancer (data not shown). When we examined associations between meat mutagens and breast cancer risk by fruit and vegetable intake (Table 5), there was some suggestion of an inverse association between MDM intake and total breast cancer among those with high fruit/vegetable intake whereas participants in the low fruit/vegetable category appeared to have a slightly increased risk of total breast cancer, however none of the observed associations reached statistical significance ( $p$ -interaction=0.07) (Table 5). We also examined associations after stratification by smoking status (never, past, current) and postmenopausal hormone use (never, past, current). Greater intake of MeIQx but not PhIP, DiMeIQx or MDM intake was associated with a significantly decreased risk of total breast cancer among current smokers and current postmenopausal hormone users (highest vs. lowest quintile: current smokers: RR 0.59, 95% CI: 0.36–0.99); current hormone users: 0.79 (95% CI: 0.64–0.97), whereas no associations were seen for past/never smokers or past/never hormone users (data not shown). Associations between HCAs and MDM and total breast cancer were not modified by age, BMI or physical activity. Furthermore, associations between HCA/MDM intake and total breast cancer were similar to those reported in table 3 when cases diagnosed within the first two years of follow-up were excluded.

## Discussion

Results from this large prospective cohort study do not support a positive association between higher intakes of meat mutagens and risk of breast cancer among postmenopausal women. There was some suggestion of a lower risk of estrogen receptor negative and progesterone receptor negative (ER–/PR–) breast cancer with higher intakes of HCAs, but none of the associations were statistically significant. Data from animal studies have long suggested an enhancing role of HCAs in mammary carcinogenesis (2,3). In addition to their mutagenic and carcinogenic properties (30) recent animal data also suggest that PhIP, the most abundant HCA in human diet, may also possess estrogenic activity (31) and can promote the secretion of prolactin (32). Endogenous hormones especially estrogen and progesterone and more recently also prolactin have been hypothesized to play a role in breast carcinogenesis and findings from both epidemiological as well as animal studies have provided compelling support for this hypothesis (33).

However, only a limited number of epidemiological studies including three recent prospective studies have examined the associations between HCA intake and breast cancer risk (5,7–13)

Here you repeated many things once you mentioned in the first sentence (for example, prospective, postmenopausal etc - so I deleted all)

Consistent with our results two of the three recent prospective studies have found no evidence for an association between HCA intake and risk of postmenopausal breast cancer (10,11). The AARP cohort using the Charred Database (11) included 3,818 breast cancer after 8 years of follow up. No associations between red, white, processed meat intake as well as meat cooked at high temperature, HCA and total mutagenic activity and risk of breast cancer were found. Results were similar after examining associations separately by hormone receptor status or by fruit and vegetable intake. No evidence for an association between HCAs and risk of breast cancer were found in a Swedish study including 430 cases with a mean follow-up of 10.4 years. That study used a HCA database developed in Sweden (10).

Contrary to the two studies and our study another recent study (13) in the Prostate, Lung, Colorectal and Ovarian Cancer Screening Trial (PLCO) reported a weak positive association between higher intake of red meat and MeIQx and risk of breast cancer (comparing highest vs. lowest quintile of intake: red meat: RR=1.23 (95% CI=1.00–1.51, ptrend=0.22); MeIQx: RR=1.26 (95% CI: 1.03–1.55); p trend=0.12).. In addition, the positive associations between red meat and breast cancer were more pronounced among ER+/PR+ breast cancer. This study also used the CHARRED database to calculate HCA intake. “Reasons for the inconsistent findings between the 4 prospective studies are unclear but one possible reason may be possible variations in genetic polymorphisms of enzymes involved in metabolism of HCAs (34,35) or timing of exposure (also see below under limitations).” I would delete this part also because you summarize below and the results aren't that inconsistent (3 null and 1 weak positive)”

Again, I deleted many things repeated below (e.g., case-control, significantly etc) Findings from case-control studies are inconsistent; and contrary to the hypothesized mechanisms, some case-control studies even observed inverse associations between breast cancer and some individual HCAs. For example, one study with 114 cases with benign breast disease found inverse association between higher PhIP intake and risk of breast cancer which the authors attributed to higher white meat intake which was also inversely associated with breast cancer risk (8). However MeIQx and DiMeIQx intake was not associated with breast cancer risk. (8) (could delete this sentence). In another study by Steck et al., (5) recent HCA intake was not associated with risk of postmenopausal breast cancer, but higher intake of MeIQx and DiMeIQx was associated with an approximately 40% decreased risk of premenopausal breast cancer. In addition, higher lifetime intakes of grilled and smoked meats were also associated with increased risk of postmenopausal breast cancer especially among those with low fruit and vegetable intake but were not associated with premenopausal breast cancer risk. In one hospital based study on 352 cases from Uruguay both MeIQx and PhIP (DiMeIQx was not examined in that study) were associated with increased risk of postmenopausal but not premenopausal breast cancer (7). In a nested case-control study of 273 cases in the Iowa Women's Health Study, a large cohort of postmenopausal women, PhIP, but not MeIQx and DiMeIQx, was associated with increased risk of breast cancer (9). In summary, while case-control studies show inconsistent results, two of the three prospective studies conducted so far have found no evidence for an association between HCA and/or MDM intake and breast cancer (10,11). In the third prospective study, there was some evidence for a statistically significant increased risk with higher intake of MeIQx intake and some suggestion of a non-significantly increased risk with higher DiMeIQx intake (13).

Contrary to the hypothesized mechanisms we found some suggestion of a lower risk of estrogen receptor negative and progesterone receptor negative (ER-/PR-) breast cancer with higher intakes of DiMeIQx and PhIP, but none of the associations were statistically significant. Reasons for these findings are unclear. Besides chance one possible reason could be that higher HCA intake may be a marker for higher intake of white meat as PhIP levels in cooked chicken are quite high ([www.charred.cancer.gov](http://www.charred.cancer.gov)), however in our study neither overall white meat intake nor PhIP from white meat were associated with ER-/PR- breast cancer.

One limitation of our study is that we did not examine genetic polymorphisms of enzymes involved in metabolism of HCAs. HCAs are not mutagenic compounds *per se* but need to undergo metabolism to exert their potential mutagenic effects. On the other hand HCAs can also be deactivated via detoxification pathways (34,35). Xenobiotic metabolic enzymes involved in activation or deactivation of HCAs include cytochrome P450 (CYP), N-acetyltransferase (NAT), sulfotransferase (SULT) and glutathione S-transferase (GST)

(34,35). Fruits and vegetables containing isothiocyanates which can induce Phase I and II metabolism enzymes may also affect metabolism of HCAs and modify the effect of HCAs (28,29). In human feeding studies substantial inter-individual variation with regard to urinary excretion of metabolites of MeIQx and PhIP have been observed, indicating considerable inter-individual variation with regard to the metabolism of HCAs (36,37). Intra-individual variation with regard to genetic polymorphisms of xenobiotic metabolic enzymes may in part explain the lack of positive associations between HCA and MDM intake and risk of breast cancer in our study. However, so far findings from epidemiological studies that have assessed possible interactions between NAT1 and NAT2 genotypes and well done meats and/or HCA intake with regard to breast cancer risk are inconsistent (8,12,38,39).

There are also some other limitations inherent to our study design. First, we cannot rule out misclassification of exposure, as we do not know how well HCA intake estimated from our cooking method questionnaires correlates with true exposure of these mutagens at the breast tissue levels. In addition, estimation of HCA intake was based on a limited number of cooking method questions. However, the aforementioned set of questions were developed based on findings from a previous pilot study conducted to establish the group of questions that could optimally predict the intake of HCAs in the NHS (17). Furthermore, we did not take into account possible modifications to cooking methods that may affect the actual amount of HCAs ingested. For example, flipping meat during the cooking process (40), microwaving (41) or marinating (42) meat prior to cooking have been shown to modify the amount of HCAs produced. Secondly, exposure was assessed at one time point only, i.e. in 1996, which may not reflect changes in cooking methods over time. However, it is very unlikely that changes in cooking methods due to underlying disease could have influenced observed associations considerably because exclusion of cases diagnosed within the first 2 years of follow-up yielded similar results for total breast cancer. Thirdly, participants in this study were only followed for up to 10 years which may not be long enough to see an effect of HCAs on breast cancer risk. Fourthly, we did not examine premenopausal breast cancer as an outcome. In a recent study using data from the Nurses Health Study II (NHS II), a large cohort of younger female nurses residing in the U.S., higher red meat intake was significantly associated with elevated risk of ER+/PR+ premenopausal breast cancer (14). These findings suggest that red meat intake as a risk factor may either differ by menopausal and/or hormone receptor status or exposure earlier in life may be the more relevant exposure. Supporting the latter hypothesis are studies that have shown that breast cancer initiation may occur earlier in life (43–45). In a recent study from the NHS II higher red meat intake during adolescence estimated using information from a validated high school FFQ was significantly associated with higher risk of premenopausal breast cancer and results were similar after adjusting for meat intake during adulthood (46). Therefore assessment of HCA intake in later stages of adulthood may not represent the relevant exposure which may in part explain the null associations between HCA intake and postmenopausal breast cancer observed in our and two other prospective studies (10,11).

In conclusion, in this large cohort study of postmenopausal women higher consumption of mutagens from meats cooked at higher temperature and longer duration was not associated with increased breast cancer risk. Our findings regarding HCAs and ER–/PR– breast cancer warrant further evaluation.

## Acknowledgments

Grants: This work was supported by a grant from the American Institute for Cancer Research as well as grant CA 55075 from the National Cancer Institute, National Institutes of Health



## References

1. Sugimura T, Wakabayashi K, Nakagama H, Nagao M. Heterocyclic amines: Mutagens/carcinogens produced during cooking of meat and fish. *Cancer Sci.* 2004; 95:290–9. [PubMed: 15072585]
2. Felton JS, Knize MG, Salmon CP, Malfatti MA, Kulp KS. Human exposure to heterocyclic amine food mutagens/carcinogens: relevance to breast cancer. *Environ Mol Mutagen.* 2002; 39:112–8. [PubMed: 11921178]
3. Snyderwine EG. Mammary gland carcinogenesis by food-derived heterocyclic amines: metabolism and additional factors influencing carcinogenesis by 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP). *Environ Mol Mutagen.* 2002; 39:165–70. [PubMed: 11921185]
4. Dai Q, Shu XO, Jin F, et al. Consumption of animal foods, cooking methods, and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev.* 2002; 11:801–8. [PubMed: 12223422]
5. Steck SE, Gaudet MM, Eng SM, et al. Cooked meat and risk of breast cancer--lifetime versus recent dietary intake. *Epidemiology.* 2007; 18:373–82. [PubMed: 17435448]
6. Zheng W, Gustafson DR, Sinha R, et al. Well-done meat intake and the risk of breast cancer. *J Natl Cancer Inst.* 1998; 90:1724–9. [PubMed: 9827527]
7. De Stefani E, Ronco A, Mendilaharsu M, Guidobono M, Deneo-Pellegrini H. Meat intake, heterocyclic amines, and risk of breast cancer: a case-control study in Uruguay. *Cancer Epidemiol Biomarkers Prev.* 1997; 6:573–81. [PubMed: 9264269]
8. Delfino RJ, Sinha R, Smith C, et al. Breast cancer, heterocyclic aromatic amines from meat and N-acetyltransferase 2 genotype. *Carcinogenesis.* 2000; 21:607–15. [PubMed: 10753193]
9. Sinha R, Gustafson DR, Kulldorff M, et al. 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine, a carcinogen in high-temperature-cooked meat, and breast cancer risk. *J Natl Cancer Inst.* 2000; 92:1352–4. [PubMed: 10944558]
10. Sonestedt E, Ericson U, Gullberg B, et al. Do both heterocyclic amines and omega-6 polyunsaturated fatty acids contribute to the incidence of breast cancer in postmenopausal women of the Malmo diet and cancer cohort? *Int J Cancer.* 2008; 123:1637–43. [PubMed: 18636564]
11. Kabat GC, Cross AJ, Park Y, et al. Meat intake and meat preparation in relation to risk of postmenopausal breast cancer in the NIH-AARP diet and health study. *Int J Cancer.* 2009; 124:2430–5. [PubMed: 19165862]
12. Mignone LI, Giovannucci E, Newcomb PA, et al. Meat consumption, heterocyclic amines, NAT2, and the risk of breast cancer. *Nutr Cancer.* 2009; 61:36–46. [PubMed: 19116874]
13. Ferrucci LM, Cross AJ, Graubard BI, et al. Intake of meat, meat mutagens, and iron and the risk of breast cancer in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Br J Cancer.* 2009; 101:178–84. [PubMed: 19513076]
14. Cho E, Chen WY, Hunter DJ, et al. Red meat intake and risk of breast cancer among premenopausal women. *Arch Intern Med.* 2006; 166:2253–9. [PubMed: 17101944]
15. Colditz GA, Rosner BA, Chen WY, Holmes MD, Hankinson SE. Risk factors for breast cancer according to estrogen and progesterone receptor status. *J Natl Cancer Inst.* 2004; 96:218–28. [PubMed: 14759989]
16. Colditz GA. The nurses' health study: a cohort of US women followed since 1976. *J Am Med Womens Assoc.* 1995; 50:40–4. [PubMed: 7722205]
17. Byrne C, Sinha R, Platz EA, et al. Predictors of dietary heterocyclic amine intake in three prospective cohorts. *Cancer Epidemiol Biomarkers Prev.* 1998; 7:523–9. [PubMed: 9641497]
18. Salvini S, Hunter DJ, Sampson L, et al. Food-based validation of a dietary questionnaire: the effects of week-to-week variation in food consumption. *Int J Epidemiol.* 1989; 18:858–67. [PubMed: 2621022]
19. Willett WC, Sampson L, Stampfer MJ, et al. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol.* 1985; 122:51–65. [PubMed: 4014201]
20. Willett, WC. Issues in analysis and presentation of dietary data. In: Willett, WC., editor. *Nutritional epidemiology.* New York: Oxford University Press; 1998. p. 321-345.
21. National Cancer Institute. CHARRED (Computerized Heterocyclic Amine Database Resource for Research in Epidemiologic Disease). <http://www.charredcancer.gov>

22. National Cancer Institute. CHARRED (Computerized Heterocyclic Amine Database Resource for Research in Epidemiologic Disease). <http://www.charredcancer.gov/nhshtml>
23. Sinha R. An epidemiologic approach to studying heterocyclic amines. *Mutat Res.* 2002; 506–507:197–204.
24. Ames BN, McCann J, Yamasaki E. Methods for detecting carcinogens and mutagens with the Salmonella/mammalian-microsome mutagenicity test. *Mutat Res.* 1975; 31:347–64. [PubMed: 768755]
25. Knize MG, Sinha R, Rothman N, et al. Heterocyclic amine content in fast-food meat products. *Food Chem Toxicol.* 1995; 33:545–51. [PubMed: 7628789]
26. Gross GA, Gruter A. Quantitation of mutagenic/carcinogenic heterocyclic aromatic amines in food products. *J Chromatogr.* 1992; 592:271–8. [PubMed: 1583097]
27. Dingley KH, Ubick EA, Chiarappa-Zucca ML, et al. Effect of dietary constituents with chemopreventive potential on adduct formation of a low dose of the heterocyclic amines PhIP and IQ and phase II hepatic enzymes. *Nutr Cancer.* 2003; 46:212–21. [PubMed: 14690798]
28. Knize MG, Kulp KS, Salmon CP, Keating GA, Felton JS. Factors affecting human heterocyclic amine intake and the metabolism of PhIP. *Mutat Res.* 2002; 506–507:153–62.
29. Steinkellner H, Rabot S, Freywald C, et al. Effects of cruciferous vegetables and their constituents on drug metabolizing enzymes involved in the bioactivation of DNA-reactive dietary carcinogens. *Mutat Res.* 2001; 480–481:285–97.
30. Sinha R, Rothman N. Exposure assessment of heterocyclic amines (HCAs) in epidemiologic studies. *Mutat Res.* 1997; 376:195–202. [PubMed: 9202756]
31. Lauber SN, Ali S, Gooderham NJ. The cooked food derived carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine is a potent oestrogen: a mechanistic basis for its tissue-specific carcinogenicity. *Carcinogenesis.* 2004; 25:2509–17. [PubMed: 15319301]
32. Lauber SN, Gooderham NJ. The cooked meat derived genotoxic carcinogen 2-amino-3-methylimidazo[4,5-b]pyridine has potent hormone-like activity: mechanistic support for a role in breast cancer. *Cancer Res.* 2007; 67:9597–602. [PubMed: 17909072]
33. Hankinson SE. Endogenous hormones and risk of breast cancer in postmenopausal women. *Breast Dis.* 2005; 24:3–15. [PubMed: 16917136]
34. Cross AJ, Sinha R. Meat-related mutagens/carcinogens in the etiology of colorectal cancer. *Environ Mol Mutagen.* 2004; 44:44–55. [PubMed: 15199546]
35. Turesky RJ. The role of genetic polymorphisms in metabolism of carcinogenic heterocyclic aromatic amines. *Curr Drug Metab.* 2004; 5:169–80. [PubMed: 15078194]
36. Stillwell WG, Sinha R, Tannenbaum SR. Excretion of the N(2)-glucuronide conjugate of 2-hydroxyamino-1-methyl-6-phenylimidazo[4,5-b]pyridine in urine and its relationship to CYP1A2 and NAT2 activity levels in humans. *Carcinogenesis.* 2002; 23:831–8. [PubMed: 12016157]
37. Stillwell WG, Turesky RJ, Sinha R, Tannenbaum SR. N-oxidative metabolism of 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx) in humans: excretion of the N2-glucuronide conjugate of 2-hydroxyamino-MeIQx in urine. *Cancer Res.* 1999; 59:5154–9. [PubMed: 10537291]
38. Zheng W, Deitz AC, Campbell DR, et al. N-acetyltransferase 1 genetic polymorphism, cigarette smoking, well-done meat intake, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev.* 1999; 8:233–9. [PubMed: 10090301]
39. Deitz AC, Zheng W, Leff MA, et al. N-Acetyltransferase-2 genetic polymorphism, well-done meat intake, and breast cancer risk among postmenopausal women. *Cancer Epidemiol Biomarkers Prev.* 2000; 9:905–10. [PubMed: 11008907]
40. Tran NL, Salmon CP, Knize MG, Colvin ME. Experimental and simulation studies of heat flow and heterocyclic amine mutagen/carcinogen formation in pan-fried meat patties. *Food Chem Toxicol.* 2002; 40:673–84. [PubMed: 11955674]
41. Felton JS, Fultz E, Dolbear FA, Knize MG. Effect of microwave pretreatment on heterocyclic aromatic amine mutagens/carcinogens in fried beef patties. *Food Chem Toxicol.* 1994; 32:897–903. [PubMed: 7959444]
42. Salmon CP, Knize MG, Felton JS. Effects of marinating on heterocyclic amine carcinogen formation in grilled chicken. *Food Chem Toxicol.* 1997; 35:433–41. [PubMed: 9216741]

43. Ariazi JL, Haag JD, Lindstrom MJ, Gould MN. Mammary glands of sexually immature rats are more susceptible than those of mature rats to the carcinogenic, lethal, and mutagenic effects of N-nitroso-N-methylurea. *Mol Carcinog.* 2005; 43:155–64. [PubMed: 15924352]
44. Colditz GA, Frazier AL. Models of breast cancer show that risk is set by events of early life: prevention efforts must shift focus. *Cancer Epidemiol Biomarkers Prev.* 1995; 4:567–71. [PubMed: 7549816]
45. Russo IH, Russo J. Role of hormones in mammary cancer initiation and progression. *J Mammary Gland Biol Neoplasia.* 1998; 3:49–61. [PubMed: 10819504]
46. Linos E, Willett WC, Cho E, Colditz G, Frazier LA. Red meat consumption during adolescence among premenopausal women and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev.* 2008; 17:2146–51. [PubMed: 18669582]

Table 1

Baseline characteristics in the Nurses Health Study population by lowest and highest quintiles of heterocyclic amine (HCA) and meat derived mutagenic activity (MDM) intake

	PhIP (ng/day)		MeIQx (ng/day)		DiMeIQx (ng/day)		Meat derived mutagenic activity (revertant colonies/day)	
	Q1 (14.3)	Q5 (201)	Q1 (3.4)	Q5 (39.1)	Q1 (0.27)	Q5 (5.5)	Q1 (834)	Q5 (7372)
Mean age (yrs)	64.7	61.3	63.4	62.8	64.0	62.3	64.2	62.1
Mean pack years smoking among smokers	24.0	25.9	22.1	29.0	22.9	27.6	23.5	26.8
Mean age at first birth	22.8	22.8	23.0	22.8	23.0	22.9	22.9	22.8
Age at menarche <12 y (%)	49.1	49.2	50.0	47.8	50.0	48.4	49.4	49.0
Parity, >=3 children (%)	58.5	61.8	57.5	62.7	57.1	62.7	58.6	61.1
Height (inches)	64.5	64.5	64.5	64.5	64.5	64.5	64.5	64.5
BMI (kg/m <sup>2</sup> )	25.7	27.1	25.3	27.4	25.5	27.2	25.5	27.1
Physical activity (METs)	18.8	18.3	20.9	15.9	20.1	17.0	19.2	17.8
Family history of breast cancer in mother/sisters (%)	14.8	14.2	15.1	14.0	15.0	14.4	15.0	14.4
History of benign breast disease (%)	31.4	30.7	33.2	29.3	32.7	29.0	32.1	30.1
Mean daily intake <sup>†</sup> (94 FFQ)								
Calories (kcal)	1602	1813	1526	1911	1541	1858	1557	1845
Total fat (gm)	48.4	62.0	42.4	71.5	44.0	67.0	46.3	64.4
Total iron (mg)	20.5	20.7	20.8	20.6	20.6	20.4	20.1	20.8
Folate (mcg)	488	485	505	466	501	472	486	483
Alcohol (gm)	3.8	6.1	4.2	5.4	4.2	5.5	4.3	5.3
Processed meat (serv/day)	0.07	0.10	0.04	0.14	0.05	0.12	0.06	0.11
Chicken +turkey (serv/day)	0.32	0.44	0.36	0.38	0.36	0.40	0.31	0.43
Red meat (serv/day)	0.11	0.18	0.05	0.27	0.06	0.24	0.08	0.21

\* Standardized for age in 1996 except for age and age at first birth; Q1 = lowest quintile, Q5 = highest quintile, number in parenthesis denote median value in each quintile (ng/day for heterocyclic amines and revertant colonies/day for meat derived mutagenic activity)

<sup>†</sup> Mean daily intakes of nutrients are energy-adjusted

<sup>‡</sup> If not noted otherwise numbers denote mean values

Abbreviations:

MeIQx=2-amino-3,8-dimethylimidazo [4,5-f] quinoxaline;

DiMeIQx=2-amino-3,4,8-trimethylimidazo [4,5-f];

PhIP=2-amino-1-methyl-6-phenylimidazo [4,5-b] pyridine

BMI=Body mass index

MET=Metabolic equivalent hours per week

**Table 2**  
Mean intake of HCA and MDM by meat type in women in the Nurses Health Study

	PhIP			MeIQx			DiMeIQx			Meat derived mutagenic activity		
	Mean (SD) (ng/day)	Percentage (%)	Percentage (%)	Mean (SD) (ng/day)	Percentage (%)	Percentage (%)	Mean (SD) (ng/day)	Percentage (%)	Percentage (%)	Mean (SD) (revertants/day)	Percentage (%)	Percentage (%)
All sources	92.1 (94.5)	100	100	18.5 (19.3)	100	100	2.4 (2.7)	100	100	3604 (3355)	100	100
Pan-fried chicken	6.1 (14.7)	6.6	4.5	0.8 (1.9)	4.5	1.0	0.02 (0.10)	1.0	1.0	473 (1170)	13.0	13.0
Broiled chicken	17.8 (30.7)	19.3	1.9	0.4 (0.7)	1.9	0.1	0.003 (0.11)	0.1	0.1	764 (1347)	21.2	21.2
Grilled chicken	37.0 (62.5)	40.2	5.5	1.0 (2.0)	5.5	34.4	0.84 (1.74)	34.4	34.4	687 (1096)	19.1	19.1
Grilled steak	21.2 (37.8)	23.0	13.9	2.6 (4.2)	13.9	0	0	0	0	476 (990)	13.2	13.2
Pan-fried steak	5.0 (17.3)	5.4	19.8	3.7 (10.9)	19.8	12.9	0.32 (1.14)	12.9	12.9	372 (1746)	10.3	10.3
Homemade beef gravy	0.84 (3.4)	0.9	10.8	2.0 (5.8)	10.8	9.0	0.22 (0.90)	9.0	9.0	214 (668)	5.9	5.9
Pan-fried bacon	4.2 (8.6)	4.6	20.5	3.8 (7.7)	20.5	18.1	0.44 (0.90)	18.1	18.1	234 (477)	6.5	6.5
Pan-fried hamburger	0	0	23.0	4.3 (4.2)	23.0	24.5	0.60 (0.59)	24.5	24.5	385 (381)	10.7	10.7

Abbreviations:

SD=Standard deviation

MeIQx=2-amino-3,8-dimethylimidazo [4,5-f] quinoxaline;

DiMeIQx=2-amino-3,4,8-trimethylimidazo [4,5-f];

PhIP=2-amino-1-methyl-6-phenylimidazo [4,5-b] pyridine



NIH-PA Author Manuscript

NIH-PA Author Manuscript

NIH-PA Author Manuscript

children and age at first birth 25-30 years, postmenopausal hormone use (never, current, past), weight change 1996-2006 (<-2 kg, -2 to +2 kg, +2.1 kg to +5 kg, +5.1 kg to +10 kg, +10.1 to +20 kg, +20.1 kg to +25 kg, >=+25.1 kg), total calorie and alcohol intake (continuous)



Relative risk of breast cancer by combinations of receptor status and quintiles of HCA, MDM and meat intake in women in the Nurses Health Study (1996–2006)

Table 4

	Quintile HCA Intake (ng/day)/Mutagenic Activity (revertants/day)					P trend
	1	2	3	4	5	
<b>Estrogen positive/Progesterone positive</b>						
<b>Total red meat (1994–2002 FFQ)</b>						
Cases	281	174	244	237	238	
Multivariate RR	1	0.89 (0.73–1.08)	0.91 (0.76–1.08)	0.92 (0.77–1.10)	0.91 (0.75–1.10)	0.46
<b>Total white meat (1994–2002 FFQ)</b>						
Cases	295	168	256	233	222	
Multivariate RR	1	1.22 (1.01–1.49)	1.08 (0.91–1.28)	0.90 (0.76–1.08)	0.97 (0.81–1.17)	0.15
<b>Processed meat (1994–2002 FFQ)</b>						
Cases/Person-years	237	239	194	237	267	
Multivariate RR <sup>†</sup>	1	1.03 (0.86–1.24)	0.90 (0.74–1.09)	0.94 (0.78–1.13)	1.06 (0.87–1.28)	0.54
<b>Total MeIQx</b>						
Cases	242	219	226	244	243	
Multivariate RR <sup>†</sup>	1	0.88 (0.73–1.06)	0.89 (0.74–1.08)	0.97 (0.81–1.17)	0.93 (0.77–1.12)	0.88
<b>Total PhIP</b>						
Cases	247	223	228	252	224	
Multivariate RR <sup>†</sup>	1	0.90 (0.75–1.08)	0.92 (0.76–1.10)	1.01 (0.84–1.21)	0.89 (0.74–1.08)	0.58
<b>Total DiMeIQx</b>						
Cases	242	211	234	238	249	
Multivariate RR <sup>†</sup>	1	0.84 (0.70–1.01)	0.94 (0.79–1.13)	0.96 (0.80–1.15)	0.98 (0.82–1.18)	0.56
<b>Meat derived mutagenic activity</b>						
Cases	221	235	244	247	227	
Multivariate RR <sup>†</sup>	1	1.05 (0.87–1.26)	1.09 (0.91–1.31)	1.10 (0.91–1.33)	0.99 (0.82–1.20)	0.77
<b>Estrogen negative/Progesterone negative</b>						
<b>Total red meat (1994–2002 FFQ)</b>						
Cases	68	46	62	57	62	
Multivariate RR	1	0.99 (0.68–1.45)	1.00 (0.71–1.43)	1.02 (0.71–1.47)	1.05 (0.72–1.53)	0.79

		Quintile HCA Intake (ng/day)/Mutagenic Activity (revertants/day)					P trend
		1	2	3	4	5	
<b>Total white meat (1994–2002 FFQ)</b>							
Cases		85	31	59	66	54	
Multivariate RR	†	1	0.77 (0.50–1.17)	0.93 (0.66–1.31)	0.86 (0.62–1.20)	0.88 (0.61–1.26)	0.53
<b>Processed meat (1994–2002 FFQ)</b>							
Cases		65	56	60	52	62	
Multivariate RR	†	1	0.86 (0.60–1.24)	0.99 (0.69–1.42)	0.77 (0.53–1.12)	0.90 (0.61–1.31)	0.61
<b>Total MeIQx</b>							
Cases		67	63	56	54	55	
Multivariate RR	†	1	0.91 (0.64–1.29)	0.83 (0.58–1.19)	0.78 (0.54–1.13)	0.79 (0.54–1.15)	0.23
<b>Total PhIP</b>							
Cases		69	68	51	61	46	
Multivariate RR	†	1	1.02 (0.73–1.43)	0.76 (0.52–1.09)	0.94 (0.66–1.34)	0.73 (0.50–1.08)	0.12
<b>Total DiMeIQx</b>							
Cases		62	67	63	58	45	
Multivariate RR	†	1	1.06 (0.75–1.51)	1.01 (0.71–1.45)	0.92 (0.64–1.33)	0.74 (0.50–1.10)	0.06
<b>Meat derived mutagenic activity</b>							
Cases		62	61	60	55	57	
Multivariate RR	†	1	0.97 (0.68–1.39)	0.96 (0.67–1.38)	0.91 (0.63–1.32)	0.93 (0.64–1.36)	0.70

\* Numbers in parenthesis next to RR are 95% confidence intervals

† Multivariate models adjusted for same covariates as denoted in table 3

Table 5

Relative risk of total breast cancer by HCA and MDM intake and fruit and vegetable intake in women in the Nurses Health Study (1996–2006)

	Low Fruit/Vegetable (<5.25 serving/day)	High Fruit/Vegetable (>=5.25 serving/day)	P-interaction
<b>Total MeIQx</b>			
Cases (lowest/highest quintile)	167/199	257/189	
Multivariate RR <sup>†</sup>	1.08 (0.87–1.35)	0.81 (0.66–1.00)	0.12
<b>Total PhIP</b>			
Cases (lowest/highest quintile)	181/153	244/197	
Multivariate RR <sup>†</sup>	1.03 (0.82–1.29)	0.77 (0.63–0.93)	0.45
<b>Total DiMeIQx</b>			
Cases (lowest/highest quintile)	176/173	235/200	
Multivariate RR <sup>†</sup>	0.95 (0.76–1.18)	0.92 (0.76–1.12)	0.86
<b>Meat derived mutagenic activity</b>			
Cases (lowest/highest quintile)	180/177	224/205	
Multivariate RR <sup>†</sup>	1.16 (0.94–1.44)	0.85 (0.70–1.03)	0.07

\* Numbers in parenthesis next to RR are 95% confidence intervals

<sup>†</sup> Multivariate models adjusted for same covariates as denoted in table 3