

RESEARCH ARTICLE

Open Access

Understanding dynamics using sensitivity analysis: caveat and solution

Thanneer M Perumal¹, Rudiyanto Gunawan^{2*}

Abstract

Background: Parametric sensitivity analysis (PSA) has become one of the most commonly used tools in computational systems biology, in which the sensitivity coefficients are used to study the parametric dependence of biological models. As many of these models describe dynamical behaviour of biological systems, the PSA has subsequently been used to elucidate important cellular processes that regulate this dynamics. However, in this paper, we show that the PSA coefficients are not suitable in inferring the mechanisms by which dynamical behaviour arises and in fact it can even lead to incorrect conclusions.

Results: A careful interpretation of parametric perturbations used in the PSA is presented here to explain the issue of using this analysis in inferring dynamics. In short, the PSA coefficients quantify the integrated change in the system behaviour due to persistent parametric perturbations, and thus the dynamical information of when a parameter perturbation matters is lost. To get around this issue, we present a new sensitivity analysis based on impulse perturbations on system parameters, which is named impulse parametric sensitivity analysis (iPSA). The inability of PSA and the efficacy of iPSA in revealing mechanistic information of a dynamical system are illustrated using two examples involving switch activation.

Conclusions: The interpretation of the PSA coefficients of dynamical systems should take into account the persistent nature of parametric perturbations involved in the derivation of this analysis. The application of PSA to identify the controlling mechanism of dynamical behaviour can be misleading. By using impulse perturbations, introduced at different times, the iPSA provides the necessary information to understand how dynamics is achieved, i.e. which parameters are essential and when they become important.

Background

Parametric sensitivity analysis (PSA) has become a must have tool in the computational systems biologists' arsenal. In most applications of this analysis, one computes sensitivity coefficients or metrics, which generally reflect the ratios between the change in a biological model output and the perturbation on system parameters that cause this change. Depending on the magnitude of the perturbations, sensitivity analyses can be classified into local (infinitesimal perturbation) and global (finite perturbation). Regardless of these classes, the interpretation of the sensitivity metrics is intuitive; parameters with large sensitivity magnitude are deemed to be important and hence considered to be the controlling factors in the system functional regulation.

Consequently, one of the common uses of PSA in systems biology is to infer the importance of cellular processes or pathways and to provide mechanistic explanations for biological behaviour [1-5].

On a separate note, dynamics is a prominent feature of many important biological processes (e.g., oscillations in cell cycle and circadian rhythm [6,7], switching behaviour in programmed cell death [8], and adaptation in chemotaxis [9]). Cellular homeostatic regulation, despite the name, relies on an active dynamical response, in which orchestrated events take place in response to internal and external stimuli. Thus, understanding cellular dynamics has become a prime concern in systems biology, in which mathematical modelling coupled with quantitative analysis have been used to gain insights on the mechanisms that give rise to and control the dynamic behaviour [1-5]. These insights can provide the molecular targets for altering system dynamic behaviour,

* Correspondence: rudi.gunawan@chem.ethz.ch

²Institute for Chemical and Bioengineering, ETH Zurich, Switzerland
Full list of author information is available at the end of the article

such as in finding treatment for diseases or in (re)engineering of cellular systems.

While there are many choices of mathematical frameworks for dynamic modelling, ordinary differential equations (ODEs) are the most commonly used modelling paradigm in systems biology and have been used to describe a wide range of biological systems. In addition, ODEs are amenable to many standard quantitative and theoretical analyses, including sensitivity analysis and bifurcation analysis, for which many off-the-shelf software packages exist that provide an integrated and user-friendly computational platform for model simulations and analyses (e.g., MATLAB [10] and XPPAUT [11]). The PSA of ODE models can be readily done using software packages such as SimBiology toolbox of MATLAB [12], PottersWheel [13], Gepasi [14], Copasi [15], JDesigner/Jarnac [16], JSim [17], BioSens [18], SBML-SAT [19], and SensSB [20]. These and other software for sensitivity analysis have been summarized in the review articles by Alves. *et al.* [21] and Klipp. *et al.* [22].

Sensitivity analysis of ODE models is well established in the science and engineering literature [23-32]. In systems biology, PSA has found wide applications, such as for model calibration and identifiability, model validation and reduction, identification of bottlenecking processes, elucidation of mechanisms of complex cellular behaviour, and investigation of cellular robustness [30,33]. A few notable examples of PSA applications in dynamic biological models include programmed cell death [34-39], budding yeast cell cycle control [6], IL-6 signalling pathway [1], circadian rhythm models [7,40,41], and coupled MAPK and PI3K signal transduction pathway [42]. In many applications, PSA is used to generate parameter ranking based on the magnitude of sensitivity coefficients, either taken at a specific time or using consolidated sensitivity metrics, such as time-integral or average or norm of sensitivity coefficients [34,43,44]. The parameter ranking is subsequently used to conclude about the mechanism or property (such as robustness) of the biological system behaviour [1-5].

In this article, we show that the dynamical aspects of cellular functional regulation cannot be inferred from the sensitivity coefficients of PSA, neither directly nor as consolidated sensitivity metrics. More importantly, the corresponding parameter rankings from PSA can give erroneous inference about the controlling mechanisms. Briefly, the reason stems from the fact that in PSA, perturbations are introduced on system parameters, which are time-invariant or static. In other words, these parametric perturbations are persistent and their effects on the system behaviour are integrated over time. Therefore, while PSA can indicate which parameter perturbations are important, it does not point to when these perturbations matter. This problem is illustrated using

local PSA of two examples: a synthetic network and a model of programmed cell death [37]. Although the illustration here was done using local sensitivity analysis, the same issue generally applies to global PSA.

To overcome this issue, a new parametric sensitivity analysis is developed in this work. This analysis differs from the classical PSA in the manner of which perturbations are introduced on model parameters, specifically using impulses, and thus is named impulse parametric sensitivity analysis (iPSA). By analyzing the consequence of impulse parameter perturbations introduced at different times, the iPSA provides time-varying, mechanistic explanation on how system dynamics is carried out. The new insights from the iPSA are demonstrated using the same two examples mentioned above.

Results and discussion

Simple network model

To illustrate the shortcoming of local PSA in analyzing system dynamics, consider a simple six state model involving three reactions with Michealis-Menten (MM) kinetics, as shown in Figure 1(a) (model parameters, rate kinetics and initial concentrations are given in Additional File 1: Supplementary Table S1). In this network, the activation of x_6 followed a switch-like




Figure 1 A simple network model. (a) A simple network with 6 states and 4 reactions. Straight arrows connect substrate to product and the dotted arrows indicate enzymatic activity. Details of model equations and parameter values are given in Table S1 (see Additional file 1: Supplementary Table S1). (b) Activation of x_6 under a constant stimulus of $x_1 = 1$: complete network (+), indirect pathway knock-out (*) and direct pathway knockout (x).

dynamics in response to the stimulus x_1 , as illustrated in Figure 1(b) (nominal). The model describes two pathways that contribute to x_6 activation: a direct x_2 pathway and an indirect x_2, x_3 , and x_5 pathway.

In this example, *in silico* knock-out experiments were performed by removing each pathway individually in order to assess the dominance of one pathway over the other in x_6 activation. Both full network and knock-out (KO) simulations were performed under a stimulus of x_1 ($t_0 = 0$) = 1. As illustrated in Figure 1(b), while the initial x_6 activation in the indirect pathway KO remained the same as that of the original model, the switch-like activation was much less pronounced. On the other hand, the original switching behaviour was preserved in the direct pathway KO, but the switching time was delayed due to a slower initial activation. Taken together, these KO simulations suggested that the x_6 activation is mainly accomplished through the indirect pathway, while the direct pathway contributes mainly to the initial x_6 activation.


Parametric sensitivity analysis for dynamical systems: A caveat

Local parametric sensitivity analysis was also used to study the pathway dominance in this simple network. Mathematically, the parametric sensitivity coefficient is given as

$$S_{ij} = \frac{\partial x_i(t)}{\partial p_j(\tau)} \quad (1)$$

where x_i is the i -th state in an ODE model and p_j is the j -th kinetic parameter of an ODE model (for a detailed description of sensitivity coefficient derivation, see Methods). These sensitivity coefficients describe the change in system output (state trajectory) at time t with respect to (an infinitesimal) perturbation on the system parameter values at time τ . Here, the PSA was performed for the same stimulus $x_1(0) = 1$ with $\tau = 0$ and the sensitivity coefficients were computed for the time range of 0-15 time units. The term *local* sensitivity analysis refers to the fact that the results will depend on the nominal parameter values around which the derivatives in Eq. (1) are calculated.

The sensitivities of x_6 with respect to all model parameters are ranked in Figure 2 using consolidated sensitivity metrics, i.e. infinite norm [44] (Figure 2(a)), FIM [43] (Figure 2(b)), and time-integral [34] (Figure 2(c)), and using sensitivity magnitudes at switching time ($t = 7.12$ time units; Figure 2(d)). The conclusion from these rankings was the same: (1) the largest sensitivity was associated with the kinetics of x_1 conversion to x_2 and (2) the direct pathway (r_2) parameters have larger sensitivities than those from the indirect pathway (r_4),




suggesting larger influence of the direct pathway on the x_6 activation. Hence, the conclusion from the PSA is in direct contradiction with the findings from *in silico* KO experiments.

This discrepancy can be explained by taking a closer look at the way parametric sensitivity coefficients in eqn. (1) are calculated:

$$S_{ij}(t, \tau) = \int_{\tau}^t \dot{S}_{ij}(\hat{t}) d\hat{t} = \int_{t_0}^t \dot{S}_{ij}(\hat{t}) H(\hat{t} - \tau) d\hat{t} \quad (2)$$

where $\tau = t_0$ is the usual perturbation time, $\dot{S}_{ij}(\hat{t})$ is the time derivative of sensitivity coefficient $S_{ij}(\hat{t})$ (see Methods for detail) and $H(t)$ is the Heaviside step function. In this case, since model parameters are static or time-invariant, the parametric perturbations in the PSA consist of step changes in the parameter values, as depicted in Figure 3(a). Hence, the sensitivity coefficients at time t represent an integrated or accumulated change in the states from τ to t due to a persistent parameter change started at time τ (see Figure 3(b)). Indeed, substituting the full equation of $\dot{S}(t)$ (see Methods) in eqn. (2) gives

$$S_{ij}(t, \tau) = \int_{t_0}^t \left[J_{i,[1..n]} S_{[1..n],j}(\hat{t}) + \frac{\partial f_i}{\partial p_j}(\hat{t}) \right] H(\hat{t} - \tau) d\hat{t} \quad (3)$$



Here, one sees two terms in the integrand that contribute to the sensitivity coefficients at time t : (1) the first is related to the (integrated) sensitivities that are carried over from the initial perturbation time τ and (2) the second accounts for the instantaneous rate changes due to the parametric perturbations that still persist at time t . Thus, in the PSA, a large sensitivity magnitude of $S_{ij}(t, \tau)$ indicates the importance of the j -th parameter in time window of τ and t , during which the perturbation is applied to the system. Hence, the use of these coefficients to infer the dynamical importance of parameters is inappropriate and can even be misleading.


For the reason above, the PSA of the simple network model gave an incorrect conclusion regarding direct versus indirect pathway activating x_6 . As seen in the *in silico* KO experiments, the direct pathway regulates the initial activation of x_6 , while the actual switching is carried out by the indirect pathway. In the PSA of this model ($\tau = 0$), the early importance of the direct pathway and also the reaction r_1 persisted beyond the initial

times in the sensitivity coefficients due to the aforementioned integrated effect. In this case, the importance of the indirect pathway was not apparent from the parameter sensitivity rankings in the background of large (integrated) sensitivities with respect to r_1 and the direct pathway. Correspondingly, the use of any time-consolidated sensitivity metrics will only worsen this problem.

Impulse parametric sensitivity analysis (iPSA)

As the problem with local PSA mentioned above is rooted from the persistent parameter perturbations, which is also done in global PSA [45], a new sensitivity analysis is formulated here that introduces impulse perturbations to model parameters as shown in Figure 3(c-d). The corresponding impulse sensitivity coefficients $iS_{ij}(t, \tau)$ reflect the change in the i -th state x_i at time t due to an impulse perturbation on the j -th parameter p_j at time τ (see Methods for the derivation and definition). Since impulse perturbations on parameters cause an immediate state changes at the perturbation time (see Methods), the inference of dynamical parametric importance can be obtained from impulse sensitivities by varying the time of perturbation. However, as with the local PSA, impulse perturbations are also *local* in nature and thus the impulse sensitivities will depend on the nominal parameter values. The global equivalent of iPSA can be formulated using pulse perturbations and is currently under development. Finally, from time-varying impulse perturbations, the iPSA can give answer to the following questions about state dynamics: which are the important parameters and when do they become important?

The iPSA was also applied to the simple network model under the same stimulus and for the same time range as above. Since the impulse perturbations are delivered at different times, the impulse sensitivity $iS_{ij}(t, \tau)$ has two-time dependence, with respect to the time of perturbation (τ) and the time of observation (t). Figure 4 shows the iPSA sensitivity coefficients of x_6 at the end of simulation time ($t = 15$ time units), with respect to the four most important parameters at different perturbation times (for complete iPSA sensitivities, see Additional File 1: Supplementary Figure S1). In agreement with the KO simulations and in contrast to the findings from PSA, the impulse sensitivities gave support to the dominance of the indirect pathway. Specifically, the results showed that x_6 activation: (1) is initiated by r_1 (high initial sensitivity to parameter $kf1$); (2) is accomplished mainly by the direct pathway prior to the switching time (sensitivity to r_2 is higher than to r_4 during these times); and (3) is subsequently carried by the indirect pathway (highest sensitivity to r_4 during switching times). A higher resolution analysis using a heat map of the complete iPSA coefficients with t and τ




between 0 and 15 time units gave the same conclusion (see Additional file 1: Supplementary Figure S2).


Fas-induced cell death model of human Jurkat T-cells

In the second case study, we consider a more complex biological network taken from the modelling of programmed cell death in Jurkat cancer T-cells (see Figure 5). The model equations and parameters were identified from experimental data [37] (see Additional file 1: Supplementary Table S2 for detailed reaction rates and parameters). In this network, the cell death is decided

by the cleaving of procaspase-3 into caspase-3 [46], in response to FasL stimulus. The model simulation showed that caspase-3 is switched ON at around $t = 6000$ seconds (see Figure 5, inset) and like the simple network above, there exist two activating pathways: the direct caspase-8 (type-I) and the indirect mitochondria-dependent pathway (type-II). As in the previous case study, the classical PSA and iPSA were applied to this network to elucidate the pathway dominance.

The classical and iPSA analyses were calculated under a constant FasL stimulation (FasL = 2 nM) over the time range of 10,000 seconds. The rankings of the important parameters that control caspase-3 according to the PSA are shown in Figure 6, using consolidated metrics: infinite norm [44](Figure 6(a)), FIM [43] (Figure 6(b)), and time integral [34](Figure 6(c)), and using the sensitivity magnitudes at switching time (Figure 6(d)) (see Additional file 1: Supplementary Figure S3 for detailed sensitivity rankings). From these rankings, one could not obtain any definitive conclusion regarding the dominance of one pathway over the other. On the contrary, the iPSA sensitivity of caspase-3 in Figure 7 clearly supported a type-II dependent caspase-3 switching with an early type-I dependent activation, in agreement with two previous analyses of this model using the Green's function matrix [47] and model reduction [48]. In Figure 7, the high caspase-3 sensitivities with respect to impulse perturbations to upstream reactions were expected during the initial times, as these served as the early cell death signal response. The cleaving of procaspase-3 was carried out by caspase-8 directly (r_5 of type-







1) before $t = 4000$ seconds, after which the mitochondrial pathway (r_{14} of type-II) became the main route of activating caspase-3 (see Additional file 1: Supplementary Figures S4 and S5 for more detail).

The discrepancy between the PSA and iPSA results can again be explained in the context of persistent versus impulse perturbations. As seen in the PSA parameter rankings in Figure 6 and following the insights offered by the iPSA in Figure 7, the effect of perturbing early response processes, including type-I pathway, was integrated over time in the PSA. For example, the highest ranked parameters in the PSA were associated with the first three reactions, r_1 to r_3 . Such integration masked the dynamical importance of different parameters in this analysis. The conclusion from the iPSA is in agreement with the simulations of KOs of type-I and type-II pathways in Figure 8. While type-II knock-out could describe the caspase-3 activation at early times, it failed to capture the switching behaviour. On the other hand, the type-I knock-out was able to reproduce the switching of caspase-3, albeit with a short delay.

The two examples above illustrate the problem of using the classical PSA in identifying the controlling mechanisms of a dynamical system. Of course, this does not mean that the PSA of dynamical models is incorrect, but rather the interpretation of the sensitivity coefficients should be carefully managed. In particular, a large sensitivity magnitude with respect to a parameter suggests the importance of this parameter in the time period between the perturbation time τ and the state observation time t . In contrast, the iPSA is developed with dynamics in mind, where the impact of a single perturbation on the system is realized only at the perturbation time and



subsequently they are delivered at varying perturbation times. By doing so, the iPSA coefficients can elucidate the way system dynamics $\mathbf{x}(t)$ is achieved, by indicating which and when parameters or processes are essential. Because of the persistent nature of perturbations used in the PSA, it is still not possible to reproduce the conclusions of the iPSA by varying the time of perturbation (see Additional file 1: Supplementary Figures S6 and S7).

Conclusions

While classical parametric sensitivity analysis provides a powerful tool to understand the parametric dependence of biological behaviour, its suitability in inferring mechanisms of dynamic behaviour has not been properly addressed. The two case studies here illustrated the caveat of using local PSA for such purpose. The issue mainly arose from the information needed to do this inference, where one needs to know not only which parameters are critical, but also when they matter. However, the persistent parametric perturbations in standard PSA are incapable of providing this information as the sensitivity coefficients represent an integrated effect. A new sensitivity analysis, called impulse parametric sensitivity analysis (iPSA), was developed with dynamical systems in mind. In particular, the iPSA makes use of local impulse perturbations introduced at different times to produce the necessary information for understanding dynamics. The application of iPSA to the case studies was able to correctly pinpoint the mechanisms responsible for dynamical system behaviour, while local PSA failed in these cases. Since the discrepancy between PSA and iPSA arises from a fundamental difference in the manner of which parametric perturbations are realized (i.e. persistent vs. impulse), the same caveat and solution can be generalized to the global PSA, in which the perturbations are no longer infinitesimal.

Methods

Mathematical Models

Ordinary differential equation models of dynamic systems can generally be written as:

$$\begin{aligned} \frac{d\mathbf{x}(t, \hat{\mathbf{p}})}{dt} &= f(\mathbf{x}, \hat{\mathbf{p}}) \\ \mathbf{x}(t_0, \hat{\mathbf{p}}) &= \mathbf{x}_0 \end{aligned} \tag{4}$$

In biological models, the state $\mathbf{x} \in \mathbb{R}^n$ is typically the concentration vector of biomolecular species, such as mRNAs and proteins, while the function f is the constitutive, often nonlinear, rate equation. The right hand side of the ODE captures the generation and consumption of biomolecules due to a variety of processes in the cell (e.g. transcription, translation, phosphorylation and

dephosphorylation, etc), the rates of which depend on a set of kinetic parameters that are consolidated in the vector $\hat{\mathbf{p}} \in \mathbb{R}^m$. Since the initial conditions \mathbf{x}_0 can be treated in the same way as model parameters, the aggregate vector $\mathbf{p} \in \mathbb{R}^{m+n}$ is used here to denote the combined parameters and initial conditions, i.e. $\mathbf{p} = [\hat{\mathbf{p}}^T \ \mathbf{x}_0^T]^T$.

Local parametric sensitivity analysis (PSA)

The effect of parameter perturbations can be written in a Taylor series expansion:

$$\begin{aligned} x_i(t, \mathbf{p} + \Delta\mathbf{p}) &= x_i(t, \mathbf{p}) + \sum_{j=1}^m \frac{\partial x_i(t, \mathbf{p})}{\partial p_j} \Delta p_j \\ &+ \frac{1}{2} \sum_{l=1}^m \sum_{j=1}^m \frac{\partial^2 x_i(t, \mathbf{p})}{\partial p_l \partial p_j} \Delta p_l \Delta p_j + \dots \end{aligned} \tag{5}$$

where the partial derivatives $\partial x_i / \partial p_j$'s are the first-order sensitivity coefficients, describing the linear change in the state x_i at any time t with respect to an infinitesimal perturbation to the parameter p_j . In general, the parametric perturbation can be introduced at any time τ ($t_0 \leq \tau \leq t$) [27], i.e.

$$\begin{aligned} S_{ij}(t, \tau) &= \frac{\partial x_i(t)}{\partial p_j(\tau)} \\ S_{ij}(t, \tau) &= \frac{\text{change in } i\text{-th state at time } t}{\text{pert. on } j\text{-th parameter at time } \tau} \end{aligned} \tag{6}$$

but in the PSA, the perturbation time τ is commonly taken to be the initial time t_0 . Hence, the argument τ is typically dropped out of eqn. (6) and the sensitivity coefficients only carry a single time dependence on the observation time t [1-3,5,30,44,49,50]. The higher order sensitivity coefficients in the Taylor series expansion are less commonly computed, and hence the focus of the current work is only on the first-order sensitivities. Because the magnitude of perturbations are infinitesimally small, the sensitivity coefficients will depend on the nominal or baseline parameter values, and thus the classical PSA is considered a local analysis.

The sensitivity coefficients of an ODE model can be computed by directly differentiating the model eqn. (4) with respect to parameters \mathbf{p} , giving the differential equation for the sensitivity coefficients as:

$$\begin{aligned} \frac{d}{dt} \mathbf{S}(t) &= \frac{\partial f}{\partial \mathbf{x}} \mathbf{S}(t) + \frac{\partial f}{\partial \mathbf{p}}; \\ \mathbf{S}(\tau) &= [\mathbf{0}^{n \times m} \ \mathbf{I}^{n \times n}] \end{aligned} \tag{7}$$

where $\partial f / \partial \mathbf{x}$ is also known as the Jacobian matrix and $\mathbf{0}^{n \times m}$ and $\mathbf{I}^{n \times n}$ are $n \times m$ zero and $n \times n$ identity

matrices, respectively. The parametric sensitivity coefficients in eqn. (7) need to be solved simultaneously with the ODE model in eqn. (4), which is called the direct method [28]. As the state and parameter values may span a large range of magnitudes, normalized sensitivity values are often used to compare among states and parameters and to generate parameter ranking, which is given by:

$$\bar{S}_{i,j}(t) = \frac{\partial x_i(t)}{\partial p_j} \frac{p_j}{x_i(t)} = \frac{\partial \log x_i(t)}{\partial \log p_j} \quad (8)$$

In most systems biology applications of PSA, parameter rankings are generated from the sensitivity coefficients, either directly or using some consolidated sensitivity metrics. In this article, four sensitivity metrics are used to rank parameters based on the PSA results:

$$\begin{aligned} [S_{\text{inf}}]_{ij} &= \max_k (|S_{ij}(t_k)|), \\ [S_{\text{FIM}}]_{ij} &= (\text{FIM})_{jj}, \quad \left(\text{FIM} = \sum_{t_k=t_0}^t \mathbf{S}_{i[1..n]}^T(t_k) \mathbf{S}_{i[1..n]}(t_k) \right) \\ [S_{\text{int}}]_{ij} &= \int_{t_0}^t S_{ij}(\hat{t}) d\hat{t}, \\ [S_{t_k}]_{ij} &= S_{ij}(t = t_k), \end{aligned} \quad (9)$$

where the indices i and j again denote the i -th state and j -th parameter, and S_{inf} , S_{FIM} , S_{int} and S_{t_k} are the sensitivity metrics based on infinite norm [44], Fisher information matrix [43], time integral [34] and sensitivity magnitude at a particular time, respectively.

Impulse parametric sensitivity analysis (iPSA)

The derivation of the iPSA coefficient follows the illustration in Figure 3(e). The sensitivity coefficients of iPSA are constructed by quantifying the ratio between the change in the state x_i at time t and the causative pulse perturbation of size $\Delta p_j / \Delta \tau$ for a duration of Δt , which is applied to the parameter p_j at time τ , in the limit Δp_j and $\Delta \tau$ tending to zero. The first step of the derivation is to quantify the change in all states \mathbf{x} at the end of the pulse perturbation, i.e. at time $\tau + \Delta \tau$, using the Taylor series expansion:

$$\Delta \mathbf{x}(\tau + \Delta \tau) = S_{[1..n],j}(\tau + \Delta \tau, \tau) \frac{\Delta p_j}{\Delta \tau} + O(\Delta p_j^2) \quad (10)$$

In the next step, the change $\Delta \mathbf{x}(\tau + \Delta \tau)$ is translated to the change in the state x_i at time t using the Green's function matrix (GFM) $\mathbf{S}^x(t, \tau + \Delta \tau)$ [47]. The (i, j) -th element of the GFM represents the sensitivity of the state x_i at time t to alteration in the state x_j at some previous time τ , i.e.

$$S_{i,j}^x(t, \tau) = \frac{\partial x_i(t)}{\partial x_j(\tau)}. \quad (11)$$

Thus, the change $\Delta x_i(t)$ due to the pulse perturbation is given by

$$\Delta x_i(t) = S_{i,[1..n]}^x(t, \tau + \Delta \tau) \Delta \mathbf{x}(\tau + \Delta \tau). \quad (12)$$

Subsequently, substitution of eqn. (10) in eqn. (12) gives

$$\Delta x_i(t) = S_{i,[1..n]}^x(t, \tau + \Delta \tau) S_{[1..n],j}(\tau + \Delta \tau, \tau) \frac{\Delta p_j}{\Delta \tau} + O(\Delta p_j^2) \quad (13)$$

Then, taking Taylor series expansion of the parametric sensitivities around the time τ and dividing both sides by Δp_j , one arrives with:

$$\begin{aligned} \Delta x_i(t) &= S_{i,[1..n]}^x(t, \tau + \Delta \tau) [S_{[1..n],j}(\tau, \tau) \\ &\quad + \dot{S}_{[1..n],j}(\tau, \tau) \Delta \tau + O(\Delta \tau^2)] \frac{\Delta p_j}{\Delta \tau} + O(\Delta p_j^2) \\ \frac{\Delta x_i(t)}{\Delta p_j} &= S_{i,[1..n]}^x(t, \tau + \Delta \tau) [\dot{S}_{[1..n],j}(\tau, \tau) \\ &\quad + O(\Delta \tau)] + O(\Delta p_j) \\ \frac{\Delta x_i(t)}{\Delta p_j} &= S_{i,[1..n]}^x(t, \tau + \Delta \tau) \left[\frac{\partial f}{\partial p_j}(\tau) \right. \\ &\quad \left. + O(\Delta \tau)] + O(\Delta p_j) \end{aligned} \quad (14)$$

Finally, taking the limit as $\Delta p_j, \Delta \tau \rightarrow 0$ such that the pulse perturbation becomes an impulse, the iPSA coefficient is obtained as:

$$iS_{i,j}(t, \tau) = S_{i,[1..n]}^x(t, \tau) \frac{\partial f}{\partial p_j}(\tau) \quad (15)$$

Note that at $t = \tau$, the iPSA coefficient reduces to

$$iS_{i,j}(\tau, \tau) = \frac{\partial f_i}{\partial p_j}(\tau) \quad (16)$$

since $\mathbf{S}^x(\tau, \tau) = \mathbf{I}$. In other words, the impulse parameter perturbation causes an immediate change in the state \mathbf{x} at time τ . By rewriting eqn. (15) as:

$$iS_{i,j}(t, \tau) = S_{i,[1..n]}^x(t, \tau) iS_{i,j}(\tau, \tau) \quad (17)$$

one can further see that the impact of this impulse perturbation takes effect only at the perturbation time τ and that the consequence on the state trajectory is equivalent to perturbing the states themselves, similar to the GFM analysis. Like in the PSA, the iPSA coefficients should be normalized for comparison and parameter ranking purposes, according to:

$$\bar{i}S_{ij}(t, \tau) = iS_{ij}(t, \tau) \frac{p_j}{x_i(t)} \quad (18)$$

Additional material

Additional file 1: Supplementary Material. Detailed results of iPSA and PSA, including model equations and parameters, of the simple network model and Fas-induced cell death model

List of abbreviations

KO: Knock-Out; ODE: Ordinary Differential Equations; PSA: Parametric Sensitivity Analysis; iPSA: Impulse Parametric Sensitivity Analysis.

Acknowledgements

TMP is supported by the Singapore Millennium Foundation scholarship.

Author details

¹Department of Chemical and Biomolecular Engineering, National University of Singapore, Singapore. ²Institute for Chemical and Bioengineering, ETH Zurich, Switzerland.

Authors' contributions

TMP performed the experiments and wrote the manuscript. RG supervised the study and wrote the manuscript. All authors have read and approved the final manuscript.

Received: 24 September 2010 Accepted: 15 March 2011

Published: 15 March 2011

References

1. Chu Y, Jayaraman A, Hahn J: **Parameter sensitivity analysis of IL-6 signalling pathways.** *IET Syst Biol* 2007, **1**:342-352.
2. Ihekwaba AE, Broomhead DS, Grimley RL, Benson N, Kell DB: **Sensitivity analysis of parameters controlling oscillatory signalling in the NF-kappaB pathway: the roles of IKK and IkkappaBalpha.** *Syst Biol (Stevenage)* 2004, **1**:93-103.
3. Ihekwaba AE, Broomhead DS, Grimley R, Benson N, White MR, Kell DB: **Synergistic control of oscillations in the NF-kappaB signalling pathway.** *Syst Biol (Stevenage)* 2005, **152**:153-160.
4. Adler P, Peterson H, Agius P, Reimand J, Vilo J: **Ranking Genes by Their Co-expression to Subsets of Pathway Members.** *Annals of the New York Academy of Sciences* 2009, **1158**:1-13.
5. Iwamoto K, Tashima Y, Hamada H, Eguchi Y, Okamoto M: **Mathematical modeling and sensitivity analysis of G1/S phase in the cell cycle including the DNA-damage signal transduction pathway.** *Biosystems* 2008, **94**:109-117.
6. Lovrics A, Zsély IGy, Csikász-Nagy A, Zádor J, Turányi T, Novák B: **Analysis of a budding yeast cell cycle model using the shapes of local sensitivity functions.** *International Journal of Chemical Kinetics* 2008, **40**:710-720.
7. Jin Y, Peng X, Liang Y, Ma J: **Uniform design-based sensitivity analysis of circadian rhythm model in Neurospora.** *Computers and Chemical Engineering* 2008, **32**:1956-1962.
8. Zhang T, Brazhnik P, Tyson JJ: **Computational analysis of dynamical responses to the intrinsic pathway of programmed cell death.** *Biophys J* 2009, **97**:415-434.
9. Hansen CH, Endres RG, Wingreen NS: **Chemotaxis in Escherichia coli: a molecular model for robust precise adaptation.** *PLoS Comput Biol* 2008, **4**:e1.
10. MATLAB; <http://www.mathworks.com/>.
11. XPPAUT; <http://www.math.pitt.edu/~bard/xpp/xpp.html>.
12. Schmidt H, Jirstrand M: **Systems Biology Toolbox for MATLAB: a computational platform for research in systems biology.** *Bioinformatics* 2006, **22**:514-515.
13. Maiwald T, Timmer J: **Dynamical modeling and multi-experiment fitting with PottersWheel.** *Bioinformatics* 2008, **24**:2037-2043.
14. Mendes P: **GEPASI: a software package for modelling the dynamics, steady states and control of biochemical and other systems.** *Comput Appl Biosci* 1993, **9**:563-571.
15. Hoops S, Sahle S, Gauges R, Lee C, Pahle J, Simus N, Singhal M, Xu L, Mendes P, Kummer U: **COPASI—a COMplex PATHway Simulator.** *Bioinformatics* 2006, **22**:3067-3074.
16. JDesigner/Jarnac; <http://sbw.kgi.edu/>.
17. JSim; <http://www.physiome.org/jsim/>.
18. BioSens; <http://www.chemengr.ucsb.edu/~ceweb/faculty/doyle/biosens/BioSens.htm>.
19. Zi Z, Zheng Y, Rundell AE, Klipp E: **SBML-SAT: a systems biology markup language (SBML) based sensitivity analysis tool.** *BMC Bioinformatics* 2008, **9**:342.
20. Rodriguez-Fernandez M, Banga JR: **SensSB: a software toolbox for the development and sensitivity analysis of systems biology models.** *Bioinformatics* 2006, **26**:1675-1676.
21. Alves R, Antunes F, Salvador A: **Tools for kinetic modeling of biochemical networks.** *Nat Biotechnol* 2006, **24**:667-672.
22. Klipp E, Liebermeister W, Helbig A, Kowald A, Schaber J: **Systems biology standards—the community speaks.** *Nat Biotechnol* 2007, **25**:390-391.
23. Saltelli A, Chan K, Scott EM: *Sensitivity Analysis: Gauging the Worth of Scientific Models* John Wiley & Sons, Ltd.; 2000.
24. Saltelli A, Ratto M, Tarantola S, Campolongo F: **Sensitivity analysis for chemical models.** *Chem Rev* 2005, **105**:2811-2827.
25. Saltelli A, Tarantola S, Campolongo F: **Sensitivity Analysis as an Ingredient of Modeling.** *Stat Sci* 2000, **15**:377-395.
26. Saltelli A, Tarantola S, Campolongo F, Ratto M: *Sensitivity Analysis in Practice: A Guide to Assessing Scientific Models* John Wiley & Sons, Ltd.; 2004.
27. Turányi T: **Sensitivity analysis of complex kinetic systems. Tools and applications.** *Journal of Mathematical Chemistry* 1990, **5**:203-248.
28. Varma A, Morbidelli M, Wu H: *Parametric Sensitivity in Chemical Systems* Cambridge University Press, Cambridge, UK; 1999.
29. Frey HC, Patil SR: **Identification and review of sensitivity analysis methods.** *Risk Anal* 2002, **22**:553-578.
30. Ingalls B: **Sensitivity analysis: from model parameters to system behaviour.** *Essays Biochem* 2008, **45**:177-193.
31. Marino S, Hogue IB, Ray CJ, Kirschner DE: **A methodology for performing global uncertainty and sensitivity analysis in systems biology.** *J Theor Biol* 2008, **254**:178-196.
32. Rabitz H, Kramer M, Dacol D: **Sensitivity analysis in chemical kinetics.** *Annual review of Physical Chemistry* 1983, **34**:419-461.
33. Stelling Jr, Sauer U, Szallasi Z, Doyle FJ, Doyle J: **Robustness of cellular functions.** *Cell* 2004, **118**:675-685.
34. Bentele M, Lavrik I, Ulrich M, Stober S, Heermann DW, Kalthoff H, Krammer PH, Eils R: **Mathematical modeling reveals threshold mechanism in CD95-induced apoptosis.** *J Cell Biol* 2004, **166**:839-851.
35. Eissing T, Allgower F, Bullinger E: **Robustness properties of apoptosis models with respect to parameter variations and intrinsic noise.** *Syst Biol (Stevenage)* 2005, **152**:221-228.
36. Hua F, Cornejo MG, Cardone MH, Stokes CL, Lauffenburger DA: **Effects of Bcl-2 levels on Fas signaling-induced caspase-3 activation: molecular genetic tests of computational model predictions.** *J Immunol* 2005, **175**:985-995.
37. Hua F, Hautaniemi S, Yokoo R, Lauffenburger DA: **Integrated mechanistic and data-driven modelling for multivariate analysis of signalling pathways.** *J R Soc Interface* 2006, **3**:515-526.
38. Shoemaker JE, Doyle III FJ: **Identifying Fragilities in Biochemical Networks: Robust Performance Analysis of Fas Signaling-Induced Apoptosis.** *Biophys J* 2008.
39. Aldridge BB, Haller G, Sorger PK, Lauffenburger DA: **Direct Lyapunov exponent analysis enables parametric study of transient signalling governing cell behaviour.** *Syst Biol (Stevenage)* 2006, **153**:425-432.
40. Stelling Jr, Gilles ED, Doyle FJ: **Robustness properties of circadian clock architectures.** *Proc Natl Acad Sci USA* 2004, **101**:13210-13215.
41. Gunawan R, Doyle FJ: **Phase sensitivity analysis of circadian rhythm entrainment.** *J Biol Rhythms* 2007, **22**:180-194.
42. Hu D, Yuan JM: **Time-dependent sensitivity analysis of biological networks: coupled MAPK and PI3K signal transduction pathways.** *J Phys Chem A* 2006, **110**:5361-5370.

43. Yue H, Brown M, He F, Jia J, Kell DB: **Sensitivity analysis and robust experimental design of a signal transduction pathway system.** *International Journal of Chemical Kinetics* 2008, **40**:730-741.
44. Yue H, Brown M, Knowles J, Wang H, Broomhead DS, Kell DB: **Insights into the behaviour of systems biology models from dynamic sensitivity and identifiability analysis: a case study of an NF-kappaB signalling pathway.** *Mol Biosyst* 2006, **2**:640-649.
45. Saltelli A: *Global sensitivity analysis: the primer* Chichester, England: John Wiley; 2008.
46. Reed JC, Doctor KS, Godzik A: **The domains of apoptosis: a genomics perspective.** *Sci STKE* 2004 2004, re9.
47. Perumal TM, Wu Y, Gunawan R: **Dynamical analysis of cellular networks based on the Green's function matrix.** *J Theor Biol* 2009, **261**:248-259.
48. Okazaki N, Asano R, Kinoshita T, Chuman H: **Simple computational models of type I/type II cells in Fas signaling-induced apoptosis.** *J Theor Biol* 2008, **250**:621-633.
49. Yamada S, Shiono S, Joo A, Yoshimura A: **Control mechanism of JAK/STAT signal transduction pathway.** *FEBS Lett* 2003, **534**:190-196.
50. Zi Z, Cho KH, Sung MH, Xia X, Zheng J, Sun Z: **In silico identification of the key components and steps in IFN-gamma induced JAK-STAT signaling pathway.** *FEBS Lett* 2005, **579**:1101-1108.

doi:10.1186/1752-0509-5-41

Cite this article as: Perumal and Gunawan: Understanding dynamics using sensitivity analysis: caveat and solution. *BMC Systems Biology* 2011 5:41.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

