



Published in final edited form as:

*Schizophr Res.* 2011 February ; 125(2-3): 143–151. doi:10.1016/j.schres.2010.11.007.

## Social Cognition in Psychosis: Multidimensional Structure, Clinical Correlates, and Relationship With Functional Outcome

Francesco Mancuso<sup>a,b,c</sup>, William P. Horan<sup>a,b</sup>, Robert S. Kern<sup>a,b</sup>, and Michael F. Green<sup>a,b</sup>

<sup>a</sup>Department of Psychiatry and Biobehavioral Sciences, Geffen School of Medicine at University of California, Los Angeles, USA

<sup>b</sup>VA Greater Los Angeles Healthcare System, VISN22 MIRECC, USA

<sup>c</sup>Department of Psychiatry, University of Naples SUN, Naples, Italy

### Abstract

Social cognitive impairments are common, detectable across a wide range of tasks, and appear to play a key role in explaining poor outcome in schizophrenia and related psychotic disorders. However, little is known about the underlying factor structure of social cognition in people with psychotic disorders due to a lack of exploratory factor analyses using a relatively comprehensive social cognitive assessment battery. In a sample of 85 outpatients with psychosis, we examined the factor structure and clinical/functional correlates of eight indexes derived from five social cognition tasks that span the domains of emotional processing, social perception, attributional style, and Theory of Mind. Exploratory factor analysis revealed three factors with relatively low inter-correlations that explained a total of 54% of the variance: (1) Hostile attributional style, (2) Lower-level social cue detection, and (3) Higher-level inferential and regulatory processes. None of the factors showed significant correlations with negative symptoms. Factor 1 significantly correlated with clinical symptoms (positive, depression-anxiety, agitation) but not functional outcome, whereas Factors 2 and 3 significantly correlated with functional outcome (functional capacity and real-world social and work functioning) but not clinical symptoms. Furthermore, Factor 2 accounted for unique incremental variance in functional capacity, above and beyond non-social neurocognition (measured with MATRICS Consensus Cognitive Battery) and negative symptoms. Results suggest that multiple separable dimensions of social cognition can be identified in psychosis, and these factors show distinct patterns of correlation with clinical features and functional outcome.

---

© 2010 Elsevier B.V. All rights reserved.

**Corresponding author:** William P. Horan, Ph.D., UCLA Department of Psychiatry & Biobehavioral Sciences, 300 UCLA Medical Plaza, Suite 2255, Los Angeles, CA 90095-6968, 310.206.8181 (voice), 310.206.3651 (fax), horan@ucla.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

#### Disclosures

None

#### Contributions

Dr. Green, Horan, and Kern designed the study and wrote the protocol. Drs. Horan and Mancuso undertook the statistical analysis, and Dr. Mancuso wrote the first draft of the manuscript. All authors contributed to and have approved the final manuscript.

#### Conflict of Interest

None

## Keywords

schizophrenia; social cognition; neurocognition; positive symptoms; negative symptoms; functional outcome

---

## 1. Introduction

Social cognition has emerged as a high priority topic for research in schizophrenia and related psychotic disorders that may help explain poor outcomes. Social cognition is a broad, multifaceted construct that refers to the cognitive and emotional functions required to understand and predict other people's mental states and behavior (Adolphs, 2009; Ochsner, 2008). Research has documented impairments across a diverse array of social cognitive processes, most commonly emotional processing, social knowledge/perception, attributional style, and Theory of Mind (Green & Horan, 2010). There have been two distinct goals of social cognitive research in schizophrenia: One devoted to understanding the nature of specific clinical symptoms (e.g., relations to paranoia or thought control) and another devoted to social cognition's role in functional outcome. Despite the proliferation of research in this promising area, several fundamental issues remain largely unexplored (Green et al., 2005; 2008).

One question concerns the underlying structure of social cognition in schizophrenia. Although there is a general consensus that social cognition is empirically and neurobiologically separable from (though related to) non-social neurocognition (Green et al., 2008; Fett et al., in press), it is not known whether the social cognitive assessments used in schizophrenia reflect a single factor or a cluster of separable factors. A few studies focusing on a narrow selection of social cognition measures suggest the presence of separable factors. For example, an exploratory factor analysis of neurocognitive and social cognitive measures found that indices of Theory of Mind, attributional bias, and agency detection loaded on three different factors (van Hooren et al., 2008). Another study using confirmatory factor analyses found that a four-factor model provided a good fit to measures of social cognition and social behavior (affect recognition, Theory of Mind, egocentricity, and rapport) (Bell et al., 2009). Along these lines, emotional intelligence shows a multidimensional structure in people with psychotic disorders that differs from healthy control subjects (Eack et al., 2009). The current study aimed to help clarify the basic structure of social cognition by examining performance across a wide range of social cognitive tasks.

A second question concerns social cognition's relation to clinical symptoms and functional outcome. Although understanding clinical symptoms has been one key motivation to study social cognition in schizophrenia, the literature provides a generally mixed picture. While recent modeling studies suggest that social cognition is separable from negative symptoms (Rassovsky et al., in press; Sergi et al., 2007), some studies report associations with particular social cognitive tests (e.g., Kohler et al., 2009). Relations to positive symptoms (e.g., thought disorder, hallucinations) are similarly inconsistent (Corcoran et al., 2008; Shamay-Tsoory et al., 2007; Woodward et al., 2009), though there has been somewhat greater consistency for associations between attributional style and paranoid delusions or beliefs (Bentall et al., 2001; Combs et al., 2007, 2009; Fornells-Ambrojo and Garety, 2009; Kinderman and Bentall, 1996). Regarding linkages to functional outcome, most studies have focused on emotion and social perception (Couture et al., 2006; Fett et al., in press). In addition, many studies include a limited assessment of functional outcome and often do not distinguish between functional capacity (what one can do on competence measures) and real-world functioning (what one actually does in the community), which is increasingly

recognized as a critical distinction (Bowie, 2006). The current study aimed to address these issues by incorporating comprehensive assessments of symptoms and functioning.

A third question is whether social cognition has “added value” in explaining functional outcome above and beyond neurocognition and clinical symptoms. Notably, at the level of simple correlations, a recent meta-analysis reported that social cognition has generally stronger relations to functional outcome than does neurocognition (Fett et al., in press). A few studies have found that social cognition (mainly emotion and social perception) does account for unique variance in outcome above and beyond neurocognition (Horan et al., in press). The current study used a relatively comprehensive social cognitive battery to further address whether social cognition plays a unique role in explaining functional outcome.

The primary goal of this research was to evaluate the factor structure of social cognition in a sample of outpatients with psychotic disorders. We used five different tests that cover the four most commonly investigated domains of social cognition in schizophrenia. Secondary goals were to: (1) examine correlations between the derived factor(s) and neurocognition, symptoms, and functional outcome, and (2) investigate whether the social cognitive factor(s) uniquely account for incremental variance in functional outcome, above and beyond neurocognition and symptoms.

## 2. Methods

### 2.1 Participants

Eighty-five outpatients were recruited from the VA Greater Los Angeles Healthcare System (VAGLAHS) and local community mental health centers. Patients met DSM-IV criteria for schizophrenia, schizoaffective disorder, or psychosis NOS (not secondary to substance use disorder) as determined by medical records and consultation with treating psychiatrists. Subjects were clinically stable (no psychiatric hospitalizations in the past 6 months, same antipsychotic medication for past 3 months). Exclusion criteria were current or past neurological disorder (e.g., epilepsy), mental retardation, or substance use disorder within the past month. Antipsychotic medication type and dose were left to the discretion of the treating physician. All participants had the capacity to give informed consent and provided written informed consent after all procedures were fully explained in accordance with procedures approved by the Institutional Review Board at the VAGLAHS. The participants were enrolled in a clinical trial comparing psychosocial interventions for social cognition and neurocognition (Horan et al., submitted); the current study used data from the baseline assessments.

### 2.2 Social cognitive assessment

1. **Emotional processing** was assessed with two tests: (a) The *Facial Emotion Identification Test* (FEIT), in which subjects view 56 digital pictures of faces from the Ekman (2004) picture set and select which emotion is expressed (happy, sad, angry, afraid, surprised, disgusted or neutral). The index of accuracy is the total number of correct items. (b) The *Managing Emotions* subtest of the Mayer-Salovey-Caruso Emotional Intelligence Test (MSCEIT) (Mayer et al., 2002; 2003) comprises two 2 subscales that examine the regulation of emotions in oneself and in one's relationships with others. These subscales include vignettes of various situations, along with ways to cope with the emotions depicted in these vignettes. Subjects were required to indicate the effectiveness of each solution, ranging from one (very ineffective) to five (very effective). A total score was derived using the MSCEIT General Consensus method.

2. Social Perception was assessed with The Half-Profile of Nonverbal Sensitivity (PONS) (Ambady et al., 1995; Rosenthal et al., 1979). The 110 scenes in this videotape-based measure last two seconds and contain facial expressions, voice intonations, and/or bodily gestures of a Caucasian female. After watching each scene, participants select which of two labels better describes a situation that would generate the social cue(s). The index of accuracy is the total number of correct items.
3. Attributional style was assessed with The Ambiguous Intentions Hostility Questionnaire (AIHQ) (Combs et al., 2007). This particular measure of attributional style focuses on a person's tendency to over-attribute hostile intentions to others and to respond to others in a hostile manner. Subjects read a series of vignettes describing social situations and answer questions about the intentions of the characters and how subjects themselves would respond to the situation. Following Combs et al. (2007), we examined scores for ambiguous situations only and computed three summary scores: Hostility bias, Aggression bias, and a composite "Blame" score (average of Intentionality, Anger, and Blame item ratings). The Hostility and Aggression bias scores were independently scored by two blinded raters (ICC's for both bias scores were .85+).
4. Theory of Mind was assessed with The Awareness of Social Inference Test (TASIT) – Part 3 (McDonald et al., 2002) is a videotape measure of Theory of Mind that contains 16 scenes depicting lies or sarcasm, with two or three actors appearing in each one. In each scene, a prologue/epilogue or camera edit provides information to the participant about the nature of the conversational exchange. After each scene, participants answer four types of forced-choice (yes/no) questions about the characters' communicative intentions, whether they want the literal or non-literal meaning of their message to be believed, their beliefs and knowledge about the situation, and their emotional state. Summary scores for Lies and Sarcasm were calculated (Kern et al., 2009). The detection of lies can be made by viewing the camera shot or from the prologue/epilogue accompanying each scene, which explicitly reveals the true state of affairs to one of the characters but not the other. To detect sarcasm, this supplemental information is helpful but not sufficient to make an accurate determination; subjects must also attend to and accurately process subtle changes in paralinguistic and other social cues from the characters' conversational exchange.

Higher scores on the social cognitive tasks indicate better performance with the exception of AIHQ. The AIHQ produces bias scores in which higher and lower scores indicate higher or lower levels of bias, respectively, toward attributing hostile intentions and blaming others in ambiguous social situations.

### 2.3 Neurocognitive assessment

The MATRICS Consensus Cognitive Battery (MCCB) (Nuechterlein and Green 2006) was used to assess general cognitive performance. It includes tests that assess seven domains of neurocognition including speed of processing, attention/vigilance, working memory, verbal learning, visual learning, reasoning and problem solving, and social cognition (Kern et al., 2008; Nuechterlein et al., 2008). Because the goal of the study was to look at specialized measures of social cognition and neurocognition separately, we used a modified composite score that did not include social cognition domain, but instead was based on the average T-scores from the six remaining domains. Higher scores indicate better performance.

## 2.4 Symptom assessment

1. Expanded 24-item *Brief Psychiatric Rating Scale* (BPRS) (Lukoff et al., 1986; Overall and Gorham, 1962) was used to evaluate psychiatric symptoms during the previous month. The current study used four subscales: *Positive*, *Negative*, *Depression-Anxiety*, and *Agitation* (Kopelowicz et al., 2008).
2. *Scale for the Assessment of Negative Symptoms* (SANS) (Andreasen, 1984) was used to assess negative symptoms during the month preceding the interview: *Affective flattening*, *Alogia*, *Anhedonia-Asociality*, and *Avolition-Apathy*. All BPRS and SANS interviewers were trained to a minimum intraclass correlation coefficient of 0.80 using established procedures (Ventura et al., 1993, 1998) by the Treatment Unit of the VISN 22 MIRECC and participated in an on-going quality assurance program. Higher ratings on the BPRS and SANS indicate more severe levels of symptoms.

## 2.5 Functional outcome assessment

1. Functional capacity was assessed using two scales: (a) The *UCSD Performance-based Skills Assessment* (UPSA) (Patterson et al., 2001), which involves role-play tests with props that are administered as simulations of events in the areas of general organization, finance, social/communications, transportation, and household chores. An overall summary score was used. (b) The *Maryland Assessment of Social Competence* (MASC) (Bellack et al., 1994), a measure of subjects' ability to solve common problems in an interpersonal context, consists of four 3-min role play scenarios, including involving initiating conversation with a casual acquaintance, two involving negotiation and compromise, and one involving standing up for one's rights. The interactions were videotaped and subsequently scored by specially trained and blinded raters. The Overall Effectiveness rating (a composite measure of the ability to maintain focus and achieve the goal of the scenario) was used.
2. Real-world functioning was assessed with The Community Adjustment Form (Stein and Test, 1980) is a semi-structured interview that evaluate real-world functioning. This measure covers 17 domains of psychosocial functioning and community adjustment. Information derived from the interview was used complete the *Role Functioning Scale* (RFS) for the domains of work functioning, independent living, and social functioning (McPheeters, 1984). The three RFS subscales moderately inter-correlated from .19 to .36; we therefore opted to evaluate them separately rather than compute a composite score. Higher scores on the UPSA, MASC, and RFS indicate better functioning.

## 2.7 Statistical analysis

Analyses were conducted in three stages: (1) The structure of the eight social cognitive indexes was assessed through correlational analyses followed by an exploratory factor analysis. We used the Maximum Likelihood extraction method because, in general, it provides better estimates than other approaches (Fabringar et al., 1999). We selected the number of factors using the scree test, which is superior to other commonly used methods of determining the number of factors (Velicer and Jackson, 1990). We chose an oblique rotation algorithm since we expected that any factors identified would likely be inter-correlated (rather than orthogonal); the individual variables from the social cognitive tests are believed to tap an over-arching social cognition dimension. (2) Correlates of the underlying dimensions derived from the factor analysis were examined. Indexes were assigned to no more than one factor based on the strength of their loadings. Factor scores were computed by first converting each social cognitive index to a z-score and then

calculating the average of the z-scored indexes that corresponded to each factor. These factor scores were then correlated with measures of neurocognition, symptoms, and functional outcome. (3) The incremental validity of social cognition in predicting functional outcome was examined with regression analyses, focusing only on the social cognitive factor scores and specific functional outcome measures that demonstrated significant inter-correlations. Multiple regression analyses examined whether the social cognitive factor scores predicted significant incremental variance ( $R^2$ ) in outcome after accounting for neurocognition and any symptoms that showed significant correlations to the relevant factors. All tests were two-tailed and results were regarded as significant below the 5% level of probability.

### 3. Results

#### 3.1 Sample characteristics and descriptive statistics

Demographic and clinical variables are presented in Table 1. This sample was chronically ill and predominantly male. Symptom levels were generally in the low to moderate range. Table 2 displays descriptive statistics for each of the social cognitive, neurocognitive, and functional outcome variables. Mean scores resemble those found in prior studies of stabilized outpatient samples (e.g., Horan et al., 2009; Roberts and Penn, 2009).

#### 3.2 Correlational and factor analysis of the social cognitive measures

Table 3 presents the correlations among the eight social cognitive indexes. The magnitude of correlations was highly variable, ranging from  $-.09$  to  $.64$ . Scores on the AIHQ showed minimal correlations with other measures. There was also a notable absence of significant correlation between the Sarcasm and Lies indexes from the TASIT.

In a factor analysis of these eight indexes (Table 4), the eigenvalue-greater-than-one rule and screen plot converged on a three-factor solution that accounted for 53.6% of the total variance. All the AIHQ indexes loaded on the first factor, which was labeled “Hostile attributional style”. Factor 2 comprised Facial Emotion Identification, PONS, and TASIT Lie scores. These tasks share a relatively low level reliance on perceiving cues and understanding information directly presented to subjects in the task stimuli, and the factor was labeled “Lower-level social cue detection”. The third factor comprised TASIT Sarcasm and MSCEIT Managing emotions, which require relatively high-level processes involved in incorporating and utilizing socio-emotional information and knowledge that is not directly present in the stimuli. This factor was labeled, “Higher-level inferential and regulatory processing”. The factors weakly intercorrelated: Factor 1 correlated  $-.07$  with Factor 2 and  $.08$  with Factor 3. Factor 2 correlated  $.26$  with Factor 3.

#### 3.3 Correlations with neurocognition, symptoms, and functional outcome

Correlations between the three social cognitive factors scores and the other study measures are shown in Table 5. There were some commonalities across the factors: None of the factors showed any correlations with negative symptoms, either assessed with the BPRS or the SANS. The factors, however, showed different patterns in other respects. Higher scores on Factor 1 (i.e., greater tendency to attribute hostility and blame to others) showed significant correlations with clinical symptoms (positive, depression-anxiety, and agitation on the BPRS) but not with functional capacity or real-world functioning. In contrast, Factors 2 and 3 showed the opposite pattern: no significant associations with clinical symptoms, but several significant relationships to functional capacity and real-world functioning. Another difference is that, although each social cognitive factor significantly correlated with the modified MCCB composite, the magnitude was significantly larger for Factor 2 compared to either Factor 1 or Factor 3 (both  $p$  values  $< .05$ , using Steiger’s  $Z$ ). Similarly, Factor 2



tended to show larger correlations with functional capacity compared to the other factors, but the comparisons only reached significance for the contrast of Factor 2 versus Factor 1 on the UPSA ( $p < .01$ ).

### 3.4 Incremental validity analyses

Factors 2 and 3 were carried forward into two incremental validity analysis because of their significant relationships to functional capacity (UPSA, MASC) and real-world functioning (RFS Work, RFS Social) in the previous correlational analysis. We evaluated whether neurocognitive and symptom levels (SANS, BPRS) significantly correlated with these functional outcome variables and should therefore also be included in the incremental validity analyses. Modified MCCB scores significantly correlated with UPSA ( $r = .54, p < .01$ ), MASC ( $r = .26, p < .05$ ), and Work functioning ( $r = .25, p < .05$ ). SANS total scores significantly correlated with UPSA ( $r = -.23, p < .05$ ), MASC ( $r = -.38, p < .01$ ), Independent Living ( $r = -.30, p < .01$ ), and Social functioning ( $r = -.51, p < .01$ ). BPRS total scores were not significantly correlated with any of the functional outcome measures (all  $r$ 's  $< .20$ ). MCCB and SANS totals scores were therefore included in the following incremental validity analyses.

Two sets of multiple regression analyses were conducted – one set for Factor 2 and another for Factor 3. In these analyses, each functional outcome index served as the dependent variable with three blocks of predictor variables entered in the following order: Block 1 = modified MCCB; Block 2 = SANS; Block 3= Either Factor 2 or Factor 3. Results for the Factor 2 analyses are shown in Table 6. For the UPSA, after accounting for neurocognition and negative symptoms Factor 2 accounted for an additional 8% ( $R^2$  change) of the variance, which was statistically significant. Similarly, for the MASC, Factor 2 accounted for a significant additional 9% of the variance. Factor 2 did not account for significant incremental variance in RFS Work or Social functioning, though there was a non-significant trend for Social functioning ( $p = .067$ ).

Results for Factor 3 are presented in Table 7. After accounting for neurocognition and negative symptoms, Factor 3 did not contribute significant incremental variance for any of the functional outcome variables. There was a trend-level finding for the MASC ( $p = .077$ ).

## 4. Discussion

Exploratory factor analysis of variables from five tests covering a wide range of social cognitive processes yielded three interpretable and weakly intercorrelated factors in outpatients with psychosis. The factors demonstrated evidence of external validity in their differential patterns of correlation with neurocognitive, clinical symptom, and functional outcome measures. The Hostile attributional style factor significantly correlated with clinical symptoms (positive, depression/anxiety, agitation) but not functional outcome, whereas The Lower-level social cue detection and Higher-level inferential and regulatory processes factors significantly correlated with functional outcome (functional capacity and real-world social and work functioning) but not clinical symptoms. Furthermore, aspects of social cognition had added value in predicting functional capacity above and beyond non-social neurocognition and symptoms. A multidimensional conceptualization of social cognition can provide a useful organizational and guiding framework for this rapidly growing area of research in psychosis.

This is the first study, to our knowledge, to conduct an exploratory factor analysis on social cognitive domains that are commonly studied in major psychopathology. The three Hostile attributional style indexes (AIHQ Hostility, Aggression and Blame) loaded on the first factor, which showed little to no correlation with the other social cognitive variables. The

second factor was labeled “Lower-level social cue detection” reflecting basic emotion detection skills (recognition of emotions in faces), low level of cognitive processing (interpretation of non-verbal information transmitted by others), and first order mental representation (detection of lies) (Peskin et al., 1996; Sullivan et al., 1995). This factor showed the highest correlation with neurocognition. The third factor was labeled “Higher-level inferential and regulatory processing” and was reflected in more refined emotional skills (ability to manage subjective emotional states), higher social cognitive functions and second order mental representation (detection of sarcasm) (Winner and Leekam, 1991; Happe, 1993). These findings support the value of considering social cognition as a multidimensional construct with hierarchically distinct lower-level and higher-level abilities (Ochsner, 2008).

Our results also showed that detection of lies and sarcasm loaded on separate factors. Several studies of school-age children have demonstrated that comprehension of deceit and irony (e.g., sarcasm) are qualitatively distinct abilities (Bara et al., 1999; Bosco and Bucciarelli, 2008; Lee and Katz, 1998). Comprehension of lies/deceit is acquired before sarcasm/irony and is based on a less complex inferential chain (Bucciarelli et al., 2003; Winner et al., 1988). We found that detection of sarcasm and managing emotions loaded on the same factor, suggesting that comprehension of sarcasm requires refined emotional skills such as empathic appreciation of the listener's emotional state (Shamay-Tsoory et al., 2005). This distinction between lies and sarcasm may have been enhanced by the way that the TASIT evaluates detection of lies – it provides all of the information about deceit in the scene and does not require much inference. A more subtle test of lie detection may have yielded a different factor structure.

In the current study, only certain aspects of social cognition showed significant relationships with clinical symptoms. No correlations were found between the three social cognitive factors and negative symptoms suggesting that social cognition and negative symptoms are largely separate constructs (Rassovsky et al., in press; Sergi et al., 2007). Also, some authors have demonstrated that the correlations between social cognition and negative symptoms can be attributed to confounding variables such as intellectual deficits or duration of illness (Langdon 2002; Pousa et al., 2008). Only the Hostile attributional style factor showed significant relations to other types of symptoms. Similar to other studies, higher tendencies to blame and respond with hostility in ambiguous social situations significantly correlated with positive symptoms (e.g., An et al., 2010; Combs et al., 2009; Janssen et al., 2006). These attributional tendencies also showed more general linkages to higher levels of depression/anxiety and agitation. Thus, the social cognitive domain of Hostile attributional style was more closely tied to indicators of clinical symptom state than the other factors and showed no significant relations to any aspect of functional outcome.

The Lower-level social cue detection and Higher-level inferential and regulatory processes factors were distinguished from the Hostile attributional style factor by a different pattern of external correlates, namely, significant relations with both functional capacity and real-world functioning but no significant relations to positive, depression/anxiety, or agitation. These findings converge with growing evidence that various aspects of social cognition show meaningful relations to both competence and performance measures of functional outcome (Couture et al., 2006; Horan et al., in press). This study also explored the associations between functional outcome and social cognition versus neurocognition, another key correlate of functional outcome. Notably, Lower-level social cue detection factor demonstrated a significantly larger correlation with neurocognition than did Higher-level inferential and regulatory processes factor, providing some evidence of differential relations between these social cognitive factors and external variables. In line with prior studies (e.g., Couture et al., 2006; Pan et al., 2009), the Lower-level social cue detection



factor accounted for additional variance in functional capacity above and beyond neurocognition and negative symptoms (8% for UPSA and 9% for MASC), demonstrating the “added value” of social cognition. However, we did not find evidence for incremental validity in real-world functioning, although other studies have (Brekke et al., 2005; Poole et al., 2000; Vauth et al., 2004). As discussed in recent integrative models of functional outcome (Bowie et al., 2008, Horan et al., 2010, Rassovsky et al., in press), neurocognitive and social cognitive competence appear more proximal to functional capacity than to real-world functioning, rendering unique relations to functional capacity simpler to demonstrate. Relations to the more distal outcome of real-world functioning can be more challenging to detect due to the various personal (e.g., motivation, self-efficacy) and socio-environmental (e.g., disability policies, cultural factors) variables that impact how one performs in the community.

The current study provides evidence for the multidimensional structure of social cognition in outpatients with psychosis. These findings should be considered in the context of several methodological factors that may limit generalizability, including a largely VA-based sample (mostly older male patients with a slightly skewed ethnic distribution), enrolling patients with three different clinically-determined psychotic diagnoses, and a lack of control for medication effects. Also, the generally low to moderate level of symptoms in our sample may have limited our ability to detect significant relations to symptoms. Nevertheless, a better understanding of the factor structure of social cognition in schizophrenia was a stated goal of a NIMH consensus meeting on this topic (Green et al., 2008) and has implications for how we interpret findings in this area. As mentioned above, social cognition has been studied in schizophrenia to better understand clinical symptoms and to better understand daily functioning. The results of these analyses suggest that social cognition in general serves both of these purposes, but specific factors serve one or the other. Even the well-documented relationships between social cognition and neurocognition are more characteristic of one factor (lower-level processes) than others, suggesting that Hostile attributional style and Higher-level inferential and regulatory processes factors involve other types of determinants. Hence, these results provide an initial step to help parse and organize this complex and rapidly-developing area.

## Acknowledgments

The authors thank Gerhard Helleman for statistical consultation and Mark McGee, Cory Tripp, Poorang Nori, and Lisa Mancini for assistance with data collection.

### Role of Funding Source

Funding for this project came from a Veterans Affairs Merit Grant and NIMH grant MH043292 (both to Dr. Green), and the VA VISN-22 Mental Illness Research Education Clinical Center. The VA had no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

## References

- Adolphs R. The social brain: Neural basis of social knowledge. *Annu. Rev. Psychol.* 2009; 60:693–716. [PubMed: 18771388]
- Ambady N, Hallahan M, Rosenthal R. On judging and being judged accurately in zero-acquaintance situations. *J. Pers. Soc. Psychol.* 1995; 69:519–529.
- An SK, Kang JI, Park JY, Kim KR, Lee SY, Lee E. Attribution bias in ultra-high risk for psychosis and first-episode schizophrenia. *Schizophr. Res.* 2010; 118(1–3):54–61. [PubMed: 20171849]
- Andreasen, NC. The scale for the assessment of negative symptoms (SANS). Iowa City, IA: The University of Iowa; 1984.

- Bara, BG.; Bosco, FM.; Bucciarelli, M. Simple and complex speech acts: what makes the difference within a developmental perspective?. In: Hahn, M.; Stoness, SC., editors. Proceedings of the XXI Conference of the Cognitive Science Society; Mahwah, NJ. Lawrence Erlbaum Associates; 1999. p. 55-60.
- Bell M, Tsang HW, Greig TC, Bryson GJ. Neurocognition, social cognition, perceived social discomfort, and vocational outcomes in schizophrenia. *Schizophr. Bull.* 2009; 35:738–747. [PubMed: 18245058]
- Bellack AS, Sayers M, Mueser KT, Bennett M. Evaluation of social problem solving in schizophrenia. *J. Abnorm. Psychol.* 1994; 103:371–378. [PubMed: 8040506]
- Bentall RP, Corcoran R, Howard R, Blackwood N, Kinderman P. Persecutory delusions: a review and theoretical integration. *Clin. Psychol. Rev.* 2001; 21:1143–1192. [PubMed: 11702511]
- Bosco FM, Bucciarelli M. Simple and complex deceptions and ironies. *J. Pragmat.* 2008; 40:583–607.
- Bowie CR, Reichenberg A, Patterson TL, Heaton RK, Harvey PD. Determinants of real-world functional performance in schizophrenia subjects: Correlations with cognition, functional capacity, and symptoms. *Am. J. Psychiatry.* 2006; 163:418–425. [PubMed: 16513862]
- Bowie CR, Leung WW, Reichenberg A, McClure MM, Patterson TL, Heaton RK, Harvey PD. Predicting schizophrenia patients real-world behavior with specific neuropsychological and functional capacity measures. *Biol. Psychiatry.* 2008; 63:505–511. [PubMed: 17662256]
- Brekke JS, Kay DD, Kee KS, Green MF. Biosocial pathways to functional outcome in schizophrenia. *Schizophr. Res.* 2005; 80:213–225. [PubMed: 16137859]
- Bucciarelli M, Colle L, Bruno G, Bara BG. How children comprehend speech acts and communicative gestures. *J. Pragmat.* 2003; 35:207–241.
- Combs DR, Penn DL, Wicher M, Waldheter E. The Ambiguous Intentions Hostility Questionnaire (AIHQ): A new measure for evaluating hostile social-cognitive biases in paranoia. *Cogn. Neuropsychiatry.* 2007; 12(2):128–143. [PubMed: 17453895]
- Combs DR, Penn DL, Michael CO, Basso MR, Wiedeman R, Siebenmorgan M, Tiegreen J, Chapman D. Perceptions of hostility by persons with and without persecutory delusions. *Cogn. Neuropsychiatry.* 2009; 14(1):30–52. [PubMed: 19214841]
- Corcoran R, Rowse G, Moore R, Blackwood N, Kinderman P, Howard R, Cummins S, Bentall RP. A transdiagnostic investigation of 'theory of mind' and 'jumping to conclusions' in patients with persecutory delusions. *Psychol. Med.* 2008; 38(11):1577–1583. [PubMed: 18005499]
- Couture SM, Penn DL, Roberts DL. The functional significance of social cognition in schizophrenia: a review. *Schizophr. Bull.* 2006; 32 Suppl 1:S44–S63. [PubMed: 16916889]
- Eack SM, Pogue-Geile MF, Greeno CG, Keshavan MS. Evidence of factorial variance of the Mayer-Salovey-Caruso Emotional Intelligence Test across schizophrenia and normative samples. *Schizophr. Res.* 2009; 114(1–3):105–109. [PubMed: 19501486]
- Ekman, P. Subtle Expression Training Tool (SETT) and Micro Expression Training Tool (METT): Paul Ekman. 2004. [www.paulekman.com](http://www.paulekman.com).
- Fabrigar LR, Wegener DT, MacCallum RC, Strahan EJ. Evaluating the use of exploratory factor analysis in psychological research. *Psychol. Methods.* 1999; 4:272–299.
- Fett AK, Viechtbauer W, Dominguez MD, Penn DL, van Os J, Krabbendam L. The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: A meta-analysis. *Neurosci. Biobehav. Rev.* (in press).
- Fornells-Ambrojo M, Garety PA. Attributional biases in paranoia: the development and validation of the Achievement and Relationships Attributions Task (ARAT). *Cogn. Neuropsychiatry.* 2009; 14:87–109. [PubMed: 19370434]
- Green MF, Horan WP. Social cognition in schizophrenia. *Curr. Dir. Psychol. Sci.* 2010; 19:243–248.
- Green MF, Olivier B, Crawley JN, Penn DL, Silverstein S. Social cognition in schizophrenia: recommendations from the measurement and treatment research to improve cognition in schizophrenia new approaches conference. *Schizophr. Bull.* 2005; 31(4):882–887. [PubMed: 16135561]
- Green MF, Penn DL, Bentall R, Carpenter WT, Gaebel W, Gur RC, Kring AM, Park S, Silverstein SM, Heinssen R. Social cognition in schizophrenia: an NIMH workshop on definitions,

- assessment, and research opportunities. *Schizophr. Bull.* 2008; 34(6):1211–1220. [PubMed: 18184635]
- Happé FGE. Communicative competence and theory of mind in autism: a test of relevance theory. *Cognition.* 1993; 48:101–119. [PubMed: 8243028]
- Horan WP, Rasseovsky Y, Kern RS, Lee J, Wynn JK, Green MF. Further Support For the Role of Dysfunctional Attitudes in Models of Real-World Functioning in Schizophrenia. *J. Psychiatr. Res.* 2010; 44:499–505. [PubMed: 20006849]
- Horan, WP.; Kern, RF.; Olivier, PO.; Green, MF. Neurocognition, social cognition, and functional outcome in schizophrenia. In: Gaebel, W., editor. *Schizophrenia -Current Science and Clinical Practice.* (in press)
- Horan WP, Kern RS, Shokat-Fadai K, Sergi MJ, Wynn JK, Green MF. Social cognitive skills training in schizophrenia: an initial efficacy study of stabilized outpatients. *Schizophr Res.* 2009; 107:47–54. [PubMed: 18930378]
- Horan WP, Kern RS, Tripp C, Helleman G, Wynn JK, Bell M, Marder SR, Green MF. Efficacy and Specificity of Social Cognitive Skills Training for Outpatients with Psychotic Disorders. (submitted).
- Janssen I, Versmissen D, Campo JA, Myin-Germeys I, van Os J, Krabbendam L. Attributional style and psychosis: Evidence for an externalizing bias in patients but not in individuals at high risk. *Psychol. Med.* 2006; 36:771–778. [PubMed: 16563206]
- Kern RS, Nuechterlein KH, Green MF, Baade LE, Fenton WS, Gold JM, Keefe RS, Mesholam-Gately R, Mintz J, Seidman LJ, Stover E, Marder SR. The MATRICS Consensus Cognitive Battery, part 2: co-norming and standardization. *Am. J. Psychiatry.* 2008; 165:214–220. [PubMed: 18172018]
- Kern RS, Green MF, Fiske AP, Kee KS, Lee J, Sergi MJ, Horan WP, Subotnik KL, Sugar CA, Nuechterlein KH. Theory of mind deficits for processing counterfactual information in persons with chronic schizophrenia. *Psychol. Med.* 2009; 39:645–654. [PubMed: 18694537]
- Kinderman P, Bentall RP. A new measure of causal locus: the internal, personal, and situational attributions questionnaire. *Pers. Individ. Dif.* 1996; 20:261–264.
- Kohler CG, Walker JB, Martin EA, Healey KM, Moberg PJ. Facial Emotion Perception in Schizophrenia: A Meta-analytic Review. *Schizophr Bull.* 2009 Vol and N!!!
- Kopelowicz A, Ventura J, Liberman RP, Mintz J. Consistency of Brief Psychiatric Rating Scale factor structure across a broad spectrum of schizophrenia patients. *Psychopathology.* 2008; 41(2):77–84. [PubMed: 18033976]
- Langdon R, Coltheart M, Ward PB, Catts SV. Disturbed communication in schizophrenia: the role of poor pragmatics and poor mind-reading. *Psychol. Med.* 2002; 32(7):1273–1284. [PubMed: 12420896]
- Lee C, Katz A. The differential role of ridicule in sarcasm and irony. *Metaphor and Symbol.* 1998; 13:1–15.
- Lukoff D, Nuechterlein KH, Ventura J. Manual for the Expanded Brief Psychiatric Rating Scale (BPRS). *Schizophr. Bull.* 1986; 12:594–602.
- Mayer, JD.; Salovey, P.; Caruso, DR. Mayer–Salovey–Caruso Emotional Intelligence Test (MSCEIT): User's Manual. Toronto, Ontario: Multi-Health Systems, Inc.; 2002.
- Mayer JD, Salovey P, Caruso DR, Sitarenios G. Measuring emotional intelligence with the MSCEIT V2.0. *Emotion.* 2003; 3:97–105. [PubMed: 12899321]
- McDonald, S.; Flanagan, S.; Rollins, J. The Awareness of Social Inference Test. Suffolk, UK: Thames Valley Test Company, Ltd.; 2002.
- McPheeters HL. Statewide mental health outcome evaluation: a perspective of two southern states. *Community Ment. Health J.* 1984; 20:44–55. [PubMed: 6723258]
- Nuechterlein, KH.; Green, MF. MATRICS Consensus Cognitive Battery. Los Angeles: MATRICS Assessment, Inc.; 2006.
- Nuechterlein KH, Green MF, Kern RS, Baade LE, Barch DM, Cohen JD, Essock S, Fenton WS, Frese FJ 3rd, Gold JM, Goldberg T, Heaton RK, Keefe RS, Kraemer H, Mesholam-Gately R, Seidman LJ, Stover E, Weinberger DR, Young AS, Zalcman S, Marder SR. The MATRICS Consensus Cognitive Battery, part 1: test selection, reliability, and validity. *Am. J. Psychiatry.* 2008; 165:203–213. [PubMed: 18172019]

- Ochsner KN. The social-emotional processing stream: five core constructs and their translational potential for schizophrenia and beyond. *Biol. Psychiatry*. 2008; 64:48–61. [PubMed: 18549876]
- Overall JE, Gorham DR. The Brief Psychiatric Rating Scale. *Psychol. Rep.* 1962; 10:799–812.
- Pan YJ, Chen SH, Chen WJ, Liu SK. Affect recognition as an independent social function determinant in schizophrenia. *Compr. Psychiatry*. 2009; 50:443–452. [PubMed: 19683615]
- Patterson TL, Goldman S, McKibbin CL, Hughs T, Jeste DV. UCSD Performance-Based Skills Assessment: development of a new measure of everyday functioning for severely mentally ill adults. *Schizophr. Bull.* 2001; 27(2):235–245. [PubMed: 11354591]
- Peskin J. Guise and guile: Children's understanding of narratives in which the purpose of pretense in deception. *Child Dev.* 1996; 67:1735–1751.
- Poole JH, Tobias FC, Vinogradov S. The functional relevance of affect recognition errors in schizophrenia. *J. Int. Neuropsychol. Soc.* 2000; 6(6):649–658. [PubMed: 11011511]
- Pousa E, Duñó R, Brébion G, David AS, Ruiz AI, Obiols JE. Theory of mind deficits in chronic schizophrenia: evidence for state dependence. *Psychiatry Res.* 2008; 158(1):1–10. [PubMed: 18166230]
- Roberts DL, Penn DL. Social cognition and interaction training in outpatients with schizophrenia: A preliminary study. *Psychiatry Res.* 2009; 166:141–147. [PubMed: 19272654]
- Rassovsky Y, Horan WP, Lee J, Sergi MJ, Green MF. Pathways between early visual processing and functional outcome in schizophrenia. *Psychol. Med.* (In press).
- Rosenthal, R.; Hall, JA.; DiMatteo, MR.; Rogers, PL.; Archer, D. Sensitivity to nonverbal communication: The PONS test. Baltimore: Johns Hopkins University Press; 1979.
- Sergi MJ, Rassovsky Y, Widmark C, Reist C, Erhart S, Braff DL, Marder SR, Green MF. Social cognition in schizophrenia: relationships with neurocognition and negative symptoms. *Schizophr. Res.* 2007; 90(1–3):316–324. [PubMed: 17141477]
- Shamay-Tsoory SG, Tomer R, Aharon-Peretz J. The neuroanatomical basis of understanding sarcasm and its relationship to social cognition. *Neuropsychology*. 2005; 19(3):288–300. [PubMed: 15910115]
- Shamay-Tsoory SG, Shur S, Barcai-Goodman L, Medlovich S, Harari H, Levkovitz Y. Dissociation of cognitive from affective components of theory of mind in schizophrenia. *Psychiatry Res.* 2007; 149(1–3):11–23. [PubMed: 17107716]
- Stein LI, Test MA. Alternatives to mental hospital treatment: I. Conceptual model treatment program and clinical evaluation. *Arch. Gen. Psychiatry*. 1980; 37:392–397. [PubMed: 7362425]
- Sullivan K, Winner E, Hopfield N. How children tell a lie from a joke: The role of second-order mental state attributions. *Br. J. Dev. Psychol.* 1995; 13:191–204.
- van Hooren S, Versmissen D, Janssen I, Myin-Germeys I, à Campo J, Mengelers R, van Os J, Krabbendam L. Social cognition and neurocognition as independent domains in psychosis. *Schizophr. Res.* 2008; 103(1–3):257–265. [PubMed: 18434094]
- Vauth R, Rüschi N, Wirtz M, Corrigan PW. Does social cognition influence the relation between neurocognitive deficits and vocational functioning in schizophrenia? *Psychiatry Res.* 2004; 128(2): 155–165. [PubMed: 15488958]
- Velicer WF, Jackson DN. Component analysis versus common factor-analysis – some further observations. *Multivariate Behav. Res.* 1990; 25(1):97–114.
- Ventura J, Green MF, Shaner A, Liberman RP. Training and quality assurance with the brief psychiatric rating scale: 'The Drift Busters'. *Int. J. Methods Psychiatr. Res.* 1993; 3:221–224.
- Ventura J, Liberman RP, Green MF, Shaner A. Training and quality assurance with the structured clinical interview for DSM-IV. *Psychiatry Res.* 1998; 79:163–173. [PubMed: 9705054]
- Winner, E.; Levy, J.; Kaplan, J.; Rosenblatt, E. Children's understanding of nonliteral language. In: Gardner, H.; Perkins, D., editors. *Art, Mind, and Education: Research from Project Zero*. Urbana, IL: University of Illinois Press; 1988.
- Winner E, Leekman S. Distinguishing irony from deception: understanding the speaker's second-order intention. *Br. J. Dev. Psychol.* 1991; 9:257–270.

Woodward TS, Mizrahi R, Menon M, Christensen BK. Correspondences between theory of mind, jumping to conclusions, neuropsychological measures and the symptoms of schizophrenia. *Psychiatry Res.* 2009; 170(2–3):119–123. [PubMed: 19906438]

**Table 1**

Demographic and clinical characteristics (N=85).

	%	M (SD)
<b>Age</b>		48.50 (8.6)
<b>Education</b>		
Participant		12.88 (1.84)
Father		12.85 (3.4)
Mother		12.48 (3.6)
<b>Age of onset</b>		22.26 (8.36)
<b>Duration of illness</b>		25.96 (10.85)
<b>Sex (male)</b>	89.4	
<b>Ethnicity</b>		
White	30.6	
Latino	12.9	
Asian	2.4	
Black	52.9	
Other	1.2	
<b>Diagnosis</b>		
Schizophrenia	68.2	
Schizoaffective	22.4	
Psychosis NOS	9.4	
<b>Symptoms</b>		
BPRS Positive		1.92 (0.75)
BPRS Depression-Anxiety		1.98 (0.81)
BPRS Negative		2.0 (0.91)
BPRS Agitation		1.24 (0.37)
BPRS Total		1.78 (0.42)
SANS Affect		1.83 (1.29)
SANS Alogia		0.65 (0.98)
SANS Apathy		3.05 (1.03)
SANS Anhedonia		2.62 (1.13)
SANS Total		1.58 (0.07)

Notes: BPRS = Brief Psychiatric Rating Scale; SANS = Scale for the Assessment of Negative Symptoms.



**Table 2**Descriptive statistics for the social cognitive, neurocognitive, and functional outcome measures ( $n = 85$ )

	Mean	SD
<b>FEIT</b>	39.79	7.02
<b>PONS</b>	76.41	7.29
<b>AIHQ Hostility</b>	1.80	.53
<b>AIHQ Aggression</b>	1.85	.52
<b>AIHQ Blame</b>	2.97	.96
<b>TASIT Lies</b>	24.58	4.07
<b>TASIT Sarcasm</b>	20.94	5.90
<b>MSCEIT- ME</b>	38.61	10.20
<b>Modified MCCB</b>	37.74	7.39
<b>UPSA</b>	73.97	10.52
<b>MASC</b>	3.30	.78
<b>RFS Work</b>	1.94	1.37
<b>RFS Independent living</b>	4.15	1.60
<b>RFS Social</b>	3.61	1.89

Notes: FEIT = Facial Emotion Identification Test; PONS = Half-Profile of Nonverbal Sensitivity; AIHQ = Ambiguous Intentions Hostility Questionnaire – Aggression, Hostility and Blame scores, respectively; TASIT = The Awareness of Social Inference Test; MSCEIT ME= Mayer-Salovey-Caruso Emotional Intelligence Test - Managing Emotions branch; MCCB = MATRICS Consensus Cognitive Battery - modified composite score; UPSA= UCSD Performance-Based Skills Assessment; MASC = Maryland Assessment of Social Competence; RFS = Role Functioning Scale.

Table 3

Correlations among the social cognitive measures

	FEIT	PONS	AIHQ Hostility	AIHQ Aggression	AIHQ Blame	TASIT Lies	TASIT Sarcasm	MSCEIT - ME
FEIT	1							
PONS	.64***	1						
AIHQ Hostility	-.16	-.15	1					
AIHQ Aggression	.07	.04	.31**	1				
AIHQ Blame	-.09	-.08	.60***	.52***	1			
TASIT Lies	.43***	.44***	-.09	.01	.03	1		
TASIT Sarcasm	.27*	.25*	.20	-.11	.09	.17	1	
MSCEIT - ME	.35***	.28*	.12	-.06	-.04	.31**	.41***	1

Notes: FEIT = Facial Emotion Identification Test; PONS = Half-Profile of Nonverbal Sensitivity; AIHQ = Ambiguous Intentions Hostility Questionnaire – Aggression, Hostility and Blame scores, respectively; TASIT = The Awareness of Social Inference Test; MSCEIT ME= Mayer-Salovey-Caruso Emotional Intelligence Test – Managing Emotions branch.

\* p&lt; 0.05;

\*\* p&lt; 0.01;

\*\*\* p&lt; 0.001

**Table 4**

Factor analysis of the social cognitive measures

	Factor 1	Factor 2	Factor 3
<b>FEIT</b>	-.07	<b>.80</b>	.26
<b>PONS</b>	-.06	<b>.80</b>	.23
<b>AIHQ Hostility</b>	<b>.65</b>	-.20	.35
<b>AIHQ Aggression</b>	<b>.64</b>	.07	-.14
<b>AIHQ Blame</b>	<b>.88</b>	-.08	.12
<b>TASIT Lies</b>	.03	<b>.56</b>	.25
<b>TASIT Sarcasm</b>	.05	.28	<b>.68</b>
<b>MSCEIT - ME</b>	-.04	.38	<b>.57</b>

Notes: Items assigned to each factor appear in bold. FEIT = Facial Emotion Identification Test; PONS = Half- Profile of Nonverbal Sensitivity; AIHQ = Ambiguous Intentions Hostility Questionnaire; TASIT = The Awareness of Social Inference Test; MSCEIT ME = Mayer-Salovey-Caruso Emotional Intelligence Test - Managing Emotions branch.

**Table 5**

Correlations between social cognitive factors and neurocognition, symptoms, and functional outcome

	<b>Factor 1</b>	<b>Factor 2</b>	<b>Factor 3</b>
<b><u>Neurocognition</u></b>			
Modified MCCB	.22*	.51***	.29**
<b><u>Symptoms</u></b>			
BPRS Positive	.28*	.04	.09
BPRS Depression-Anxiety	.30**	.06	.07
BPRS Negative	-.02	.17	-.15
BPRS Agitation	.22*	-.16	.17
BPRS Total	.37**	.03	.07
SANS Affect	-.08	.11	-.14
SANS Alogia	.01	-.12	-.18
SANS Apathy	.03	-.21	-.20
SANS Anhedonia	.05	-.18	.09
SANS Total	.02	-.05	-.20
<b><u>Functional outcome</u></b>			
UPSA	.11	.52***	.34**
MASC	.10	.38**	.29*
RFS Work	.05	.26*	.21
RFS Independent living	-.07	.09	.10
RFS Social	-.12	.24*	.25*

Notes: MCCB = MATRICS Consensus Cognitive Battery - modified composite score; BPRS = Rating Scale; SANS = Scale for the Assessment of Negative Symptoms; UPSA= UCSD Performance- Assessment; MASC = Maryland Assessment of Social Competence; RFS = Role Functioning Scale.

\* p< 0.05;

\*\* p< 0.01;

\*\*\* p< 0.001

**Table 6**

Regression analyses for incremental prediction of functional outcome: Factor 2 (Lower-level social cue detection)

Dependent variable	Predictor	R <sup>2</sup>	R <sup>2</sup> Change	F Change
UPSA	<i>MCCB</i>	.29		32.07***
	<i>SANS</i>	.32	.03	3.75 <sup>†</sup>
	<i>Factor 2</i>	.40	.08	10.74**
MASC	<i>MCCB</i>	.06		4.79*
	<i>SANS</i>	.19	.13	11.19***
	<i>Factor 2</i>	.27	.09	8.40**
RFS Work	<i>MCCB</i>	.07		5.90*
	<i>SANS</i>	.09	.03	2.20
	<i>Factor 2</i>	.11	.01	1.27
RFS Social	<i>MCCB</i>	.03		2.56
	<i>SANS</i>	.28	.25	27.52***
	<i>Factor 2</i>	.31	.03	3.45 <sup>†</sup>

Notes: UPSA= UCSD Performance-Based Skills Assessment; MASC = Maryland Assessment of Social Competence; RFS = Role Functioning Scale; MCCB = MATRICS Consensus Cognitive Battery - modified composite score; SANS = Scale for the Assessment of Negative Symptoms.

\* p<0.05;

\*\* p<0.01;

\*\*\* p<0.001;

<sup>†</sup> p < .10.

**Table 7**

Regression analyses for incremental prediction of functional outcome: Factor 3 (Higher-level inferential and regulatory processes)

Dependent variable	Predictor	R <sup>2</sup>	R <sup>2</sup> Change	F Change
UPSA	<i>MCCB</i>	.29		32.07***
	<i>SANS</i>	.32	.03	3.75 <sup>†</sup>
	<i>Factor 3</i>	.35	.03	3.21 <sup>†</sup>
MASC	<i>MCCB</i>	.06		4.79*
	<i>SANS</i>	.19	.13	11.19***
	<i>Factor 3</i>	.21	.03	2.41
RFS Work	<i>MCCB</i>	.07		5.90*
	<i>SANS</i>	.09	.03	2.20
	<i>Factor 3</i>	.11	.01	1.25
RFS Social	<i>MCCB</i>	.03		2.56
	<i>SANS</i>	.28	.25	27.52***
	<i>Factor 3</i>	.29	.01	1.59

Notes: UPSA= UCSD Performance-Based Skills Assessment; MASC= Maryland Assessment of Social Competence; RFS = Role Functioning Scale; MCCB = MATRICS Consensus Cognitive Battery - modified composite score; SANS = Scale for the Assessment of Negative Symptoms.

\* p < 0.05;

\*\* p < 0.01;

\*\*\* p < 0.001;

<sup>†</sup> p < .10.