



Published in final edited form as:

Clin Psychol (New York). 2010 December ; 17(4): 293–306. doi:10.1111/j.1468-2850.2010.01221.x.

Comorbidity of Anxiety and Depression in Youth: Implications for Treatment and Prevention

Judy Garber and
Vanderbilt University

V. Robin Weersing
SDSU/UCSD Joint Doctoral Program in Clinical Psychology

Judy Garber: judy.garber@vanderbilt.edu

Abstract

The high level of concurrent and sequential comorbidity between anxiety and depression in children and adolescents may result from (a) substantial overlap in both the symptoms and items used to assess these putatively different disorders, (b) common etiological factors (e.g., familial risk, negative affectivity, information processing biases, neural substrates) implicated in the development of each condition, and (c) negative sequelae of anxiety conferring increased risk for the development of depression. Basic research on their various common and unique etiological mechanisms has guided the development of efficacious treatments for anxiety and depressive disorders in youth. Potential processes through which the successful treatment of childhood anxiety might prevent subsequent depression are described.

Anxiety and depression frequently co-occur both concurrently and sequentially in children and adolescents, and one often increases the risk of the other over time. The most common anxiety diagnoses in youth are separation anxiety disorder (SAD), social anxiety disorder (SOC), generalized anxiety disorder (GAD), and specific phobia (SP); depression diagnoses include major depressive disorder (MDD) and dysthymic disorder (DD). These various anxiety and depressive disorders appear together across generations, clustering strongly within families. Unpacking the whole of the internalizing comorbidity literature is beyond the scope of this paper (see Axelson & Birmaher, 2001; Brady & Kendall, 1992; Seligman & Ollendick, 1998). Rather, we address several key questions regarding comorbid anxiety and depression in youth. What is the extent of their concurrent and sequential overlap? What may account for the observed patterns of comorbidity between these disorders? What are the implications of their comorbidity for the treatment and prevention of anxiety and depression?

Extent of Concurrent and Sequential Comorbidity

Anxiety and mood symptoms and disorders in youth are distressing, impairing, and prevalent (e.g., Costello et al., 2003), interfere with interpersonal relationships and academic achievement, and increase the risk of suicide and other psychopathology (e.g., Gould et al., 1998; Rohde, Lewinsohn, & Seeley, 1994). Moreover, such negative effects may propagate forward into adulthood (Rohde et al., 1994; Weissman et al., 1999). The prognosis for comorbid anxiety and depression is worse than either condition alone, with higher risk of recurrence, longer duration, increased suicide attempts, greater impairment, less favorable response to treatment, and greater utilization of mental health services (Birmaher et al., 1996; Ezpeleta, Domenech, & Angold, 2006).

An estimated 15% to 20% of youth in the United States meet criteria for any anxiety disorder (i.e., SAD: 2.8% to 8%, SP: 10%, SOC: 7%, panic disorder: 1%–3%, GAD: 4.3%) (Beesdo, Knapp, & Pine, 2009), and approximately 1 in 5 youth will have an episode of MDD by age 18 (Lewinsohn et al., 1993). The level of comorbidity between these common problems is substantial, as evidenced by both high correlations between dimensional measures of anxious and depressive symptoms (e.g., Cole, Peeke, Martin, Truglio, & Seroczynski, 1997; Stark & Laurent, 2001) and diagnostic comorbidity rates as high as 75% in some clinical samples (Sorensen, Nissen, Mors, & Thomsen, 2005; Weersing, Gonzalez, Campo, & Lucas, 2008).

Interestingly, this high degree of comorbidity does not appear to be symmetrical. In community samples, 25% to 50% of youth with depression also meet criteria for an anxiety disorder, whereas only 10% to 15% of those with a primary anxiety disorder have a concurrent depressive disorder (Angold et al., 1999; Axelson & Birmaher, 2001; Costello et al., 2003). Thus, youth with primary depressive disorders tend to have comorbid anxiety more often than do those with primary anxiety disorders have comorbid depression (Merikangas & Avenevoli, 2002; Ollendick, Shortt, & Sander, 2005).

Several factors might explain this apparent imbalance. First, subsyndromal levels of symptoms often have not been assessed in studies of comorbidity. That is, children diagnosed with anxiety disorders may have concurrent depressive symptoms even if they do not meet full criteria for a depressive diagnosis, and these co-occurring, subthreshold symptoms may account for the apparent link between anxiety and subsequent depressive disorder in adolescence. Indeed, sub-diagnostic depressive symptoms have been found to be a more reliable predictor of subsequent depressive disorders than symptoms of either separation or social anxiety (Keenan, Feng, Hipwell, & Klostermann, 2009).

Second, anxiety disorders are quite heterogeneous; the extent of comorbidity with depression depends on which anxiety symptoms and disorders are assessed (Avenevoli, Stolar, Li, Dierker, & Merikangas, 2001; Chaplin, Gillham, & Seligman, 2009; Moffitt et al., 2007). For example, whereas panic disorder does not consistently predict subsequent depression, SOC and GAD in childhood tend to be associated with depression during adolescence (Bittner et al., 2007; Keenan et al., 2009). Thus, although there is considerable comorbidity among anxiety disorders (Angold et al., 1999), combining anxiety disorders together likely distorts the apparent strength and direction of the relation between particular anxiety disorders and depression.

Third, the degree of comorbidity varies by age and developmental period. Whereas anxiety is more prevalent during childhood, depression increases during adolescence (e.g., Cohen et al., 1993; Woodward & Fergusson, 2001). Youth with comorbid anxiety and depression tend to be older than those with either disorder alone (Brady & Kendall, 1992; Merikangas & Avenevoli, 2002). This may be due, in part, to differences in the structure and differentiation of affect across development. For example, in young children (third graders), anxiety and depression form a unified, indistinguishable construct, whereas in older children (sixth graders), a dual-factor or tripartite model is more common (Cole, Truglio, & Peeke, 1997). Thus, higher rates of comorbid anxiety and depressive disorders tend to be found in adolescents than children (Ollendick et al., 2005).

Finally, whereas concurrent comorbidity of anxiety and depressive disorders is substantial and relatively undisputed (Brady & Kendall, 1992; Karlson et al., 2006), less clear is the extent and direction of “sequential comorbidity;” that is, when one disorder reliably precedes the other (Angold et al., 1999). Comorbid anxiety and depression may have strong effects on one another such that the presence of anxiety symptoms may lead to an increase

in depressive symptoms and vice versa (Bittner et al., 2007; Goodwin, Fergusson, & Horwood, 2004); most studies of sequential comorbidity have focused on anxiety as the predictor and depression as the outcome, rather than the reverse. In general, evidence indicates that anxiety symptoms and disorders in childhood often precede the onset of depressive disorders in adolescence and young adulthood (Chorpita & Daleiden, 2002; Pine, Cohen, Gurley, Brook, & Ma, 1998), particularly for girls (Breslau, Schultz, & Peterson, 1995; Chaplin et al., 2009; Keenan & Hipwell, 2005), and may contribute to the increased risk of depression in females (Bittner et al., 2004).

Less evidence exists of depression preceding anxiety, however (e.g., Axelson & Birmaher, 2001; Orvaschel, Lewinsohn, & Seeley, 1995). Only a few balanced investigations have been conducted in which individuals with mood disorders are followed longitudinally to assess the onset of an anxiety disorder as well as the reverse (Gallerani, Garber, & Martin, 2010; Keenan et al., 2009; Pine et al., 1998). For example, Pine and colleagues (1998) reported that MDD during adolescence significantly predicted a five-fold increased risk of GAD in young adulthood.

What Accounts for Comorbidity between Anxiety and Depression

Several nonmutually exclusive explanations for the observed comorbidity between anxiety and depression have been suggested, some of which are methodological and others more substantive. We highlight here some of the most frequently suggested reasons for such comorbidity, and later discuss their implications for intervention.

Item and Symptom Overlap

One methodological explanation for the high correlation between self-report measures of anxiety and depression is that the items on these scales are quite similar or even identical. Investigations that have excluded all overlapping items, however, have found that the correlations of the abbreviated measures were still significant ($r = .34$) (Stark & Laurent, 2001), and that such modified versions of the self-report measures only reduced the shared variance (52%–72%) in trait constructs by about 13% (Cole, Truglio, & Peeke, 1997).

A related, but more complex, reason for the observed comorbidity between anxiety and depression is that the symptoms that define these disorders are similar. For example, social avoidance can characterize both SOC and MDD, although the underlying reason for and function of the avoidance may differ. The clearest example of symptom overlap is between GAD and MDD, both of which include fatigue, sleep disturbance, concentration difficulties, and irritability (in children). The proposed revision for DSM-V eliminates these specific symptoms from the criteria that define GAD, which may be one way to reduce comorbidity. The validity of these new symptom criteria for GAD remains to be demonstrated in adults as well as children.

Underlying Negative Affectivity

An important substantive explanation for the association between anxiety and depression is that they have shared etiological influences. In particular, some of the overlap between anxious and depressive symptoms likely is due to a common underlying latent risk factor of general negative affectivity (NA) or negative emotionality (Barlow, 2000; Clark & Watson, 1991), which is related to the personality construct of neuroticism (Watson, 2000). NA represents the extent to which an individual feels distress (e.g., upset, sad, angry, guilty, worried) and not calm, relaxed, or peaceful. According to the tripartite model (Clark & Watson, 1991), high NA characterizes both anxiety and depression, whereas low positive affect (PA) and loss of interest or pleasure are unique to depression, and somatic tension and physiological hyperarousal (PH) are distinct features of anxiety. Removing variance

attributable to general NA reduces the correlation between anxiety and depression and facilitates their discrimination on the remaining unique features (i.e., PA, PH).

Support for the tripartite model has been found in clinical and nonclinical samples of children and adolescents (Laurent & Ettelson, 2001). Particularly robust is the finding that general NA accounts for much of the variance shared by depression and anxiety in youth (Chorpita, Daleiden, Moffitt, Yim, & Umemoto, 2000; Tully, Zajac & Venning, 2009). Evidence for the physiological arousal component has been more mixed, however (e.g., Chorpita, 2002; Jacques & Mash, 2004; Lonigan, Hooe, David, & Kistner, 1999). Watson (2005) proposed a reconfiguration of mood and anxiety disorders into “distress” and “fear” factors, which together comprise a higher order internalizing factor (Slade & Watson, 2006). Accordingly, GAD, MDD, DD, and post-traumatic stress disorder (PTSD) cluster with the distress disorders, whereas panic disorder, SOC, and SPs comprise the fear disorders. Some evidence that GAD is more closely linked with depression as compared to anxiety has been found in children (Higa-McMillan, Smith, Chorpita & Hayashi, 2008; Lahey et al., 2008), although evidence that GAD is more closely aligned with other anxiety disorders than to depression also has been reported (e.g., Beesdo, Pine, Lieb, & Wittchen, 2010).

Shared Familial Risk

Family studies consistently reveal that offspring of depressed parents are at high risk for developing early-onset anxiety disorders as well as depression, and offspring of anxious parents are at risk to develop early-onset depression (Warner, Mufson, Weissman, 1995; Weissman, Warner, Wickramaratne, Moreau, & Olfson, 1997). Familial risk may be a marker for genetic and/or shared environmental mechanisms.

Twin studies indicate that a common genetic influence likely accounts for some of the covariation between anxious and depressive symptoms (Hudziak, Rudiger, Neale, Heath, & Todd, 2000; Thapar & McGuffin, 1997). An endophenotype common to both pediatric anxiety and depression may be “temperament” such as negative affectivity (as discussed earlier), stress reactivity, affect dysregulation, behavioral inhibition, and harm avoidance (e.g., Caspi, Moffitt, Newman, & Silva, 1996; Kelvin, Goodyer, & Altham, 1996). Moreover, consistent with the notion of heterotypic continuity, the shared genetic liability may be differentially expressed as anxiety earlier and depression later in development (Eaves, Silberg, & Erkanli, 2003). In particular, the common genetic/biological diathesis may result in anxiety or depression depending on the timing of the environmental events. In genetically vulnerable children, stressors that occur in childhood may produce anxiety, whereas those occurring during adolescence may lead to depression. The developmental progression from anxiety to depression may reflect a “readiness” (Kovacs & Devlin, 1998; p. 54) to show certain physiological aspects of anxiety (e.g., agitation, hyperarousal) earlier in development, and certain other physiological (e.g., vegetative symptoms) and cognitive (e.g., rumination) aspects of depression later.

Although genes likely play a significant role in the etiology and comorbidity of anxiety and depression, copious evidence also implicates parenting behaviors in the intergenerational transmission of anxiety and depression (e.g., McLeod, Weisz, & Wood, 2007; McLeod, Wood, & Weisz, 2007). For example, parental rejection and control are positively correlated with both anxiety and depression in children, with rejection being more strongly associated with depression and control more associated with anxiety (Rapee, 1997). Such parenting behaviors have been found to characterize both anxious and depressed mothers who tend to exhibit less warmth and more controlling behaviors toward their children (e.g., Lovejoy, Graczyk, O’Hare, & Neuman, 2000; Whaley, Pinto, & Sigman, 1999). In addition, insecure attachment early in childhood has been linked with both anxiety and depression later (Davila, Ramsay, Stroud, & Steinberg, 2005).

Similar Information Processing Biases and Neural Substrates

Negative cognitions and information processing errors such as catastrophizing, rumination, and worry characterize both anxiety and depression (Dozios & Beck, 2008; Martin & Tesser, 1996). Maladaptive interpretations of negative social events in particular are cognitive biases associated with both social anxiety and depression (Wilson & Rapee, 2005). Similar to depressed individuals, people high in social anxiety tend to make internal, stable, and global attributions for failures in interpersonal situations (Alfano, Joiner, & Perry, 1994; Hope, Gansler, & Heimberg, 1989), although negative inferential style interacts with stress to specifically predict depression and not anxiety in adolescents (Hankin, 2008).

Data from performance-based studies indicate that anxious and depressed youth also share information-processing biases in attention toward threat and interpretation of ambiguous situations as negative or threatening (e.g., Bogels & Zigterman, 2000; Ladouceur et al., 2005). The cognitive processes of individuals with anxiety and depression tend to be primed to attend more quickly and readily to threat (Coles, Turk, & Heimberg, 2007; Gotlib, Krasnoperova, Neubauer, Joorman, 2004). Attention toward and interpretation of threat is assumed to worsen symptoms of internalizing psychopathology (Dalgleish et al., 2003; Lonigan, Vasey, Phillips, & Hazen, 2004), and in turn, cognitive symptoms of anxiety and depression presumably then worsen attention to and interpretation of negative stimuli (Brozovich & Heimberg, 2008).

Finally, similar neural-circuitry dysfunction related to emotional modulation of perception and behavior has been found in individuals with either anxiety or depressive disorders (e.g., Phillips, Drevets, Rauch, & Lane, 2003; Thomas et al., 2001). For example, adolescents with anxiety or mood disorders have been found to exhibit both common and distinct functional neural correlates (i.e., amygdala responses) during face processing, depending on the specific attention and emotion states engaged (Beesdo et al., 2009). In particular, Beesdo and colleagues reported that both anxious and depressed youth showed greater amygdala activation than healthy controls when viewing fearful faces, whereas disorder specificity emerged during passive viewing of emotional stimuli. Thus, anxiety and depressive disorders in youth involve many complex commonalities, as well as distinguishable amygdala-related biases, information processing patterns, temperaments, family environments, and genetic vulnerabilities.

Anxiety as a Risk for Depression

As noted earlier, anxiety and depression show substantial sequential comorbidity, with anxiety symptoms and disorders often preceding the onset of mood disorders (e.g., Avenevoli et al. 2001; Kim-Cohen et al., 2003; Pine et al. 1998). Accordingly, anxiety itself may be a causal risk factor (Kraemer et al., 1997) for the development of depression (Bittner et al., 2004; Wittchen, Beesdo, Bittner, & Goodwin, 2003), and the experience of childhood anxiety may be both the source of high sequential comorbidity over development and of elevated rates of concurrent anxiety and depression in adolescence.

Childhood anxiety may have depressogenic effects through a variety of mechanisms. For example, social anxiety likely increases individuals' affiliative behaviors as well as their attempts to avoid negative evaluations from others (e.g., Alden & Taylor, 2004; Leary & Kowalski, 1995). This may result in dysfunctional social behaviors that are intended to be protective, but actually end up increasing the likelihood of the very rejection they are trying to avoid. Anxiety-driven social withdrawal and isolation can intensify peer rejection, resulting in feelings of loneliness, low self-worth, and sadness (Gazelle & Ladd, 2003), particularly for girls for whom interpersonal relationships are especially important (Rose & Rudolph, 2006). Similarly, avoidance of expressing emotions due to fears of rejection and

humiliation has been found to mediate the relation between social anxiety and changes in depressive symptoms (Grant, Beck, Farrow, & Davila, 2007). Socially anxious individuals who inhibit their expression of negative emotions may feel devalued, sad, and resentful, and over time, become depressed. Furthermore, individuals with social anxiety tend to make negative inferences about the meaning of adverse social events for their future and self-worth (Stopa & Clark, 2000), which have been identified as cognitive vulnerabilities to depression (Abramson, Metalsky, & Alloy, 1989).

Summary

The high level of comorbidity between anxiety and depression in youth may be the result of three non-exclusive factors: (a) substantial overlap in both the symptoms and items used to assess these putatively different disorders, (b) common etiological factors implicated in the development of each condition, and (c) negative sequelae of anxiety conferring increased risk for the development of depression. The first explanation has important implications for the nosology of disorders, creation of valid assessments, and the design and conduct of research with internalizing youth. The second and third explanations may guide the development of treatments for youth with internalizing problems and the prevention of depression in adolescents and adults.

Implications for Intervention

Treatment

Evidence-based treatments are typically designed to target the precipitating and maintaining factors of disorders to bring about symptom remission and functional improvement. To the extent that anxiety and depression in youth share common etiological underpinnings, the efficacious treatments developed for these disorders likely share common features and mechanisms of action. Examination of the intervention literature indicates that this is largely what appears to have transpired. For both depression and anxiety, the best practice pharmacotherapy is with selective serotonin reuptake inhibitors (SSRIs) (e.g., TADS, 2004; Walkup et al., 2008). Generally, higher doses of SSRIs are required for therapeutic effects on depression than anxiety (with the notable exception of dosing for obsessive compulsive disorder). The same agents, however, appear to have similar benefits and adverse event profiles across youth with anxiety or depression (Bridge et al., 2007).

The primary psychosocial intervention for both anxiety and depression is cognitive behavioral therapy (CBT; see Compton et al., 2004; Kendall, 2010), with positive effects for CBT reported across the anxiety disorders and for mild to moderate levels of depressive disorder (for severe depression, the combination of CBT and SSRI may be warranted, see Brent et al., 2008). The CBT interventions that treat these conditions contain similar elements, although the programs differ in their complexity and number of strategies employed. For example, various CBT manuals for anxiety and depression include problem solving, assertiveness training, cognitive restructuring, family communication skills training, relaxation, exposure, pleasant activity scheduling, and behavioral activation (Weersing, 2004). Across all these techniques, the different CBT manuals share a core focus on (a) the interplay between thoughts, feelings, and behaviors, and (b) within this framework, teaching adaptive responses to stress and coping with negative emotionality.

In addition to the surface similarity of effective interventions for anxious and depressed youth, interventions for one condition may have beneficial spill-over effects on comorbid symptoms of the other disorder. Several randomized controlled trials examining the effects of individual, group, and family CBT and exposure-based treatments for anxiety disorders in children have found significant reductions in self-reported depressive symptoms as well

(Barrett Dadds, & Rapee, 1996; Kendall et al., 1997; Kendall et al 2008; Kendall, Safford, Flannery-Schroeder, & Webb, 2004; Manassis et al., 2002; Silverman et al., 1999). A comprehensive meta-analysis of the effects of psychological therapy for depression in children and adolescents examined whether such treatments affected other conditions also (Weisz, McCarty, & Valeri, 2006). To examine this specificity question, Weisz and colleagues compared the effect sizes (ES) for measures of depression with those for anxiety symptoms across ten studies that had assessed both. They found that depression treatments produced a significant reduction in anxiety symptoms (ES = 0.39) that was only marginally lower than that found for depressive symptoms (ES = 0.57). Thus, finding that anxiety treatments reduce depressive symptoms and depression treatments beneficially affect anxiety is consistent with the view that anxiety and depression in youth are closely associated empirically (e.g., Achenbach & Rescorla, 2001) and likely share common risk factors such as negative affectivity (e.g., Clark & Watson, 1991; Laurent & Ettelson, 2001).

The presence of one disorder, however, sometimes reduces the efficacy of the treatment of the other. If they were essentially the same condition, then why wouldn't the interventions affect the comorbid disorder as well? Some depression treatment studies have found that comorbid anxiety predicts a worse outcome (Curry et al., 2006; Emslie et al., 1998; Vostanis, Feehan, & Grattan, 1998), although other studies have shown that the presence of anxiety predicted a more positive outcome of CBT for depression (Brent et al., 1998; Rohde et al., 2001). Results of investigations of whether depression reduces the efficacy of anxiety treatments also have been mixed. A significant relation between depressive symptoms and a less favorable response to anxiety treatment has been found in some studies (e.g., Berman, Weems, Silverman, & Kurtines, 2000), but not others (Southam-Gerow, Kendall, & Weersing, 2001). Thus, interventions targeting one disorder will not necessarily successfully treat the other comorbid condition, even if similar treatments are effective for each diagnosis on its own.

Taken together, these results have led to the development of interventions designed to treat comorbid anxiety and depression or, more ambitiously, to successfully target anxiety, depression, or comorbid anxiety and depression within a single treatment protocol. Such work is still in the pilot stage of implementation and testing, but two approaches have emerged. One program with youth stems from adult work developing a unified theory of emotional disorders (Barlow, Allen, & Choate, 2004), and emphasizes the core cognitive biases and avoidance of negative emotions common to both anxiety and depression (Ehrenreich, Goldstein, Wright, & Barlow, 2009). A second approach is a more behavioral intervention focusing on graded engagement in activities designed to increase positive affect and reduce the functional impairment associated with internalizing symptoms in youth (Weersing, Gonzalez, Campo, & Lucas, 2008). The technique of "graded engagement" combines aspects of behavioral activation for depression, exposure techniques for anxiety, and problem solving skills to target the maladaptive responses to stress and negative emotions seen across anxiety and depression. Notably, this intervention targets both threshold and subthreshold levels of comorbidity between anxiety and depression, and the functional impact that these symptoms have on each other. For example, depressed youth may have difficulty developing rewarding and mood enhancing relationships if they also have a history of social anxiety. Thus, although similar interventions are efficacious with both anxiety and depressive disorders, treatments need to address their unique features as well.

Prevention

Whereas treatment aims to reduce existing symptoms and disorders, the goal of prevention is to decrease the likelihood of the onset of a disorder or a worsening of symptoms, particularly among individuals at risk based on their having current subsyndromal symptoms

(i.e., indicated prevention) or other risk factors such as parental psychopathology or exposure to stress (i.e., selected prevention). Although less extensive, research on the prevention of anxiety or depression parallels the treatment literature in several ways (for reviews, see Horowitz & Garber, 2006; Neil & Christensen, 2009; Stice et al., 2009). First, interventions aimed at preventing anxiety and depression use many of the same procedures. For example, some depression prevention programs explicitly include anxiety reducing techniques such as relaxation, anxiety management, and stress inoculation (e.g., Clarke, Hawkins, Murphy, & Sheeber, 1993; Hains & Ellmann, 1994). Second, the presence of one disorder sometimes moderates the effect of the preventive intervention on the other. For instance, some depression prevention programs have been found to be better for children with higher baseline levels of emotional arousal (Hains & Ellmann, 1994) or clinical anxiety (Lowry-Webster, Barrett, & Dadds, 2001; Lowry-Webster, Barrett, & Lock, 2003).

Third, interventions aimed at preventing one disorder sometimes affect the other as well. For example, the effects of the Penn Prevention program have been found to be as strong or even stronger in reducing anxious as compared to depressive symptoms (Gillham et al., 2006; Roberts, Kane, Thomson, Bishop, & Hart, 2003). Gillham and colleagues suggested that the stronger effects on anxiety may have been because the anti-anxiety skills were easier to learn compared to the skills that more directly targeted depression. It also is possible that including strategies aimed at reducing anxiety dilutes the effects of the overall intervention on depression.

In contrast, anxiety prevention programs have been found to have only modest effects on depressive symptoms. For instance, the FRIENDS program, a universal school-based intervention designed to prevent internalizing problems through enhancing problem-solving and coping skills, has been found to significantly reduce anxiety. A reduction in depressive symptoms, however, was found only at the 12-month evaluation and not at 24 or 36 months (Barrett, Farrell, Ollendick, & Dadds, 2006; Lock & Barrett, 2003; Lowry-Webster et al., 2001).

Treating Anxiety to Prevent Depression

Given that childhood anxiety often precedes and is a risk factor for depression (Costello et al., 2003; Wittchen et al., 2003), a potentially useful intervention strategy may be to treat anxiety as a means of preventing subsequent depression (Bienvenu & Ginsburg, 2007; Flannery-Schroeder, 2006). Although some treatments for anxiety also produce significant decreases in depressive symptoms (e.g., Barrett et al., 1996; Kendall et al., 2004; Manassis et al., 2002), the strategy of preventing depression by successfully treating anxiety in youth has yet to be tested in a way that disentangles the baseline levels of both anxiety and depression. Several important questions remain regarding the treatment and prevention of anxiety and depression, respectively. Do treatments for anxiety disorders prevent the onset of subsequent depressive disorders in addition to reducing concurrent depressive symptoms? If successful treatment of anxiety prevents subsequent depression, how enduring are these effects? For whom does the treatment of anxiety prevent depression? What are the optimal implementation parameters, such as dose and timing? Finally, what accounts for the depression prevention effects of treatments for anxiety in children? That is, through what mechanisms do these significant effects occur?

Table 1 outlines several ways that treating anxiety might be sufficient to prevent depression without directly implementing depression prevention strategies. If successful, then treating anxiety could be an efficient and cost effective way to prevent depression, at least among anxious children. We suggest at least four different ways this might occur: (a) Anxiety is a probabilistic risk factor for the development of depression. Removing anxiety may eliminate

this risk, although the underlying processes are not specified. (b) The therapeutic activities that target anxiety (e.g., cognitive reappraisal; coping) may generalize to depression. Children might learn to apply the skills acquired for dealing with their anxiety to other situations and emotions, without necessarily being explicitly taught to do so. The extent to which such transfer of learning occurs will partially depend on the child's level of cognitive development. (c) If anxiety is a *causal* risk factor for depression, then reducing anxiety should affect the processes (e.g., avoidance) that account for this link, thereby preventing depression. For example, anxiety leads some children to avoid social situations and hence miss out on positively reinforcing activities, thereby increasing their chances of depression. Similarly, anxiety may produce awkward social behaviors that result in rejection and subsequent depression. If treatment directly eliminates such anxiety and the associated mediating pathways, then depression might be prevented. (d) If anxiety and depression share common mechanism(s) such as negative affectivity, then treatments for anxiety that reduce the common risk factor(s) might prevent depression as well.

Finally, if treating anxiety is *not* sufficient to prevent depression, then it may be necessary to supplement anxiety treatments with intervention techniques specifically aimed at preventing depression. Although several successful depression prevention programs exist (see Brunwasser, Gillham, & Kim, 2009; Horowitz & Garber, 2006; Stice et al., 2009 for reviews), studies are needed that explicitly test the efficacy of combining depression prevention strategies, either simultaneously or sequentially, with treatments for anxiety (see bottom of Table 1).

RESEARCH RECOMMENDATIONS

1. Determine what are the most efficacious and cost-effective ways of providing interventions that treat and prevent both anxiety and depression.
2. Conduct randomized clinical trials with sufficiently long follow-up intervals to determine if treating anxiety significantly decreases the likelihood of subsequent depression.
3. Identify specific mechanisms that link the different anxiety disorders to depression, and test interventions that directly target these mechanisms.
4. Conduct treatment-dismantling studies to identify the specific components of the treatments that contribute to symptom reduction and improved functioning.

Acknowledgments

This work was supported in part from grants from NIMH (R01MH064735; R01MH064503) and the William T. Grant Foundation.

References

- Abramson LY, Metalsky GI, Alloy LB. Hopelessness depression: A theory-based subtype of depression. *Psychological Review*. 1989; 96:358–372.
- Achenbach, T.; Rescorla, L. Manual for the ASEBA school-age forms and profiles. Burlington: University of Vermont Research Center for Children, Youth, & Families; 2001.
- Alden LE, Taylor CT. Interpersonal processes in social phobia. *Clinical Psychology Review*. 2004; 24:857–882. [PubMed: 15501559]
- Alfano MS, Joiner TE, Perry M. Attributional style: A mediator of the shyness–depression relationship? *Journal of Research in Personality*. 1994; 28:287–300.

- Angold A, Costello EJ, Erkanli A. Comorbidity. *Journal of Child Psychology, Psychiatry, and Allied Disciplines*. 1999; 40:57–87.
- Avenevoli S, Stolar M, Li J, Dierker L, Merikangas KR. Comorbidity of depression in children and adolescents: Models and evidence from a prospective high-risk family study. *Biological Psychiatry*. 2001; 49:1071–1081. [PubMed: 11430849]
- Axelson DA, Birmaher B. Relation between anxiety and depressive disorders in childhood and adolescence. *Depression and Anxiety*. 2001; 14:67–78. [PubMed: 11668659]
- Barlow DH. Unraveling the mysteries of anxiety and its disorders from the perspective of emotion theory. *American Psychologist*. 2000; 55:1247–1263. [PubMed: 11280938]
- Barlow DH, Allen LB, Choate ML. Toward a unified treatment for the emotional disorders. *Behavior Therapy*. 2004; 35:205–230.
- Barrett PM, Dadds MR, Rapee RM. Family treatment of childhood anxiety: A controlled trial. *Journal of Consulting and Clinical Psychology*. 1996; 64:333–342. [PubMed: 8871418]
- Barrett PM, Farrell LJ, Ollendick TH, Dadds M. Long-term data from an Australian FRIENDS universal prevention trial. *Journal of Clinical Child and Adolescent Psychology*. 2006; 35:333–342.
- Beesdo K, Knappe S, Pine DS. Anxiety and anxiety disorders in children and adolescents: developmental issues and implications for *DSM-V*. *Psychiatric Clinics of North America*. 2009; 32:483–524. [PubMed: 19716988]
- Beesdo K, Lau JY, Guyer AE, McClure-Tone EB, Monk CS, Nelson EE, Fromm SS, Goldwin M, Wittchen H-U, Leibenluft E, Ernst M, Pine DS. Common and distinct amygdala-function perturbations in depressed vs. anxious adolescents. *Archives of General Psychiatry*. 2009; 66:275–285. [PubMed: 19255377]
- Beesdo K, Pine DS, Lieb R, Wittchen HU. Incidence and risk patterns of anxiety and depressive disorders and categorization of generalized anxiety disorder. *Archives of General Psychiatry*. 2010; 67:47–57. [PubMed: 20048222]
- Berman SL, Weems CF, Silverman WK, Kurtines WM. Predictors of outcome in exposure-based cognitive and behavioral treatments for phobic and anxiety disorders in children. *Behavior Therapy*. 2000; 31:713–731.
- Bienvenu OJ, Ginsburg GS. Prevention of anxiety disorders. *International Review of Psychiatry*. 2007; 19:647–654. [PubMed: 18092242]
- Birmaher B, Ryan ND, Williamson DE, Brent DA, Kaufman J, Dahl RE, et al. Childhood and adolescent depression: A review of the past 10 years, Part I. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1996; 35:1427–1439. [PubMed: 8936909]
- Bittner A, Egger HL, Erkanli A, Costello EJ, Foley DL, Angold A. What do childhood anxiety disorders predict? *Journal of Child Psychology and Psychiatry*. 2007; 48:1174–1183. [PubMed: 18093022]
- Bittner A, Goodwin RD, Wittchen HU, Beesdo K, Höfler M, Lieb R. What characteristics of primary anxiety disorders predict subsequent major depressive disorder? *Journal of Clinical Psychiatry*. 2004; 65:618–626. [PubMed: 15163247]
- Bogels SM, Zigterman D. Dysfunctional cognitions in children with social phobia, separation anxiety disorder, and generalised anxiety disorder. *Journal of Abnormal Child Psychology*. 2000; 28:205–211. [PubMed: 10834771]
- Brady EU, Kendall PC. Comorbidity of anxiety and depression in children and adolescents. *Psychological Bulletin*. 1992; 111:244–255. [PubMed: 1557475]
- Brent DA, Emslie G, Clarke G, Wagner KD, Asarnow JR, Keller M, et al. Switching to another SSRI or to Venlafaxine with or without cognitive behavioral therapy for adolescents with SSRI-resistant depression: The TORDIA randomized controlled trial. *Journal of the American Medical Association*. 2008; 299:901–913. [PubMed: 18314433]
- Brent DA, Kolko DJ, Birmaher B, et al. Predictors of treatment efficacy in a clinical trial of three psychosocial treatments for adolescent depression. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1998; 37:906–914. [PubMed: 9735610]
- Breslau N, Schultz L, Peterson E. Sex differences in depression: A role for preexisting anxiety. *Psychiatry Research*. 1995; 58:1–12. [PubMed: 8539307]

- Bridge JA, Iyengar S, Salary CB, Barbe R, Birmaher B, Pincus HA, Ren L, Brent DA. Clinical response and risk for reported suicidal ideation and suicide attempts in pediatric antidepressant treatment: A meta-analysis of randomized controlled trials. *Journal of the American Medical Association*. 2007; 297:1683–1696. [PubMed: 17440145]
- Brozovich F, Heimberg RG. An analysis of post-event processing in social anxiety disorder. *Clinical Psychology Review*. 2008; 28:891–903. [PubMed: 18294745]
- Brunwasser SM, Gillham JE, Kim ES. A meta-analytic review of the Penn Resiliency Program's effect on depressive symptoms. *Journal of Consulting and Clinical Psychology*. 2009; 77:1042–1054. [PubMed: 19968381]
- Caspi A, Moffitt TE, Newman DL, Silva PA. Behavioral observations at age 3 years predict adult psychiatric disorders: longitudinal evidence from a birth cohort. *Archives of General Psychiatry*. 1996; 53:1033–1039. [PubMed: 8911226]
- Chaplin TM, Gillham JB, Seligman MEP. Gender, anxiety, and depressive symptoms: A longitudinal study of early adolescents. *Journal of Early Adolescence*. 2009; 29:307–327. [PubMed: 19756209]
- Chorpita BF. The tripartite model and dimensions of anxiety and depression: An examination of structure in a large school sample. *Journal of Abnormal Child Psychology*. 2002; 30:177–190. [PubMed: 12002397]
- Chorpita BF, Daleiden EL. Tripartite dimensions of emotion in a child clinical sample: measurement strategies and implications for clinical utility. *Journal of Consulting and Clinical Psychology*. 2002; 70:1150–1160. [PubMed: 12362965]
- Chorpita BF, Daleiden EL, Moffitt C, Yim L, Umemoto LA. Assessment of tripartite factors of emotion in children and adolescents I: Structural validity and normative data of an affect and arousal scale. *Journal of Psychopathology and Behavioral Assessment*. 2000; 22:141–160.
- Clark LA, Watson D. Tripartite model of anxiety and depression: Psychometric evidence and taxonomic implications. *Journal of Abnormal Psychology*. 1991; 100:316–336. [PubMed: 1918611]
- Clarke GN, Hawkins W, Murphy M, Sheeber LB. School-based primary prevention of depressive symptomatology in adolescents: Findings from two studies. *Journal of Adolescent Research*. 1993; 8:183–204.
- Cohen P, Cohen J, Brook J. An epidemiological study of disorders in late childhood and adolescence-II: Persistence of disorders. *Journal of Child Psychology and Psychiatry*. 1993; 34:869–877. [PubMed: 8408372]
- Cole DA, Peeke LA, Martin JM, Truglio R, Seroczynski AD. A longitudinal look at the relation between depression and anxiety in children and adolescents. *Journal of Consulting and Clinical Psychology*. 1997; 106:586–597.
- Cole DA, Truglio R, Peeke L. Relation between symptoms of anxiety and depression in children: a multitrait-multimethod multigroup assessment. *Journal of Consulting and Clinical Psychology*. 1997; 65:110–119. [PubMed: 9103740]
- Coles ME, Heimberg RG, Schofield CA. Interpretation of facial expressions and social anxiety: Specificity and source of biases. *Cognition and Emotion*. 2008; 22:1159–1173.
- Compton SN, March JS, Brent D, Albano AM, Weersing VR, Curry J. Cognitive-behavioral psychotherapy for anxiety and depressive disorders in children and adolescents: An evidence-based medicine review. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2004; 43:930–959. [PubMed: 15266189]
- Costello EJ, Mustillo S, Erkanli A, Keeler G, Angold A. Prevalence and development of psychiatric disorders in childhood and adolescence. *Archives of General Psychiatry*. 2003; 60:837–844. [PubMed: 12912767]
- Curry J, Rohde P, Simons A, et al. Predictors and moderators of acute outcome in the Treatment for Adolescents with Depression Study (TADS). *Journal of American Academy of Child and Adolescent Psychiatry*. 2006; 45:1427–1439.
- Dalgleish T, Taghavi R, Neshat-Doost H, Moradi A, Canterbury R, Yule W. Patterns of processing bias for emotional information across clinical disorders: A comparison of attention, memory, and prospective cognition in children and adolescents with depression, generalized anxiety, and

- posttraumatic stress disorder. *Journal of Clinical Child and Adolescent Psychology*. 2003; 32:10–21. [PubMed: 12573928]
- Davila, J.; Ramsay, M.; Stroud, CB.; Steinberg, S. Attachment as vulnerability to the development of psychopathology. In: Hankin, BL.; Abela, JRZ., editors. *Development of psychopathology: A vulnerability-stress perspective*. Thousand Oaks, CA: Sage; 2005. p. 215-242.
- Dozois, DJA.; Beck, AT. Cognitive schemas, beliefs and assumptions. In: Dobson, KS.; Dozois, DJA., editors. *Risk factors in depression*. Oxford: Elsevier; 2008. p. 121-143.
- Eaves L, Silberg J, Erkanli A. Resolving multiple epigenetic pathways to adolescent depression. *Journal of Child Psychology and Psychiatry*. 2003; 44:1006–1014. [PubMed: 14531583]
- Ehrenreich JT, Goldstein CR, Wright LR, Barlow DH. Development of a unified protocol for the treatment of emotional disorders in youth. *Child & Family Behavior Therapy*. 2009; 31:20–37. [PubMed: 19617930]
- Emslie GJ, Rush AJ, Weinberg WA, et al. Fluoxetine in child and adolescent depression: Acute and maintenance treatment. *Depression and Anxiety*. 1998; 7:32–39. [PubMed: 9592630]
- Ezpeleta L, Domenech JM, Angold A. A comparison of pure and comorbid CD/ODD and depression. *Journal of Child Psychology and Psychiatry*. 2006; 47:704–712. [PubMed: 16790005]
- Flannery-Schroeder EC. Reducing anxiety to prevent depression. *American Journal of Preventive Medicine*. 2006; 31:S136–S142. [PubMed: 17175408]
- Gallerani CM, Garber J, Martin NC. The temporal relation between depression and comorbid psychopathology in adolescents at varied risk for psychopathology. *Journal of Child Psychology and Psychiatry*. 2010; 51:242–249. [PubMed: 19874429]
- Gazelle H, Ladd GW. Anxious solitude and peer exclusion: A diathesis-stress model of internalizing trajectories in childhood. *Child Development*. 2003; 74:257–278. [PubMed: 12625449]
- Gillham JE, Reivich KJ, Freres DR, Lascher M, Litzinger S, Shatté A, et al. School-based prevention of depression and anxiety symptoms in early adolescence: A pilot of a parent intervention component. *School Psychology Quarterly*. 2006; 21:323–348.
- Goodwin RD, Fergusson DM, Horwood LJ. Early anxious/withdrawn behaviors predict later internalizing disorders. *Journal of Child Psychology and Psychiatry*. 2004; 45:874–883. [PubMed: 15056317]
- Gotlib IH, Krasnoperova E, Neubauer Yue D, Joorman J. Attentional biases for negative interpersonal stimuli in clinical depression. *Journal of Abnormal Psychology*. 2004; 113:127–135.
- Gould MS, King R, Greenwald S, Fisher P, Schwab-Stone M, Kramer R, et al. Psychopathology associated with suicidal ideation and attempts among children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1998; 37:915–923. [PubMed: 9735611]
- Grant DM, Beck JG, Farrow SM, Davila J. Do interpersonal features of social anxiety influence the development of depressive symptoms? *Cognition & Emotion*. 2007; 21:646–663.
- Hains AA, Ellman SW. Stress inoculation training as a preventive intervention for high school youths. *Journal of Cognitive Psychotherapy: An International Quarterly*. 1994; 8:219–232.
- Hankin BL. Cognitive vulnerability-Stress model of depression during adolescence: Investigating depressive symptom specificity in a multi-wave prospective study. *Journal of Abnormal Child Psychology*. 2008; 36:999–1014. [PubMed: 18437551]
- Higa-McMillan CK, Smith RL, Chorpita BF, Hayashi K. Common and unique factors associated with DSM-IV-TR internalizing disorders in children. *Journal of Abnormal Child Psychology*. 2008; 36:1279–1288. [PubMed: 18607718]
- Hope DA, Gansler DA, Heimberg RG. Attentional focus and causal attributions in social phobia: Implications from social psychology. *Clinical Psychology Review*. 1989; 9:49–60.
- Horowitz JL, Garber J. The prevention of depressive symptoms in children and adolescents: A meta-analytic review. *Journal of Consulting and Clinical Psychology*. 2006; 74:401–415. [PubMed: 16822098]
- Hudziak JJ, Rudiger LP, Neale MC, Heath AC, Todd RD. A twin study of inattentive, aggressive, and anxious/depressed behaviors. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2000; 39:469–476. [PubMed: 10761349]

- Jacques HAK, Mash EJ. A test of the tripartite model of anxiety and depression in elementary and high school boys and girls. *Journal of Abnormal Child Psychology*. 2004; 32:13–25. [PubMed: 14998108]
- Keenan K, Feng X, Hipwell A, Klostermann S. Depression begets depression: comparing the predictive utility of depression and anxiety symptoms to later depression. *Journal of Child Psychology and Psychiatry*. 2009; 50:1167–1175. [PubMed: 19344385]
- Keenan K, Hipwell AE. Preadolescent clues to understanding depression in girls. *Clinical Child and Family Psychology Review*. 2005; 8:89–105. [PubMed: 15984082]
- Kelvin RG, Goodyer IM, Altham PM. Temperament and psychopathology amongst siblings of probands with depressive and anxiety disorders. *Journal of Child Psychology and Psychiatry*. 1996; 37:543–550. [PubMed: 8807434]
- Kendall PC, Flannery-Schroeder E, Panichelli-Mindel SM, Southam-Gerow M. Therapy for youths with anxiety disorders: a second randomized clinical trial. *Journal of Consulting and Clinical Psychology*. 1997; 65:366–380. [PubMed: 9170760]
- Kendall PC, Hudson J, Gosch E, Flannery-Schroeder E, Suveg C. Cognitive behavioral therapy for anxiety disordered youth: A randomized clinical trial evaluating child and family modalities. *Journal of Consulting and Clinical Psychology*. 2008; 76:282–297. [PubMed: 18377124]
- Kendall PC, Safford S, Flannery-Schroeder E, Webb A. Child anxiety treatment: outcomes in adolescence and impact on substance use and depression at 7.4-year follow-up. *Journal of Consulting and Clinical Psychology*. 2004; 72:276–287. [PubMed: 15065961]
- Kim-Cohen J, Caspi A, Moffitt TE, Harrington H, Milne BJ, Poulton R. Prior juvenile diagnoses in adults with mental disorder: Developmental follow-back of a prospective-longitudinal cohort. *Archives of General Psychiatry*. 2003; 60:709–717. [PubMed: 12860775]
- Kovacs M, Devlin B. Internalizing disorders in childhood. *Journal of Child Psychology and Psychiatry Allied Disciplines*. 1998; 39:47–63.
- Kraemer HC, Kazdin AE, Offord DR, Kessler RC, Jensen PS, Kupfer DJ. Coming to terms with the terms of risk. *Archives of General Psychiatry*. 1997; 54:337–343. [PubMed: 9107150]
- Lahey BB, Rathouz PJ, Van Hulle C, Urbano RC, Krueger RF, Applegate B, et al. Testing structural models of DSM-IV symptoms of common forms of child and adolescent psychopathology. *Journal of Abnormal Child Psychology*. 2008; 36:187–206. [PubMed: 17912624]
- Ladouceur CD, Dahl RE, Williamson DE, Birmaher B, Ryan ND, Casey BJ. Altered emotional processing in pediatric anxiety, depression, and comorbid anxiety-depression. *Journal of Abnormal Child Psychology*. 2005; 33:165–177. [PubMed: 15839495]
- Laurent J, Ettelson R. An examination of the tripartite model of anxiety and depression and its application to youth. *Clinical Child and Family Psychology Review*. 2001; 4:209–230. [PubMed: 11783739]
- Leary, MR.; Kowalski, RM. The self-presentation model of social phobia. In: Heimberg, RG.; Liebowitz, MRM.; Hope, DA.; Schneier, FR., editors. *Social phobia: Diagnosis, assessment, and treatment*. New York: Guilford Press; 1995. p. 94-112.
- Lewinsohn PM, Hops H, Roberts RE, Seeley JR, Andrews JA. Adolescent psychopathology: I. Prevalence and incidence of depression and other DSM-III-R disorders in high school students. *Journal of Abnormal Psychology*. 1993; 102:133–144. [PubMed: 8436689]
- Lock S, Barrett PM. A longitudinal study of developmental differences in universal preventive intervention for child anxiety. *Behaviour Change*. 2003; 20:183–199.
- Lonigan CJ, Hooe ES, David CF, Kistner JA. Positive and negative affectivity in children: confirmatory factor analysis of a two-factor model and its relation to symptoms of anxiety and depression. *Journal of Consulting and Clinical Psychology*. 1999; 67:374–386. [PubMed: 10369058]
- Lonigan CJ, Vasey MW, Phillips BM, Hazen RA. Temperament, anxiety, and the processing of threat-relevant stimuli. *Journal of Clinical Child and Adolescent Psychology*. 2004; 33:8–20. [PubMed: 15028537]
- Lovejoy MC, Graczyk PA, O'Hare E, Neuman G. Maternal depression and parenting: A meta-analytic review. *Clinical Psychology Review*. 2000; 20:561–592. [PubMed: 10860167]

- Lowry-Webster HM, Barrett PM, Dadds MR. A universal prevention trial of anxiety and depressive symptomatology in childhood: Preliminary data from an Australian study. *Behaviour Change*. 2001; 18:36–50.
- Lowry-Webster HM, Barrett PM, Lock S. A universal prevention trial of anxiety symptomatology during childhood: Results at one-year follow-up. *Behaviour Change*. 2003; 20:25–43.
- Manassis K, Mendlowitz SL, Scapillato D, Avery D, Fiksenbaum L, Freire M, et al. Group and individual cognitive-behavioral therapy for childhood anxiety disorders. A randomized trial. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2002; 41:1423–1430. [PubMed: 12447028]
- Martin, LL.; Tesser, A. *Striving and feeling: Interactions among goals, affect, and self-regulation*. Hillsdale, NJ: Erlbaum; 1996.
- McLeod BD, Weisz JR, Wood JJ. Examining the association between parenting and childhood depression: A meta-analysis. *Clinical Psychology Review*. 2007; 27:986–1003. [PubMed: 17449154]
- McLeod BD, Wood JJ, Weisz JR. Examining the association between parenting and childhood anxiety: A meta-analysis. *Clinical Psychology Review*. 2007; 27:155–172. [PubMed: 17112647]
- Merikangas, K.; Avenevoli, S. Epidemiology of mood and anxiety disorders in children and adolescents. In: Tsuang, MT.; Tohen, M., editors. *Textbook in Psychiatric Epidemiology*. 2. New York, NY: Wiley; 2002. p. 657-704.
- Moffitt TE, Caspi A, Harrington H, Milne BJ, Melchior M, Goldberg D, Poulton R. Generalized anxiety disorder and depression: childhood risk factors in a birth cohort followed to age 32. *Psychological Medicine*. 2007; 37:441–452. [PubMed: 17201999]
- Neil AL, Christensen H. Efficacy and effectiveness of school-based prevention and early intervention programs for anxiety. *Clinical Psychology Review*. 2009; 29:208–215. [PubMed: 19232805]
- Ollendick, TH.; Shortt, AL.; Sander, JB. Internalizing disorders of childhood and adolescence. In: Maddux, JE.; Winstead, BA., editors. *Psychopathology: Foundations for a contemporary understanding*. Mahwah, NJ: Lawrence Erlbaum; 2005. p. 353-376.
- Orvaschel H, Lewinsohn PM, Seeley JR. Continuity of psychopathology in a community sample of adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1995; 34:1525–1535. [PubMed: 8543521]
- Phillips ML, Drevets W, Rauch SL, Lane R. Neurobiology of emotion perception, II: Implications for major psychiatric disorders. *Biological Psychiatry*. 2003; 54:515–528. [PubMed: 12946880]
- Pine DS, Cohen P, Gurley D, Brook J, Ma Y. The risk for early-adulthood anxiety and depressive disorders in adolescents with anxiety and depressive disorders. *Archives of General Psychiatry*. 1998; 55:56–64. [PubMed: 9435761]
- Rapee RM. Potential role of childrearing practices in the development of anxiety and depression. *Clinical Psychology Review*. 1997; 17:47–67. [PubMed: 9125367]
- Roberts C, Kane R, Thomson H, Bishop B, Hart B. The prevention of depressive symptoms in rural school children: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*. 2003; 71:622–628. [PubMed: 12795585]
- Rohde P, Clarke GN, Lewinsohn PM, et al. Impact of comorbidity on a cognitive-behavioral group treatment for adolescent depression. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40:795–802. [PubMed: 11437018]
- Rohde P, Lewinsohn PM, Seeley JR. Are adolescents changed by an episode of major depression? *Journal of the American Academy of Child and Adolescent Psychiatry*. 1994; 33:1289–1298. [PubMed: 7995795]
- Rose A, Rudolph KD. A review of sex differences in peer relationship processes: Potential trade-offs for the emotional and behavioral development of girls and boys. *Psychological Bulletin*. 2006; 132:98–131. [PubMed: 16435959]
- Seligman LD, Ollendick TH. Comorbidity of anxiety and depression in children and adolescents: An integrative review. *Clinical Child and Family Psychology Review*. 1998; 1:125–144. [PubMed: 11324302]
- Silverman WK, Kurtines WM, Ginsburg GS, Weems CF, Rabian B, Serafini LT. Contingency management, self-control, and education support in the treatment of childhood phobic disorders:

- a randomized clinical trial. *Journal of Consulting and Clinical Psychology*. 1999; 67:675–687. [PubMed: 10535234]
- Slade T, Watson D. The structure of common *DSM-IV* and ICD-10 mental disorders in the Australian general population. *Psychological Medicine*. 2006; 36:1593–1600. [PubMed: 16882356]
- Sorensen MJ, Nissen JB, Mors O, Thomsen PH. Age and gender differences in depressive symptomatology and comorbidity: An incident sample of psychiatrically admitted children. *Journal of Affective Disorders*. 2005; 84:85–91. [PubMed: 15620389]
- Stark KD, Laurent J. Joint factor analysis of the children's depression inventory and the revised children's manifest anxiety scale. *Journal of Clinical Child Psychology*. 2001; 30:552–567. [PubMed: 11708242]
- Stice E, Shaw H, Bohon C, Marti CN, Rohde P. A meta-analytic review of depression prevention programs for children and adolescents: Factors that predict magnitude of intervention effects. *Journal of Consulting & Clinical Psychology*. 2009; 77:486–503. [PubMed: 19485590]
- Southam-Gerow MA, Kendall PC, Weersing VR. Examining outcome variability: correlates of treatment response in a child and adolescent anxiety clinic. *Journal of Clinical Child Psychology*. 2001; 30:422–436. [PubMed: 11501258]
- Stopa L, Clark DM. Social phobia and interpretation of social events. *Behavior Research and Therapy*. 2000; 38:273–283.
- Thapar A, McGuffin P. Anxiety and depressive symptoms in childhood: A genetic study of comorbidity. *Journal of Child Psychology and Psychiatry Allied Disciplines*. 1997; 38:651–656.
- Thomas KM, Drevets WC, Dahl RE, Ryan ND, Birmaher B, Eccard CH, Axelson D, Whalen PJ, Casey BJ. Amygdala response to fearful faces in anxious and depressed children. *Archives of General Psychiatry*. 2001; 58:1057–1063. [PubMed: 11695953]
- Treatment for Adolescents with Depression Study (TADS) Team. Fluoxetine, cognitive-behavioral therapy, and their combination for adolescents with depression: Treatment for Adolescents with Depression Study (TADS) randomized controlled trial. *Journal of the American Medical Association*. 2004; 292:807–820. [PubMed: 15315995]
- Tully PJ, Zajac IT, Venning AJ. The structure of anxiety and depression in a normative sample of younger and older Australian adolescents. *Journal of Abnormal Child Psychology*. 2009; 37:717–726. [PubMed: 19238534]
- Vostanis P, Feehan C, Grattan E. Two-year outcome of children treated for depression. *European Child and Adolescent Psychiatry*. 1998; 7:12–18. [PubMed: 9563808]
- Walkup JT, Albano AM, Piacentini J, Birmaher B, Compton SN, Sherrill JT, et al. Cognitive behavior therapy, sertraline, or a combination in childhood anxiety. *The New England Journal of Medicine*. 2008; 359:2753–2766. [PubMed: 18974308]
- Warner V, Mufson L, Weissman MM. Offspring at high and low risk for depression and anxiety: mechanisms of psychiatric disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1995; 34:786–797. [PubMed: 7608053]
- Watson, D. *Mood and temperament*. New York: Guilford Press; 2000.
- Watson D. Rethinking the mood and anxiety disorders: A quantitative hierarchical model for *DSM-V*. *Journal of Abnormal Psychology*. 2005; 114:522–536. [PubMed: 16351375]
- Weersing, VR. The psychotherapy manual: A tool for many tasks. In: Shaffer, D., editor. *Issues in the conduct and interpretation of studies of adolescent depression*; Invited plenary session conducted at the annual meeting of the American Academy of Child and Adolescent Psychiatry; Washington, D. C: 2004 October.
- Weersing VR, Gonzalez A, Campo JV, Lucas AN. Brief behavioral therapy for pediatric anxiety and depression: Piloting an integrated treatment approach. *Cognitive and Behavioral Practice*. 2008; 15:126–139.
- Weissman MM, Warner V, Wickramaratne P, Moreau D, Olfson M. Offspring at risk: Early-onset major depression and anxiety disorders over a decade. *Archives of General Psychiatry*. 1997; 54:932–940. [PubMed: 9337774]
- Weissman MM, Wolk S, Wickramaratne P, Goldstein RB, Adams P, Greenwald S, et al. Children with prepubertal-onset major depressive disorder and anxiety grown up. *Archives of General Psychiatry*. 1999; 56:794–801. [PubMed: 12884885]

- Weisz JR, McCarty CA, Valeri SM. Effects of psychotherapy for depression in children and adolescents: A meta-analysis. *Psychological Bulletin*. 2006; 132:132–149. [PubMed: 16435960]
- Whaley SE, Pinto A, Sigman M. Characterizing interactions between anxious mothers and their children. *Journal of Consulting and Clinical Psychology*. 1999; 67:826–836. [PubMed: 10596505]
- Wilson JK, Rapee RM. The interpretation of negative social events in social phobia with versus without comorbid mood disorder. *Anxiety Disorders*. 2005; 19:245–274.
- Wittchen HU, Beesdo K, Bittner A, Goodwin RD. Depressive episodes—evidence for a causal role of primary anxiety disorders? *European Psychiatry*. 2003; 18:384–393. [PubMed: 14680714]
- Woodward LJ, Fergusson DM. Life course outcomes of young people with anxiety disorders in adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2001; 40:1086–1093. [PubMed: 11556633]

Table 1

Treating Anxiety to Prevent Depression

Type	Description	Processes
	Treating anxiety is sufficient to prevent depression.	
Unspecified Risk Factor	Simply removing anxiety is sufficient to prevent depression. Treatment doesn't explicitly involve treating or preventing depression directly.	Remove anxiety; processes unspecified (could be biology, cognitions, etc.)
Generalization	Treatment for anxiety generalizes, so skills learned serve to prevent later depression	Coping, cognitive restructuring, exposure
Causal Risk Factor	Treating anxiety directly addresses mechanism(s) that lead to depression	Avoidance, social skills. Anxiety as a causal risk factor for depression.
Common Mechanism(s)	Treatment of anxiety directly addresses common mechanism(s)	Reducing the common factor such as NA eliminates both
	Treating anxiety is not sufficient to prevent depression.	
Simultaneous	Treat anxiety and add strategies for preventing depression	add depression prevention
Sequential	Treat anxiety first, then add depression prevention strategies	add depression prevention