



Published in final edited form as:

J Med Chem. 2010 October 14; 53(19): 7048–7060. doi:10.1021/jm1006676.

Synthesis and Biological Evaluation of Bivalent Ligands for the CB1 Receptor

Yanan Zhang^{*,†}, Anne Gilliam[†], Rangan Maitra[†], M. Imad Damaj[‡], Julianne M. Tajuba[†], Herbert H. Seltzman[†], and Brian F. Thomas^{*,†}

[†]Research Triangle Institute, Research Triangle Park, NC 27709

[‡]Department of Pharmacology and Toxicology, Virginia Commonwealth University, Richmond, VA 23298

Abstract

Dimerization or oligomerization of many G protein-coupled receptors, including the CB1 receptor, is now widely accepted and may have significant implications towards medications development targeting these receptor complexes. A library of bivalent ligands composed of two identical CB1 antagonist pharmacophores derived from SR141716 linked by spacers of various lengths were developed. The affinities of these bivalent ligands at CB1 and CB2 receptors were determined using radiolabeled binding assays. Their functional activities were measured using GTP- γ -S accumulation and intracellular calcium mobilization assays. The results suggest that the nature of the linker and its length are crucial factors for optimum interactions of these ligands at CB1 receptor binding sites. Finally, selected bivalent ligands (**5d** and **7b**) were able to attenuate the antinociceptive effects of the cannabinoid agonist CP55,940 in a rodent tail-flick assay. These novel compounds as probes will enable further evaluation of CB1 receptor dimerization and oligomerization, its functional significance, and may prove useful in the development of new therapeutic approaches to G protein-coupled receptor mediated disorders.

Introduction

The endocannabinoid system (ECS) is comprised of the CB1 and CB2 receptors, their endogenous ligands (endocannabinoids), and the proteins involved in endocannabinoid synthesis and inactivation, as well as the intracellular signaling pathways affected by endocannabinoids.¹ Increasing evidence suggest that the endocannabinoid system is critically involved in a variety of physiological and pathological conditions. More importantly, modulation of the endocannabinoid system may hold therapeutic promise in a wide range of disparate diseases such as pain, inflammatory diseases, peripheral vascular disease, appetite enhancement or suppression, and locomotor disorders.² Most of the actions exerted by exogenous cannabinoids or endocannabinoid in the brain are mediated by the CB1 receptor, which belongs to the G-protein-coupled receptors (GPCRs) super family, the largest class of cell surface receptors. GPCRs, including the class A family of which the CB1 receptor is a member, are attractive targets for medication development.

While GPCRs were traditionally considered monomeric, it is now well accepted that many GPCRs, including the CB1 receptor,^{3,4} exist on the cell membrane as homo- and hetero-

^{*} To whom correspondence should be addressed. For Y.Z.: Tel: 919-541-1235. Fax: 919-541-6499. yzhang@rti.org. For B.F.T.: Tel: 919-541-6552. Fax: 919-541-6499. bft@rti.org.

Supporting Information Available: HPLC data of target compounds (**5a-f**, **6a-d**, **7a-f** and **8a-d**). This material is available free of charge via the Internet at <http://pubs.acs.org>.

dimers or higher-order oligomers.⁵ Moreover, receptor oligomerization is often essential for receptor function (e.g., the GABAB receptor)⁶, and can also modulate ligand interaction, activation, signal transduction, and internalization.⁷⁻¹¹ For example, it has been proposed that a μ - δ opioid receptor heterodimer is the fundamental signaling unit that mediates opioid tolerance and dependence through specific signal transducer(s) that recognize and couple to the heterodimer but not to μ -receptor monomers/homomers.¹² In an analogous fashion, modulation of the CB1 receptor dimers or oligomers may offer novel opportunities to uniquely target and manipulate function of the endocannabinoid system.

The importance of GPCR dimerization and oligomerization *in vivo* remains to be elucidated and exploited, largely due to a lack of selective pharmacological tools and immunological reagents. Among various efforts to modulate the GPCR oligomers, bivalent ligands, which are defined as two pharmacophores linked by spacers, represent a unique and promising approach and may provide such a tool.^{13, 14} Bivalent ligands, provided they have suitable functional affinity at the monomeric receptor, are expected to selectively bind with greatly enhanced affinity to ligand recognition sites on heterodimers and oligomers due to the small containment volume for the second pharmacophore after the binding of the first one and the formation of thermodynamically more stable complexes. At the same time, bivalent ligands may display unique properties since they interact with more than one receptor simultaneously. Indeed, bivalent ligands have been developed for variety of G protein-coupled receptor targets, including opioids,^{13, 15} adrenergic,^{16, 17} dopamine,¹⁸ serotonin^{19, 20} and muscarinic receptors.^{21, 22} These bivalent ligands have been shown to be able to selectively target homo- or heterodimers and display unique pharmacological properties as compared to their monomeric subunits. However, to the best of our knowledge, there are no bivalent ligands developed for the CB1 receptor to date.

Here we present our efforts in the design and synthesis of symmetrical bivalent ligands targeting CB1 receptor dimers. The bivalent ligands confer two identical core structure of 1,5-diarylpyrazole derived from **1** (SR141716, or rimonabant, Figure 1) joined by a variety of linkers. Compound **1** was initially reported by Sanofi-Recherche as a highly potent and selective CB1 receptor antagonist/inverse agonist. It was the first drug to selectively block both the *in vitro* and *in vivo* effects of cannabinoids that are mediated by the CB1 receptor. Compound **1** was approved for the treatment of obesity in Europe before its recent withdrawal from the market due to undesirable psychological effects. This compound also shows great promise in many potential therapeutic applications including smoking addiction, drug and alcohol dependence, cognitive disorders, inflammation and arthritis.^{23, 24} By developing bivalent ligands with **1** as the pharmacophore, we aim to affect the binding affinities of these ligands to cannabinoid receptor monomers/dimers and perhaps alter their efficacies or signal transduction pathways as antagonists/inverse agonists. We hereby describe the synthesis and preliminary pharmacological examination of a series of bivalent ligands that possess linkers of various lengths, and describe the results with respect to the optimal linker length for affinity, and their related pharmacological activity in two signal transduction pathways. For comparative purposes, corresponding monovalent ligands were synthesized to evaluate the contribution of the presence of the linkers to activity. In doing so, these bivalent ligands enhance our understanding of the structure and function of CB1 receptor dimers.

Results and Discussions

Bivalent ligand design

In order to focus our efforts on the efficient development of bivalent ligands, we selected a prototypical CB1 receptor inverse agonist, SR141716 (**1**). In addition to the high affinity and potency at the CB1 receptor *in vitro* and *in vivo*, the structure-activity relationships on this

class of compounds have been well studied and documented. This allowed for an informed selection of appropriate positions to attach the linkers to the molecules without likely altering their activity significantly. It also permitted efficient synthesis following known procedures with minimal modifications. In particular, SAR results on this structure class indicate that the 3-carboxamide position generally tolerates the replacement of the 1-aminopiperidyl group with a variety of substituents including alkyl groups and aromatic groups (**2**, Figure 1).²⁵⁻²⁷ Therefore, bivalent ligands linked through the 3-position were initially developed.

A series of bivalent ligands with linkers of varying lengths were synthesized and evaluated in efforts to optimize the linker for bridging of the receptors dimers. The optimal linker lengths, or the distances between the binding sites on neighboring receptors in receptor dimers or oligomers, have been reported on a number of GPCRs. Molecular modeling studies based on the crystal structure of rhodopsin suggested a distance between the individual receptors to be in the vicinity of $\sim 35\text{\AA}$, although the receptor dimer was in a head-to-tail orientation.²⁸ Similarly, molecular modeling on the opioid receptor suggested that the distance between the recognition sites of either the interlocking or contact dimers with a TM5,6-interface is $\sim 27\text{\AA}$, while it's greater ($\sim 32\text{\AA}$) in dimers with TM4,5-interface.¹³ However, during their studies on opioid bivalent ligands, Portoghese and co-workers discovered that optimal activity was obtained when spacers are in the range of 22\AA (~ 19 atoms).²⁹ Neumeyer and coworkers found that bivalent ligands for the opioid receptors having spacers containing 10 methylene units or less displayed the highest affinities.^{30, 31} More recently, a series of adenosine A_{2A} antagonist/dopamine D_2 agonist bivalent ligands were developed where linkers ranged between 26 and 66 atoms.³² Interestingly, affinities of the bivalent ligands to both receptors stayed almost identical with the elongation of the linkers. The authors indicated that linkers with 26 atoms were of sufficient length to allow the bivalent ligands to bind to receptor dimers according to receptor docking experiments, and suggested that the lack of correlation between binding affinity and linker length might be due to the high flexibility of the mixed peptide/polyethylene glycol linkers. Based on these findings and others, linkers between 5 and 23 atoms were initially examined in our laboratory to determine optimal linker length.

Three types of linkers have been considered in the design of the bivalent ligands. The first class investigated were polyethylene glycol linkers. The second category is composed of small peptides (Figure 1). These two classes of linkers are readily available and have been employed in bivalent ligand development by a number of groups.^{13, 22, 33} These linkers are not only readily available but also offer the advantage of gradually increasing the linker length. However, our preliminary results with these two types of linkers failed to show promise. Both compounds **3** and **4** had low affinity in radiolabeled binding and were inactive in GTP- γ -S and calcium assays (data not shown). This is consistent with literature results that suggest hydrophobic groups are generally preferred at this 3-carboxamide position.^{25, 26, 34} Linkers composed of alkylamines were also examined (Figure 2). The selection of these hydrophobic molecules was based on the SAR studies in our laboratory and also by Wiley and co-workers that indicate substitution of the 3-carboxamide with hydrocarbons usually retains or sometimes even improves the affinity or antagonist activity of **1**.^{25, 26, 34} Accordingly, a series of alkyl triamines were selected to construct the bivalent ligands (**5a-f**, Figure 2). A protonatable nitrogen atom was introduced in the middle of the chain in order to reduce the incremental increases in hydrophobicity upon elongation of the alkyl chains. This nitrogen not only provides symmetry of the bivalent ligands, it also facilitates the construction of long alkyl linkers. Additionally, the N-methyl series of analogs (**6a-d**) were prepared to examine the possible hydrogen bonding effects of the alkyl amine linker.

Chemistry

Compound **3** was obtained by coupling between the pyrazole carboxylic acid (**9**), which was readily prepared from commercially available 4-chloropropiophenone in three steps following the procedure developed in our laboratory,^{35, 36} and 1,11-diamino-3,6,9-trioxaundecane using BOP as the coupling agent (Scheme 1). In the preparation of **4**, acid **9** was coupled to glycine methyl ester hydrochloride under standard coupling conditions that employed HOBt, EDCI and triethylamine in THF to give the methyl ester (**10**) in almost quantitative yield. Hydrolysis of **10** in methanolic sodium hydroxide at room temperature furnished **11** in quantitative yield. Coupling of **11** with glycine methyl ester hydrochloride under identical conditions as that of **9**, followed by mild hydrolysis (LiOH, MeOH/THF/H₂O), provided **12** in excellent yield. Finally, reaction of **12** with excess ethylenediamine furnished **4**.

The route to bivalent ligands **5a-f** and monovalent ligands **7a-f** required the use of the N-H triamine linkers **18a-f**. While **18a-b** were commercially available, **18c-f** were prepared in our laboratory as shown in Scheme 2. For **18c-d**, the starting bromoalkyl nitriles (**15c-d**) were commercially available. Intermediates **15e** (m = 8) and **15f** (m = 10) in the preparation of **18e-f** needed to be synthesized from dibromides **13** and **14**, respectively. This was readily accomplished by displacing one of the bromides in these dibromoalkanes with a cyano group using sodium cyanide in DMSO. Thereafter, bis-alkylation of benzylamine with bromides **15c-f** in the presence of potassium carbonate in 1-butanol or dimethylformamide (DMF) furnished amines **16c-f** in excellent yields. Reduction of these nitriles was readily accomplished by hydrogenation catalyzed with Raney Nickel to give **17c-f**. Another hydrogenation using palladium on carbon removed the benzyl groups to afford triamines **18c-f**, which were generally of sufficient purity and were used in the following step without further purification. It is worth noting that the sequence of the hydrogenations was important and debenylation followed by reduction of nitriles failed to give the desired products in satisfactory yields.

The coupling of acid **9** and triamines **18a-f** was then attempted using several methods in order to furnish the bivalent ligands. Initial trials on the coupling employing the acid chloride or activation of acid **9** with agents such as chloroformates or BOP all failed to display selectivity and products with acylation at all three amino sites were obtained as the primary products. Eventually carbonyldiimidazole (CDI) appeared to provide satisfactory selectivity, and the desired products **5a-f**, where acylation occurred at the two primary amino sites, were obtained in reasonable yields (Scheme 3). Under similar conditions, acylation at only one of the primary amino groups could be readily accomplished with the employment of excess triamines **18a-f** to provide **7a-f**.

Following a procedure analogous to that of scheme 2, the N-methyl triamine linkers were prepared as depicted in Scheme 4. Bis-alkylation of methylamine in methanol or methylamine hydrochloride with bromides **15c-f** in ethanol under microwave conditions or heated in sealed pressure tubes provided **19a-d** in almost quantitative yields. Hydrogenation catalyzed by palladium on carbon provided the triamines (**20a-d**) in excellent yields. Similar to Scheme 3, the N-methyl bivalent ligands **6a-d** and the monovalent controls **8a-d** were obtained by coupling reactions between acid **9** and amines **20a-d** using CDI.

Pharmacology

Affinity of bivalent ligands

All the target compounds were evaluated in competition binding assays using both rat whole brain membrane preparations and cells stably transfected with either the human CB1 or CB2 receptors. The receptor binding affinities were determined in competitive displacement

assays using radioligands [³H]-**1** and [³H]-CP55,940. The results are summarized in Tables 1 and 2.

Most of the bivalent ligands displayed nM affinity at the CB1 receptor, albeit somewhat lower than the parent compound **1**. Similarly to **1**, all bivalent ligands and monovalent controls also showed reasonable selectivity for the CB1 receptor over the CB2 receptor, displaying little or no affinity at the CB2 receptor. Noticeably, all compounds exhibited higher affinity (2-3 fold) for the CB1 receptor in the displacement of [³H]**1** than the structurally different [³H]CP55,940. This is in agreement with observations previously reported by Wiley and coworkers where CP55,940 derivatives were usually better ligands in displacement of radiolabeled CP55,940 than **1**.³⁷

Interestingly, the binding affinity of the N-H bivalent ligand series at the CB1 receptor appeared to be sensitive to the length of the linkers (Table 1). Specifically, the affinity initially increased with increasing linker length, and then decreased as the linker was extended. The peak affinity was obtained with **5d** (n = 7), where the linker is composed of 15 atoms, against both radioligands (12.3 nM vs. [³H]CP55,940 and 4.41 nM vs. [³H]**1**). Such an initial increase, followed by a subsequent decrease in affinity, is consistent with observations in the bivalent opioid ligands made by the Portoghese and Neumeyer groups.^{13, 30, 38, 39} Thus, it is hypothesized that linker length is critical for the ability of the bivalent ligands to bind two CB1 receptors simultaneously, as insufficient length would not permit bridging and spacers of excessive length would reduce bridging due to increased confinement volume. Such a transition suggests that bridging of vicinal receptors by bivalent ligands occurs most efficiently with optimal linker length. This hypothesis is also supported by the different pattern that was observed for the corresponding monovalent controls in this series (**7a-f**, Table 1), where the affinity increased as the spacer became longer, with the most potent compound determined to be **7f** (n = 11, 23 atoms). The observation that monovalent ligands display comparable or even higher affinity than the corresponding bivalent ligands (**5e** vs. **7e** and **5f** vs. **7f**) has also been previously reported in bivalent ligands for other GPCRs including opioids, 5-HT₄ and GnRHR.^{19, 30, 33, 40} One possibility is that the linker may represent an additional recognition site and only one pharmacophore is needed when the spacer is of sufficient length. Most significantly, the highest affinity bivalent ligand (**5d**), where the linker is composed of 15 atoms, displays higher affinity (~4 fold for both radioligands) than the corresponding monovalent control (**7d**), indicating that the presence of the second pharmacophore increases the ability for the compounds to bind to the receptor, possibly by simultaneously occupying vicinal recognition sites of neighboring receptors in the receptor dimer. Again, this affinity enhancement of bivalent ligands over their corresponding monovalent ligands has been widely observed in previous studies on the opioid receptors, although it is often a modest (~ 2 fold) difference.³⁰ The optimal linker length of 15 atoms in the present study is consistent with the range reported for bivalent ligands developed for other GPCRs.^{29, 30, 41, 42}

A similar trend in affinity was also observed in the bivalent ligands of the N-Me series (**6a-d**, Table 2) with respect to their ability to compete for [³H]CP55,940 and [³H]**1** binding. Affinity initially increased with linker length, with **6a** (n = 5) and **6b** (n = 7) displaying the greatest ability in displacing either [³H]**1** or [³H]CP55,940. Slightly different than the N-H monovalent ligand series, the binding affinity of **8a-d** at the CB1 receptor initially increased and then remained relatively constant with the elongation of the linker, with **8c** and **8d** displaying almost identical K_i's. Interestingly, when the linkers are of the same length, the affinities of the bivalent ligands from both the N-H and N-Me series are relatively similar; indicating that the presence of the N-methyl group did not appear to interfere with the interaction of the bivalent ligands with the receptors.

Inverse agonist/antagonist activity of bivalent ligands

All compounds were examined in vitro using both [³⁵S]GTP- γ -S accumulation and intracellular calcium mobilization assays to characterize their efficacy, inverse agonist activity, and apparent affinity (pA_2). The results are shown in table 3 and 4.

In the [³⁵S]GTP- γ -S assay using whole rat brain, most dimers and monomers appeared to act as weak inverse agonists. Similar to **1**, most compounds required μ M concentrations to show inverse agonist activity in hCB1 transfectants, and the change from basal activity was relatively modest (<25% decrease in basal binding under the conditions used). However, most compounds potently shifted the concentration-response curve of the agonist CP55,940, indicating that they were high affinity antagonists. Significantly, the pA_2 values against CP55,940 stimulated [³⁵S]-GTP- γ -S binding in hCB1 cells were correlated with their K_i values in both the N-H and N-Me series. Specifically, the pA_2 values first increased and then decreased for the bivalent ligands and always increased for the monovalent ligands. The bivalent ligand with the highest apparent affinity in the N-H series was **5d** whereas **6b** showed the highest pA_2 value in the N-Me series. The higher potencies of the bivalent ligand **5d** and **6b** than the monovalent ligands **7d** and **8b**, respectively, is consistent with the binding of these bivalent ligands to CB1 receptor dimers.

All the compounds were also tested using a calcium mobilization assay as a measure of CB1 receptor function and again showed nM potency. The same trend of an initial increase followed by a subsequent decrease in potency with increasing linker length was observed for bivalent ligands in both series (**5a-f** and **6a-d**), with the exception of **6a**, whereas the potency generally increased and stayed consistent for the monovalent ligands (**7a-f** and **8a-d**). **5c** and **5d** showed the greatest potency in N-H bivalent ligands and **6a** was the most potent N-Me bivalent ligand. However, no potency enhancement at the optimal linker (15 atoms) was observed in this assay between the bivalent and monovalent ligands (**5d** vs. **7d** and **6b** vs. **8b**). While data from the three in vitro assays trended well, differences were noted between K_i , pA_2 and K_e values. This is not surprising as different endpoints and biological systems were experimentally employed for ligand characterization.

Tail-flick Studies

The bivalent ligands with the highest affinity and potency, **5d** and **6b**, and their monovalent controls, **7d** and **8b**, were evaluated for their ability to block the antinociceptive effects of the cannabinoid agonist CP55,940 in a rodent tail-flick assay. In the experiment, the tail was exposed to 55°C warm water and the amount of time taken for the animal to move (flick) its tail away from the heat was recorded. Test compounds or vehicle were administered i.p. at 10 mg/kg i.p. to male mice 30 minutes prior to the administration of vehicle or 1.5 mg/kg CP55,940. Tail-flick times were measured 30 minutes after treatment with CP55,940. Antinociceptive response was calculated as percentage of maximum possible effect.

As shown in Figure 3, a single 10 mg/kg i.p. dose of the bivalent ligand **6b** and its monomeric control **8b** could significantly attenuate the antinociceptive response to CP55,940. However, the N-H analogs (**5d** and **7d**) were considerably less active. It remains to be determined if the differences in potencies in vivo can be attributed to differences in biodistribution or metabolism of the various ligands. Nevertheless, these results suggest that these bivalent compounds, despite their high molecular weight, are able to attenuate nociceptive responses by central and/or peripheral mechanisms by antagonizing the CB1 receptor complexes.

Conclusions

The concept of homo- and heterodimerization has opened new potential avenues for the development of drugs targeted at GPCRs. One emerging approach is to employ bivalent ligands that specifically bind to these receptor dimers. Ideally, bivalent ligands with linkers of optimal length will bind to receptor dimers with greatly enhanced affinity due to the formation of thermodynamically stable complexes. Indeed, significant progress has been made in a number of GPCRs including opioids,^{13, 15} adrenergic,^{16, 17} dopamine,¹⁸ serotonin^{19, 20} and muscarinic receptors.^{21, 22} Most significantly, much success has been recently achieved by Portoghesi and co-workers with bivalent opioid ligands *in vivo*.^{42, 43} In particular, μ -opioid (MOP) agonist/ δ -opioid (DOP) antagonist bivalent ligands were shown to be potent analgesics after systemic administration, but did not produce the tolerance or dependence seen with traditional monovalent opioid analgesics.¹² However, to the best of our knowledge, there are no bivalent ligands developed for the CB1 receptor to date. It is now well established that this receptor is a viable target to treat various indications including smoking addiction, drug and alcohol dependence, metabolic syndrome, cancer, fibrosis and inflammation. The consequences of altered cellular function as a result of dimerization and oligomerization of CB1 receptors are being explored. Availability of specific probes to understand the physiological importance of such interactions hold the key to better understand the role of cannabinoid signaling in the context of health and disease.

In the present study, we synthesized a library of symmetrical bivalent ligands containing two SR141716 moieties joined by aminoalkyl linkers. All the target compounds were evaluated in radiolabeled binding assays at the CB1 and CB2 receptors, functional [³⁵S]GTP- γ -S accumulation assay and functional calcium mobilization assay. In all three assays, a clear trend could be detected where the bivalent ligands showed initially increased and then decreased affinity/activity with elongation of the linkers, whereas the monovalent ligands generally continued increasing or stayed consistent once the linker length was sufficiently long. The fact that the bivalent ligands showing higher affinity and activity than their respective monovalent controls (**5d** vs. **7d** and **6b** vs. **8b**) when the spacer was comprised of 15 atoms in both the radiolabeled binding and [³⁵S]GTP- γ -S assay, respectively, strongly suggests bridging of neighboring receptors. It is worth noting that, although only moderate affinity or potency enhancement was observed for the bivalent over the monovalent ligands in our binding or functional assays, additional evidence in support of this hypothesis has been reported using techniques such as FRET as demonstrated by Russo and coworker in 5-HT₄ receptors.¹⁹ Finally, selected bivalent ligands (**5d** and **7b**) and the corresponding monovalent controls (**6d** and **8b**) were able to attenuate the antinociceptive effects of the cannabinoid agonist CP55,940 in the tail-flick assay. Taken together, further evaluation of this bivalent ligand approach of the CB1 receptor dimerization or oligomerization is clearly needed and may serve as the basis for the ultimate development of new medications.

Experimental

Chemistry

Reactions were conducted under N₂ atmospheres using oven-dried glassware. All solvents and chemicals used were reagent grade. Anhydrous tetrahydrofuran, dichloromethane, and N,N-dimethylformamide (DMF) were purchased from Aldrich and used as such. Unless otherwise mentioned, all reagents and chemicals were purchased from commercial vendors and used as received. Flash column chromatography was carried out on a Teledyne ISCO CombiFlash Companion system using RediSep Rf prepacked columns. Purity and characterization of compounds were established by a combination of HPLC, TLC, gas chromatography mass spectrometry (GC-MS), and NMR analytical techniques described below. ¹H and ¹³CNMR spectra were recorded on a Bruker Avance DPX-300 (300 MHz)

spectrometer and were determined in CHCl₃-d or MeOH-d₄ with tetramethylsilane (TMS) (0.00 ppm) or solvent peaks as the internal reference unless otherwise noted. Chemical shifts are reported in ppm relative to the solvent signal, and coupling constant (J) values are reported in hertz (Hz). Thin-layer chromatography (TLC) was performed on EMD precoated silica gel 60 F254 plates, and spots were visualized with UV light or I₂ detection. Low-resolution mass spectra were obtained using a Waters Alliance HT/Micromass ZQ system (ESI). High-resolution mass spectra were obtained in the Mass Spectrometry Laboratory, Department of Chemistry, University of Michigan. All test compounds were greater than 95% pure as determined by HPLC on an Agilent 1100 system using an Agilent Zorbax SB-Phenyl, 2.1x150 mm, 5 μm column with gradient elution using the mobile phases (A) H₂O containing 0.1% CF₃COOH and (B) MeCN. A flow rate of 0.5 mL/min was used for **5a-f** and **7a-d** and 1.0 mL/min for **6a-f** and **8a-d**.

15c-d and **18a-b** were purchased from Aldrich and were used as such.

5-(4-chlorophenyl)-N-{13-[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazol-3-yl]-13-oxo-3,6,9-trioxa-12-azatridec-1-yl}-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide (3)

Benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (BOP) (116mg, 0.262 mmol) was added to a solution of acid **9** (100 mg, 0.262 mmol) in 15 mL of THF. After 5 min, 1,11-diamino-3,6,9-trioxaundecane (30 mg, 0.157 mmol) was added. The reaction was stirred at room temperature for 1h. The solvent was removed and the resulting slurry was diluted with saturated aqueous NaHCO₃ and extracted with ethyl acetate (2 × 30mL). The combined organic layers were washed with brine and dried. The residue was purified on silica using MeOH-CHCl₃-NH₄OH and EtOAc to give **3** (95mg, 78.9%) as a solid. ¹H NMR (CDCl₃) δ 2.38 (s, 6H), 3.58 (m, 16H), 7.05 (d, J = 9.0, 4H), 7.28 (m, 8H), 7.41 (s, 2H). MS: C₂₅H₂₉Cl₃N₄O₄, [M+H]⁺ 555.2.

Methyl N-[[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazol-3-yl]carbonyl]glycinate (10)

To a solution of acid **9** (2g, 5.24 mmol) in 60 mL of CH₂Cl₂ was added sequentially HOBt (0.78g, 5.76 mmol), EDCI (1.1g, 5.76 mmol) and glycine methyl ester hydrochloride (0.66g, 5.24 mmol). The reaction was stirred at room temperature for 15 min before Et₃N was added. The reaction was stirred for 12h. The reaction was diluted with CH₂Cl₂ (100 mL) and washed with 1N HCl, NaHCO₃ and then brine. The organic layer was dried with Na₂SO₄ and concentrate to give **10** as a white solid. ¹H NMR (CDCl₃) δ 2.36 (s, 3H), 3.78 (s, 3H), 4.22 (d, J = 5.7, 2H), 7.05 (d, J = 6.6, 2H), 7.30 (m, 4H), 7.40 (t, J = 3.0, 1H), 7.43 (s, 1H).

The product was of sufficient purity and was used in the next step without further purification.

N-[[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazol-3-yl]carbonyl]glycine (11)

A solution of **10** in 30 mL of MeOH and 30 mL of 2N NaOH was stirred at room temperature for 16h. The solvent was concentrated *in vacuo* and the resulting solution was washed with ether. The aqueous solution was acidified with 6N HCl and then extracted with EtOAc (3 × 100 mL). The combined organic layers were washed with water, brine and dried with Na₂SO₄. The solvent was removed *in vacuo* to give **11** (2.09g, 90.9% over both steps). ¹H NMR (CDCl₃) δ 2.35 (s, 3H), 4.26 (d, J = 6.0, 2H), 7.06 (d, J = 9.0, 2H), 7.31 (m, 4H), 7.42 (s, 1H), 7.58 (t, 3.0, 1), 10.78 (bs, 1H).

***N*-[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazol-3-yl]carbonyl]glycylglycine (**12**)**

Following the procedure for the preparation of **10**, **11** was coupled to glycine methyl ester hydrochloride to provide the methyl ester in 69.5% yield. ¹H NMR (CDCl₃) δ 2.36 (s, 3H), 3.75 (s, 3H), 4.07 (d, J = 6.0, 2H), 4.16 (d, J = 6.0, 2H), 6.77 (t, J = 3.0, 1H), 7.07 (d, J = 6.0, 2H), 7.29 (m, 4H), 7.43 (s, 1H), 7.53 (t, J = 3.0, 1H).

A solution of the above methyl ester (200mg, 0.40 mmol) and LiOH (25mg, 1.2 mmol) in 10 mL of THF-MeOH (3:1) and 2 mL of water was stirred at room temperature for 12h. The reaction was acidified with 3N HCl and extracted with EtOAc (2 × 25 mL). The combined organic layers were washed with water, brine and dried with Na₂SO₄. The solvent was evaporated to give **12** (175mg, 88.4%) as a white solid. ¹H NMR (CDCl₃) δ 2.31 (s, 3H), 3.78 (s, 2H), 4.09 (s, 2H), 7.19 (d, J = 8.4, 2H), 7.37 (d, J = 8.4, 2H), 7.46 – 7.56 (m, 3H).

***N*-[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazol-3-yl]carbonyl]glycyl-*N*-(2-aminoethyl)glycinamide (**4**)**

To a solution of **12** (140mg, 0.28 mmol) in THF at room temperature was added BOP (125.0mg, 0.28 mmol) and ethylenediamine (9.0 μL, 1.41 mmol). The mixture was stirred for 15 min before Et₃N was added. The reaction was stirred for 12h. The reaction was quenched with water and extracted with EtOAc (2 × 40 mL). The combined organic layers were washed with 1N HCl, saturated NaHCO₃ and brine and then dried with Na₂SO₄. The solvent was removed and the residue was purified on silica gel using MeOH-CHCl₃-NH₄OH and EtOAc to give **4** (35mg, 34.5%) as a solid. ¹H NMR (CDCl₃) δ 2.31 (s, 3H), 2.87 (d, J = 6.0, 2H), 3.34 (d, J = 6.0, 2H), 3.96 (d, J = 6.0, 2H), 4.08 (d, J = 6.0, 2H), 7.06 (m, 3H), 7.20 – 7.40 (m, 5H), 7.42 (s, 1H), 7.80 (t, J = 3.0, 1H). MS: C₂₃H₂₃Cl₃N₆O₃, [M+H]⁺ 537.4.

9-Bromononanenitrile (15e**)**

Sodium cyanide (3.6g, 73.5 mmol) was added in portions to a solution of 1,8-dibromooctane (**13**) (20g, 73.5 mmol) in 50 mL of DMSO at 60°C. After 30 min, the reaction was stopped and allowed to cool to room temperature. The reaction was diluted with 200 mL of diethyl ether and 200 mL of hexane and then washed with water (2 × 50 mL). The organic layer was separated, dried with sodium sulfate and concentrated. The resulting slurry was purified on silica using MeOH/CH₂Cl₂ (1:9) to give **15e** (7.16g, 44.7%) as a colorless oil. ¹H NMR (CDCl₃) δ 1.35 (t, J = 7.5, 3H), 1.82 (s, 3H), 4.32 (q, J = 7.5, 2H), 7.24 (d, J = 9.0, 1H), 7.37 (d, J = 7.5, 1H), 7.48 (s, 1H).

11-Bromoundecanenitrile (15f**)**

15f (6.95g, 42.4%) was obtained from 1,10-dibromodecane (**14**) (20g, 66.6 mmol) as an oil. ¹H NMR (CDCl₃) δ 1.15 – 1.50 (m, 12H), 1.65 (m, 2H), 1.84 (m, 2H), 3.34 (t, J = 7.5, 2H), 3.41 (t, J = 6.0, 2H).

5,5'-(Benzylimino)dipentanenitrile (16c**)**

A mixture of benzylamine (0.5g, 4.67 mmol), potassium carbonate (1.94g, 14.0 mmol) and potassium iodide (0.27g, 1.63 mmol) was heated to 115°C. A solution of 5-bromopentanenitrile in 1-butanol was added drop wise. The resulting mixture was kept at 115°C for 20h. The reaction was allowed to cool to room temperature and then filtered. The solid was washed with diethyl ether (2 × 30 mL). The combined organic layers were extracted with 3N HCl (2 × 20 mL). The aqueous layer was washed with ether and basified with sodium carbonate. The resulting solution was then extracted with ether (3 × 40 mL). The combined organic layers were dried with Na₂SO₄ and concentrated to give **16c** (1.02g,

81.1%) as an oil. $^1\text{H NMR}$ (CDCl_3) δ 1.50 – 1.70 (m, 8H), 2.26 (t, J = 6.6, 4H), 2.42 (t, J = 6.3, 4H), 3.51 (s, 2H), 7.15 – 7.32 (m, 5H).

The product was off sufficient purity and used in the next step without further purification.

7,7'-(Benzylimino)diheptanenitrile (16d)

16d (1.43g, 94.1%) was obtained from benzylamine (0.5g, 4.67 mmol) and 7-bromoheptanenitrile (1.86g, 9.80 mmol) as a colorless oil. $^1\text{H NMR}$ (CDCl_3) δ 1.20 – 1.65 (m, 16H), 2.29 (t, J = 6.9, 4H), 2.39 (t, J = 6.9, 4H), 3.52 (s, 2H), 7.29 (m, 5H).

9,9'-(Benzylimino)dinonanenitrile (16e)

16e (1.79g, 100%) was obtained from benzylamine (0.5g, 4.67 mmol) and **15e** (2.04g, 9.33 mmol) as an oil. $^1\text{H NMR}$ (CDCl_3) δ 1.20 – 1.70 (m, 24H), 2.20 – 2.45 (m, 8H), 3.51 (s, 2H), 2.24 (m, 5H).

11,11'-(Benzylimino)diundecanenitrile (16f)

16f (2.0g, 84.8%) was obtained from benzylamine (0.7g, 6.53 mmol) and **15f** (3.38g, 13.72 mmol). $^1\text{H NMR}$ (CDCl_3) δ 1.20 – 1.50 (m, 24H), 1.65 (m, 4H), 1.83 (m, 4H), 2.35 (t, J = 7.2, 4H), 2.90 (m, 4H), 4.16 (s, 2H), 7.43 (m, 3H), 7.64 (m, 2H).

N-(5-aminopentyl)pentane-1,5-diamine (18c)

A suspension of **16c** (0.5g, 1.86 mmol) and Raney Nickel (0.5g) in ethanol (40 mL), THF (10 mL) and 2N sodium hydroxide (8 mL) was stirred under hydrogen (50 psi) for 20h. The suspension was filtered through Celite and concentrated. The resulting slurry was diluted with water (40 mL) and then extracted with CH_2Cl_2 (2×50 mL). The combined organic layers were washed with brine and dried over Na_2SO_4 to give **17c** as an off-white oil. **17c** was used in the next step without purification.

A suspension of **17c** (0.35g, 1.25mmol), 10% palladium on carbon (40 mg) in ethanol (15 mL) and acetic acid (5 mL) was stirred under 50 psi hydrogen for 3h. The suspension was filtered through Celite and the filtrate was concentrated. To the resulting slurry was added 2N NaOH (20 mL) and extracted with CH_2Cl_2 (2×30 mL). The combined organic layers were washed with water and dried with Na_2SO_4 . The solvent was evaporated in vacuo to give **18c** (228 mg, 91.1% over both steps) as a clear oil. $^1\text{H NMR}$ (CDCl_3) δ 1.15 – 1.55 (m, 12H), 2.53 (t, J = 6.9, 4H), 2.62 (t, J = 6.6, 4H).

N-(7-aminoheptyl)heptane-1,7-diamine (18d)

Following the procedure for the synthesis of **18c**, **18d** (0.85g, 94.6%) was obtained from **16d** (1.2g, 3.69 mmol). $^1\text{H NMR}$ (CDCl_3) δ 1.15 – 1.50 (m, 20H), 2.52 (t, J = 7.2, 4H), 2.61 (t, J = 6.9, 4H).

N-(9-aminononyl)nonane-1,9-diamine (18e)

Following the procedure for the synthesis of **18c**, **18e** (0.425g, 77.5%) was obtained from **16e** (0.7g, 1.83 mmol). $^1\text{H NMR}$ (CDCl_3) δ 1.20 – 1.55 (m, 28H), 2.57 (t, J = 7.2, 4H), 2.66 (t, J = 6.9, 4H).

N-(11-aminoundecyl)undecane-1,11-diamine (18f)

Following the procedure for the synthesis of **18c**, **18f** (0.34g, 84.1%) was obtained from **16f** (0.5g, 1.14 mmol). $^1\text{H NMR}$ (CDCl_3) δ 1.10 – 1.50 (m, 36H), 2.56 (t, J = 7.5, 4H), 2.62 (t, J = 7.2, 4H).

***N,N*-(iminodiethane-2,1-diyl)bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (5a)**

A solution of acid **9** (0.2g, 0.52 mmol), carbonyldiimidazole (85mg, 0.52 mmol) in 5 mL of CH₂Cl₂ was stirred at room temperature for 1h. TLC showed the complete consumption of the starting material. Triamine **18a** (28 μL, 0.26 mmol) was added and the reaction was allowed to stir at room temperature for 3h. The reaction was diluted with CH₂Cl₂ and washed sequentially with NaHCO₃, water and brine. The solution was dried with Na₂SO₄ and concentrated. The resulting slurry was purified on silica using CHCl₃-MeOH-NH₄OH (80:18:2) and EtOAc to give **5a** (0.13g, 61.1%) as a white solid. ¹H NMR (CDCl₃) δ 2.35 (s, 6H), 2.90 (t, J = 6.0, 4H), 3.52 (dt, J₁ = J₂ = 6.0, 4H), 7.05 (d, J = 6.0, 4H), 7.22 – 7.30 (m, 10H), 7.40 (s, 2H). HRMS: C₃₈H₃₁Cl₆N₇O₂, [M+H]⁺ calcd 828.0749, found 828.0768.

***N,N*-(iminodipropane-3,1-diyl)bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (5b)**

Following the procedure for the preparation of **5a**, **5b** was obtained from **18b** in 49.5% yield. ¹H NMR (CDCl₃) δ 1.75 (m, 4H), 2.35 (s, 6H), 2.72 (t, J = 6.0, 4H), 3.45 (dt, J₁ = 9.0, J₂ = 6.0, 4H), 7.05 (d, J = 6.0, 4H), 7.28 (m, 8H), 7.40 (s, 2H), 7.54 (t, J = 3.0, 2H). HRMS: C₄₀H₃₅Cl₆N₇O₂, [M+H]⁺ calcd 856.1062, found 856.1076.

***N,N*-(iminodipentane-5,1-diyl)bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (5c)**

Following the procedure for the preparation of **5a**, **5c** was obtained from **18c** in 42.5% yield. ¹H NMR (CDCl₃) δ 1.37 – 1.64 (m, 12H), 2.37 (s, 6H), 2.59 (t, J = 6.0, 4H), 3.41 (dt, J₁ = 9.0, J₂ = 6.0, 4H), 6.96 (t, J = 3.0, 2H), 7.06 (d, J = 6.0, 4H), 7.26 (m, 8H), 7.43 (s, 2H). HRMS: C₄₄H₄₃Cl₆N₇O₂, [M+H]⁺ calcd 912.1688, found 912.1679.

***N,N*-(iminodiheptane-7,1-diyl)bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (5d)**

Following the procedure for the preparation of **5a**, **5d** was obtained from **18d** in 55.6% yield. ¹H NMR (CDCl₃) δ 1.34 – 1.70 (m, 20H), 2.37 (s, 6H), 2.56 (t, J = 6.0, 4H), 3.41 (dt, J₁ = 6.9, J₂ = 6.6, 4H), 6.95 (t, J = 3.0, 2H), 7.05 (d, J = 6.0, 4H), 7.29 (m, 8H), 7.43 (s, 2H). HRMS: C₄₈H₅₁Cl₆N₇O₂, [M+H]⁺ calcd 968.2314, found 968.2321.

***N,N*-(iminodinonane-9,1-diyl)bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (5e)**

Following the procedure for the preparation of **5a**, **5e** was obtained from **18e** in 42.8% yield. ¹H NMR (CDCl₃) δ 1.26 – 1.59 (m, 28H), 2.37 (s, 6H), 2.60 (t, J = 6.0, 4H), 3.40 (dt, J₁ = 6.6, J₂ = 6.0, 4H), 6.95 (t, J = 3.0, 2H), 7.28 (m, 8H), 7.43 (s, 2H). HRMS: C₅₂H₅₉Cl₆N₇O₂, [M+H]⁺ calcd 1024.2940, found 1024.2972.

***N,N*-(iminodiundecane-11,1-diyl)bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (5f)**

Following the procedure for the preparation of **5a**, **5f** was obtained from **18f** in 38.9% yield. ¹H NMR (CDCl₃) δ 1.25 – 1.61 (m, 36H), 2.37 (s, 6H), 2.61 (t, J = 7.5, 4H), 3.40 (dt, J₁ = 6.9, J₂ = 6.6, 4H), 6.96 (t, J = 6.0, 2H), 7.06 (d, J = 8.4, 4H), 7.29 (m, 8H), 7.42 (s, 2H). HRMS: C₅₆H₆₇Cl₆N₇O₂, [M+H]⁺ calcd 1080.3566, found 1080.3590.

***N*-{2-[(2-aminoethyl)amino]ethyl}-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (7a)**

A solution of acid **9** (50 mg, 0.13 mmol) and carbonyldiimidazole (21mg, 0.13 mmol) in 5 mL of CH₂Cl₂ was stirred at room temperature for 30 min. TLC showed the complete consumption of starting acid. This suspension was then added to a solution of triamine **18a** (41 mg, 0.39 mmol) drop wise. The resulting clear solution was allowed to stir at room temperature for 3h. The reaction was diluted with CH₂Cl₂ and washed sequentially with NaHCO₃, water and brine. The solution was dried with Na₂SO₄ and concentrated. The resulting slurry was purified on silica using MeOH-CHCl₃-NH₄OH (80:18:2) and EtOAc to give **5a** (47 mg, 89.0%) as a solid. ¹H NMR (CDCl₃) δ 2.36 (s, 3H), 2.72 (t, J = 5.7, 2H), 2.80 (t, J = 4.8, 2H), 2.87 (t, J = 6.0, 2H), 3.53 (dt, J₁ = J₂ = 6.0, 2H), 7.05 (d, J = 6.0, 2H), 7.28 (m, 5H), 7.42 (s, 1H). HRMS: C₂₁H₂₂Cl₃N₅O, [M+H]⁺ calcd 466.0968, found 466.0971.

***N*-{3-[(3-aminopropyl)amino]propyl}-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (7b)**

Following the procedure for the preparation of **7a**, **7b** was obtained from **18b** in 67.9% yield. ¹H NMR (CDCl₃) δ 1.64 (m, 2H), 1.79 (m, 2H), 2.37 (s, 3H), 2.67 – 2.77 (m, 6H), 3.53 (dt, J₁ = 5.7, J₂ = 5.4, 2H), 7.05 (d, J = 8.4, 2H), 7.30 (m, 4H), 7.42 (s, 1H), 7.56 (m, 1H). HRMS: C₂₃H₂₆Cl₃N₅O, [M+H]⁺ calcd 494.1281, found 494.1284.

***N*-{5-[(5-aminopentyl)amino]pentyl}-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (7c)**

Following the procedure for the preparation of **7a**, **7c** was obtained from **18c** in 44.9% yield. ¹H NMR (CDCl₃) δ 1.36 – 1.60 (m, 12H), 2.37 (s, 3H), 2.59 – 2.72 (m, 6H), 3.43 (dt, J₁ = 9.3, J₂ = 6.6, 2H), 6.95 (t, J = 3.0, 1H), 7.06 (d, J = 8.4, 2H), 7.29 (m, 4H), 7.43 (s, 1H). HRMS: C₂₇H₃₄Cl₃N₅O, [M+H]⁺ calcd 550.1907, found 550.1909.

***N*-{7-[(7-aminoheptyl)amino]heptyl}-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (7d)**

Following the procedure for the preparation of **7a**, **7d** was obtained from **18d** in 35.2% yield. ¹H NMR (CDCl₃) δ 1.26 – 1.62 (m, 20H), 2.38 (s, 3H), 2.56 (m, 4H), 2.67 (t, J = 6.6, 2H), 3.41 (dt, J₁ = 6.9, J₂ = 6.6, 2H), 6.96 (t, J = 3.0, 1H), 7.06 (d, J = 6.9, 2H), 7.29 (m, 4H), 7.43 (s, 1H). HRMS: C₃₁H₄₂Cl₃N₅O, [M+H]⁺ calcd 606.2533, found 606.2536.

***N*-{9-[(9-aminononyl)amino]nonyl}-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (7e)**

Following the procedure for the preparation of **7a**, **7e** was obtained from **18e** in 47.2% yield. ¹H NMR (CDCl₃) δ 1.18 – 1.62 (m, 28H), 2.38 (s, 3H), 2.58 (t, J = 7.2, 4H), 2.67 (t, J = 6.9, 2H), 3.40 (dt, J₁ = 6.9, J₂ = 6.6, 2H), 6.96 (t, J = 3.0, 1H), 7.28 (m, 4H), 7.43 (s, 1H). HRMS: C₃₅H₅₀Cl₃N₅O, [M+H]⁺ calcd 662.3159, found 662.3150.

***N*-{11-[(11-aminoundecyl)amino]undecyl}-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (7f)**

Following the procedure for the preparation of **7a**, **7f** was obtained from **18f** in 46.7% yield. ¹H NMR (CDCl₃) δ 1.20 – 1.65 (m, 36H), 2.38 (s, 3H), 2.58 (t, J = 7.2, 4H), 2.67 (t, J = 6.9, 2H), 3.40 (dt, J₁ = 6.9, J₂ = 6.6, 2H), 6.95 (t, J = 3.0, 1H), 7.05 (d, J = 8.4, 2H), 7.28 (m, 4H), 7.43 (s, 1H). HRMS: C₃₉H₅₈Cl₃N₅O, [M+H]⁺ calcd 718.3785, found 718.3784.

5,5'-(Methylimino)dipentanenitrile (19a)

A pressure tube equipped with a mixture of methyl amine hydrochloride (1g, 16 mmol), **15c** (3.7 mL, 32 mmol), potassium carbonate (4.4g, 32 mmol) and potassium iodide (0.53g, 3.2 mmol) in 20 mL of ethanol was heated to 110°C for 16 hours. The reaction was cooled to room temperature and the solvent was removed. The resulting slurry was partitioned between ethyl acetate and water. The organic layer was washed with brine and dried. The resulting residue was purified by chromatography on silica gel to give **19a** (1.24g, 40.1%) as an off-white oil. ¹H NMR (CDCl₃) δ 1.55 – 1.85 (m, 8H), 2.19 (s, 3H), 2.38 (m, 8H).

7,7'-(Methylimino)diheptanenitrile (19b)

19b was synthesized from **15d** in 48.8% yield following procedure for **19a**. ¹H NMR (CDCl₃) 1.25 -1.70 (m, 16H), 2.18 (s, 3H), 2.33 (m, 8H).

9,9'-(Methylimino)dinonanenitrile (19c)

19c was synthesized from **15e** in 55.0% yield following procedure for **19a**. ¹H NMR (CDCl₃) 1.25 – 1.50 (m, 20H), 1.65 (m, 4H), 2.19 (s, 3H), 2.31 (m, 8H).

11,11'-(Methylimino)diundecanenitrile (19d)

19d was synthesized from **15f** in 42.5% yield following procedure for **19a**. ¹H NMR (CDCl₃) 1.20 – 1.50 (m, 24H), 1.66 (m, 4H), 1.85 (m, 4H), 2.36 (t, J = 3.9, 4H), 2.78 (s, 3H), 3.01 (t, J = 8.4, 4H).

N-(5-aminopentyl)-N-methylpentane-1,5-diamine (20a)

20a was synthesized from **19a** in 59.0% yield following procedure for **17c**. ¹H NMR (CDCl₃) δ 1.15 – 1.55 (m, 12H), 2.20 (s, 3H), 2.31 (t, J = 7.5, 4H), 2.69 (t, J = 6.9, 4H).

N-(7-aminoheptyl)-N-methylheptane-1,7-diamine (20b)

20b was synthesized from **19b** in 73.8% yield following procedure for **17c**. ¹H NMR (CDCl₃) δ 1.20 – 1.50 (m, 24H), 2.18 (s, 3H), 2.29 (t, J = 7.5, 4H), 2.67 (t, J = 6.9, 4H).

N-(9-aminononyl)-N-methylnonane-1,9-diamine (20c)

20c was synthesized from **19c** in 99.0% yield following procedure for **17c**. ¹H NMR (CDCl₃) δ 1.20 – 1.55 (m, 28H), 2.19 (s, 3H), 2.29 (t, J = 7.8, 4H), 2.67 (t, J = 6.9, 4H).

N-(11-aminoundecyl)-N-methylundecane-1,11-diamine (20d)

20d was synthesized from **19d** in 79.0% yield following procedure for **17c**. ¹H NMR (CDCl₃) δ 1.20 – 1.50 (m, 36H), 2.19 (s, 3H), 2.29 (t, J = 7.8, 4H), 2.67 (t, J = 6.9, 4H).

N,N-[(methylimino)dipentane-5,1-diyl]bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide] (6a)

Following the procedure for the preparation of **5a**, **6a** was obtained from **20a** in 50.2% yield. ¹H NMR (CDCl₃) δ 1.37 – 1.65 (m, 12H), 2.24 (s, 3H), 2.38 (m, 10H), 3.41 (dt, J₁ = 6.6, J₂ = 6.0, 4H), 7.00 (t, J = 3.0, 2H), 7.06 (d, J = 6.0, 4H), 7.28 (m, 8H), 4.23 (s, 2H). HRMS: C₄₅H₄₅Cl₆N₇O₂, [M+H]⁺ calcd 926.1844, found 926.1866.

N,N-[(methylimino)diheptane-7,1-diyl]bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide] (6b)

Following the procedure for the preparation of **5a**, **6b** was obtained from **20b** in 40.3% yield. ¹H NMR (CDCl₃) δ 1.26 -1.60 (m, 20H), 2.18 (s, 3H), 2.28 (t, J = 6.0, 4H), 2.37 (s,

6H), 3.41 (dt, $J_1 = 6.6$, $J_2 = 6.0$, 4H), 6.95 (t, $J = 3.0$, 2H), 7.06 (d, $J = 6.0$, 4H), 7.28 (m, 8H), 7.42 (s, 2H). HRMS: $C_{49}H_{53}Cl_6N_7O_2$, $[M+H]^+$ calcd 982.2470, found 982.2482.

***N,N*-[(methylimino)dinonane-9,1-diyl]bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (6c)**

Following the procedure for the preparation of **5a**, **6c** was obtained from **20c** in 53.5% yield. 1H NMR ($CDCl_3$) δ 1.28 – 1.50 (m, 24H), 1.50 – 1.65 (m, 4H), 2.19 (s, 3H), 2.29 (t, $J = 7.8$, 4H), 2.38 (s, 6H), 3.40 (dt, $J_1 = 6.9$, $J_2 = 6.6$, 4H), 6.96 (t, $J = 3.0$, 2H), 7.06 (d, $J = 8.4$, 4H), 7.29 (m, 8H), 7.43 (s, 2H). HRMS: $C_{53}H_{61}Cl_6N_7O_2$, $[M+H]^+$ calcd 1038.3096, found 1038.3094.

***N,N*-[(methylimino)diundecane-11,1-diyl]bis[5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (6d)**

Following the procedure for the preparation of **5a**, **6d** was obtained from **20d** in 43.8% yield. 1H NMR ($CDCl_3$) δ 1.20 – 1.65 (m, 36H), 2.29 (s, 3H), 2.41 (m, 10H), 3.42 (dt, $J_1 = 9.0$, $J_2 = 6.0$, 4H), 6.94 (t, $J = 3.0$, 2H), 7.05 (d, $J = 9.0$, 4H), 7.28 (m, 8H), 7.42 (s, 2H). HRMS: $C_{57}H_{69}Cl_6N_7O_2$, $[M+H]^+$ calcd 1094.3722, found 1094.3715.

***N*-[5-(5-aminopentyl)(methylamino)pentyl]-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (8a)**

Following the procedure for the preparation of **7a**, **8a** was obtained from **20a** in 60.8% yield. 1H NMR ($CDCl_3$) δ 1.22 – 1.70 (m, 12H), 2.20 (s, 3H), 2.32 (t, $J = 7.5$, 4H), 2.37 (s, 3H), 2.69 (t, $J = 6.9$, 2H), 2.42 (dt, $J_1 = 6.9$, $J_2 = 6.6$, 2H), 6.99 (t, $J = 3.0$, 1H), 7.06 (d, $J = 6.6$, 2H), 7.29 (m, 4H), 7.43 (s, 1H). HRMS: $C_{28}H_{36}Cl_3N_5O$, $[M+H]^+$ calcd 564.2064, found 564.2073.

***N*-[7-(7-aminoheptyl)(methylamino)heptyl]-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (8b)**

Following the procedure for the preparation of **7a**, **8b** was obtained from **20b** in 29.5% yield. 1H NMR ($CDCl_3$) δ 1.20 – 1.65 (m, 20H), 2.19 (s, 3H), 2.29 (t, $J = 7.8$, 4H), 2.38 (s, 3H), 2.67 (t, $J = 6.6$, 2H), 2.41 (dt, $J_1 = 6.9$, $J_2 = 6.6$, 2H), 6.95 (t, $J = 3.0$, 1H), 7.06 (d, $J = 6.9$, 2H), 7.28 (m, 4H), 7.43 (s, 1H). HRMS: $C_{32}H_{44}Cl_3N_5O$, $[M+H]^+$ calcd 620.2690, found 620.2687.

***N*-[9-(9-aminononyl)(methylamino)nonyl]-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (8c)**

Following the procedure for the preparation of **7a**, **8c** was obtained from **20c** in 34.9% yield. 1H NMR ($CDCl_3$) δ 1.20 – 1.65 (m, 28H), 2.19 (s, 3H), 2.29 (t, $J = 9.0$, 4H), 2.38 (s, 3H), 2.67 (s, 3H), 3.41 (dt, $J_1 = 9.0$, $J_2 = 6.0$, 2H), 6.95 (t, $J = 3.0$, 1H), 7.05 (d, $J = 9.0$, 2H), 7.28 (m, 4H), 7.43 (s, 1H). HRMS: $C_{36}H_{52}Cl_3N_5O$, $[M+H]^+$ calcd 676.3316, found 676.3312.

***N*-[11-(11-aminoundecyl)(methylamino)undecyl]-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (8d)**

Following the procedure for the preparation of **7a**, **8d** was obtained from **20b** in 40.6% yield. 1H NMR ($CDCl_3$) δ 1.20 – 1.60 (m, 36H), 2.20 (s, 3H), 2.29 (t, $J = 7.8$, 4H), 2.38 (s, 3H), 3.40 (dt, $J_1 = 6.9$, $J_2 = 6.6$, 2H), 6.95 (t, $J = 3.0$, 1H), 7.07 (d, $J = 6.6$, 2H), 7.29 (m, 4H), 7.43 (s, 1H). HRMS: $C_{40}H_{60}Cl_3N_5O$, $[M+H]^+$ calcd 732.3942, found 732.3947.

Receptor Binding Assays

CB1 and CB2 Receptor Binding Assays

The CB1 receptor binding assay involved membranes isolated from a HEK-293 expression system whereas the CB2 receptor was expressed in CHO-K1 cells (Sigma-Aldrich Chemical Co., St. Louis, Mo). The methods used for performing binding assays in transfected cells expressing human CB1 or CB2 receptors were similar to those previously described for rat brain membrane preparations.^{25, 35} Binding was initiated with the addition of 40 μ g of cell membrane proteins to assay tubes containing [³H]CP-55,940 (ca. 130 Ci/mmol) or [³H]-1 (ca. 22.4 Ci/mmol), a test compound (for displacement studies), and a sufficient quantity of buffer (50 mM Tris•HCl, 1 mM EDTA, 3 mM MgCl₂, 5 mg/mL BSA, pH 7.4) to bring the total incubation volume to 0.5 mL. All assays were performed in polypropylene test tubes. In the displacement assays, the concentrations of [³H]CP-55,940 and [³H]-1 7.2 nM and 20 nM, respectively. Nonspecific binding was determined by the inclusion of 10 μ M unlabeled CP-55,940, or **1**. All cannabinoid analogs were prepared by suspension in buffer A from a 1 mg/mL ethanol stock. Following incubation at 30 °C for 1 h, binding was terminated by vacuum filtration through GF/C glass fiber filter plates (Packard, Meriden, CT, pretreated in buffer B for at least 1 h) in a 96-well sampling manifold (Millipore, Bedford, MA). Reaction vessels were washed twice with 4 mL of ice cold buffer (50 mM Tris•HCl, 1 mg/mL BSA). The filter plates were air-dried and sealed on the bottom. Liquid scintillate was added to the wells and the top sealed. After incubating the plates in cocktail for at least 2 h, the radioactivity present was determined by liquid scintillation spectrometry. Assays were done in duplicate, and results represent combined data from three to six independent experiments. Saturation and displacement data were analyzed by unweighted nonlinear regression of receptor binding data. For displacement studies, curve-fitting and IC₅₀ calculation were done with GraphPad Prism (GraphPad Software, Inc., San Diego, CA), which fits the data to one and two-site models and compares the two fits statistically.

GTP- γ -[³⁵S] assay

GTP- γ -[³⁵S] assays were performed to determine the ability of target compounds to shift the binding curves of the agonist CP55,940 or **1**. Reaction mixtures consisted of either CP55,940 (2.5 pM to 25 μ M) or **1** (10 pM to 100 μ M), 20 μ M GDP, and 100 pM GTP- γ -[³⁵S] in 50 mM Tris•HCl, pH 7.4, 1 mM EDTA, 5 mM MgCl₂, 100 mM NaCl, and 1 mg/mL BSA. The effects of these compounds on agonist binding were compared at concentrations of 1, 10, and 100 nM vs. reactions with no antagonist in a final reaction mixture volume of 0.5 mL. Binding was determined using membrane preparations as previously described. Data analysis was performed using global nonlinear regression analysis of the dose–response curves (Prism, GraphPad), and pA₂ values were calculated. The calculations were performed with the slope of the Schild line constrained to 1, as well as unconstrained, and an *F*-test (*P* < 0.05) was used to determine the best model.

Calcium mobilization assay

Calcium mobilization was performed in CHO cells co-expressing G α 16 protein and the human CB1 receptor cDNAs. Activation of CB1 receptor leads to coupling of this receptor to the promiscuous G α 16 protein and consequent mobilization of intracellular calcium. In the assay, the apparent agonist dissociation equilibrium constants (K_e) value of each compound was determined by running a 6-point half-log CP55,940 concentration response curve in the presence and absence of a single concentration of antagonist.⁴⁴ The concentration of antagonist was chosen such that it caused at least a 2-fold increase (shift to the right) in the CP55,940 curve but did not exceed 10 μ M to retain pharmacological relevance. A three-parameter logistic equation was fit to the concentration response data with Prism (GraphPad Software; San Diego, CA) to calculate K_e. These values were

reported as the mean \pm SEM from at least three independent experiments. **1** was employed as the positive control (antagonist) for inhibition of CB1 activity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institute of Health/National Institute on Drug Abuse (NIDA) Grants DA019217 (B.F.T.), DA026582 (Y.Z.) and NIAAA grant AA017514 (R.M.).

References

1. De Petrocellis L, Cascio MG, Di Marzo V. The endocannabinoid system: a general view and latest additions. *Br J Pharmacol*. 2004; 141(5):765–74. [PubMed: 14744801]
2. Pacher P, Batkai S, Kunos G. The endocannabinoid system as an emerging target of pharmacotherapy. *Pharmacol Rev*. 2006; 58(3):389–462. [PubMed: 16968947]
3. Mackie K. Cannabinoid receptor homo- and heterodimerization. *Life Sci*. 2005; 77(14):1667–73. [PubMed: 15978631]
4. Wager-Miller J, Westenbroek R, Mackie K. Dimerization of G protein-coupled receptors: CB1 cannabinoid receptors as an example. *Chem Phys Lipids*. 2002; 121(1-2):83–9. [PubMed: 12505693]
5. George SR, O'Dowd BF, Lee SP. G-protein-coupled receptor oligomerization and its potential for drug discovery. *Nat Rev Drug Discov*. 2002; 1(10):808–20. [PubMed: 12360258]
6. Lee SP, O'Dowd BF, Ng GY, Varghese G, Akil H, Mansour A, Nguyen T, George SR. Inhibition of cell surface expression by mutant receptors demonstrates that D2 dopamine receptors exist as oligomers in the cell. *Mol Pharmacol*. 2000; 58(1):120–8. [PubMed: 10860933]
7. Franco R, Casado V, Cortes A, Mallol J, Ciruela F, Ferre S, Lluís C, Canela EI. G-protein-coupled receptor heteromers: function and ligand pharmacology. *Br J Pharmacol*. 2008; 153 1:S90–8. [PubMed: 18037920]
8. Rios CD, Jordan BA, Gomes I, Devi LA. G-protein-coupled receptor dimerization: modulation of receptor function. *Pharmacol Ther*. 2001; 92(2-3):71–87. [PubMed: 11916530]
9. Minneman KP. Heterodimerization and surface localization of G protein coupled receptors. *Biochem Pharmacol*. 2007; 73(8):1043–50. [PubMed: 17011524]
10. Milligan G. G protein-coupled receptor dimerization: function and ligand pharmacology. *Mol Pharmacol*. 2004; 66(1):1–7. [PubMed: 15213289]
11. Milligan G. G-protein-coupled receptor heterodimers: pharmacology, function and relevance to drug discovery. *Drug Discov Today*. 2006; 11(11-12):541–9. [PubMed: 16713906]
12. Daniels DJ, Lenard NR, Etienne CL, Law PY, Roerig SC, Portoghese PS. Opioid-induced tolerance and dependence in mice is modulated by the distance between pharmacophores in a bivalent ligand series. *Proc Natl Acad Sci U S A*. 2005; 102(52):19208–13. [PubMed: 16365317]
13. Portoghese PS. From models to molecules: opioid receptor dimers, bivalent ligands, and selective opioid receptor probes. *J Med Chem*. 2001; 44(14):2259–69. [PubMed: 11428919]
14. Messer WS Jr. Bivalent ligands for G protein-coupled receptors. *Curr Pharm Des*. 2004; 10(17):2015–20. [PubMed: 15279542]
15. Peng X, Neumeyer JL. Kappa receptor bivalent ligands. *Curr Top Med Chem*. 2007; 7(4):363–73. [PubMed: 17305578]
16. Zheng W, Lei L, Lalchandani S, Sun G, Feller DR, Miller DD. Yohimbine dimers exhibiting binding selectivities for human α 2a- versus α 2b-adrenergic receptors. *Bioorg Med Chem Lett*. 2000; 10(7):627–30. [PubMed: 10762040]
17. Lalchandani SG, Lei L, Zheng W, Suni MM, Moore BM, Liggett SB, Miller DD, Feller DR. Yohimbine dimers exhibiting selectivity for the human α 2C-adrenoceptor subtype. *J Pharmacol Exp Ther*. 2002; 303(3):979–84. [PubMed: 12438517]

18. Tamiz AP, Bandyopadhyay BC, Zhang J, Flippen-Anderson JL, Zhang M, Wang CZ, Johnson KM, Tella S, Kozikowski AP. Pharmacological and behavioral analysis of the effects of some bivalent ligand-based monoamine reuptake inhibitors. *J Med Chem.* 2001; 44(10):1615–22. [PubMed: 11334571]
19. Russo O, Berthouze M, Giner M, Soulier JL, Rivail L, Sicsic S, Lezoualc'h F, Jockers R, Berque-Bestel I. Synthesis of specific bivalent probes that functionally interact with 5-HT(4) receptor dimers. *J Med Chem.* 2007; 50(18):4482–92. [PubMed: 17676726]
20. Meltzer PC, Kryatova O, Pham-Huu DP, Donovan P, Janowsky A. The synthesis of bivalent 2beta-carbomethoxy-3beta-(3,4-dichlorophenyl)-8-heterobicyclo[3.2.1]octane s as probes for proximal binding sites on the dopamine and serotonin transporters. *Bioorg Med Chem.* 2008; 16(4):1832–41. [PubMed: 18053732]
21. Christopoulos A, Grant MK, Ayoubzadeh N, Kim ON, Sauerberg P, Jeppesen L, El-Fakahany EE. Synthesis and pharmacological evaluation of dimeric muscarinic acetylcholine receptor agonists. *J Pharmacol Exp Ther.* 2001; 298(3):1260–8. [PubMed: 11504829]
22. Rajeswaran WG, Cao Y, Huang XP, Wroblewski ME, Colclough T, Lee S, Liu F, Nagy PI, Ellis J, Levine BA, Nocka KH, Messer WS Jr. Design, synthesis, and biological characterization of bivalent 1-methyl-1,2,5,6-tetrahydropyridyl-1,2,5-thiadiazole derivatives as selective muscarinic agonists. *J Med Chem.* 2001; 44(26):4563–76. [PubMed: 11741475]
23. Beardsley PM, Thomas BF. Current evidence supporting a role of cannabinoid CB1 receptor (CB1R) antagonists as potential pharmacotherapies for drug abuse disorders. *Behav Pharmacol.* 2005; 16(5-6):275–96. [PubMed: 16148435]
24. Bifulco M, Grimaldi C, Gazzero P, Pisanti S, Santoro A. Rimonabant: just an antiobesity drug? Current evidence on its pleiotropic effects. *Mol Pharmacol.* 2007; 71(6):1445–56. [PubMed: 17327463]
25. Francisco ME, Seltzman HH, Gilliam AF, Mitchell RA, Rider SL, Pertwee RG, Stevenson LA, Thomas BF. Synthesis and structure-activity relationships of amide and hydrazide analogues of the cannabinoid CB(1) receptor antagonist N-(piperidinyl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide (SR141716). *J Med Chem.* 2002; 45(13):2708–19. [PubMed: 12061874]
26. Thomas BF, Francisco ME, Seltzman HH, Thomas JB, Fix SE, Schulz AK, Gilliam AF, Pertwee RG, Stevenson LA. Synthesis of long-chain amide analogs of the cannabinoid CB1 receptor antagonist N-(piperidinyl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide (SR141716) with unique binding selectivities and pharmacological activities. *Bioorg Med Chem.* 2005; 13(18):5463–74. [PubMed: 15994087]
27. Dyck B, Goodfellow VS, Phillips T, Grey J, Haddach M, Rowbottom M, Naeve GS, Brown B, Saunders J. Potent imidazole and triazole CB1 receptor antagonists related to SR141716. *Bioorg Med Chem Lett.* 2004; 14(5):1151–4. [PubMed: 14980654]
28. Palczewski K, Kumasaka T, Hori T, Behnke CA, Motoshima H, Fox BA, Le Trong I, Teller DC, Okada T, Stenkamp RE, Yamamoto M, Miyano M. Crystal structure of rhodopsin: A G protein-coupled receptor. *Science.* 2000; 289(5480):739–45. [PubMed: 10926528]
29. Portoghese PS, Larson DL, Sayre LM, Yim CB, Ronsisvalle G, Tam SW, Takemori AE. Opioid agonist and antagonist bivalent ligands. The relationship between spacer length and selectivity at multiple opioid receptors. *J Med Chem.* 1986; 29(10):1855–61. [PubMed: 3020244]
30. Neumeyer JL, Zhang A, Xiong W, Gu XH, Hilbert JE, Knapp BI, Negus SS, Mello NK, Bidlack JM. Design and synthesis of novel dimeric morphinan ligands for kappa and micro opioid receptors. *J Med Chem.* 2003; 46(24):5162–70. [PubMed: 14613319]
31. Peng X, Knapp BI, Bidlack JM, Neumeyer JL. Synthesis and preliminary in vitro investigation of bivalent ligands containing homo- and heterodimeric pharmacophores at mu, delta, and kappa opioid receptors. *J Med Chem.* 2006; 49(1):256–62. [PubMed: 16392810]
32. Soriano A, Ventura R, Molero A, Hoen R, Casado V, Cortes A, Fanelli F, Albericio F, Lluís C, Franco R, Royo M. Adenosine A2A receptor-antagonist/dopamine D2 receptor-agonist bivalent ligands as pharmacological tools to detect A2A-D2 receptor heteromers. *J Med Chem.* 2009; 52(18):5590–602. [PubMed: 19711895]

33. Bongers KM, van den Berg RJ, Heitman LH, IJ AP, Oosterom J, Timmers CM, Overkleeft HS, van der Marel GA. Synthesis and evaluation of homo-bivalent GnRHR ligands. *Bioorg Med Chem*. 2007; 15(14):4841–56. [PubMed: 17517510]
34. Wiley JL, Jefferson RG, Grier MC, Mahadevan A, Razdan RK, Martin BR. Novel pyrazole cannabinoids: insights into CB(1) receptor recognition and activation. *J Pharmacol Exp Ther*. 2001; 296(3):1013–22. [PubMed: 11181936]
35. Zhang Y, Burgess JP, Brackeen M, Gilliam A, Mascarella SW, Page K, Seltzman HH, Thomas BF. Conformationally constrained analogues of N-(piperidinyl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide (SR141716): design, synthesis, computational analysis, and biological evaluations. *J Med Chem*. 2008; 51(12):3526–39. [PubMed: 18512901]
36. Seltzman HH, Carroll FI, Burgess JP, Wyrick CD, Burch DF. Tritiation of SR141716 by metallation-iodination-reduction: tritium-proton nOe study. *Journal of Labelled Compounds & Radiopharmaceuticals*. 2002; 45(1):59–70.
37. Wiley, JL.; Wang, P.; Mahadevan, A.; Razdan, RK.; Martin, BR. *Pyrazole Cannabinoids With Non-CB1, Non-CB2 Activity*. International Cannabinoid Research Society 19th Symposium; St Charles, Illinois, USA. 2009.
38. Portoghese PS, Edward E. Smitsman-Bristol-Myers Squibb Award Address. The role of concepts in structure-activity relationship studies of opioid ligands. *J Med Chem*. 1992; 35(11):1927–37. [PubMed: 1317919]
39. Neumeyer JL, Peng X, Knapp BI, Bidlack JM, Lazarus LH, Salvadori S, Trapella C, Balboni G. New opioid designed multiple ligand from Dmt-Tic and morphinan pharmacophores. *J Med Chem*. 2006; 49(18):5640–3. [PubMed: 16942040]
40. Bongers KM, van den Berg RJ, Knijnenburg AD, Heitman LH, Ijzerman AP, Oosterom J, Timmers CM, Overkleeft HS, van der Marel GA. Synthesis and evaluation of homodimeric GnRHR antagonists having a rigid bis-propargylated benzene core. *Bioorg Med Chem*. 2008; 16(7):3744–58. [PubMed: 18282756]
41. Bhushan RG, Sharma SK, Xie Z, Daniels DJ, Portoghese PS. A bivalent ligand (KDN-21) reveals spinal delta and kappa opioid receptors are organized as heterodimers that give rise to delta(1) and kappa(2) phenotypes. *J Med Chem*. 2004; 47(12):2969–72. [PubMed: 15163177]
42. Daniels DJ, Kulkarni A, Xie Z, Bhushan RG, Portoghese PS. A bivalent ligand (KDAN-18) containing delta-antagonist and kappa-agonist pharmacophores bridges delta2 and kappa1 opioid receptor phenotypes. *J Med Chem*. 2005; 48(6):1713–6. [PubMed: 15771416]
43. Xie Z, Bhushan RG, Daniels DJ, Portoghese PS. Interaction of bivalent ligand KDN21 with heterodimeric delta-kappa opioid receptors in human embryonic kidney 293 cells. *Mol Pharmacol*. 2005; 68(4):1079–86. [PubMed: 16006595]
44. Kosterlitz HW, Lees GM, Wallis DI, Watt AJ. Non-specific inhibitory effects of morphine-like drugs on transmission in the superior cervical ganglion and guinea-pig isolated ileum. *Br J Pharmacol*. 1968; 34(3):691P–692P.




Figure 1.
SR141716 (**1**) and 3- substituted analogs with alkyl or polar linkers




Figure 2.
Bivalent and monovalent ligands with triamine linkers







Figure 3. Blockade of the antinociceptive effect of CP55,940 by bivalent and monovalent ligands. Results are expressed as the percent maximal possible effect (% MPE, where % MPE = [(test _ control)/(maximum latency _ control) · 100]) as defined by a 10 sec cut-off for the noxious stimulus. Significant differences ($p < 0.05$) from vehicle-CP55,940 treated controls denoted with an asterisk (*).

**Scheme 1.**

Synthesis of compounds **3** and **4**.


Reagents and conditions: a). Benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (BOP), 1,11-diamino-3,6,9-trioxaundecane, THF; b). Glycine methyl ester hydrochloride, HOBt, EDCI, Et₃N, CH₂Cl₂; c). NaOH, MeOH/H₂O; d). LiOH, THF/MeOH/H₂O; e). Ethylenediamine, HOBt, EDCI, Et₃N, CH₂Cl₂.

**Scheme 2.**Synthesis of N-H triamine linkers **18c-f**Reagents and conditions: (a). NaCN, K₂CO₃, DMSO; (b). benzylamine, K₂CO₃, 1-butanol or DMF, 100° C; (c). H₂, Raney Nickel, ethanol, 2N. NaOH; (d). H₂, Pd/C, ethanol.

**Scheme 3.**

Synthesis of bivalent ligands **5a-f** and monovalent ligands **7a-f**

Reagents and Conditions: CDI (0.5 equiv. for **5a-f**, 3 equiv. for **7a-f**), CH_2Cl_2 .

**Scheme 4.**

Synthesis of N-Me bivalent ligands **6a-d** and monovalent ligands **8a-d**

Reagents and Conditions: (a). Methylamine in methanol or methylamine hydrochloride, K_2CO_3 , ethanol, microwave or heated in pressure tube; (b). H_2 , Pd/C, ethanol; (c). CDI (0.5 equiv. for **6a-d**, 3 equiv. for **8a-d**), CH_2Cl_2 .

Table 1
Binding affinities of N-H bivalent ligands **5a-f** and **7a-f** against [³H]-CP55,940 and [³H]-1

Cmpd	n	Linker (atoms)	Displacement Assay vs. Tritiated Ligands: Ki (nM) in hCB1 (n=2)			Displacement vs. ³ H-CP55,940: Ki (nM) in hCB2		
			³ H-CP55,940	SEM	³ H-1	SEM	SEM	CBI/CB2
1			6.18	1.2	1.18	0.1	313	50.6
5a	2	5	229	75.0	94.0	8.00	1285	5.6
5b	3	7	174	1.0	41.9	5.40	496	2.9
5c	5	11	68.1	12.6	30.4	4.40	451	6.6
5d	7	15	12.3	1.10	4.41	0.34	553	45.0
5e	9	19	54.1	16.3	57.4	44.7	a	>46
5f	11	23	99.3	35.8	37.0	4.55	a	>25
7a	2	5	1225	359	506	56.5	a	>2
7b	3	7	a	-	a	-	a	-
7c	5	11	349	36.5	230	4.50	a	>7
7d	7	15	46.7	1.85	19.5	1.35	622	13.3
7e	9	19	14.0	2.10	5.44	0.62	419	29.9
7f	11	23	4.56	0.83	2.30	0.20	305	66.9

^a Ki > highest standard of 2500 nM

Table 2

Binding affinities of N-Me bivalent ligands **6a-d** and monovalent ligands **8a-d** against [³H-CP55,940] and [³H-1]

Compd	n	Linker (atoms)	Displacement Assay vs. Tritiated Ligands: Ki (nM) in hCB1 (n=2)			Displacement vs. ³ H-CP55,940: Ki (nM) in hCB2		
			³ H-CP55,940	SEM	³ H-1	SEM	SEM	CBI/CB2
6a	5	11	38.7	4.0	6.35	1.07	1037	26.8
6b	7	15	17.3	0.45	27.5	1.90	683	39.5
6c	9	19	247	29.0	94.1	25.9	a	>10
6d	11	23	1885	450	1292	524	a	>1.3
8a	5	11	162	22.5	88.8	0.45	a	>15
8b	7	15	37.5	4.45	15.5	0.40	1934	51.6
8c	9	19	10.0	1.27	6.12	1.06	265	26.5
8d	11	23	14.7	5.27	5.51	1.81	426	29.0

^a Ki > highest standard of 2500 nM

Table 3

Functional assessment of the alkyl N-H series of bivalent ligands **5a-f** and monovalent ligands **6a-f** at the CB1 receptor

Cmpd	Linker	GTP- γ -S Assay in Rat Brain		GTP- γ -S Assay in hCB1		GTP- γ -S Assay in hCB1		Calcium Assay	
		EC50 (nM)	E _{max}	EC50 (nM)	E _{max}	P _{A2} in hCB1	+/- 95% confidence limits	K _e (nM)	SEM
1		56305	-37.8	ND	-	8.59	0.08	1.1	0.12
5a	5	b	-35.9	237	-25.5	7.08	0.52	2702	411
5b	7	718	-37.2	84.2	-31.9	7.41	0.27	1304	279
5c	11	b	-29.6	33.0	8.5	7.53	0.28	476	69
5d	15	1193	-25.0	179	-38.8	8.08	0.24	567	64
5e	19	1.34	10.4	27.9	6.5	7.76	0.60	4165	1142
5f	23	b	-12.2	a	-	7.56	0.35	d	-
7a	5	7243	-22.9	ND	-	c	c	d	-
7b	7	b	-34.8	ND	-	c	c	d	-
7c	11	2222	-40.7	ND	-	6.12	0.32	5399	1105
7d	15	3279	-52.8	ND	-	7.69	0.46	502.6	225
7e	19	2393	-65.2	ND	-	8.25	0.28	31.6	3.1
7f	23	3616	-29.2	ND	-	8.50	0.26	146.5	19.1

^aEC50 > highest standard of 25,000 nM;

^bEC50 > highest standard of 10,000 nM;

^cDoes not converge, unable to calculate value;

^dno shift observed at 10,000 nM;

ND: not done.

Table 4

Functional assessment of the N-Me series of bivalent ligands **6a-d** and monovalent ligands **8a-d** at the CB1 receptor

Cmpd	Linker	GTP- γ -S Assay in Rat Brain		GTP- γ -S Assay in hCB1		GTP- γ -S Assay in hCB1		Calcium Assay	
		EC50 (nM)	E _{max}	EC50 (nM)	E _{max}	PA ₂ in hCB1	+/- 95% confidence limits	K _e (nM)	SEM
6a	11	1.78	-3.8	59.8	-13.1	7.96	0.22	107.7	20.8
6b	15	1460	-40.8	2238	-32.7	8.37	0.24	478.3	14.3
6c	19	b	-	161	-23.0	7.53	0.25	c	-
6d	23	29.3	17.3	a	-	6.30	0.32	c	-
8a	11	3037	-53.3	ND	-	6.51	0.77	2873	418
8b	15	4367	-62.0	ND	-	7.66	0.55	219.8	2.8
8c	19	3404	-67.9	ND	-	8.34	0.25	140.9	29.1
8d	23	8768	-49.3	ND	-	7.63	0.65	205.5	2.9

^a EC₅₀ > highest standard of 25,000 nM;

^b EC₅₀ > highest standard of 10,000 nM;

^c no shift observed at 10,000 nM