



Published in final edited form as:

*Biol Psychiatry*. 2009 October 15; 66(8): 795–800. doi:10.1016/j.biopsych.2009.05.018.

## Alcohol Consumption Indices of Genetic Risk for Alcohol Dependence

**Julia D. Grant, Arpana Agrawal, Kathleen K. Bucholz, Pamela A.F. Madden, Michele L. Pergadia, Elliot C. Nelson, Michael T. Lynskey, Richard D. Todd, Alexandre A. Todorov, Narelle K. Hansell, John B. Whitfield, Nicholas G. Martin, and Andrew C. Heath**

Midwest Alcoholism Research Center, Department of Psychiatry, Washington University School of Medicine, St. Louis, Missouri, USA (JDG, AA, KKB, PAFM, MLP, ECN, MTL, RDT [deceased], AAT, ACH); and the Queensland Institute of Medical Research, Brisbane, Australia (NKH, JBW, NGM)

### Abstract

**Background**—Previous research has reported a significant genetic correlation between heaviness of alcohol consumption and alcohol dependence (AD), but this association might be driven by the influence of AD on consumption rather than the reverse. We test the genetic overlap between AD symptoms and a heaviness of consumption measure among individuals who do not have AD. A high genetic correlation between these measures would suggest that a continuous measure of consumption may have a useful role in the discovery of genes contributing to dependence risk.

**Methods**—Factor analysis of 5 alcohol use measures was used to create a measure of heaviness of alcohol consumption. Quantitative genetic analyses of interview data from the 1989 Australian Twin Panel (n=6257 individuals;  $M=29.9$  years) assessed the genetic overlap between heaviness of consumption, DSM-IV AD symptoms, DSM-IV AD symptom clustering, and DSM-IV alcohol abuse.

**Results**—Genetic influences accounted for 30–51% of the variance in the alcohol measures and genetic correlations were 0.90 or higher for all measures, with the correlation between consumption and dependence symptoms among non-dependent individuals estimated at 0.97 (95% CI: 0.80–1.00).

**Conclusions**—Heaviness of consumption and AD symptoms have a high degree of genetic overlap even among non-dependent individuals in the general population, implying that genetic influences on dependence risk in the general population are acting to a considerable degree through heaviness of use, and that quantitative measures of consumption will likely have a useful role in the identification of genes contributing to AD.

### Keywords

alcohol dependence; heaviness of consumption; heritability; genetic overlap; twins; gene identification

---

Address reprint requests to Julia D. Grant, Ph.D., Washington University School of Medicine, Dept. of Psychiatry, 660 S. Euclid, Campus Box 8134, St. Louis, MO 63110, USA; grantj@psychiatry.wustl.edu.

### DISCLOSURES

The authors have no financial or other potential conflicts of interest.

## INTRODUCTION

Extensive family, twin and adoption study literatures have documented the strong familial transmission of alcoholism, and its genetic underpinnings (1–4). Large-sample studies with widely varying operationalizations of alcoholism have been remarkably consistent in documenting moderately strong genetic contributions to variation in alcohol dependence (AD), typically accounting for 40–60% of variation in risk. This literature has generated gene-discovery efforts through genetic linkage methods (5–12), with successful follow-up of positive linkage regions (13–15), and through genomewide association studies of alcoholism (16,17). However, with the exception of genetic analyses of individual alcoholism symptoms (18,19), the specific aspects of AD symptomatology that best define its core heritable phenotype(s) has received inadequate attention. Alcohol abuse, for example, is commonly ignored as genetically uninformative, though the evidentiary basis for this assumption is weak (20). Despite widespread reliance on a binary dependence measure, evidence supporting a continuum of alcohol problems has been found repeatedly in both clinically ascertained (21) and general community (22–24) samples. From such analyses have come proposals for a quasi-continuous characterization of AD symptoms for DSM-V (25,26).

Research (27) has documented strong genetic influence on alcohol consumption patterns in general community samples (28), with reports of a high genetic correlation between heaviness of consumption and AD risk (29). Unfortunately, the existing literature cannot exclude the possibility that this association is driven by the influence of AD onset on progression of drinking patterns rather than vice versa, although statistical methods can estimate the genetic correlation between AD risk and consumption with dependence effects on consumption excluded (30). Confirmation of a high genetic correlation between nearly continuous consumption measures and AD risk would suggest that such measures may be useful in the discovery of genes contributing to dependence risk.

In this paper, we use the power of the twin study design to address three questions: (i) the plausibility of a dimensional model for the genetic transmission of AD risk; (ii) the evidence for cotransmission of AD versus alcohol abuse when abuse is defined hierarchically (31); and (iii) the evidence for at least partial genetic cotransmission of heaviness of consumption and AD risk among individuals without AD.

## METHODS AND MATERIALS

### Samples

We make use of data from a young adult Australian twin cohort (3) for genetic analyses, and from spouses (32) of a second older twin sample (32,33) for corroboration of factor analyses and test-retest reliability assessments.

**Australian Young Adult Twin Panel (“1989 Cohort”)**—Twin pairs for this volunteer sample were born 1964–1971, were identified in 1980–1982 through mass media appeals and school systems, were raised together, and included a broad spectrum of sociodemographic groups (34). As described elsewhere (3,34), twins were not assessed as children, but completed a mailed questionnaire in 1989 and a telephone diagnostic interview between 1996 and 2000. Of the 8020 twins (4010 pairs) identified in childhood, 6257 individuals (78.0%) were interviewed as young adults (2761 complete pairs: 698 MZ female, 494 MZ male, 513 DZ female, 395 DZ male, and 661 DZ unlike-sexed pairs; and 735 twin singletons, N=371 female). Mean age at interview was 30.0 years for women (N=3454) and 29.9 years for men (N=2803).

**Australian Spouse Sample**—In 1994–1997, spouses/partners (1430 females,  $M=43.5$  years; 2384 males,  $M=48.5$  years, [32]) of an older twin cohort (“1981 Cohort”, [33]) completed a telephone diagnostic interview. The same assessments used with the “1989 Twin Cohort” were used with this spouse sample, allowing for examination of the consistency of factor loadings in a somewhat older sample. A subset of individuals from the spouse cohort ( $n=665$ ) completed a second interview in 2003–2005 for one of several coordinated studies, allowing for estimation of the long-term stability of the measures. Data from respondents targeted for a general population study of large sibships ( $n=5485$ , including  $n=264$  previously interviewed spouses) were used to estimate factor scoring coefficients which were then applied to an additional  $n=401$  spouses who completed a second interview for one of two non-population-based studies (35).

## Measures

All participants completed a telephone-administered adaptation of the SSAGA, a semi-structured diagnostic interview designed for genetic studies of alcoholism (36). Interviews were completed by lay interviewers who received 2-weeks of basic training in telephone-interviewing and continuing in-service training (33). Interviewers were supervised by a Master’s-level Clinical Psychologist, and interviews were audio-taped for quality-control purposes unless permission to do so was refused. The SSAGA has been shown to have excellent within-center and across-center AD reliability (kappas: 0.84–0.89, [36]) and test-retest reliability (tetrachoric correlation=0.77, [33]). Alcohol use and lifetime DSM-IV alcohol abuse and dependence questions were asked of all respondents who had either used alcohol regularly (defined as having consumed alcohol at least once a month for 6 months or more) or who reported having gotten drunk. Heaviness of consumption was assessed through 2 lifetime indices and 3 indices from the 12-month period of heaviest use. Lifetime indices were maximum drinks consumed in a 24-hour period (log-transformed) and maximum tolerance (maximum drinks consumed before becoming drunk, or maximum drinks consumed before feeling any effect for those who had never been drunk; log-transformed). Indices from the 12-month period of heaviest use were typical weekly consumption (typical number of drinks consumed per drinking occasion multiplied by typical frequency of consuming alcohol; log-transformed), frequency of heavy drinking (5 or more drinks on a single occasion), and frequency of drinking to intoxication. Our primary analysis focused on the total number of lifetime AD symptoms endorsed, with a separate conditional clustering variable indicating whether at least 3 AD symptoms had ever occurred within a 12-month period. Per DSM-IV criteria, alcohol abuse was defined hierarchically (i.e., only among respondents who did not have AD).

## Statistical Analysis

Simple descriptive statistics were used to summarize rates of alcohol use and problems in the 1989 twin cohort. Factor analysis of the 5 alcohol consumption measures, conducted in SAS (37) separately for men and women and for the 1989 twin and 1981 spouse samples, was used to generate a heaviness of consumption factor score for each individual. For joint analyses with categorical outcomes, factor scores were collapsed into a 7-level measure to ease computational burden.

To address the relationships between genetic influences on AD symptomatology, AD symptom clustering (whether 3 or more symptoms occurred within a 12-month period; undefined in those with 0–2 AD symptoms), alcohol abuse (undefined in those with 3 or more AD symptoms), and heaviness of consumption (undefined in those with 3 or more AD symptoms), we used an approach described by Heath *et al.* (30). In a twin analysis containing an unconditional phenotype (e.g. AD symptomatology) and a conditional phenotype (e.g. heaviness of consumption, defined only in unaffected individuals) and using

the assumption of underlying normally distributed risk ('liability') distributions, they showed that although a full bivariate genetic model is underidentified if the unconditional phenotype is binary, this problem no longer applies if the unconditional phenotype includes at least two non-zero categories for which the conditional phenotype is defined (e.g. consumption is defined in those reporting 0, 1 or 2 AD symptoms).

We first fit a standard univariate genetic model to a 5-level AD symptom count measure (0, 1, 2, 3, or 4 or more symptoms) using Mx (38), to confirm the plausibility of the assumption that the liability distribution underlying symptom count has a bivariate normal joint distribution in twin pairs. We then extended this model to the multivariate case by adding (a) the 7-level alcohol consumption factor score (set to missing in individuals with 3 or more AD symptoms), (b) a variable indicating whether the respondent reported clustering of 3 or more AD symptoms in the same 12 month period (defined only in those reporting 3 or more AD symptoms), and (c) DSM-IV alcohol abuse (defined only in those reporting 0–2 AD symptoms), thereby estimating the genetic, shared environmental, and non-shared environmental contributions to, and correlations between, these variables.

## RESULTS

### Sample Characteristics

As shown in Table 1, the young adult Australian twin cohort is characterized by near universal alcohol use and a high prevalence of heavy drinking and of DSM-IV alcohol dependence and abuse.

### Alcohol consumption factor score

For the twin sample, a single factor model adequately accounted for the covariation among the consumption measures, based on Eigenvalues and scree plots. All five items had substantial factor loadings, with loadings of similar magnitude in the young adult twin cohort (0.69–0.92 for women; 0.68–0.93 for men) and the older spouse cohort (factor loadings: 0.59–0.83 for women; 0.57–0.93 for men). In all cases, typical consumption during the period of heaviest use had the highest factor loading and frequency of intoxication during the period of heaviest use the lowest. Eight-year test-retest correlations in the spouse cohort were 0.76 in women and 0.78 in men.

As shown in Table 2, twin factor score varied as a function of genetic risk, with the highest factor scores among AD individuals, intermediate factor scores for those who did not have AD but had a cotwin with AD, and the lowest factor scores among pairs concordant for no AD. The lower factor score for non-AD individuals with an AD MZ cotwin compared to AD individuals suggests either overlapping environmental influence on the factor score and AD, or an impact of AD on heaviness of consumption.

### Genetic analyses

Fitting multiple threshold models to two-way twin pair contingency tables for AD symptom count confirmed that the observed data met the assumption of an underlying normal liability distribution in like-sex male (Chi-square=29.24, d.f.=30, p=0.51) and female (Chi-square=32.61, d.f.=30, p=0.34) twin pairs, and for the 5-group analysis including unlike-sex pairs (Chi-square=75.05, d.f.=75, p=0.48).

Table 3 summarizes variance component estimates for the quadrivariate genetic model, with genetic and environmental correlations between the variables summarized in Table 4. Major genetic conclusions are: (a) moderate heritability of AD symptom count (39%), with a genetic correlation with the temporal clustering variable approaching unity ( $r_G=0.99$ ); (b)

high heritability of heaviness of consumption (50%), with a high genetic correlation with AD symptom count ( $r_G=0.97$ ); (c) high heritability of alcohol abuse (51% in non-AD individuals), with a high genetic correlation with AD symptoms ( $r_G=0.96$ ). Non-shared environmental correlations were intermediate in magnitude and, for conditional phenotypes, had broad confidence intervals. Shared environmental parameters were all non-significant. As shown in Table 5, the genetic correlations between the consumption factor score (set to missing for those with 3 or more AD symptoms) and individual DSM-IV AD symptoms ranged from 0.67 to 0.95, with complete genetic overlap a possibility for 6 of the 7 symptoms.

## DISCUSSION

Using data from a general community twin sample, we have shown that a composite alcohol factor score is highly reliable (see also Agrawal *et al.* [39]), moderately heritable (50%), and has a high genetic correlation with AD symptomatology ( $r_G=0.97$ ), indicating that genetic influences on dependence risk and consumption overlap considerably in the general population. We previously noted a smaller genetic association with confidence intervals that excluded unity ( $r_G=0.63$ , CI: 0.53–0.72 [29]). However, the previous analyses assessed consumption at the time of interview and included consumption assessments for AD individuals.

The excellent reliability and substantial heritability of our quantitative consumption factor score and its high genetic correlation with AD suggest that the factor score will be useful in the identification of genes contributing to AD. While it is possible that binary diagnostic AD measures may play a role in the discovery of genes contributing to heaviness of consumption, continuous measures such as consumption are considerably more powerful predictors than binary ones, and are therefore likely to be especially useful for detecting the numerous small effects that contribute to AD risk. Prior studies using diagnostic assessments of AD have been limited by loss of power (e.g., through use of affecteds-only) or by heterogeneity among the unaffected individuals. Use of a quantitative measure circumvents these challenges by allowing for assessment of genomic effects across the range of liability to dependence. The factor score also represents an improvement over AD symptom count measures, which are highly skewed. Our study also suggests that quantitative indices of alcohol consumption may allow investigators to indirectly examine genetic and genomic effects on AD vulnerability *even in the absence of full diagnostic data*, thereby greatly augmenting sample sizes and even further increasing power to detect modest allelic effects (40,41). It is hoped that future gene-discovery efforts using quantitative measures will be better able to identify the effects of polymorphisms that act across a range of vulnerability to alcohol use disorder, as well as their interplay with environmental influences.

Genes that influence alcohol metabolism undoubtedly play some role in contributing to differences in alcohol consumption in this European ancestry sample. It is well known that the ALDH2 locus, which is polymorphic in individuals of Asian ancestry but not those of European ancestry, is associated with differences in both AD and alcohol consumption (42,43). Although analysis of SNPs across the ADH gene have confirmed significant effects of ADH gene variants in non-Jewish individuals of European ancestry, both on metabolism (44) and on multiple indices of heaviness of consumption (45), the variance in consumption level accounted for was small.

Our findings also have implications for the use of DSM-based alcohol measures. Despite the concerns surrounding alcohol abuse (46), abuse was substantially heritable (51%) and had a genetic correlation with dependence that did not differ significantly from unity. Thus, abuse may provide information on individuals at genetic risk of AD who would be missed by using

dependence exclusively. We found less support for a focus on symptom temporal clustering: it was endorsed by 88% of women and 90% of men with 3 or more symptoms, had only modest heritability (30%), and had a genetic correlation of unity with AD symptom count, suggesting it added little information beyond that provided by dependence symptoms.

Some may question the inclusion of tolerance in our factor score creation. However, our absolute tolerance measure is quite distinct from the DSM tolerance criterion which is relative (i.e., tolerance is a change in consumption before feeling an effect). Additional analyses indicated a correlation of 0.98 between our current 5-item factor score and a reduced 4-item factor, suggesting that the two factor scores are highly comparable.

Our conclusions must be tempered by the recognition that our sample was relatively young and almost entirely of European ancestry. In addition, some may view the inclusion of large numbers of only moderate AD cases as a limitation (47). We have argued elsewhere (48), however, that moderate dependence is associated with many substantively important outcomes, including reproductive delay (49), and marital breakdown, family conflict, and adverse environmental exposures including physical and sexual abuse (48). Additionally, moderate dependence is itself an important transition in the progression to severe dependence. What we cannot exclude from our analyses, because of relatively small numbers of severely dependent cases, is the likelihood that additional genetic factors contribute to severe dependence risk above and beyond those associated with heaviness of drinking. In the long term, only the progressive and parallel identification of genes contributing to risk of dependence in community ascertained versus severe clinic samples will clarify this question.

## Acknowledgments

We thank the twins and their spouses for their participation and cooperation, the project coordinators and interviewers under the supervision of Dixie J. Statham, and the data managers under the supervision of John Pearson and David Smyth. This work was supported by grants to Drs. Andrew C. Heath (AA010249, AA007728, AA011998, AA013321), Nicholas G. Martin (AA013326), Kathleen K. Bucholz (DA014363, AA012640), Pamela A.F. Madden (DA012854), Richard D. Todd (AA013320), Arpana Agrawal (ABMRF/Foundation for Alcohol Research, DA023668), Elliot C. Nelson (AA013446), Michele L. Pergadia (DA019551), and Michael T. Lynskey (DA018660).

## References

1. Heath, AC.; Slutske, WS.; Madden, PAF. Gender differences in the genetic contribution to alcoholism risk and to alcohol consumption patterns. In: Wilsnack, RW.; Wilsnack, SC., editors. *Gender and Alcohol: Individual and Social Perspectives*. New Brunswick, NJ: Rutgers Center of Alcohol Studies; 1997. p. 114-149.
2. Kendler, KS.; Prescott, CA. *Genes, Environment and Psychopathology: Understanding the Causes of Psychiatric and Substance Use Disorders*. New York: Guilford Press; 2006.
3. Knopik VS, Heath AC, Madden PAF, Bucholz KK, Slutske WS, Nelson EC, et al. Genetic effects on alcohol dependence risk: Re-evaluating the importance of psychiatric and other heritable risk factors. *Psychol Med*. 2004; 34:1519–1530. [PubMed: 15724882]
4. Nurnberger JI Jr, Wiegand R, Bucholz K, O'Connor S, Meyer ET, Reich T, et al. A family study of alcohol dependence: Coaggregation of multiple disorders in relatives of alcohol-dependent probands. *Arch Gen Psychiatry*. 2004; 61:1246–1256. [PubMed: 15583116]
5. Foroud T, Bucholz KK, Edenberg HJ, Goate A, Newman RJ, Porjesz B, et al. Linkage of an alcoholism-related severity phenotype to chromosome 16. *Alcohol Clin Exp Res*. 1998; 22:2035–2042. [PubMed: 9884148]
6. Foroud T, Edenberg HJ, Goate A, Rice J, Flury L, Koller DL, et al. Alcoholism susceptibility loci: Confirmation studies in a replicate sample and further mapping. *Alcohol Clin Exp Res*. 2000; 24:933–945. [PubMed: 10923994]



7. Hansell, NK.; Agrawal, A.; Whitfield, JB.; Morley, KI.; Gordon, SD.; Lind, PA., et al. Can we identify genes for heaviness of alcohol use in samples ascertained for heterogeneous purposes?. (in review)
8. Hansell, NK.; Agrawal, A.; Whitfield, JB.; Morley, KI.; Gordon, SD.; Lind, PA., et al. Linkage analysis of alcohol dependence symptom score in a community sample: Effects of exclusion criteria. (in review)
9. Hill SY, Shen S, Zezza N, Hoffman EK, Perlin M, Allan W. A genome wide search for alcoholism susceptibility genes. *Am J Med Genet B Neuropsychiatr Genet.* 2004; 128B:102–113. [PubMed: 15211641]
10. Kendler KS, Kuo PH, Todd Webb B, Kalsi G, Neale MC, Sullivan PF, et al. A joint genomewide linkage analysis of symptoms of alcohol dependence and conduct disorder. *Alcohol Clin Exp Res.* 2006; 30:1972–1977. [PubMed: 17117961]
11. Prescott CA, Sullivan PF, Kuo PH, Webb BT, Vittum J, Patterson DG, et al. Genomewide linkage study in the Irish affected sib pair study of alcohol dependence: Evidence for a susceptibility region for symptoms of alcohol dependence on chromosome 4. *Mol Psychiatry.* 2006; 11:603–611. [PubMed: 16534506]
12. Reich T, Edenberg HJ, Goate A, Williams JT, Rice JP, van Eerdewegh P, et al. Genome-wide search for genes affecting risk for alcohol dependence. *Am J Med Genet B: Neuropsychiatr Genet.* 1998; 81:207–215.
13. Edenberg HJ, Dick DM, Xuei X, Tian H, Almasy L, Bauer LO, et al. Variations in GABRA2, encoding the alpha 2 subunit of the GABA(A) receptor, are associated with alcohol dependence and with brain oscillations. *Am J Hum Genet.* 2004; 74:705–714. [PubMed: 15024690]
14. Edenberg JH, Foroud T. The genetics of alcoholism: Identifying specific genes through family studies. *Addiction Biol.* 2006a; 11:386–396.
15. Edenberg HJ, Xuei X, Chen HJ, Tian H, Wetherill LF, Dick DM, et al. Association of alcohol dehydrogenase genes with alcohol dependence: A comprehensive analysis. *Hum Mol Genet.* 2006b; 15:1539–1549. [PubMed: 16571603]
16. Johnson C, Drgon T, Liu Q-R, Walther D, Edenberg H, Rice J, et al. Pooled association genome scanning for alcohol dependence using 104,268 SNPs: Validation and use to identify alcoholism vulnerability loci in unrelated individuals from the Collaborative Study on the Genetics of Alcoholism. *Am J Med Genet B Neuropsychiatr Genet.* 2006; 141:844–853. [PubMed: 16894614]
17. Uhl GR, Drgon T, Johnson C, Fatusin OO, Liu Q-R, Contoreggi C, et al. Higher order” addiction molecular genetics: Convergent data from genome-wide association in humans and mice. *Biochem Pharmacol.* 2008; 75:98–111. [PubMed: 17764662]
18. Johnson EO, van den Bree MBM, Pickens RW. Indicators of genetic and environmental influence in alcohol-dependent individuals. *Alcohol Clin Exp Res.* 1996; 20:67–74. [PubMed: 8651465]
19. Slutske WS, True WR, Scherrer JF, Heath AC, Bucholz KK, Eisen SA, et al. The heritability of alcoholism symptoms: “Indicators of genetic and environmental influence in alcohol-dependent individuals” revisited. *Alcohol Clin Exp Res.* 1999; 23:759–769. [PubMed: 10371393]
20. Prescott CA, Kendler KS. Genetic and environmental contributions to alcohol abuse and dependence in a population-based sample of male twins. *Am J Psychiatry.* 1999; 156:34–40. [PubMed: 9892295]
21. Bucholz KK, Heath AC, Reich T, Hesselbrock VM, Kramer JR, Nurnberger JI Jr, et al. Can we subtype alcoholism? A latent class analysis of data from relatives of alcoholics in a multicenter family study of alcoholism. *Alcohol Clin Exp Res.* 1996; 20:1462–1471. [PubMed: 8947326]
22. Heath AC, Bucholz KK, Slutske WS, Madden PAF, Dinwiddie SH, Dunne MP, et al. The assessment of alcoholism in surveys of the general community: what are we measuring? Some insights from the Australian twin panel survey. *Int Rev Psychiatry.* 1994; 6:295–307.
23. Lynskey MT, Nelson EC, Neuman RJ, Bucholz KK, Madden PAF, Knopik VS, et al. Limitations of DSM-IV operationalizations of alcohol abuse and dependence in a sample of Australian twins. *Twin Res Hum Genet.* 2005; 8:574–584. [PubMed: 16354499]
24. Saha TD, Chou SP, Grant BF. Toward an alcohol use disorder continuum using item response theory: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychol Med.* 2006; 36:931–941. [PubMed: 16563205]

25. Helzer JE, Kraemer HC, Krueger RF. The feasibility and need for dimensional psychiatric diagnoses. *Psychol Med.* 2006a; 36:1671–1680. [PubMed: 16907995]
26. Helzer JE, van den Brink W, Guth SE. Should there be both categorical and dimensional criteria for the substance use disorders in DSM-V? *Addiction.* 2006b; 101 (Suppl 1):17–22. [PubMed: 16930157]
27. Partanen, J.; Bruun, K.; Markkanen, T. *Inheritance of Drinking Behavior.* Helsinki: Finnish Foundation for Alcohol Studies; 1966.
28. Heath, AC. Genetic influences on drinking behavior in humans. In: Begleiter, H.; Kissin, B., editors. *Alcohol and Alcoholism, Vol. 1, The Genetics of Alcoholism.* New York: Oxford University Press, Inc; 1995. p. 82-121.
29. Whitfield JB, Zhu G, Madden PAF, Neale MC, Heath AC, Martin NG. The genetics of alcohol intake and of alcohol dependence. *Alcohol Clin Exp Res.* 2004; 28:1153–1160. [PubMed: 15318113]
30. Heath AC, Martin NG, Lynskey MT, Todorov AA, Madden PAF. Estimating two-stage models for genetic influences on alcohol, tobacco, or drug use initiation and dependence vulnerability in twin and family data. *Twin Res.* 2002; 5:113–124. [PubMed: 11931689]
31. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* 4. Washington, DC: American Psychiatric Association; 1994.
32. Grant JD, Heath AC, Bucholz KK, Madden PAF, Agrawal A, Statham DJ, et al. Spousal concordance for alcohol dependence: Evidence for assortative mating or spousal interaction effects? *Alcohol Clin Exp Res.* 2007; 31:717–728. [PubMed: 17378921]
33. Heath AC, Bucholz KK, Madden PAF, Dinwiddie SH, Slutske WS, Bierut LJ, et al. Genetic and environmental contributions to alcohol dependence risk in a national twin sample: Consistency of findings in men and women. *Psychol Med.* 1997; 27:1381–1396. [PubMed: 9403910]
34. Hansell NK, Agrawal A, Whitfield JB, Morley KI, Zhu G, Lind PA, et al. Long-term stability and heritability of alcohol consumption and dependence. *Twin Res Hum Genet.* 2008; 11:287–304. [PubMed: 18498207]
35. Saccone SF, Pergadia ML, Loukola A, Broms U, Montgomery GW, Wang JC, et al. Genetic linkage to chromosome 22q12 for a heavy-smoking quantitative trait in two independent samples. *Am J Hum Genet.* 2007; 80:856–866. [PubMed: 17436240]
36. Bucholz KK, Cadoret R, Cloninger CR, Dinwiddie SH, Hesselbrock VM, Nurnberger JI Jr, et al. A new, semi-structured psychiatric interview for use in genetic linkage studies: A report on the reliability of the SSAGA. *J Stud Alcohol.* 1994; 55:149–158. [PubMed: 8189735]
37. SAS Institute. *SAS/STAT User's Guide, Version 8.* Cary, NC: SAS Institute Inc; 2000.
38. Neale, MC.; Boker, SM.; Xie, G.; Maes, HH. *Mx: Statistical Modeling.* 6. VCU Box 900126, Richmond, VA 23298: Department of Psychiatry; 2003.
39. Agrawal A, Grant JD, Littlefield AK, Waldron M, Pergadia ML, Lynskey MT, et al. Developing a quantitative measure of alcohol consumption for genomic studies on prospective cohorts. *J Stud Alcohol Drugs.* 2009; 70:157–168. [PubMed: 19261227]
40. Houlston RS, Webb E, Broderick P, Pittman AM, DiBernardo MC, et al. COGENT Study. Meta-analysis of genome-wide association data identifies four new susceptibility loci for colorectal cancer. *Nat Genet.* 2008; 40:1426–1435. [PubMed: 19011631]
41. Cooper JD, Smyth DJ, Smiles AM, Plagnol V, Walker NM, Allen JE, et al. Meta-analysis of genome-wide association study data identifies additional type 1 diabetes risk loci. *Nat Genet.* 2008; 40:1399–1401. [PubMed: 18978792]
42. Heath, AC.; Madden, PAF.; Bucholz, KK.; Nelson, EC.; Todorov, A.; Price, RK., et al. Genetic and environmental risks of dependence on alcohol, tobacco, and other drugs. In: Plomin, R.; DeFries, JC.; Craig, IW.; McGuffin, P., editors. *Behavioral Genetics in the Postgenomic Era.* Washington, DC: American Psychological Association; 2003.
43. Higuchi S, Matsushita S, Muramatsu T, Murayama M, Hayashida M. Alcohol and aldehyde dehydrogenase genotypes and drinking behavior in Japanese. *Alcohol Clin Exp Res.* 1996; 20:493–497. [PubMed: 8727242]



44. Birley AJ, James MR, Dickson PA, Montgomery GW, Heath AC, Martin NG, et al. ADH single nucleotide polymorphism associations with alcohol metabolism *in vivo*. *Hum Mol Genet.* 2009; 18:1533–1542. [PubMed: 19193628]
45. Macgregor S, Lind PA, Bucholz KK, Hansell NK, Madden PAF, Richter MM, et al. Associations of ADH and ALDH2 gene variation with self report alcohol reactions, consumption and dependence: an integrated analysis. *Hum Mol Genet.* 2009; 18:580–593. [PubMed: 18996923]
46. Hasin D, Hatzenbuehler ML, Keyes K, Ogburn E. Substance use disorders: Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) and International Classification of Diseases, tenth edition (ICD-10). *Addiction.* 2006; 101 (Suppl 1):59–75. [PubMed: 16930162]
47. Babor TF, Caetano R. Subtypes of substance dependence and abuse: Implications for diagnostic classification and empirical research. *Addiction.* 2006; 101 (Suppl 1):104–110. [PubMed: 16930166]
48. Heath, AC.; Lynskey, MT.; Waldron, M. Child and adolescent substance use and substance use disorders. In: Rutter, M.; Taylor, EA., editors. *Rutter's Child and Adolescent Psychiatry.* 5. Oxford, UK: Blackwell; 2008.
49. Waldron M, Heath AC, Bucholz KK, Madden PAF, Martin NG. Alcohol dependence and reproductive onset: Findings in two Australian twin cohorts. *Alcohol Clin Exp Res.* 2008; 32:1–10. [PubMed: 17949471]

**Table 1**

Frequency distribution of alcohol consumption items and abuse/dependence symptoms in the Australian Young Adult Twin Panel ("1989 Cohort")

	Women (%)	Men (%)
Full Sample (n=3454 women, n=2803 men):		
Lifetime Abstainers	0.9	1.2
Of non-abstainers (n=3422 women, n=2769 men):		
Ever regular drinkers <sup>a</sup>	87.3	94.1
Ever been intoxicated	88.0	96.1
Of reg. drinkers/ever intox. (n=3207 women, n=2701 men):		
Maximum drinks consumed in 24 hours (lifetime)		
1-4	10.1	1.7
5-10	47.0	11.0
11-15	22.3	17.9
16-20	8.9	16.1
21-25	5.8	16.1
26-30	2.4	13.0
31+	3.5	24.2
Maximum tolerance (lifetime) <sup>b</sup>		
≤ 2	9.6	2.7
3-4	24.4	8.3
5-6	31.0	19.9
7-8	15.5	19.8
9-10	10.8	17.0
11-14	5.4	18.2
15+	3.2	14.1
Typical # of drinks per occasion (heaviest period)		
1-2	30.6	16.7
3-4	31.6	27.1
5-6	20.9	22.2
7-8	8.9	14.1
9-11	5.2	8.9
12+	2.8	11.0
Frequency consumed alcohol (heaviest period)		
One day per month or less often	16.6	7.9
2-3 days per month	12.0	6.6
1 day per week	20.7	12.2
2 days per week	22.9	22.2
3-4 days per week	17.4	28.1
5-7 days per week	10.4	23.0

	Women (%)	Men (%)
Frequency had 5+ drinks in a single occasion (heaviest period)		
Never	18.8	5.0
At least once, but less than 1 time per month	14.7	8.5
1–3 times per month	18.2	12.6
1 time per week	18.9	16.8
2 times per week	16.8	23.8
3–7 times per week	12.6	33.3
Frequency drank to intoxication (heaviest period)		
Never	13.2	5.5
At least once, but less than 1 time per month	35.0	21.2
1–3 times per month	22.1	22.0
1 time per week	15.2	19.9
2 times per week	9.5	18.8
3–7 times per week	5.0	12.5
Alcohol dependence symptoms		
0	30.5	15.0
1	28.0	23.8
2	22.4	25.4
3	9.8	15.9
4+	9.3	19.9
Alcohol dependence clustering (of those with 3 or more Sx)	88.0	89.8
Alcohol abuse (of those with 0–2 dependence Sx)	7.1	18.3

<sup>a</sup> consumed alcohol at least once a month for 6 or more months;

<sup>b</sup> maximum drinks consumed before becoming drunk/feeling any effect

**Table 2**

Alcohol factor score as a function of respondent and cotwin dependence history, by zygosity and gender in the Australian Young Adult Twin Panel (“1989 Cohort”). Categories for non-dependent respondents are ordered by predicted genetic risk (see Methods)

	# of respondents	Mean Factor Score (s.d.)
Female Twins (n=3050)		
Respondent AD	530	1.00 (0.84)
Respondent not AD, MZ cotwin AD	118	0.32 (0.86)
Respondent not AD, DZ female cotwin AD	111	0.17 (0.75)
Respondent not AD, DZ male cotwin AD	151	0.08 (0.72)
Same-sex pair concordant no AD	1759	-0.29 (0.83)
Unlike sex pair concordant no AD	381	-0.26 (0.82)
Male Twins (n=2493)		
Respondent AD	852	0.65 (0.70)
Respondent not AD, MZ cotwin AD	125	0.08 (0.80))
Respondent not AD, DZ female cotwin AD	60	-0.11 (0.87)
Respondent not AD, DZ male cotwin AD	144	-0.04 (0.75)
Unlike sex pair concordant no AD	383	-0.35 (0.98)
Same-sex pair concordant no AD	929	-0.41 (0.90)

**Table 3**

Proportions of variance and 95% confidence intervals for heaviness of alcohol use and alcohol abuse/dependence variables in the Australian Young Adult Twin Panel (“1989 Cohort”)

	<b>Genetic Variance</b>	<b>Shared Environmental Influence</b>	<b>Nonshared Environmental Influence</b>
AD symptom count	0.39* (0.27–0.49)	0.07 (0.00–0.16)	0.54* (0.49–0.54)
Alcohol consumption factor score <sup>a</sup>	0.50* (0.36–0.63)	0.10 (0.00–0.22)	0.40* (0.35–0.44)
AD symptom clustering <sup>b</sup>	0.30* (0.13–0.55)	0.16 (0.00–0.35)	0.55* (0.41–0.66)
Alcohol abuse diagnosis <sup>a</sup>	0.51* (0.25–0.53)	0.08 (0.00–0.29)	0.41* (0.31–0.52)

\* Indicates  $p < .05$

<sup>a</sup> undefined in those with 3 or more alcohol dependence symptoms

<sup>b</sup> undefined in those with 0–2 alcohol dependence symptoms



**Table 4**

Point estimates and 95% confidence intervals for genetic correlations (above diagonal) and nonshared environmental correlations (below diagonal) of heaviness of alcohol use and alcohol abuse/dependence variables in the Australian Young Adult Twin Panel (“1989 Cohort”)

	<b>AD symptom count</b>	<b>Consumption</b>	<b>AD symptom clustering</b>	<b>Alcohol Abuse Diagnosis</b>
AD symptom count	—————	0.97* (0.91–1.00)	0.99* (0.80–1.00)	0.96* (0.73–0.99)
Consumption <sup>a</sup>	0.53* (0.45–0.59)	—————	0.99* (0.78–1.00)	0.90* (0.74–1.00)
AD symptom clustering <sup>b</sup>	0.79* (0.69–0.90)	0.60 (–0.26–0.99)	—————	0.95* (0.56–1.00)
Alcohol Abuse Diagnosis <sup>a</sup>	0.44* (0.32–0.46)	0.39* (0.30–0.51)	0.68 (–0.42–0.98)	—————

\* Indicates  $p < .05$

<sup>a</sup> undefined in those with 3 or more alcohol dependence symptoms

<sup>b</sup> undefined in those with 0–2 alcohol dependence symptoms

**Table 5**

Genetic correlations between the alcohol consumption factor score and individual DSM-IV alcohol dependence symptoms (factor score set to missing among for those with 3 or more AD symptoms)

<b>DSM-IV Dependence Symptom</b>	<b><math>r_G</math> (95% CI)</b>
Withdrawal	0.92 (0.52–1.00)
Tolerance	0.95 (0.77–1.00)
Used More Than Intended	0.89 (0.82–1.00)
Unable to Quit/Persistent Desire to Quit	0.67 (0.56–0.89)
Spent Much Time Getting/Using	0.83 (0.67–1.00)
Reduced Activities	0.73 (0.36–1.00)
Continued Use Despite Physical/Emotional/Health Problems	0.81 (0.62–1.00)