



Published in final edited form as:

*Oper Dent.* 2011 March ; 36(1): 2–11. doi:10.2341/10-137-CR.

## Methods used by Dental Practice-Based Research Network (DPBRN) dentists to diagnose dental caries

**Dr. Valeria V. Gordan, DDS, MS, MSCI\* [Professor],**

Department of Operative Dentistry, College of Dentistry, University of Florida, Gainesville, Florida, USA.

**Dr. Joseph L Riley III, PhD [Associate Professor, ],**

Department of Community Dentistry and Behavioral Sciences, College of Dentistry, University of Florida, Gainesville, Florida, USA.

**Dr. Ricardo M. Carvalho, DDS, MS, PhD [Professor],**

Department of Prosthodontics, Faculdade de Odontologia de Bauru, Bauru, SP, Brazil

**Dr. John Snyder, DMD [private practitioner],**

Permanente Dental Associates, Portland, Oregon, USA.

**Dr. James L Sanderson Junior, DMD [private practitioner],**

Hoover, Alabama, USA.

**Dr. Mary Anderson, DDS [private practitioner], and**

Minneapolis, Minnesota, USA.

**Dr. Gregg H. Gilbert, DDS, MBA, FAAHD, FACD [Professor and Chair]**

Department of General Dental Sciences, School of Dentistry, University of Alabama at Birmingham, Birmingham, Alabama, USA.

**For The DPBRN Collaborative Group**

### Abstract

**Objectives**—To (1) identify the methods that dentists in The Dental Practice-Based Research Network (DPBRN) use to diagnose dental caries; (2) quantify their frequency of use; and (3) test the hypothesis that certain dentist and dental practice characteristics are significantly associated with their use.

**Methods**—A questionnaire about methods used for caries diagnosis was sent to DPBRN dentists who reported doing at least some restorative dentistry; 522 dentists participated. Questions included use of dental radiographs, dental explorer, laser fluorescence, air-drying, fiber optic devices, and magnification, as used when diagnosing primary, secondary/recurrent, or non-specific caries lesions. Variations on the frequency of their use were tested using multivariate analysis and Bonferroni tests.

**Results**—Overall, the dental explorer was the instrument most commonly used to detect primary occlusal caries as well as to detect caries at the margins of existing restorations. In contrast, laser fluorescence was rarely used to help diagnose occlusal primary caries. For proximal caries, radiographs were used to help diagnose 75-100% of lesions by 96% of the DPBRN dentists.

\* **Corresponding author:** University of Florida, College of Dentistry P.O. Box 100415, Gainesville, FL 32610-04415 Phone: 352 273 5846 Fax: 352 846 1643 [vgordan@dental.ufl.edu](mailto:vgordan@dental.ufl.edu).

The DPBRN Collaborative Group comprises practitioner-investigators, faculty investigators, and staff investigators who contributed to this DPBRN activity. A list of these persons is at <http://www.dpbrn.org/users/publications/Default.aspx>

Dentists who use radiographs most often to assess proximal surfaces of posterior teeth, were significantly more likely to also report providing a higher percentage of patients with individualized caries prevention ( $p = .040$ ) and seeing a higher percentage of pediatric patients ( $p = .001$ ).

**Conclusion**—Use of specific diagnostic methods varied substantially. The dental explorer and radiographs are still the most commonly used diagnostic methods.

---

## INTRODUCTION

Research has shown that significant inconsistencies exist in the diagnosis of caries.<sup>1, 2</sup> These inconsistencies are partly related to the marked variation among dentists when diagnosing dental caries, as well as the lack of sensitive and specific methods to detect and quantify carious lesions.<sup>3</sup> Previous reports indicate sensitivity values ranging from 0.77 to 1.00, and specificity values ranging from 0.45 to 0.93.<sup>4</sup>

Inaccurate diagnosis will result in either overtreatment, with placement of unnecessary restorations, or under-treatment, which will potentially miss the benefit of non-invasive types of treatment at an earlier stage of caries development.<sup>1</sup> Either circumstance will directly or indirectly impact patients' oral health status and short- and long-term cost of care. Identifying the methods being used for caries diagnosis and the factors associated with their use is important to understanding the current status of caries diagnosis in non-academic daily clinical practice. This information should be invaluable to develop consistent criteria for caries diagnosis and to target dentists who could enhance their diagnostic accuracy.

The Dental Practice-Based Research Network (DPBRN) has a valuable mixture of dental practices with diverse treatment philosophies, dental care reimbursement models, and patient populations that presents an advantageous opportunity to investigate variation in how dentists diagnose dental caries. Dentists in DPBRN have much in common with dentists at large<sup>5</sup>, while simultaneously offering substantial diversity in the characteristics of its dentists, their practices, and their patients.<sup>6</sup> DPBRN mainly comprises dentist practitioner-investigators from five regions: AL/MS: Alabama/Mississippi; FL/GA: Florida/Georgia; MN: dentists employed by HealthPartners and private practitioners in Minnesota; PDA: Permanente Dental Associates in cooperation with Kaiser Permanente Center for Health Research in Oregon and Washington; and SK: Denmark, Norway, and Sweden.<sup>7</sup> The purpose of this study was to investigate the methods that DPBRN dentists use to diagnose dental caries. The specific aims were to: (1) identify the methods that dentists in DPBRN use to diagnosis dental caries; (2) quantify their frequency of use; and (3) test the hypothesis that certain dentist and dental practice characteristics are significantly associated with their use.

## MATERIALS AND METHODS

This study queried dentists participating in DPBRN, which comprises outpatient dental practices that have affiliated to investigate research questions and to share experiences and expertise ( $n=1,166$ ). Participants in DPBRN were recruited through mass mailings to licensed dentists from the participating regions. As part of enrollment in DPBRN, all practitioner-investigators completed an Enrollment Questionnaire about their practice characteristics and themselves. An “Assessment of Caries Diagnosis and Caries Treatment” (“Caries Questionnaire”) questionnaire was sent to DPBRN dentists who reported on the Enrollment Questionnaire that they do at least some restorative dentistry ( $n=901$ ). A pilot study documented comprehension and item test-retest reliability across 15 days using a sample of 35 network dentists. All items in the final version met a test-retest reliability cutoff of  $\kappa > 0.7$ . Practitioner-investigators were asked to return the questionnaire

within three weeks. A reminder letter was sent after the third week if the questionnaire had not been returned. After an additional three weeks, a second reminder was sent. Both of these questionnaires are available at <http://www.dpbrn.org/users/publications/Supplement.aspx>.

The Caries Questionnaire contained clinical photographs and had a range of questions, including caries-related diagnostic and clinical decision-making processes, caries risk assessment, and use of prevention techniques. Five hundred thirty-two participating DPBRN dentists returned the questionnaire, which represents an overall return rate of 59%. There were no participation differences by gender, area of specialty, or years since dental school graduation. Ten surveys were completed by dentists whose practices were outside the designated regions and were not included in this study; therefore we reported data from 522 dentists.

## Measures

Table 1 presents the wording for the series of questions about the methods used to diagnose dental caries. Questions specifically investigated the use of radiographs for proximal and occlusal lesions; dental explorer for the detection of primary occlusal caries and secondary/recurrent caries at the margins of existing restorations; air-drying for primary caries on unspecified locations; laser fluorescence for primary caries on occlusal surfaces and fiber optic transillumination for proximal lesions in anterior teeth. Additionally, when responses to the use of air-drying were positive, respondents were prompted to report on the length of time used for the air-drying procedure. The response choices were 0 percent, 1-24 percent, 25-49 percent, 50-74 percent, 75-99 percent, 100 percent.

## Statistical analyses

The percentages for each caries diagnostic method were coded to the categories' median to maintain the interval nature of the data so that parametric statistics could be used: 0%=0%, 1-24%=12.5%, 25-49%=37%, 50-74%=62%, 75-99%=87%, 100%=100%. Descriptive statistics for each of the diagnostic methods were calculated.

The general linear model was used to test for differences across DPBRN region on the use of each diagnostic method. Other practice characteristics tested as independent variables in each model were type of practice, dentist gender, percentage of pediatric patients, years since dental school graduation, whether caries risk was assessed, percentage of patients who received individualized caries prevention, and percentage of patient contact time spent performing non-implant restorative procedures. "Type of practice" was categorized for each dentist as being in either: (1) a solo or small group private practice (SGP); (2) a large group practice (LGP); or (3) a public health practice (PHP). "Small" practices were defined as those that had 3 or fewer dentists. Public health practices were defined as those that receive the majority of their funding from public sources.

Other dichotomous variables were coded as follows: dentist gender, male=0 and female = 1; performing caries risk assessment =0 and not performing caries risk assessment = 1. Multivariate pair-wise comparisons using a Bonferroni correction were used to interpret region differences. Observations for some dentists may lack independence because some network dentists practice within the same office and this may influence their responses or practice patterns. However, data that could account for this possibility were not available for the statistical analysis

## RESULTS

Table 2 shows the distribution of practice characteristics for the participating dental offices. Table 3 summarizes the frequency of use of the diagnostic methods. Table 4 shows the parameter estimates for practice characteristics associated with the frequency of use of each diagnostic method ( $p < 0.1$ ).

### Practice Characteristics and Use of Diagnostic Tools

#### Radiographs

**Proximal surface of a posterior tooth:** Dentists who provide a higher percentage of patients with individualized caries prevention ( $p = 0.040$ ) and who see a higher percentage of pediatric patients ( $p = 0.001$ ) would use radiographs to assess proximal surfaces of posterior teeth on a greater percentage of patients. There was also a trend for a positive association between number of years since dental school graduation and greater use of radiographs to assess proximal surfaces of posterior teeth ( $p = 0.068$ ) (Table 4).

Dentists in the AL/MS (93%), FL/GA (95%), MN (88%), and PDA (95%) regions would use radiographs to assess proximal surfaces of posterior teeth on a significantly greater percentage of patients than dentists in the SK region (76%, all at  $p < 0.001$ ). Dentists in the FL/GA ( $p = 0.050$ ) and PDA ( $p = 0.028$ ) regions also would use radiographs to assess proximal surfaces of posterior teeth more often than dentists from the MN region would.

**Occlusal surface:** None of the practice variables were significant predictors of using radiographs to assess occlusal caries (Table 4). However, regional differences were found in which dentists in the FL/GA (63%,  $p = 0.033$ ) and PDA (69%,  $p = 0.013$ ) regions would use radiographs to assess proximal surfaces of posterior teeth on a significantly greater percentage of their patients than dentists in the SK region would (45%).

#### Use of explorers

**Occlusal surface:** Dentists in a SGP practice model (90% vs. 80%,  $p = 0.018$ ) and those who perform caries risk assessment (94% vs. 87%,  $p = 0.04$ ) use the explorer on occlusal lesions on a greater percentage of patients than dentists using a non-SGP practice model or do not use caries risk assessment.

Dentists in the AL/MS (93%), FL/GA (89%), and PDA (91%) regions would use the explorer on occlusal surfaces significantly more often on patients than dentists in the SK region (69%),  $p < 0.001$ . Dentists in the AL/MS ( $p < 0.001$ ), FL/GA ( $p = 0.015$ ), and PDA ( $p < 0.001$ ) regions would use the explorer on occlusal surfaces significantly more often than dentists in the MN region (88%).

**Margin of an existing restoration:** There was a trend for dentists in a SGP practice model (92% vs. 87%) and those who do not perform caries risk assessment (94% vs. 87%) to use the explorer on the margin of an existing restoration on a greater percentage of their patients than their respective comparison groups (both at  $p = 0.07$ ). Dentists in the AL/MS (93%,  $p = 0.031$ ) region would use the explorer on the margin of an existing restoration significantly more often than dentists from the SK region (84%).

**Other methods (Table 4)**—Laser fluorescence on occlusal surfaces was used most often by dentists who spend the highest percentage of their patient care time doing restorative procedures ( $p < 0.001$ ). There was a trend for dentists with the least number of years since dental school graduation ( $p = 0.088$ ) to use laser fluorescence more often than dentists with

longer tenure. Dentists in the FL/GA (13%,  $p=0.025$ ) region use laser fluorescence significantly more often than dentists in the AL/MS region (6%).

There was a positive association between the percentage of patients who receive individualized caries prevention and the use of air drying to help detect a carious lesion ( $< 0.001$ ). Dentists in the PDA (60%) region would be less likely to use air drying when diagnosing a caries lesion than dentists in the SK region (84%,  $p = 0.004$ ). Time of air drying presented no significant associations.

There was a positive association between the percentage of patients who receive individualized caries prevention and the use of fiber optic transillumination on proximal lesions ( $p < 0.001$ ), with no significant regional differences.

Male dentists (61% vs. 40% for females) and practices with a lower percentage of pediatric patients ( $p < 0.001$ ) were more likely to use magnification when examining a carious lesion. Dentists in the FL/GA (79%,  $p = 0.009$ ), MN (73%,  $p = 0.001$ ) and PDA (67%,  $p = 0.004$ ) regions use magnification for caries detection more often than dentists from the SK region (31%). Dentists in the FL/GA ( $p < 0.001$ ) and MN ( $p = 0.013$ ) regions also use magnification more often than dentists in the AL/MS region (49%).

## DISCUSSION

The accurate diagnosis of the presence of disease is paramount for appropriate care. Even though several methods have been described for the diagnosis of dental caries, no consistent criteria are available.<sup>8</sup> The diagnosis of occlusal caries is highly subjective, with considerable variation in the ability and experience among clinicians to appropriately diagnose and treat occlusal caries.<sup>9,10</sup> Reviews of the literature point to methodological difficulties in drawing valid comparisons between studies due to incompatible criteria and simulations.<sup>11,12</sup> It has also been concluded that no caries diagnostic method fulfills all of the ideal criteria for accurate measurements needed to plan appropriate care.<sup>11-13</sup>

In the current study, dentists would use the dental explorer, a tactile tool, as the main instrument for the detection of occlusal and recurrent dental caries. The low specificity (approximately 40 percent) reported for this diagnostic tool when used as the only diagnostic method is of concern.<sup>14-19</sup> Dentists tend to over-diagnose caries if a sharp instrument is used and sticks in any deep pit and fissure without the true evidence of caries. A “sticking” probe is not necessarily indicative of decay and may be entirely due to local anatomic features. Additionally, applying pressure with a sharp explorer has been called into question, particularly in Europe and Scandinavia, because of documented damage to surface integrity and possible implantation of organisms, both of which may increase lesion susceptibility.<sup>20, 21</sup> Although this last issue is somewhat contentious, the evidence suggests that an explorer should be used lightly or not at all on occlusal surfaces. As shown in this study some dentists from the SK region not use dental explorer at all for the detection of occlusal dental caries.

The use of magnification to assist visual examination was consistently used by almost half of the dentists in the study in the United States regions. The visual examination for evaluating carious lesions has been described and, in certain studies, results have been validated *in vitro* by sectioning the teeth after the lesions had been visually scored<sup>22</sup>; the histological features are considered as the ‘gold standard’.<sup>23</sup> The visual method for evaluation of dental caries has low sensitivity and high specificity in diagnosing occlusal caries.<sup>24-26</sup> Conversely some dentists from the SK region do not use magnification at all for dental caries detection.

This study did not address the issue of whether an individual dentist reported using more than one method to assess caries diagnosis. It is possible that the same individuals who reported that they would use the explorer in 100% of the cases could also always use magnification, and vice-versa. We did explore whether certain pairs of diagnostic techniques were more often used by certain dentists than others. However, when we calculated correlations between the frequencies of use for each of the caries diagnostic techniques, the correlations were small, ranging from  $r = 0.15$  for radiographs and explorer on occlusal lesion, to occlusal explorer and magnification  $r = 0.02$ . Apart from numerous possible combinations of the data, some studies reported on the use of combined methods to improve the accuracy of the diagnostic of caries. This approach also seems to be a matter of controversy. While the combination of three methods (*e.g.*, visual examination, radiographic examination, and laser fluorescence) has been reported to improve accuracy and provide the best treatment plan for occlusal surfaces<sup>27</sup>, others have indicated that a combination of diagnostic methods (*ca.* bitewing radiographs, electric conductance, quantitative light fluorescence, and DIAGNOdent) only slightly improved the correct diagnosis when compared to visual examination alone. Of more concern was the fact that the combined methods resulted in decreased specificity, thus leading examiners to elect more invasive treatments.<sup>28</sup>

The use of radiographs for caries detection has a long history, and it is still a widely used diagnostic method.<sup>13, 29</sup> In the current study, radiographs have been consistently and widely used for the diagnosis of proximal dental caries and less often for the diagnosis of occlusal dental caries. It is accepted that occlusal lesions are initiated on the fissure walls and can therefore be obscured by sound superficial tissue.<sup>30</sup> This seems to be a known concept among the respondents and it may explain the tendency for a lower frequency of use of radiographs as a diagnostic method for occlusal caries. Dental radiographs have been reported to be inadequate for detecting caries in the occlusal surface until the lesion is well advanced through the enamel and into the dentin.<sup>30</sup> False positives can also occur with radiographic diagnosis, and specificities of 66% to 98% have been recorded *in vitro*.<sup>31,32</sup> However, studies have reported high sensitivity values for radiographs when diagnosing proximal dental caries.<sup>14, 33-37</sup> More importantly, radiographs can be a reliable way to monitor the remineralization process of proximal dental caries if fluoride therapy or other remineralization treatment is being used. Our study has shown that 96% of the respondents would use radiographs for diagnosis of proximal lesions in more than 75% of the cases. With the exception of SK, the use of radiographs was reported to be used in more than 75% of the cases by the vast majority (>90%) of all respondents from US regions. This study did not investigate all diagnostic methods available for detection of proximal lesions in posterior teeth and, therefore, it is unclear whether respondents make use of other diagnostic methods in those cases. Improved methods for the diagnosis of proximal lesions have been reported with the use of tooth separation and impression techniques.<sup>38,39</sup> These methods accurately identify the presence of cavitation on the surface and thus allow for a more accurate decision regarding treatment.

Fiber-optic transillumination (FOTI) has been evaluated in a number of studies for detection of posterior proximal caries lesions reporting low to good sensitivity and good specificity.<sup>35, 40-43</sup> One *in vitro* study suggested that a combination of FOTI and visual examination is valid for determination of occlusal lesion depth.<sup>44</sup> As the buccal-lingual distance of anterior teeth is, in general, smaller than in posterior teeth, clinicians may rely more on visual inspection for the examination in this area. This may explain the low frequency of use of FOTI found in our study. Yet, 10% reported on using it in more than 50% of the cases.

Even though advances in technology have made available other assisting diagnostic methods, such as FOTI and DIAGNOdent, the study shows that dentists participating in the

DPBRN do not frequently use these methods. Various levels of sensitivity have been reported for DIAGNOdent<sup>34, 45-50</sup> and FOTI<sup>44, 51-53</sup> for the diagnosis of occlusal and proximal dental caries. The strength of the evidence however, is low as the available literature is insufficient to support generalizable estimates of the sensitivity and specificity studies of these diagnostic methods. Again, there was no attempt to investigate whether clinicians are using FOTI and Diagnodent as a secondary method in combination with the more traditional methods. From the lower frequencies reported, we speculate that this is the case.

Caries lesions occur in a variety of anatomic locations and have unique aspects of configuration and rate of progression. These differences make it unlikely that a single diagnostic method will have the adequate sensitivity and specificity to detect caries at all sites. Reports in the literature have concluded that conventional visual, tactile, and radiographic examinations used as individual diagnostic methods provide less than ideal diagnostic sensitivity.<sup>14-17, 54</sup>

Many of the Alabama dentists were part of pre-existing dental network that was not formed specifically for the DBPRN and this may partially explain the participation rates in this study. Additionally, the study sample is not a random sample of general dentists in the United States. Consequently, the extent to which these findings generalize to this population cannot be stated with certainty. Based on comparisons to dentists who responded to the 2004 ADA Survey of Dental Practice, DPBRN dentists have much in common with dentists at large.<sup>5-6</sup> The only key characteristic with a statistically significant difference seems to be that DPBRN dentists tend to have a more-recent year of graduation from dental school than dentists at large. DPBRN dentists certainly represent a substantial diversity with regard to practice settings, patient populations, rural-urban area of residence, and geographic locations.

Substantial differences exist among dentists regarding the clinical decision-making process, which is in part due to the lack of sensitivity and specificity provided by some caries diagnostic methods. Variation of the diagnostic methods among the different DPBRN regions is probably related to the type of participating practices. AL/MS and FL/GA participants are generally solo practitioners in fee-for-service, self-pay, or private insurance-based practices. PDA and MN comprise dentists working in large group practices, SK practitioners are mostly in public health practices, and the MN region mainly comprises large group practices with a few solo private practice practitioners. Since public health practices and some large group practices employ the practitioners, it may be that these organizations are able to use standardized training and diagnostic tools. HP Dental Group, for example, has its own guideline for caries risk assessment, called “caries guideline” which has specific instructions on assessment of caries risk and protocols for radiographic and clinical diagnosis of dental caries. In fact, MN and SK were the regions reporting the least percentage of clinicians using explorer in all cases. These are the regions more aligned with the current scientific evidence on the validity of using an explorer to detect occlusal caries.

An International Consensus Report, 20 years ago, concluded that “research emphasis should be placed on the development of improved and standardized methods for caries diagnosis”.<sup>54</sup> Today's status of caries diagnosis in dental private offices still calls for evidence-based and standardized protocols. Unfortunately, the status of caries diagnosis has not changed significantly during the past several decades. A systemic review completed recently corroborates this conclusion (Ewoldsen and Koka, 2010). Without proper protocol and consistent criteria for caries diagnosis, the cost of dental treatment could be significantly affected (Shugars and Bader, 1996). Establishing accurate methods and criteria for caries

diagnosis is fundamental to determining normative treatment needs, to reduce dental over-treatment (Bader and Shugars, 1992) and consequently reduce the cost of care.

## CONCLUSIONS

There is considerable variation in the methods used for diagnosing caries across the DPBRN regions.

1. AL/MS reported the highest frequency of use of dental explorer in 100% of the cases and MN the lowest.
2. Use of radiographs to diagnose proximal caries for all patients ranged from 63% in FL/GA to 21% in SK. Dentists who provide a higher percentage of patients with individualized caries prevention and who see a higher percentage of pediatric patients are more likely to use radiographs to assess proximal surfaces of posterior teeth.
3. Laser fluorescence on occlusal surfaces was used most often by practices that perform the highest percentage of restorative procedures. There was a trend for dentists with the least number of years since dental school graduation to use laser fluorescence more often than dentists with longer tenure.

Overall, the dental explorer and radiographs are still the main diagnostic methods for the diagnosis of dental caries.

## Acknowledgments

This investigation was supported by NIH-NIDCR grants DE-16746 and DE-16747. Opinions and assertions contained herein are those of the authors and are not to be construed as necessarily representing the views of the respective organizations or the National Institutes of Health.

## REFERENCES

1. Pine CM, ten Bosch JJ. Dynamics of and diagnostic methods for detecting small carious lesions. *Caries Research*. 1996; 30(6):381–388. [PubMed: 8946093]
2. Bader JD, Shugars DA, Bonito AJ. A systematic review of the performance of methods for identifying carious lesions. *Journal of Public Health Dentistry*. 2002; 62(4):201–13. [PubMed: 12474624]
3. Tranæus S, Shi X-Q, Angmar-Månsson B. Caries risk assessment: Methods available to clinicians for caries detection. *Community Dentistry and Oral Epidemiology*. 2005; 33:265–273. [PubMed: 16008633]
4. Espelid I, Tveit AB, Fjellveit A. Variations among dentists in radiographic detection of occlusal caries. *Caries Research*. 1994; 28:169–175. [PubMed: 8033190]
5. Makhija SK, Gilbert GH, Rindal DB, Benjamin PL, Richman JS, Pihlstrom DJ, for the DPBRN Collaborative Group. Dentists in practice-based research networks have much in common with dentists at large: evidence from The Dental Practice-Based Research Network. *General Dentistry*. 2009; 57:270–275. [PubMed: 19819818]
6. Makhija SK, Gilbert GH, Rindal DB, Benjamin PL, Richman JS, Pihlstrom DJ, Qvist V, for the DPBRN Collaborative Group. Practices participating in a dental PBRN have substantial and advantageous diversity even though as a group they have much in common with dentists at large. *BioMed Central Oral Health*. 2009; 15:9–26.
7. Gilbert GH, Williams OD, Rindal DB, Pihlstrom DJ, Benjamin PL, Wallace MC, DPBRN Collaborative Group. The creation and development of the dental practice-based research network. *Journal of the American Dental Association*. 2008; 139(1):74–81. [PubMed: 18167389]



8. National Institutes of Health. Office of Medical Application of Research. Consensus Development Conference Statement: Diagnosis and Management of Dental Caries Throughout Life. 2001. March 26-28, 2001. from: [http://odp.od.nih.gov/consensus/cons/115/115\\_intro.htm](http://odp.od.nih.gov/consensus/cons/115/115_intro.htm)
9. Bader JD, Brown JP. Dilemmas in caries diagnosis. *Journal of the American Dental Association*. 1993; 124:48–50. [PubMed: 8505450]
10. Swenson E, Hennessy B. Detection of occlusal carious lesions: an in vitro comparison of clinicians diagnostic abilities at varying levels of experience. *General Dentistry*. 2009; 51:60–66. [PubMed: 19146144]
11. Bader J, Shugars D, Bonito A. Systematic reviews of selected dental caries diagnostic and management methods. *Journal of Dental Education*. 2001; 65:960–968. [PubMed: 11699997]
12. Stookey GK, Gonzalez-Cabezas C. Emerging methods of caries diagnosis. *Journal of Dental Education*. 2001; 65:1001–1006. [PubMed: 11699969]
13. Van Rijkom HM, Verdonschot EH. Factors involved in validity measurements of diagnostic tests for approximal caries—a meta-analysis. *Caries Research*. 1995; 29:364–370. [PubMed: 8521438]
14. Chong MJ, Seow WK, Purdie DM, Cheng E, Wan V. Visual-tactile examination compared with conventional radiography, digital-radiography, and DIAGNOdent in the diagnosis of occlusal occult caries in extracted premolars. *Pediatric Dental Journal*. 2003; 25:341–349.
15. Jahangiri L, Wahlers C, Hittelman E, Matheson P. Assessment of sensitivity and specificity of clinical evaluation of cast restoration marginal accuracy compared to stereomicroscopy. *Journal of Prosthetic Dentistry*. 2005; 93:138–142. [PubMed: 15674223]
16. Lussi A. Validity of diagnostic and treatment decisions of fissure caries. *Caries Research*. 1991; 25:296–303. [PubMed: 1913769]
17. Thompson NJ, Boyer EM. Validity of oral health screening in field conditions: pilot study. *Journal of Dental Hygiene*. 2006; 80:9. [PubMed: 16734994]
18. Verdonschot EH, Bronkhorst EM, Burgersdijk RC, Konig KG, Schaeken MJ, Truin GJ. Performance of some diagnostic systems in examinations for small occlusal carious lesions. *Caries Research*. 1992; 26:59–64. [PubMed: 1568239]
19. Ricketts D, Kidds E, Weerheijm K, de Soet H. Hidden caries: what is it? Does it exist? Does it matter? *International Dental Journal*. 1997; 47:259–265. [PubMed: 9448806]
20. Ekstrand KR, Qvist V, Thylstrup A. Light microscope study of the effect of probing in occlusal surfaces. *Caries Research*. 1987; 21:368–374. [PubMed: 3475183]
21. van Dorp CS, Exterkate RA, ten cate JM. The effect of dental probing on subsequent enamel demineralization. *American Society of Dentistry for Children. Journal of Dentistry for Children*. 1988; 55:343–347. [PubMed: 3170872]
22. Ekstrand KR, Ricketts DN, Kidd EA. Reproducibility and accuracy of three methods for assessment of demineralization depth of the occlusal surface: an in vitro examination. *Caries Research*. 1997; 31:224–231. [PubMed: 9165195]
23. Machiulskiene V, Nyvad B, Baelum V. Prevalence and severity of dental caries in 12-year-old children in Kaunas, Lithuania. *Caries Research*. 1998; 32:175–180. [PubMed: 9577982]
24. Le YL, Verdonschot EH. Performance of diagnostic systems in occlusal caries detection compared. *Community Dentistry and Oral Epidemiology*. 1994; 22:187–191. [PubMed: 8070247]
25. Kidd EAM, Ricketts DNJ, Pitts NB. Occlusal caries diagnosis: a changing challenge for clinicians and epidemiologists. *Journal of Dentistry*. 1993; 21:323–331. [PubMed: 8258581]
26. Wenzel A, Larsen MJ, Fejerskov O. Detection of occlusal caries without cavitation by visual inspection, film radiograph, xeroradiographs, and digitized radiographs. *Caries Research*. 1991; 25:365–371. [PubMed: 1747887]
27. Valera FB, Pessan JP, Valera RC, Mondelli J, Percinoto C. Comparison of visual inspection, radiographic examination, laser fluorescence and their combinations on the treatment decisions for occlusal surfaces. *American Journal of Dentistry*. 2008; 21:25–29. [PubMed: 18435372]
28. Pereira AC, Eggertsson H, Martinez-Mier EA, Mialhe FL, Eckert GJ, Zero DT. Validity of caries detection on occlusal surfaces and treatment decisions based on results from multiple caries-detection methods. *European Journal of Oral Sciences*. 2009; 117:51–57. [PubMed: 19196318]
29. Pitts NB, Stamm JW. Proceedings from the international consensus workshop on caries clinical trials. *Journal of Dental Research*. 2004; 83(Special Issue C):125–128.

30. Pitts, N. Advances in radiographic detection methods and caries management rationale. In: Stookey, GK., editor. Early detection of dental caries. School of dentistry, Indiana University; Indianapolis, IN: 1996. p. 39-50.
31. Ferreira Zandona AG, Analoui M, Schemehorn BR, Eckert GJ, Stookey GK. Laser fluorescence detection of demineralization in artificial occlusal fissures. *Caries Research*. 1998; 32:31–40. [PubMed: 9438569]
32. Huysmans MC, Longbottom C, Pitts N. Electrical methods in occlusal caries diagnosis: An in vitro comparison with visual inspection and bite-wing radiography. *Caries Research*. 1998; 32:324–329. [PubMed: 9701656]
33. Pitts NB, Renson CE. Reproducibility of computer-aided image-analysis-derived estimates of the depth and area of radiolucencies in approximal enamel. *Journal of Dental Research*. 1985; 64:1221–1224. [PubMed: 3861653]
34. Tonioli MB, Bouschlicher MR, Hillis SL. Laser fluorescence detection of occlusal caries. *American Journal of Dentistry*. 2002; 15:268–273. [PubMed: 12572647]
35. Vaarkamp J, ten Bosch JJ, Verdonschot EH, Bronkhorst EM. The real performance of bitewing radiography and fiber-optic transillumination in approximal caries diagnosis. *Journal of Dental Research*. 2000; 79:1747–1751. [PubMed: 11077989]
36. Valachovic RW, Douglass CW, Berkey CS, McNeil BJ, Chauncey HH. Examiner reliability in dental radiography. *Journal of Dental Research*. 1986; 65:432–436. [PubMed: 3457049]
37. Yalcinkaya S, Kunzel A, Willers R, Thoms M, Becker J. Subjective image quality of digitally filtered radiographs acquired by the Durr Vistascan system compared with conventional radiographs. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology & Endodontics*. 2006; 101:643–651.
38. Seddon RP. The detection of cavitation in carious approximal surfaces in vivo by tooth separation, impression and scanning electron microscopy. *Journal of Dentistry*. 1989; 17:117–120. [PubMed: 2671075]
39. Mariath AA, Bressani AE, Araujo FB. Elastomeric impression as a diagnostic method of cavitation in proximal dentin caries in primary molars. *Journal of Applied Oral Science*. 2007; 15:529–533.
40. Oby-Musset AM, Cahen PM, Turlot JC, Frank R. Approximal caries diagnosis in epidemiological studies: transillumination or bitewing radiographs? *Journal de Biologie Buccale*. 1988; 16:13–17. [PubMed: 3165088]
41. Peers A, Hill FJ, Mitropoulos CM, Holloway PJ. Validity and reproducibility of clinical examination, fiber-optic transillumination, and bite-wing radiology for the diagnosis of small approximal carious lesions: an in vitro study. *Caries Research*. 1993; 27:307–311. [PubMed: 8402807]
42. Stephen KW, Russell JI, Creanor SL, Burchell CK. Comparison of fiber-optic transillumination with clinical and radiographic caries diagnosis. *Community Dentistry and Oral Epidemiology*. 1987; 15:90–94. [PubMed: 3552397]
43. Vaarkamp J, ten Bosch JJ, Verdonschot EH, Traanaeus S. Quantitative diagnosis of small approximal caries lesions utilizing wavelength-dependent fiber-optic transillumination. *Journal of Dental Research*. 1997; 76:875–882. [PubMed: 9126184]
44. Cortes DF, Ellwood RP, Ekstrand KR. An in vitro comparison of a combined FOTI/visual examination of occlusal caries with other caries diagnostic methods and the effect of stain on their diagnostic performance. *Caries Research*. 2003; 37:8–16. [PubMed: 12566633]
45. Aljehani A, Yang L, Shi XQ. In vitro quantification of smooth surface caries with DIAGNOdent and DIAGNOdent pen. *Acta Odontologica Scandinavica*. 2007; 65:60–63. [PubMed: 17354096]
46. El-Housseiny AA, Jamjoum H. Evaluation of visual, explorer, and a laser device for detection of early occlusal caries. *Journal of Clinical Pediatric Dentistry*. 2001; 26:41–48. [PubMed: 11688812]
47. Kuhnisch J, Ziehe A, Brandstad A, Heinrich-Weltzien R. An in vitro study of the reliability of DIAGNOdent measurements. *Journal of Oral Rehabilitation*. 2004; 31:895–899. [PubMed: 15369472]
48. Lussi A, Hellwig E. Performance of a new laser fluorescence device for the detection of occlusal caries in vitro. *Journal of Dentistry*. 2006; 34:467–471. [PubMed: 16431009]

49. Virajsilp V, Thermontree A, Aryatawong S, Paiboonwaraschat D. Comparison of proximal caries detection in primary teeth between laser fluorescence and bitewing radiography. *Pediatric Dental Journal*. 2005; 27:493–499.
50. Yin W, Feng Y, Hu D, Ellwood RP, Pretty IA. Reliability of quantitative laser fluorescence analysis of smooth surface lesions adjacent to the gingival tissues. *Caries Research*. 2007; 41:186–189. [PubMed: 17426397]
51. Hintze H, Wenzel A, Danielsen B, Nyvad B. Reliability of visual examination, fiber-optic transillumination and bite-wing radiography, and reproducibility of direct visual examination following tooth separation for the identification of cavitated carious lesions in contacting approximal surfaces. *Caries Research*. 1998; 32:204–209. [PubMed: 9577986]
52. Holt RD, Azevedo MR. Fibre optic transillumination and radiographs in diagnosis of approximal caries in primary teeth. *Community Dental Health*. 1989; 6:239–247. [PubMed: 2679987]
53. Schneiderman A, Elbaum M, Shultz T, Keem S, Greenebaum M, Driller J. Assessment of dental caries with digital imaging fiber-optic transillumination (DIFOTI): in vitro study. *Caries Research*. 1997; 31:103–110. [PubMed: 9118181]
54. McComb D, Tam LE. Diagnosis of occlusal caries: Part I. Conventional methods. *Journal of the Canadian Dental Association*. 2001; 67:454–457. [PubMed: 11583606]
55. Ewoldsen N, Koka S. There are no clearly superior methods for diagnosing, predicting, and noninvasively treating dental caries. *Journal of Evidence-Based Dentistry*. 2010; 10:16–17.
56. Shugars DA, Bader JD. Cost implications of differences in dentists' restorative treatment decisions. *Journal of Public Health Dentistry*. 1996; 56:219–222. [PubMed: 8906706]
57. Bader JD, Shugars DA. Understanding dentists' restorative treatment decisions. *Journal of Public Health Dentistry*. 1992; 52:102–110. [PubMed: 1564688]

**Table 1**

Questions used to assess the methods used by DPBRN dentists to diagnose dental caries.

Instructions. These questions have to do with methods that you may use to diagnose dental caries. Please circle the one number that best corresponds to your answer. Patients can vary substantially from one practice to the next, but we are interested in the patients in YOUR practice.

- 1 When you examine patients to determine if they have a caries lesion on a proximal (mesial or distal) surface, on a posterior tooth, on what percent of these patients do you use radiographs to help diagnose the lesion?
- 2 When you examine patients to determine if they have a caries lesion on the occlusal surface, on what percent of these patients do you use radiographs to help diagnose the lesion?
- 3 When you examine patients to determine if they have a primary occlusal caries lesion, on what percent of these patients do you use a dental explorer to help diagnose the lesion?
- 4 When you examine patients to determine if they have a caries lesion at the margin of an existing restoration (recurrent/secondary caries) on what percent of these patients do you use a dental explorer to help diagnose the lesion?
- 5 When you examine patients to determine if they have a primary caries lesion on the occlusal surface, on what percent of these patients do you use laser fluorescence (for example, Diagnodent®)?
- 6 When you examine patients to determine if they have a primary caries lesion, on what percent of these patients do you use air-drying to help diagnose the lesion?\*\*\*
- 7 When you examine patients to determine if they have a caries lesion on a proximal (mesial or distal) surface of an anterior tooth, on what percent of these patients do you use fiber optic transillumination to help diagnose the lesion?
- 8 When you examine patients to determine if they have a caries lesion, on what percent of these patients do you use some sort of magnification to help diagnose the lesion?

\*\*\* Respondents who reported using air-drying were also asked: Approximately how long do you dry the tooth surface? The response choices were: 1-2 seconds, 3-4 seconds, 5 seconds, more than 5 seconds **Participants had the following answering choices: 1 – Never or 0% 2 – 1 to 24% 3 – 25 to 49% 4 – 50 to 74% 5 – 75 to 99% 6 – Every time or 100%**

Note: The percentages for each method used were recoded to the categories' median to maintain the interval nature of the data so that parametric statistics could be used: 0 percent=0, 1-24 percent=12.5, 25-49 percent=37, 50-74 percent=62, 75-99 percent=87, 100 percent=100.

Table 2

Summary of dentists' and practices' characteristics

Practice variable	Total % (n)	AL/MS 57% (298)	FL/GA 19% (98)	MN 5% (28)	PDA 10% (51)	SK 9% (47)
Practice type						
Solo or small group practice	81% (422)	99% (296)	98% (96)	11% (3)	0 (0%)	57% (27)
Large group practice	14% (98)	0 (0%)	0 (0%)	89% (25)	100% (51)	0 (0%)
Public health practice	5% (51)	1% (2)	2% (2)	0 (0%)	0 (0%)	43% (20)
Gender (male)	81% (425)	86% (255)	86% (84)	68% (19)	82% (42)	53% (25)
Percent of pediatric patients	25% (SD=23)	25% (SD=21)	23% (SD=23)	23% (SD=19)	18% (SD=18)	31% (SD=33)
Years since dental school graduation (mean)	23 (SD=11)	24 (SD=11)	25 (SD=10)	19 (SD=9)	18 (SD=10)	22 (SD=12)
Caries risk assessed	70% (363)	65% (194)	56% (55)	79% (22)	98% (50)	89% (42)
Individualized caries prevention	52% (SD=34)	46% (SD=32)	53% (SD=35)	62% (SD=29)	71% (SD=29)	58% (SD=34)
Time spend in non-implant restorative services	60% (SD=19)	61% (SD=20)	59% (SD=19)	66% (SD=15)	59% (SD=14)	52 (SD=22)

**Table 3**

Frequency of use for each of the caries diagnostic methods by DPBRN dentists.

	<b>Range (Min-Max)</b>	<b>Not used at all % (n)</b>	<b>Used on all patients % (n)</b>	<b>% Patients Mean (SD)</b>
Radiograph proximal (n=522)	12-100%	0% (n=0)	50% (n=262)	92% (12)
Radiograph occlusal (n=521)*	0-100%	1% (n=1)	14% (n=71)	56% (33)
Dental explorer occlusal lesion (n=522)	0-100%	<1% (n=2)	61% (n=317)	88% (23)
Dental explorer on the margin of an existing restoration (n=520)*	0-100%	2% (n=9)	63% (n=328)	91% (16)
Laser fluorescence occlusal (n=521)*	0-100%	86% (n=450)	1% (n=4)	7% (21)
Air Drying (n=521)*	0-100%	4% (n=19)	32% (n=168)	70% (32)
Drying time in seconds (n=488)*	0-5+	----	----	1.2 (1.3)
Fiber optic proximal (n=520)*	0-100%	35% (n=180)	2% (n=9)	21% (27)
Magnification (n=521)*	0-100%	21% (n=110)	40% (n=205)	57% (43)

\* These numbers do not equal to 522 as some dentists left some of the questions unanswered.

**Table 4**

Practice characteristics associated with each diagnostic method.

	<b>B (SE)</b>	<b>SE</b>	<b>p</b>
<b>Radiograph for proximal caries</b>			
Years since dental school graduation	-.082	.047	.068
Individualized caries prevention	.030	.015	.040
Percent of pediatric patients	-.785	.228	.001
<b>Radiograph for occlusal lesions <sup>a</sup></b>			
<b>Explorer on occlusal lesions</b>			
Private practice	13.352	5.640	.018
Caries risk assessment	4.820	2.337	.040
<b>Explorer on the margin of an existing restoration</b>			
Private practice	7.692	4.198	.068
Caries risk assessment	3.067	1.678	.068
<b>Use of laser fluorescence on occlusal surfaces</b>			
Percent of non-implant restorative work	1.900	.484	< .001
Year since graduation from dental school	-.158	.092	.088
<b>Air drying when diagnosing occlusal caries</b>			
Individualized caries prevention	.209	.045	<.001
<b>Fiber optic transillumination for proximal caries</b>			
Individualized caries prevention	.141	.039	<.001
<b>Use of Magnification</b>			
Gender	-13.414	5.136	.009
Percentage of pediatric patients	-4.104	.886	<.001

This table provides parameter estimates for practice characteristic associated with the frequency of use of each diagnostic method where significance was <.10 and are adjusted for regional differences. Regional differences are reported in results section text for each diagnostic method.

<sup>a</sup>None of the practice characteristic were associated with the use of radiographs for occlusal lesions.

Private practice variable was coded as SGP=2, PHS or LGP = 1

Caries risk assessment yes=1, no=2