

Published in final edited form as:

J Alzheimers Dis. 2010 January 1; 22(3): 741–763. doi:10.3233/JAD-2010-101020.

Amyloid Beta as a Modulator of Synaptic Plasticity

Mordhwaj S Parihar¹ and Gregory J Brewer^{2,*}

¹ School of Studies in Biotechnology & Zoology, Vikram University, Ujjain, MP, India

² Kenneth Stark Endowed Chair in Alzheimer Research, Departments of Neurology and Medical Microbiology, Immunology and Cell Biology, Southern Illinois University School of Medicine, Springfield, IL, USA

Abstract

Alzheimer's disease is associated with synapse loss, memory dysfunction and pathological accumulation of amyloid beta in plaques. However, an exclusively pathological role for amyloid beta is being challenged by new evidence for an essential function of amyloid beta at the synapse. Amyloid beta protein exists in different assembly states in the central nervous system and plays distinct roles ranging from synapse and memory formation to memory loss and neuronal cell death. Amyloid beta is present in the brain of symptom-free people where it likely performs important physiological roles. New evidence indicates that synaptic activity directly evokes the release of amyloid beta at the synapse. At physiological levels, amyloid beta is a normal, soluble product of neuronal metabolism that regulates synaptic function beginning early in life. Monomeric amyloid beta 40 and 42 are the predominant forms required for synaptic plasticity and neuronal survival. With age, some assemblies of amyloid beta are associated with synaptic failure and Alzheimer's disease pathology, possibly targeting the N-methyl-D-aspartic acid (NMDA) receptor through the $\alpha 7$ -nicotinic acetylcholine receptor ($\alpha 7$ -nAChR), mitochondrial amyloid- β alcohol dehydrogenase (ABAD) and cyclophilin D. But emerging data suggests a distinction between age effects on the target response in contrast to the assembly state or the accumulation of the peptide. Both aging and beta amyloid independently decrease neuronal plasticity. Our laboratory has reported that amyloid beta, glutamate and lactic acid are each increasingly toxic with neuron age. The basis of the age-related toxicity partly resides in age-related mitochondrial dysfunction and an oxidative shift in mitochondrial and cytoplasmic redox potential. In turn, signaling through phosphorylated extracellular signal-regulated protein kinases (pERK) is affected along with an age-independent increase in phosphorylated cAMP response element-binding protein (pCREB) This review examines the long-awaited functional impact of amyloid beta on synaptic plasticity.

Keywords

Alzheimer's disease; Synapse; Aging; mitochondria; survival signaling

1. Introduction

Alzheimer's disease (AD) arises on a neuropathological background of amyloid plaques and neurofibrillary tangles (NFT) characterized by ongoing neurodegeneration in areas of brain involved in learning and memory. Synaptic pathology is an early marker of both AD and aging [1,2,3]. The best understood AD pathogenesis could be explained by a loss of plasticity [4,5] that may adversely affect dendritic ramifications, synaptic remodeling, long-

*Corresponding author: Professor Gregory J Brewer, gbrewer@siu.edu, Phone: 217-545-5230, FAX: 217-545-3227.

term potentiation (LTP), axonal sprouting, neurite extension, synaptogenesis, and neurogenesis. Plasticity, the process by which synapses modulate their strength and form new connections with other neurons plays a particularly important role in response to injury and disease [6]. Age is the most important and universal risk factor for AD perhaps because the biological capacity for plasticity decreases with age [7,8]. With aging, a combination of subclinical AD pathology, inflammation, Lewy bodies, microinfarcts and vascular lesions in the cortical and hippocampal regions contribute to late-life brain atrophy and dementia in each individual [9]. Thus, as a consequence of age, the ability to sustain learning and memory are diminished or slowed [10]. Age could interact with other variables that influence neuroplasticity in at least two ways. Conventionally, aging may shift the complex balance of amyloid beta ($A\beta$) metabolism away from the potentially neurotrophic products of α secretase processing and toward the production of neurotoxic moieties containing the intact $A\beta$ fragment [11]. Alternatively, aging could render the brain vulnerable to $A\beta$ -protein neurotoxicity [12] an age-related susceptibility [13]. $A\beta$ toxicity *in vivo* is also highly species-specific; toxicity is greater in aged rhesus monkeys than in aged marmoset monkeys, and is not significant in aged rats [12]. These results suggest that $A\beta$ neurotoxicity *in vivo* is a pathological response of the aging brain, which is more pronounced in higher order primates. Thus, longevity may contribute to the unique susceptibility of humans to AD by rendering the brain vulnerable to $A\beta$ neurotoxicity [12]. The pathological hallmark of $A\beta$ deposits appears to precede the hallmark of phosphorylated tau in the brain [14]. $A\beta$ 1-42 at high doses impairs cognitive and memory functions in mouse models of AD. But the relationship of $A\beta$ deposition to synapse loss is less clear in these models. Additionally, whether $A\beta$ deposits might be the early symptom contributing to neurodegeneration or whether synaptic pathology might be an early event preceding amyloid deposition in AD is not clear. Synaptic loss might occur early in AD and molecular biomarkers such as tau hyperphosphorylation and $A\beta$ deposits might be indicators of prolonged disease. Moreover, a central question is whether $A\beta$ plays a direct role in the neurodegenerative process in AD. There are two schools of thought on involvement of $A\beta$ in Alzheimer's disease. In one, $A\beta$ initiates the disease once produced in excess, which has motivated most AD clinical trials. Alternatively, $A\beta$ does not initiate but rather is secondary to other pathogenic events as a protective response to neuronal insult [15].

Two models have emerged to explain the role of $A\beta$ in normal and AD pathological state. According to the most popular model, oligomeric [16] and fibrillary $A\beta$ deposits [17,18] are responsible for the eventual neuronal degeneration involving disruption of glutamatergic synaptic function leading to the characteristic cognitive deficits [19,20,21,22]. The second model dictates that $A\beta$ may normally serve as a negative feedback signal that maintains neuronal activity within a normal dynamic range. *In vivo* studies on wild-type animals [23] and *in vitro* studies on wild-type [24] and knock-out [25] animals demonstrate that $A\beta$ production significantly increases with increase in communication between brain cells [26] and this increased level depresses excitatory synapses and reduces neuronal activity. $A\beta$ was proposed as a regulator of ion channel function [27] and as essential for neuronal health [28]. $A\beta$ is secreted from neurons in response to synaptic activity and that $A\beta$, in turn, down regulates synaptic transmission [29]. This negative feedback loop could operate as a physiological homeostatic mechanism to limit levels of neuronal activity [30].

A youthful role for $A\beta$ may enhance neuronal plasticity to help the remaining neural circuits compensate for lost or broken circuits and improve overall network performance and neurological function (Figure 1). Improving network activity may also help to prevent the inexorable loss of neuronal processes and cell bodies that occurs in AD and other neurodegenerative disorders. In the present paper we discuss the recent mechanistic link between $A\beta$ function and synaptic plasticity. This review focuses on the interface between a

physiological role of A β and toxicity in terms of molecular mechanisms of synaptic plasticity.

2. A β formation and assembly

The A β peptide is derived by proteolytic processing of the amyloid precursor protein [APP; 31,32], a type I integral membrane protein [33]. A β is a prominent constituent of the amyloid plaques present in AD brain [34]. In the beginning, APP is delivered to the plasma membrane where it is subjected to proteolytic processing by α -secretase. However, absent the α -secretase cleavage, APP molecules are internalized into endocytic compartments where they are subjected to cleavage by β -secretase (BACE) and γ -secretase to generate A β . Therefore, inhibition of the combined action of β - and γ -secretase that leads to A β peptide generation is regarded as a promising approach to treat AD. The Golgi apparatus and to a lesser extent the endoplasmic reticulum are also sources of a distinct population of A β peptides secreted into the extracellular space [35]. The majority of secreted A β peptides are 40 amino acids in length (A β 40), whereas a smaller fraction of A β is cleaved to produce a 42 amino acid species (A β 42). A β 42 is the main amyloid peptide that drives production of amyloid fibrils [36] in AD patients. A β in turn can be degraded by proteases such as the insulin-degrading enzyme [37,38] and neprilysin [39]. A β generated from axon-transported APP is released from presynaptic sites and subsequently accumulates close to the nerve terminal [40].

The hydrophobicity of A β 42 leads to a number of aggregation states [41]. This ability to self-associate [49,50] into different assembly states ranges from dimers to soluble oligomers to insoluble aggregates of fibrils [51]. Initially, it was assumed that toxicity was mediated by fibrillar A β similar to that present in amyloid plaques. This together with the findings that monomer is innocuous and that amyloid plaques alone cannot account for disease has led many to conclude, if it isn't fibrillar A β and it isn't A β monomer then it must be some other form of A β [52]. Soluble non-fibrillar A β assemblies [19,53] are more toxic than the fibrillar form, but as yet the specific form of A β which causes injury to neurons *in vivo* has not been identified. These oligomers have been described in cultured cells [44] as well as in APP transgenic mouse brain and human brain [45,46,47,48]. Synthetic oligomers specifically bind to synapses of hippocampal neurons [42,43]. Soluble A β (sA β) is found, at low concentrations, as a normal constituent of biological fluids [54,55,56]. As demonstrated by Piccini et al. [57], the composition as well as the aggregation and toxicity properties of soluble A β aggregates that accumulate in AD is different from those of normal aging. One such form of A β known as N-terminal truncated pyroglutamyl amyloid peptide (A β py3-42) is the predominant form of early aggregates that alter the membrane permeability, suggesting that they form pores in the membrane like other amyloidogenic peptides [57,58]. Naturally occurring A β peptides can begin to assemble *in vivo* into metastable dimers, trimers and higher oligomers while still at low nanomolar levels [44,59,60]. Immunocytochemical approaches identified different forms of A β produced with age in an APP transgenic (Tg2576) mouse model of AD [61]. Initially A β 40 and A β 42, the most predominant form of A β , occurs in its soluble form. At 6–10 months, the soluble forms of A β decrease as SDS-insoluble forms of A β 40 and A β 42 increase exponentially. SDS-resistant A β oligomers develop only in older Tg2576. After 11–12 months, A β is converted into diffuse plaques. At later ages, the A β accumulates in diffuse plaques, neuritic plaques with amyloid cores [61]. It seems reasonable that the synaptic and neuronal compromise seen at sites distant from plaques is mediated by an A β species that can readily diffuse and access the space in and surrounding the synaptic cleft [52]. The physiological level of A β is controlled by its production, degradation and clearance [23,62] while a defect in clearance leads to the accumulation of A β [52].

3. Activity-dependent A β production

Since synaptic loss correlates better with memory loss than plaque burden and people with mutations in the APP gene who make more amyloid invariably develop AD, the critical question emerges, what is the role of A β in synaptic activity and progression of the disease process? The mechanism regulating A β production and its subsequent release by neurons is closely linked with synaptic activity [29]. Thus, understanding the factors that regulate A β levels has implications for disease pathogenesis as well as for developing therapeutics. In animal models as well as in humans, the activation of muscarinic M1 acetylcholine receptors increases α -secretase cleavage of APP and consequently reduces A β levels [63,64] whereas activation of NMDARs decreases α -secretase cleavage, consequently increasing A β levels [65]. Stimulation with muscarinic agonists or activators of protein kinase C (PKC), such as phorbol esters causes the up-regulation of the α -secretase cleavage of APP [66]. Thus, regulation of the α -secretase contributes to regulation of A β peptide and toxicity in vivo [67]. Modulating synaptic transmission has been shown to alter extracellular soluble A β levels in organotypic brain slice [29]. Synaptic activity modulates interstitial fluid A β levels in vivo in APP transgenic and wild-type mice [23]. Neural activity regulates the trafficking of proteins at synaptic sites [68,69]. Thus it is possible that the induction of neural activity promotes the endocytosis of surface APP, enhancing the accessibility of APP to BACE in endosomal/recycling compartments [29]. In acute brain slices, synaptic vesicle cycling alone, in the absence of neuronal depolarization, was sufficient to drive release of A β from neurons [23]. Similar increases in A β levels were demonstrated in hippocampal seizures induced by electrical stimulation. However, decreasing synaptic transmission using tetrodotoxin (TTX) or tetanus toxin rapidly reduced interstitial fluid A β levels by 30% and 80%, respectively [26]. Cirrito et al. [26] also show that synaptic activity-induced increase in endocytosis drives more APP into the endocytic compartment, ultimately resulting in increased A β production and release. A β produced in the endocytic pathway is then brought to the cell surface where it is released into the extracellular fluid [70]. Inhibition of endocytosis reduces APP internalization and reduces A β production and release in cell lines [71]. Support for a casual link between synaptic activity and A β levels in humans comes from recent brain imaging studies of regions that contain the most metabolic activity throughout life (and presumably have the highest levels of neuronal activity) are the same regions that degenerate and accumulate A β [72]. The increased synaptic activity enhances both oligomer formation and localization at synaptic sites in rat and mouse hippocampal slices and primary human cortical neurons in culture [73]. These oligomers appear to bind to NMDARs at the synapse [74] and induce internalization of NMDAR [75] and deregulation of NMDA signaling pathways [76]. In this regard, neuronal activity not only enhances oligomer targeting but also facilitates oligomer formation at synaptic sites [73]. These findings indicate that sustained synaptic activity causes an increase in oligomeric A β which accumulates with age and leads to synaptic dysfunction and neuronal death. There seems to be a homeostasis of a normal negative feedback function under normal physiological conditions where an increased neuronal activity produces more A β ; the enhanced A β production depresses synaptic function; the depressed synaptic function will decrease neuronal activity [29]. Disturbances in this homeostatic mechanism of A β could produce the problems of AD patients. Persistently elevated neuronal activity if it is unchecked could lead to excitotoxicity [77], as well as higher levels of secreted A β peptides, which may convert to neurotoxic fibrils with consequent synaptic depression and neuronal toxicity [78,79]. Thus A β toxicity might represent a disturbance of normal function with the net balance of production *versus* clearance determining a beneficial synaptic or toxic fate.

4. A normal physiologic role for A β

A β has been extensively studied because of its association with plaques in AD brains, interference with synaptic function and possible pathogenesis of AD [60]. However, A β exists in normal individuals without any known pathology. Therefore, there has been a conscientious search for its normal physiological role in the brain, particularly its possible involvement in synaptic plasticity and neuronal survival. A β at physiological levels is essential for synaptic plasticity in normal individuals [80]. The physiological functions of A β , as well as the primary mechanisms that initiate early A β -mediated synaptic plasticity are now being revealed. We hypothesize that the protective or destructive effects of A β are determined by its relative concentration in addition to the age-related cellular environment. Physiologically low regulated concentrations of A β would play a critical function in mediating synaptic plasticity and improve cognitive functions whereas, accumulation of higher concentrations of A β together with age effects cause disruptions of this regulation with consequent synaptic dysfunction and loss, as seen in AD [76]. Other factors such as structural changes of A β due to pathological processing and/or post-translational dysregulations as well as age related changes in A β clearance cannot be excluded. The large body of evidence for activity-dependent production of A β strongly suggests a normal function for this peptide. Proposed functions of A β include control of synaptic activity and memory consolidation, trophic and neuronal survival, cholesterol transport and antioxidant functions.

4.1. Lower levels of A β modulate synaptic plasticity

It is generally believed that plasticity, such as LTP and long-term depression (LTD), is important for learning and memory. Multiple signaling pathways, including several protein kinases and phosphatases, are required for the generation of LTP and LTD [81]. These same pathways have been shown to influence *in vivo* phenomena, such as learning and memory [82]. A β is released in lower amounts in normal brains throughout life during synaptic activity and seems to be beneficial for normal brain synaptic functions [83]. A β is normally produced in the brain, where the *in vivo* concentration in the rodent brain has been estimated to be in the picomolar range [84]. Picomolar A β is present in both cerebrospinal fluid and plasma of healthy individuals throughout life [55]. Recent results indicate that A β serves an essential role at the synapse and in synaptic structure-functional plasticity critical to learning and memory. A necessary role of A β in synaptic plasticity and memory in normal brain is supported by the observation that APP knock-out (KO) mice show impaired LTP and memory [85]. The impaired synaptic plasticity and memory found in BACE1 KO mice also suggest a necessary role of A β [86]. Similarly, the necessary role of A β in synaptic plasticity and memory is seen from loss of presenillin function (γ -secretase) [87]. Puzzo et al. [83] show that picomolar concentrations (200 pM) of both A β 42 monomers and oligomers cause a marked increase in long-term potentiation, whereas high nanomolar concentrations (200 nanomolar) lead to the well established reduction of potentiation in the hippocampus. Picomolar levels of A β 42 also produce a pronounced enhancement of both reference and contextual fear memory. Thus, lower concentrations play a novel positive modulatory role on neurotransmission and memory, whereas high concentrations are associated with neuronal cell death. The lower concentrations of aged A β 42 used by Puzzo et al. [83] are close to those found in the normal brain [88,89,90,91] and shown to enhance LTP and memory. Increase in synaptic activity will increase A β production [29] while lowering synaptic activity minimizes A β production. Similarly specific stimulation of NMDA receptors promotes A β production [65] and A β in turn depresses synaptic activity. Thus indirectly A β also plays a role in suppressing excessive glutamate release. Interestingly, A β may play an important role during synapse elimination [92] and stimulation of neurogenesis in the hippocampus during brain development [93]. These studies provide

convincing evidence to show that physiological levels of A β have a role in controlling synaptic activity.

4.2. Neuronal survival

A β 42 is normally produced and secreted by cells in much lower quantities than A β 40, which represents ~90% of the total secreted A β [94]. However both species of A β are necessary for neuronal survival. Both A β 40 and A β 42 at physiological concentrations are important in neuronal survival and memory (Table 1). Physiological levels of A β also have trophic and neuroprotective actions in trophic deprived conditions [95]. Many A β has a physiological role in normal synapse function. In organotypic hippocampal slices, BACE activity is increased by synaptic activity and the resulting A β peptides depress excitatory transmission through α -amino-3-hydroxyl-5-methyl-4-isoxazole-propionate (AMPA) and N-methyl-D-aspartic acid (NMDA) receptors, suggesting a role for A β in homeostatic plasticity [29]. A β may have an important physiological role in synapse elimination during brain development [reviewed in reference 92]. Inhibition of endogenous A β production by exposure to inhibitors either of β - or γ -secretases in primary neuronal cultures caused neuronal cell death [28]. Thus targeting A β formation pharmacologically, or immunologically, could be deleterious [96]. Both A β 40 and A β 42 have been shown to be protective. Synthetic A β 42 monomers at 30–100 nM support the survival of developing neurons under conditions of trophic deprivation and protect mature neurons against excitotoxic death [97]. In cultured neural stem cells, A β 42 increased the number of newborn neurons [93]. A β 42 exhibited highly protective effects not only when combined with NMDA (100 nM), but also when applied before or after the NMDA pulse [97]. The later evidence excludes a direct interaction between A β 42 monomers and NMDARs. Monomers of A β 40 were also fully protective against NMDA toxicity. Neurotrophic function of A β 40 was obtained in a cell culture treated with picomolar levels of A β 40 [98,28]. Cells treated with such picomolar levels of A β 40 reverse the toxicity of secretase inhibition. These findings provide compelling evidence for a role for A β in neuronal survival. The pro-survival role of A β is summarized in Fig. 1.

4.2.1. Insulin like growth factor-I (IGF-I) and insulin signaling—The underlying mechanism of neuronal survival with A β is emerging. Neuronal synapses and astrocytes of memory-processing brain regions possess insulin receptors (IRs) [99] which when activated by insulin facilitate synaptic plasticity in normal brain [100]. IR and Insulin-like growth factor I (IGF-I) receptors consist of α -subunits and transmembrane β -subunits. Binding of insulin or IGF-I to the α -subunit increases the intrinsic tyrosine kinase activity of the β -subunit, and causes autophosphorylation of the β -subunit, thus triggering tyrosine phosphorylation of insulin receptor substrate (IRS)-1 and IRS-2, as well as Shc [101] as an important pathway of cell survival. To protect against A β toxicity, the tyrosine-phosphorylated sites create binding sites for various signal-transducing molecules containing Src homology-2 domain, such as phosphoinositide 3-kinase (PI3K) and growth factor receptor-bound protein 2 (Grb2), thus activating PI3K/phosphoinositide-dependent kinase 1 (PDK1)/Akt (protein kinase B)/glycogen synthase kinase (GSK)-3 α -3 β and Ras/Raf-1/mitogen-activated protein kinase/extracellular-signal regulated kinase (MEK/ERK) signaling pathways [101]. In normal brain, IGF-I and insulin promote glucose utilization, energy metabolism, and neuronal survival [102], largely through PI3K/Akt/GSK-3 β signaling [103,104]. Consistent with positive effects of insulin on synaptic plasticity [105], acute insulin treatment improved memory in rats [106] and also in normal adults and AD patients [107] by strongly activating ERK and Akt and blocking c-Jun N-terminal kinase (JNK) activation in a PI3K-dependent manner [108]. The mechanism involves many steps beginning with A β activation of IGF-1/insulin receptors by locally produced IGF-1 or,

possibly, A β monomers may bind to IGF-1/insulin receptors, as already shown for A β oligomers [109,110].

4.2.2. The PI3K/Akt signaling—PI3K, a membrane-associated second messenger protein, and its downstream kinase, Akt, are associated with neuronal survival [111] and plasticity [112] via activation of transcription pathways and protein synthesis. PI3K pathway, which required the activation of IGF-1/insulin receptors, is the most convincing pro-survival effect of A β 42 monomers. The PI3K signaling pathway is important in the transmission of survival signals in many cell types including neurons [113,114]. The PI3K-Akt signaling cascade, initiated by IRS, is phosphorylated by stimulated insulin- and IGF-receptor tyrosine kinases [115]. One of the kinases known to lie downstream of PI3K is Akt, which can be directly activated by products of PI3K [116] by promoting its phosphorylation at Ser473 and Thr308 [117]. Activated Akt, in turn, phosphorylates a wide range of substrates activating anti-apoptotic (survival) factors and inactivating pro-apoptotic factors [114,117]. Certain proapoptotic mediators, such as the transcription factor forkhead (FOXO), the tau kinase GSK-3 β , and the Bcl2 antagonist BAD proteins, are inactivated by Akt [118,119]. Akt substrates such as mammalian target of rapamycin (mTOR; Ser2448) and decreased levels of cell-cycle inhibitors (p27^{kip1}) are found in AD temporal cortex when compared to controls [120]. Akt downregulates the activities of GSK-3 α and GSK-3 β by phosphorylating the former at Ser21 and the latter at Ser9 [118]. GSK-3 α has been implicated in the production of A β peptide [117,121] while increased GSK-3 β activity has been implicated in tau hyperphosphorylation [122,123,124] and neuronal cell death [124,125]. Phosphorylation/inactivation of GSK-3 β , suppresses GSK-3 β -dependent phosphorylation of tau at residues overphosphorylated in AD and prevents apoptosis of confluent cells. Treatment of cortical neurons with A β 42 monomers increased Ser9 phosphorylation (inhibition) of the Akt substrate, GSK-3 β [97]. Inhibition of GSK-3 β promotes cell survival through a variety of mechanisms including a reduced degradation of β -catenin, which then translocates into the nucleus and activates the transcription of protective genes [126].

4.2.3. Extracellular-signal regulated kinase 1/2 signaling (ERK1 and ERK2)—The mitogen-activated protein kinase (MAPK) family of protein kinases is traditionally viewed as important kinases in transmitting extracellular membrane signals into the nucleus. The 44 kDa ERK1 and 42 kDa ERK2 are members of the MAPK superfamily that specifically respond to A β in brain cells [127]. ERK1 and ERK2 are known to be activated through dual phosphorylation by the MAPK/ERK on threonine and tyrosine in the Thr-Glu-Tyr sequence of the activation loop [128,129]. ERK signaling is critical for memory and tightly regulated by many proteins. ERKs are critical for human learning as revealed by human mental retardation syndromes [130]. They are also known to contribute to molecular information processing in dendrites, to stabilize structural changes in dendritic spines and to interact with scaffolding and structural proteins at the synapse [131]. ERK is an important neuronal marker for activity through activation by cytosolic calcium and depolarization of the membrane [132,133]. On phosphorylation and activation, ERKs phosphorylate other cytoplasmic effectors and are translocated into the nucleus where they phosphorylate transcription factors such as Myc, Fos, Jun, and Elk1 [104,134]. Direct substrates of the ERKs include two members of the RSK family of protein serine-threonine kinases, RSK1 and RSK2. The transcription factor CREB is phosphorylated on serine 133 *in vivo* by RSK2 in NGF-stimulated PC12 cells [135]. The dependence of CREB phosphorylation on activation of the ERK pathway is suggested by inhibition of A β -induced phosphorylation of CREB by piceatannol and the MEK inhibitor PD98059. Other kinases, such as protein kinase A (PKA) or Ca²⁺/calmodulin-dependent protein kinases [CAM kinases; 136] may also contribute to phosphorylation of cyclic AMP response element (CRE)-binding protein

(CREB) in response to A β but the complete inhibition of CREB phosphorylation by PD98059 suggests that the ERK pathway is the main signaling pathway elicited by A β leading to transcriptional activation through CREB [137]. These data provide a mechanism by which A β alters gene expression through the transcription factor CREB [137], possibly resulting in a ceiling of activation that limits further formation of new memories.

4.2.3. Protein kinase C—PKC is part of a multigene family of serine-threonine kinases central to many signal transduction pathways [138] with a prominent role in memory [139]. It is likely that A β -induced increases in cytosolic Ca²⁺ signals are transmitted to PKC for PKC-mediated transcriptional activation. In addition, PKC activates ERK by interacting with Ras or Raf-1 [140] to initiate CREB phosphorylation. While PKC levels decline in AD [141], their activation restores K⁺ channel function in cells from AD patients [142]. In addition, activation of PKC directly or indirectly enhances the α -processing cleavage of APP [143]. The activation of PKC has also been shown to prevent A β toxicity in rat primary hippocampal neurons [144].

4.2.4. Calcium signaling—Synaptic activity is required for neurons to survive [145] by entry of appropriate amounts of Ca²⁺ through synaptic NMDA receptors and other Ca²⁺ channels [146]. The process implicates key protein effectors, such as CaMKs, MAPK/ERKs, and CREB. Properly controlled homeostasis of calcium signaling not only supports normal brain physiology but also maintains neuronal integrity and long-term cell survival. Ca²⁺ signaling pathways can suppress apoptosis and promote survival through two mechanistically distinct processes. One process involves the PI3K/AKT signaling pathway which promotes survival [147]. The other pathway requires the generation of calcium transients in the cell nucleus which offers long-lasting neuroprotection [146,148]. Malfunctioning of calcium signaling to the cell nucleus may lead to neurodegeneration and neuronal cell death [149].

Dysregulation of intracellular calcium signaling has been implicated in the pathogenesis of Alzheimer's disease [150]. A β is known to act through multiple targets [151] including Ca²⁺ channels and various receptors in membranes. Synthetic A β binds to the calcium permeable nAChRs with high affinity [152]. A β 42 administered in the low picomolar range activates nAChRs at presynaptic nerve endings of synaptosomes [83,153]. Under normal conditions, activation of nAChRs is necessary for the A β -induced increase in synaptic plasticity and memory [23]. However, it remains to be determined whether these effects are mediated by a direct physical interaction of the peptide with the nAChR. In addition, A β enhances transmitter release by transient increase of glutamate release from the presynaptic terminal that results from brief periods of high frequency stimulation with Ca²⁺ buildup within the terminal that triggers mechanisms of short-term synaptic plasticity [154].

A β directly interacts with Ca²⁺ channels such as voltage-dependent calcium channels (VDCC) and TRP cation channels (TRPC) to produce a transient increase in Ca²⁺ necessary for synaptic plasticity and neuronal survival. A β interacts directly with the recombinant L-type Ca²⁺ channel (α 1C) subunit to increase Ca²⁺ channel protein at the cell membrane and hence increased Ca²⁺ conductance [80]. Within the TRPC subfamily, TRPC3 and 6 have been shown to protect cerebellar granule neurons against serum deprivation-induced cell death in cultures and promote neuronal survival in rat brain [155]. A neuronal survival mechanism of A β may also involve altered expression of K⁺ channels [80]. In cerebellar granule neurons, 24-h pre-incubation with 1 μ M unaggregated A β protein resulted in a 60% increase in the 'A'-type component of K⁺ current possibly reflecting Ca²⁺-mediated gene expression [156]. A full understanding of these signal transduction pathways of Ca²⁺ may lead to refined pharmacological strategies that minimize deadly effects of Ca²⁺ entry and optimize its growth- and survival-promoting properties.

4.2.5. Transcriptional activation—CREB is one of the best characterized stimulus-induced transcription factors that activate transcription of target genes in response to a diverse array of stimuli, including neuronal activity, a variety of protein kinases such as protein kinase A (PKA), MAPK/ERKs, pp90 ribosomal S6 kinase (RSK), and Ca²⁺/calmodulin-dependent protein kinases (CaMKs). These kinases all phosphorylate CREB at a particular residue, serine 133 (Ser133) which is required for CREB-mediated transcription [157]. In contrast to CaMKs, ERKs cannot directly phosphorylate CREB. Two related RSKs and mitogen- and stress-activated protein kinases (MSKs) transmit the signal from activated ERKs to CREB [158]. CREB has been shown to be involved in certain types of hippocampal LTP as well as long-term memory, neurogenesis and synaptogenesis [159,160]. Transcriptional activation of CREB recruits a multiprotein assembly called a transcriptional co-activator complex. These often include proteins with intrinsic acetyltransferase activity [161]. Among the best characterized transcriptional co-activator proteins is CREB binding protein (CBP) [162]. A role for CBP in memory storage was first demonstrated in a mouse model of Rubinstein-Taybi Syndrome [163]. There is no direct evidence indicating how lower levels of A β might initiate CREB phosphorylation principally by Ca²⁺ signaling and/or through PKA/Atk/ERK pathways. However, exceeding physiological levels of A β could deregulate Ca²⁺ signaling mechanism by excessive accumulation of Ca²⁺ in the cytoplasm and cytoplasmic organelles such as mitochondria. Since hippocampal neuronal calcium is one of the most potent signals in neuronal gene expression [149], A β -induced Ca²⁺ deregulation may lead to compromised synaptic function. Consistence with this hypothesis, AD has been associated with impaired cAMP signaling which may contribute to the pathophysiology of the disease. Levels of the activated (i.e. phosphorylated) form of CREB are reduced in AD compared to that of an age-matched healthy control group [164]. Calcium signaling to the cell nucleus is the key inducer of CREB phosphorylation on its activator site serine 133 [165]. Experiments in aged neurons show altered calcium signaling at the level of either calcium signal generation and/or calcium signal propagation [166]. These studies indicate a critical role of calcium in A β -induced synaptic activity and memory formation by regulating specific signal transduction pathways.

4.3. Cholesterol transport

High cholesterol levels have been linked to overproduction of A β and are a risk factor for AD. One of the physiological functions of A β has been suggested to control cholesterol transport [167]. Prevalence of AD is reduced among people treated with inhibitors of cholesterol biosynthesis, statins [168,169] and animal studies support these results [170]. In vitro and in vivo studies have shown that cholesterol modulates APP processing and affects APP mRNA expression [171]. Another mechanism is the increased binding of A β to ApoE4 over non-E4 alleles. ApoE is a lipid and cholesterol transport protein responsible for the efflux of cholesterol from neurons to form stable complexes both in vitro and in vivo [172]. Allele ApoE4 is a major risk factor in AD [173]. This relationship might promote synaptogenesis, since in vitro studies have demonstrated that cholesterol released by astroglia increases synaptogenesis [174,175] with resulting modulation of spike rates [176]. Together, this evidence indicates that one of the physiological functions of APP might be to control cholesterol movement across neuronal membranes [167].

4.4. Antioxidant

The three histidine residues in A β control the redox activity of iron, indicating that A β is likely to be an important antioxidant. A β 40 at 5 μ M was found to protect primary neuronal cultures from the neurotoxicity of iron [94]. Nakamura et al. [177] found that A β 40 or A β 42 inhibits Fe³⁺ or Cu⁺²-catalyzed ascorbate oxidation and hydroxyl radical generation. Nanomolar concentrations of A β can block neuronal apoptosis following oxidative damage,

which suggests that A β has a protective role against oxidative stress [178] and is essential for neuronal survival [28,94]. Monomeric A β 40 was found to protect neurons cultured in a medium containing 1.5 μ M Fe $^{2+}$ without antioxidant molecules. However, the antioxidant protection of monomeric A β 40 depends on the type of oxidant used. A β 40 inhibits cultured neurone death caused by Cu $^{2+}$, Fe $^{2+}$, and Fe $^{3+}$ but does not protect neurons against H $_2$ O $_2$ -induced damage [94]. In cerebral cortical neuronal cultures, monomeric A β 40 inhibits the reduction of Fe $^{3+}$ induced by vitamin C and the generation of superoxides and prevents lipid peroxidation induced by Fe $^{2+}$ [94]. Moreover, monomeric forms of A β 42 also exhibited antioxidant and neuroprotective effects. However, oligomeric or aggregated A β 40 and A β 42 were devoid of such antioxidant activity and their neuroprotective activity was demolished. Thus, depriving neurons of the protective activity of A β 42 monomers may also be an important factor in neurodegeneration [97]. These findings provide novel insights on a normal antioxidant role of A β and indicate that monomeric A β protects neurons by quenching metal-inducible oxygen radical generation and thereby inhibits neurotoxicity.

5. Effects of A β on dendritic spine plasticity and synaptic function

Synapse loss is the strongest anatomical correlate of the degree of clinical impairment in AD [179]. Loss of dendritic spines at the sites of excitatory synaptic transmission may be the major pathological mechanism in Alzheimer's disease. However, issues regarding the level of A β concentration, type of A β species as well as the mechanisms of its production and actions that lead to synaptic loss remained poorly understood. Continuous overproduction of A β at dendrites or axons acts locally to reduce the number and plasticity of synapses [76,180]. The majority of excitatory synapses in the brain are made on the heads of dendritic spines [181]. Initially, synapse degeneration begins at the level of dendritic spines, the loci of memory-initiating mechanisms [182,183,184]. As seen in AD and transgenic mouse AD models, significant decreases occur in spine density [185,186,187,188], molecules involved in spine signaling [189,190] and control of filamentous actin (F-actin) [191]. In a mouse model for AD, the vicinity of amyloid plaques is characterized by highly dysmorphic neurites and spine turnover [192,193] causing a net loss of spines. These abnormalities in dendritic spines develop even before appearance of clinical symptoms in AD, likely because of cognitive reserve [187]. This phenotype could be caused by A β oligomers, which have been shown to block LTP and directly induce LTD, spine loss and memory loss [50]. Soluble oligomers of A β have a direct synaptotoxic effect at nanomolar concentrations [51]. In hippocampal culture, the soluble A β produced abnormalities in spine composition, shape, and abundance that strongly support the hypothesis that soluble A β initiates toxic mechanisms in AD brains that account for synaptic damage [74]. Continued exposure to A β caused abnormal spine morphology, with induction of long thin spines, loss of spine cytoskeletal protein drebin and a significant decrease in spine density [74]. In a direct investigation of the acute effects of extracellular and intracellular A β 42 peptides on synaptic transmission, Moreno et al. [194] noticed inhibition of synaptic transmission by nanomolar concentrations of intra-axonal oligomeric A β 42, but not oligomeric A β 40 or extracellular oligomeric A β 42. Similar nanomolar levels of A β disrupt hippocampal LTP [60,195]. Importantly, physiological concentrations of A β in extracellular fluids are picomolar [196]. Thus, local dendritic and axonal abnormalities associated with amyloid deposits lead to loss of synapses and the breakage of dendrites and axons in AD [187,193]. As dendritic spines are the major connecting elements of one neuron with another in the brain, changes in spine plasticity would have a detrimental impact on disease pathogenesis and progression. Overall, the accumulation of soluble or fibrillar amyloid deposits in AD causes disruption of synaptic connections on a permanent basis and this likely contributes to the cognitive decline and memory [197]. This decreased synaptic activity leads to the elimination of synapses and loss of network activity [198,199].

The molecular mechanism of spine loss by A β is not clear. Electron microscopic studies demonstrate that oligomeric A β is localized within the synaptic compartment [200] or that it is bound to the extracellular surface of the spine suggesting that oligomeric A β may interact directly at the synapse to cause dysfunction and spine collapse [201]. Soluble A β causes abnormal expression of Arc, a synaptic memory related protein that causes abnormal spine shape and glutamate receptor trafficking [42,202]. A β treatment of cultured hippocampal neurons leads to the inactivation of PKA and persistence of its regulatory subunit PKAII α [203]. Since glutamate treatment reduces phosphorylated CREB phosphorylation and the decrease is reversed by rolipram (a phosphodiesterase inhibitor that raises cAMP and leads to the dissociation of the PKA catalytic and regulatory subunits), a similar mechanism may inhibit LTP by A β . Later studies confirmed the activation of the PKA/CREB pathway in both cultured neurons and murine hippocampal slices after inhibition of LTP by A β [204,205]. Interestingly, the toxicity of micromolar fibrillar A β on cultured neurons correlates with an age-related increase in phosphorylated extracellular signal-regulated kinase (pERK) as well as an age-independent over-activation of pCREB [7]. A β -induced activation of ERK1/2 may reduce mitochondrial respiration and ATP production by decreasing complex I activity and substrate oxidation through complex I [206]. Oligomers can also compromise synaptic function by altering the permeability of neuronal membranes and disrupting ion homeostasis [207,208]. None of these studies of action of A β on protein kinases have identified the proximal target of A β . However, these observations suggest that A β acts directly on the pathways involved in the formation of late LTP. Agents that enhance the cAMP/PKA/CREB-signaling pathway have potential for the treatment of AD [203]. These studies clearly support the emerging view that impaired synaptic function may be more important for the development of AD than neuronal cell death which occurs at later stages of the disease [199]. However, the major question of how abnormal spine dynamics and alterations in spine plasticity contribute to the disease progression in AD is still not very clear. The major challenge to prevent such loss in spine plasticity could prove invaluable for the treatment of neurodegenerative diseases.

6. Molecular targets of A β induced synaptic dysfunction

The search for a mechanism by which A β impairs synaptic plasticity has led to the identification of the cell surface receptors and signaling pathways mediating A β -induced synaptotoxicity. Cell surface interaction sites reported for A β include receptor for Advanced Glycation End products (RAGE) and NMDAR [152,209]. A β has been variously reported to directly affect the activity of NMDAR, possibly by binding to nAChRs, or intracellular mitochondrial cyclophilin D (CypD), mitochondrial A β alcohol dehydrogenase (ABAD) or certain protein kinases. Examination of the evidence for these multiple activities of A β and their affinity constants may distinguish direct binding partners from downstream effectors.

6.1. NMDA receptors

It is well known that excitatory synapses contain AMPA and NMDA ionotropic glutamate receptors as well as metabotropic type glutamate receptors (mGluRs) positioned on dendritic spines [210,211]. A β -induced synaptic dysfunction has been attributed to the synaptic removal of AMPA receptors (AMPA receptors); however, it is unclear how A β induces this loss [212]. Glutamatergic processes are strongly implicated in causing and mediating the symptoms of AD [213]. The clinical use of memantine in treatment of AD provides direct support for the involvement of NMDARs in the cognitive deficits [214]. Memantine acts as a low-to-moderate affinity open channel uncompetitive inhibitor of NMDARs at therapeutic concentrations [215,216]. A β promotes glutamatergic excitotoxicity and potently disrupts glutamatergic synapses and plasticity providing an explanation for the cognitive deficits in AD [217]. A β alters the glutamatergic transmission system by inducing marked reductions in levels of AMPA and NMDA receptors at the neuronal plasma membrane [74,75,218,219].

At physiological levels, A β selectively enhances NMDAR-mediated currents and synaptic transmission [220,221]. While increased A β application promotes endocytosis of NMDARs in cortical neurons and produces a rapid and persistent depression of NMDA-evoked currents in cortical neurons [75]. This reduction of NMDAR-dependent currents is thought to result from A β -mediated activation of nAChRs [75]. A β can also promote increased Ca²⁺ influx and elevate the levels of potentially toxic reactive oxygen species in an NMDAR-dependent manner [221,222]. In general, brief periods of high synaptic activity open NMDARs, leading to a long-lasting increase in postsynaptic AMPAR number, spine growth and LTP of synaptic transmission [223,224,225]. Snyder et al. [75] reported a pathway by which A β reduces glutamatergic transmission and NMDAR – dependent LTP. The application of A β 42 to cultured cortical neurons promoted endocytosis of NMDARs, effectively reducing the density of NMDARs at synapses. At higher concentrations, A β is known to enhance activation of NMDARs [226] and cause NMDAR agonist-induced delayed cognitive dysfunction [227]. It is apparent that excessive or inappropriate activation of NMDAR can block LTP [228,229].

Alternatively, low levels of synaptic stimulation can activate NMDARs to produce NMDA-dependent or mGluR-dependent LTD. These two forms of LTD can induce removal of postsynaptic AMPARs and loss of spines [230,231,232,233]. Interestingly, sublethal NMDAR activation increased the production and secretion of A β [65,234]. However, some of the conceptually and biomedically most important questions that have arisen from these novel insights concern the molecular mechanisms by which A β and the glutamatergic transmission system cooperate at the synapse to synergistically regulate and control synaptic transmission. Is excitotoxicity that results from excessive glutamate receptor activation the main trigger that increases the secretion of A β during synaptic transmission or is A β solely a pathologic product directing cell demise? It is possible that the increased A β production causes an increase in NMDA activation and increased NMDA activation in turn increases the A β production within limits. This process would have a negative impact on synaptic function if there were a homeostatic balance between NMDA activation and A β production. However, excess NMDA activation or excess A β generation both are harmful for synaptic plasticity that could lead to the cognitive impairment in AD.

Treatment with A β oligomers also causes reduction of post-synaptic density-95 (PSD-95), an adaptor protein that plays a critical role in synaptic plasticity and in the stabilization of AMPAR and NMDAR at synapses [218]. Dysregulation of NMDAR function causes excessive neuronal Ca²⁺ influx, oxidative stress [222], and inhibition of the Wnt/ β -catenin signaling pathway [235,236].

6.2. Nicotinic acetylcholine receptors (nAChRs)

Activation of the neuronal pentameric nAChRs is involved in diverse brain functions including synaptic plasticity and memory [237,238] and enhances transmitter release in several brain regions including the hippocampus [239], the spinal cord dorsal horn [240], the olfactory bulb, and the amygdala [241]. The increase of synaptic plasticity by A β requires activation of nAChRs [242]. Because activation of nAChRs is necessary for the A β -induced increase of synaptic plasticity and memory under normal conditions, A β may modify glutamate release with a mechanism dependent upon activation of nAChRs [83]. However, several reports of the effect of A β 42 on nAChRs are conflicting. Some studies have reported that A β 42 activates nAChRs [243,244], while others indicate that A β 42 inhibits nAChRs [245,246]. For example, physiological levels of A β can activate while toxic levels inhibit presynaptic nAChR and evoke changes in presynaptic Ca²⁺ levels in rat hippocampus and neocortex [244]. Interestingly, picomolar concentrations of A β 42 were effective in activating nAChRs while higher levels of A β produced inhibitory action. The disparity may depend on the nanomolar A β 1–42 inhibition of nicotine-induced Ca²⁺ responses while

picomolar A β 42 directly evokes sustained increases in presynaptic Ca²⁺ via nAChRs [244]. A β 42 binds to the nAChR with picomolar affinity [247]. This binding can modulate presynaptic, glutamate-mediated synaptic transmission or glutamate release, suggesting that A β 42-dependent cholinergic modulation activates signal transduction mechanisms that ultimately result in synaptic transmission and memory consolidation [248]. However, it remains to be determined whether these effects are mediated by a direct physical interaction of the A β peptide with the nAChRs. Immunohistochemical studies on human sporadic AD brains show that A β 42 and nAChR, are both present in neuritic plaques and co-localize in individual cortical neurons suggesting that A β could be tightly associated with nAChR [247]. Alternatively, A β might be responsible for regulation of nAChR function through strong binding with membrane lipids [249]. Picomolar or higher A β 42 acting through nAChRs, can elicit ERK MAPK activation in hippocampal cultures [245,250] possibly triggered by Ca²⁺ influx [243,251]. ERK is known to regulate transcription factors such as CREB and Elk-1 by phosphorylation [140], which help initiate transcription of memory-associated genes that contain their respective regulatory elements [252]. Therefore, over-activation of nAChRs and excessive Ca²⁺ influx or dysregulation of Ca²⁺ homeostasis provide a molecular mechanism for the cholinergic dysfunction that is a hallmark of AD [253,254].

6.3. Mitochondrial cyclophilin D

Mitochondria serve as direct targets of neuronal toxicity in which A β associates with the outer mitochondrial membrane, inter-membrane space, inner mitochondrial membrane, and the matrix [255]. Progressive accumulation of A β in cortical mitochondria in AD patients and also in brains from transgenic AD type mouse models suggests a role for mitochondrial A β in the pathogenesis or development of the disease. Once inside the mitochondria, A β is able to interact with a number of targets, including the mitochondrial proteins ABAD and cyclophilin-D (CypD) [256]. Opening the mitochondrial permeability transition pore (MPTP) to depolarize mitochondria and release cytochrome C may be central to mitochondrial and neuronal malfunction in AD patients [257]. CypD, an integral part of the MPTP, whose opening leads to cell death, interacts with A β peptide within the mitochondria of AD patients and a Tg mouse model of AD [258]. MPTP causes mitochondrial swelling, outer membrane rupture, release of cell death mediators and enhances production of reactive oxygen species (ROS). Computer simulation studies show that A β interacts with both ANT and CypD [259]. CypD/A β interaction causes an oxidative stress and increased MPTP opening that triggers neurodegeneration [258]. CypD-deficient cortical mitochondria are resistant to A β - and Ca²⁺-induced mitochondrial swelling and MPTP opening [257]. Adenine nucleotide translocase (ANT) is a transport protein for ADP and ATP and component of MPTP that binds directly to CypD [260]. This interaction may facilitate its anchoring in the inner membrane and disturbance of the mitochondrial membrane potential, mitochondrial swelling and cell death [259]. Interestingly, the MPTP also requires the participation of members of the Bcl-2 family proteins but a clear understanding of the interaction of A β with CypD together with both proapoptotic or antiapoptotic Bcl-2 family proteins in AD has not been made. The ability of CypD to protect neurons from A β - and oxidative stress-induced cell death and its role in improvement of synaptic and cognitive functions has been suggested to provide a new therapeutic approach for the treatment of conditions associated with AD. Together these studies provide new mechanisms for A β targets that link the MPTP to synaptic stress and the neurodegeneration seen in AD.

6.4. Mitochondrial A β -binding alcohol dehydrogenase (ABAD)

ABAD is a member of the short chain dehydrogenase reductase family in mitochondria that binds A β [261]. Binding of A β to ABAD distorts the enzyme's structure, rendering it inactive. Binding also promotes mitochondrial generation of free radicals. In neurons,

ABAD is predominately localized to mitochondria. Upon binding ABAD, A β triggers events leading to neuronal apoptosis through a mitochondrial pathway [262,263]. Mitochondrial A β levels in the 3xTg-AD mouse increase significantly at 9 months and temporally correlate with increased levels of A β binding to ABAD [264]. Interestingly, mitochondrial ABAD is upregulated in neurons from AD patients [263]. The ABAD-A β complex has been hypothesized to induce oxidant stress and mitochondrial dysfunction [265]. Increased expression of ABAD exacerbates A β -mediated mitochondrial and neuronal stress [263,266]. A β binding to ABAD causes free radical production and neuronal apoptosis [267]. Neurons cultured from transgenic mice with targeted overexpression of a mutant form of amyloid precursor protein and ABAD (Tg mAPP/ABAD) displayed spontaneous generation of hydrogen peroxide and superoxide anion, and decreased ATP, as well as subsequent release of cytochrome c from mitochondria and induction of caspase-3-like activity followed by DNA fragmentation and loss of cell viability [266]. In addition, cytochrome c oxidase (COX) activity was selectively decreased in neurons cultured from mAPP/ABAD mice. In vivo, mAPP/ABAD mice displayed reduced levels of brain ATP and COX activity, diminished glucose utilization, as well as electrophysiological abnormalities in hippocampal slices compared with mAPP mice [266]. ABAD-A β binding and enhanced generation of oxidants in brain mitochondria of transgenic mice results in exaggerating neuronal stress and impaired learning and memory [268]. Analysis of hippocampal slices from mAPP/ABAD mice were shown to display diminished LTP compared with other genotypes [266]. However, positive effects of low levels of A β have not been studied. Similar to CypD, the ABAD-A β interaction may also represent a novel treatment target against AD. Other intra-mitochondrial targets of A β remain to be discovered. These data suggest that mitochondrial ABAD, ordinarily a contributor to metabolic homeostasis, has the capacity to become a pathogenic factor in an A β rich environment.

7. Redox and phosphorylative energetic exhaustion: A β -induced mitochondrial and synaptic dysfunction

Increasing evidence indicates that the mitochondrial dysfunction is an important early factor in the development of AD-like pathology [264]. Mitochondria are known to accumulate in synapses [269,270] and mitochondrial trafficking to synapses is dynamic and regulated by synaptic activity [271]. However, increasing evidence indicates that accumulation of A β in mitochondria occurs before extracellular amyloid deposition and increases with age. A β has been found in the mitochondria of both AD brain and transgenic mouse models of AD overexpressing A β [255,256,272,]. A β has been detected in mitochondria from postmortem brain specimens of AD patients [255] and also in isolated mitochondria from the cerebral cortex of APP transgenic mice [272]. APP and its derivatives, monomeric and oligomeric forms of A β , interact with mitochondrial membranes [263,272,273,274] or mitochondrial matrix protein ABAD [264] leading to mitochondrial dysfunction. Accordingly, A β is linked to the mitochondrial malfunction observed in the Alzheimer's disease brain and mouse models of AD [255,272]. Substantial evidence indicates that mitochondria serve as direct targets for A β protein mediated neuronal toxicity.

Studies of postmortem brains from AD patients and transgenic mouse models of AD suggest that mitochondria are involved in oxidative damage induced by A β early in AD progression [reviewed in reference 275]. Lower levels of ROS are required for synaptic signaling with ROS acting as messenger molecules in the process of LTP [276]. However, high levels of ROS have been implicated as damaging toxic molecules in the age-related impairments of LTP [276,277]. Our previous work shows that ROS levels increase with age of neurons in parallel with an age-related decline in transmembrane potential [278]. As mitochondrial transmembrane potential is a driving force for cellular production of ATP, its decline in neurons will have a long term effect in many important energy driven reactions. Increased

oxidative stress, coupled with dysregulation of calcium homeostasis and resulting apoptosis of vulnerable neuronal populations, are proposed to underlie the loss of synaptic activity and associated cognitive decline [275]. From these deficiencies emerges the concept of synaptic energy exhaustion in AD, both phosphorylative (ATP) and redox (NAD[P]H) energies. Our previous work shows that hippocampal NAD(P)H and glutathione (GSH) decline with age in association with increased susceptibility to glutamate toxicity in neurons of old-age [279]. Thus, an age-related decline in neuronal reducing currency (NAD[P]H) and reducing buffer (GSH) will surely promote oxidative stress and excess ROS. It is noteworthy that in the early stages of AD, there is already a reduction in the number of mitochondria [280] and the activities of tricarboxylic acid cycle enzymes [281] and cytochrome C oxidase [282]. However, how ROS are produced at the synapse in response to A β oligomers is not fully known. Excessive ROS are locally generated in response to synaptic A β oligomer binding [222]. This ROS formation can be totally blocked by the mitochondrial uncoupler, 2,4-dinitrophenol which suggests a central role of mitochondria in A β -induced oxidative stress [222]. Many studies suggest the possible involvement of oxidative stress and calcium dysfunction in A β toxicity [283,284]. Experiments with *Caenorhabditis elegans* containing inducible A β 42 indicate that oxidative stress can precede fibrillogenesis [285]. These reports were strengthened by findings that A β toxicity at the synapse is dependent on the presence of a functional mitochondrial electron transport chain which is a principal site of ROS formation as well as a major target for their deleterious effects [286].

The question as to why brain synaptic ROS levels increase with age is uncertain, but may involve lack of use [287] followed by acute overstimulation of excitatory NMDARs that leads to excessive ROS, related to excess Ca²⁺ entry into mitochondria [288]. Dysregulation of NMDAR function induced by A β binding to neuronal synapses may lead to synaptic mitochondrial dysfunction and excessive ROS formation [222,289]. Memory mechanisms might be directly compromised by elevated ROS, which could explain the connection between AD and oxidative stress [222]. The increase in oxidative damage exhibited by synaptic mitochondria will damage synapses, affect neurotransmission and might be ultimately responsible for cognitive decline in AD patients. Taken together these studies provide convincing evidence for the concept that mitochondria have a pivotal role in A β -induced synaptic dysfunction and neuronal stress. Improved function of mitochondria is an effective way of reducing effects of aging and may inhibit neuronal cell death in AD [287].

8. Conclusions

The present review highlights some important physiological roles for A β in the CNS during normal function and AD pathogenesis. Given the important role that A β plays in various activities at the synapse, A β should not be regarded merely as a toxic factor that requires eradication to avoid dementia. There is enough evidence to suggest an essential activity-dependent role of A β in modulation of synaptic activity and neuronal survival. However, dissociation of the synaptic effects of aging from A β remains to be investigated. Several extracellular and intracellular synaptic receptors are important targets of A β oligomers. Alterations in receptor characteristics and synaptic dysfunction are crucial to the early memory deficit and cognitive decline in AD. Alterations in synaptic plasticity, mitochondrial function and neurotransmission by A β can affect activity-dependent signaling and gene expression, resulting in the disintegration of neural networks and, ultimately, the failure of neural functions [290]. These ongoing discussions provide new insights into strategies for development of AD therapy that not only reduce the amount of A β but also inhibit A β aggregation and restore mitochondrial and redox energy for synaptic function.

Acknowledgments

This work was supported in part by NIH grants R01 AG032431, R56 AG013435 and the Kenneth Stark Endowed Chair for Alzheimer Research to GJB.

Abbreviations

ABAD	mitochondrial A β -binding alcohol dehydrogenase
Ca²⁺CAM	Calcium calmodulin
CAMKIV	Calcium/calmodulin-dependent protein kinase type IV
CREB	Cyclic AMP response element (CRE)-binding protein
CBP	CREB binding protein
CypD	cyclophilin D
ERK	extracellular signal-regulated kinase
Fas	FS7-associated cell surface antigen
FOXO	forkhead transcription factor
GSK	glycogen synthase kinase
Grb2	growth factor receptor-bound protein 2
IR	insulin receptor
IRS-1	insulin receptor substrate-1
IRS-2	insulin receptor substrate-2
LTP	Long term potentiation
mitDH	mitochondrial dehydrogenases
MEK	MAPK kinase, MAPK, mitogen-activated protein kinase
MPTP	membrane permeability transition pore opening
mTOR	mammalian target of rapamycin
α7-nAChR	α 7-nicotinic acetylcholine receptor
NMDAR	N-methyl-D-aspartic acid receptor
PDK1	3-Phosphoinositide-dependent kinase 1
PI3K	phosphoinositide 3' kinase
PKB/Akt	protein kinases B
ROS	reactive oxygen species
VDCC	voltage-dependent calcium channels
TRPC	TRP cation channel

References

1. Greenough WT, West RW, DeVoogd TJ. Subsynaptic plate perforations: changes with age and experience in the rat. *Science*. 1978; 202:1096–1098. [PubMed: 715459]
2. Agnati LF, Benfenati F, Solfrini V, Biagini G, Fuxe K, Guidolin D, Carani C, Zini I. Brain aging and neuronal plasticity. *Ann N Y Acad Sci*. 1992; 673:180–186. [PubMed: 1336643]

3. Martin LJ, Pardo CA, Cork LC, Price DL. Synaptic pathology and glial responses to neuronal injury precede the formation of senile plaques and amyloid deposits in the aging cerebral cortex. *Am J Pathol.* 1994; 145:1358–1381. [PubMed: 7992840]
4. Brewer GJ. Neuronal plasticity and stressor toxicity during aging. *Exp Gerontol.* 2000; 35:1165–1183. [PubMed: 11113600]
5. Buell SJ, Coleman PD. Dendritic growth in the aged human brain and failure of growth in senile dementia. *Science.* 1979; 206:854–856. [PubMed: 493989]
6. Mesulam MM. Neuroplasticity failure in Alzheimer's disease: bridging the gap between plaques and tangles. *Neuron.* 1999; 24:521–529. [PubMed: 10595506]
7. Brewer GJ, Torricelli JR, Lindsey AL, Kunz EZ, Neuman A, Fisher DR, Joseph JA. Age-related toxicity of amyloid-beta associated with increased pERK and pCREB in primary hippocampal neurons: reversal by blueberry extract. *J Nutr Biochem.* 2010 Epub ahead of print. 10.1016/j.nutbio.2009.08.005
8. Hedden T, Gabrieli JD. Insights into the ageing mind: a view from cognitive neuroscience. *Nat Rev Neurosci.* 2004; 5:87–96. [PubMed: 14735112]
9. Fotuhi M, Hachinski V, Whitehouse PJ. Changing perspectives regarding late-life dementia. *Nat Rev Neurol.* 2009; 5:649–658. [PubMed: 19918254]
10. Lanahan A, Lyford G, Stevenson GS, Worley PF, Barnes CA. Selective alteration of long-term potentiation-induced transcriptional response in hippocampus of aged, memory-impaired rats. *J Neurosci.* 1997; 17:2876–2885. [PubMed: 9092609]
11. Pasternack JM, Palmert MR, Usiak M, Wang R, Zurcher-Neely H, Gonzalez-De Whitt PA, Fairbanks MB, Cheung T, Blades D, Heinrikson RL. Alzheimer's disease and control brain contain soluble derivatives of the amyloid protein precursor that end within the beta amyloid protein region. *Biochemistry.* 1992; 31:10936–10940. [PubMed: 1420205]
12. Geula C, Wu CK, Saroff D, Lorenzo A, Yuan M, Yankner BA. Aging renders the brain vulnerable to amyloid beta-protein neurotoxicity. *Nat Med.* 1998; 4:827–831. [PubMed: 9662375]
13. Brewer GJ. Age-related toxicity to lactate, glutamate, and beta-amyloid in cultured adult neurons. *Neurobiol Aging.* 1998; 19:561–568. [PubMed: 10192215]
14. LaFerla FM, Oddo S. Alzheimer's disease: Abeta, tau and synaptic dysfunction. *Trends Mol Med.* 2005; 11:170–176. [PubMed: 15823755]
15. Lee HG, Zhu X, Castellani RJ, Nunomura A, Perry G, Smith MA. Amyloid-beta in Alzheimer disease: the null versus the alternate hypotheses. *J Pharmacol Exp Ther.* 2007; 321:823–829. [PubMed: 17229880]
16. Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, Selkoe D. Soluble oligomers of amyloid Beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron.* 2009; 62:788–801. [PubMed: 19555648]
17. Selkoe DJ. Amyloid protein and Alzheimer's disease. *Sci Am.* 1991; 265:68–71. [PubMed: 1785042]
18. Hardy JA, Higgins GA. Alzheimer's disease: the amyloid cascade hypothesis. *Science.* 1992; 256:184–185. [PubMed: 1566067]
19. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL. Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A.* 1998; 95:6448–6453. [PubMed: 9600986]
20. Hsia AY, Masliah E, McConlogue L, Yu GQ, Tatsuno G, Hu K, Kholodenko D, Malenka RC, Nicoll RA, Mucke L. Plaque-independent disruption of neural circuits in Alzheimer's disease mouse models. *Proc Natl Acad Sci U S A.* 1999; 96:3228–3233. [PubMed: 10077666]
21. Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science.* 2002; 297:353–356. [PubMed: 12130773]
22. Walsh DM, Selkoe DJ. Deciphering the molecular basis of memory failure in Alzheimer's disease. *Neuron.* 2004; 44:181–193. [PubMed: 15450169]
23. Cirrito JR, Yamada KA, Finn MB, Sloviter RS, Bales KR, May PC, Schoepp DD, Paul SM, Mennerick S, Holtzman DM. Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron.* 2005; 48:913–922. [PubMed: 16364896]

24. Ting JT, Kelley BG, Lambert TJ, Cook DG, Sullivan JM. Amyloid precursor protein overexpression depresses excitatory transmission through both presynaptic and postsynaptic mechanisms. *Proc Natl Acad Sci U S A*. 2007; 104:353–358. [PubMed: 17185415]
25. Priller C, Bauer T, Mitteregger G, Krebs B, Kretschmar HA, Herms J. Synapse formation and function is modulated by the amyloid precursor protein. *J Neurosci*. 2006; 26:7212–7221. [PubMed: 16822978]
26. Cirrito JR, Kang JE, Lee J, Stewart FR, Verges DK, Silverio LM, Bu G, Mennerick S, Holtzman DM. Endocytosis is required for synaptic activity-dependent release of amyloid-beta in vivo. *Neuron*. 2008; 58:42–51. [PubMed: 18400162]
27. Ramsden M, Henderson Z, Pearson HA. Modulation of Ca²⁺ channel currents in primary cultures of rat cortical neurones by amyloid beta protein (1–40) is dependent on solubility status. *Brain Res*. 2002; 956:254–261. [PubMed: 12445693]
28. Plant LD, Boyle JP, Smith IF, Peers C, Pearson HA. The production of amyloid beta peptide is a critical requirement for the viability of central neurons. *J Neurosci*. 2003; 23:5531–5535. [PubMed: 12843253]
29. Kamenetz F, Tomita T, Hsieh H, Seabrook G, Borchelt D, Iwatsubo T, Sisodia S, Malinow R. APP processing and synaptic function. *Neuron*. 2003; 37:925–937. [PubMed: 12670422]
30. Reinhard C, Hebert SS, De Strooper B. The amyloid-beta precursor protein: integrating structure with biological function. *EMBO J*. 2005; 24:3996–4006. [PubMed: 16252002]
31. Selkoe DJ, Yamazaki T, Citron M, Podlisny MB, Koo EH, Teplow DB, Haass C. The role of APP processing and trafficking pathways in the formation of amyloid beta-protein. *Ann N Y Acad Sci*. 1996; 777:57–64. [PubMed: 8624127]
32. Gralle M, Ferreira ST. Structure and functions of the human amyloid precursor protein: the whole is more than the sum of its parts. *Prog Neurobiol*. 2007; 82:11–32. [PubMed: 17428603]
33. Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science*. 2002; 297:353–356. [PubMed: 12130773]
34. Glenner GG, Wong CW. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem Biophys Res Commun*. 1984; 120:885–890. [PubMed: 6375662]
35. Greenfield JP, Tsai J, Gouras GK, Hai B, Thinakaran G, Checler F, Sisodia SS, Greengard P, Xu H. Endoplasmic reticulum and trans-Golgi network generate distinct populations of Alzheimer beta-amyloid peptides. *Proc Natl Acad Sci U S A*. 1999; 96:742–747. [PubMed: 9892704]
36. Jarrett JT, Berger EP, Lansbury PT Jr. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry*. 1993; 32:693–697.
37. Leissring MA. The AbetaCs of Abeta-cleaving proteases. *J Biol Chem*. 2008; 283:29645–29649. [PubMed: 18723506]
38. Farris W, Mansourian S, Chang Y, Lindsley L, Eckman EA, Frosch MP, Eckman CB, Tanzi RE, Selkoe DJ, Guenette S. Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain in vivo. *Proc Natl Acad Sci U S A*. 2003; 100:4162–4167. [PubMed: 12634421]
39. Iwata N, Tsubuki S, Takaki Y, Shirohane K, Lu B, Gerard NP, Gerard C, Hama E, Lee HJ, Saido TC. Metabolic regulation of brain Abeta by neprilysin. *Science*. 2001; 292:1550–1552. [PubMed: 11375493]
40. Lazarov O, Lee M, Peterson DA, Sisodia SS. Evidence that synaptically released beta-amyloid accumulates as extracellular deposits in the hippocampus of transgenic mice. *J Neurosci*. 2002; 22:9785–9793. [PubMed: 12427834]
41. Selkoe DJ. Alzheimer's disease is a synaptic failure. *Science*. 2002; 298:789–791. [PubMed: 12399581]
42. Lacor PN, Buniel MC, Chang L, Fernandez SJ, Gong Y, Viola KL, Lambert MP, Velasco PT, Bigio EH, Finch CE, Krafft GA, Klein WL. Synaptic targeting by Alzheimer's-related amyloid beta oligomers. *J Neurosci*. 2004; 24:10191–10200. [PubMed: 15537891]
43. Barghorn S, Nimmrich V, Striebinger A, Krantz C, Keller P, Janson B, Bahr M, Schmidt M, Bitner RS, Harlan J, Barlow E, Ebert U, Hillen H. Globular amyloid beta-peptide oligomer - a

- homogenous and stable neuropathological protein in Alzheimer's disease. *J Neurochem.* 2005; 95:834–847. [PubMed: 16135089]
44. Podlisny MB, Ostaszewski BL, Squazzo SL, Koo EH, Rydell RE, Teplow DB, Selkoe DJ. Aggregation of secreted amyloid beta-protein into sodium dodecyl sulfate-stable oligomers in cell culture. *J Biol Chem.* 1995; 270:9564–9570. [PubMed: 7721886]
 45. Enya M, Morishima-Kawashima M, Yoshimura M, Shinkai Y, Kusui K, Khan K, Games D, Schenk D, Sugihara S, Yamaguchi H, Ihara Y. Appearance of sodium dodecyl sulfate-stable amyloid beta-protein (A β) dimer in the cortex during aging. *Am J Pathol.* 1999; 154:271–279. [PubMed: 9916941]
 46. Kawarabayashi T, Shoji M, Younkin LH, Wen-Lang L, Dickson DW, Murakami T, Matsubara E, Abe K, Ashe KH, Younkin SG. Dimeric amyloid beta protein rapidly accumulates in lipid rafts followed by apolipoprotein E and phosphorylated tau accumulation in the Tg2576 mouse model of Alzheimer's disease. *J Neurosci.* 2004; 24:3801–3809. [PubMed: 15084661]
 47. Lesne S, Koh MT, Kotilinek L, Kaye R, Glabe CG, Yang A, Gallagher M, Ashe KH. A specific amyloid-beta protein assembly in the brain impairs memory. *Nature.* 2006; 440:352–357. [PubMed: 16541076]
 48. Roher AE, Chaney MO, Kuo YM, Webster SD, Stine WB, Haverkamp LJ, Woods AS, Cotter RJ, Tuohy JM, Krafft GA, Bonnell BS, Emmerling MR. Morphology and toxicity of A β (1-42) dimer derived from neuritic and vascular amyloid deposits of Alzheimer's disease. *J Biol Chem.* 1996; 271:20631–20635. [PubMed: 8702810]
 49. Powers ET, Morimoto RI, Dillin A, Kelly JW, Balch WE. Biological and chemical approaches to diseases of proteostasis deficiency. *Annu Rev Biochem.* 2009; 78:959–991. [PubMed: 19298183]
 50. Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, Brett FM, Farrell MA, Rowan MJ, Lemere CA, Regan CM, Walsh DM, Sabatini BL, Selkoe DJ. Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med.* 2008; 14:837–842. [PubMed: 18568035]
 51. Walsh DM, Selkoe DJ. A beta oligomers - a decade of discovery. *J Neurochem.* 2007; 101:1172–1184. [PubMed: 17286590]
 52. Shankar GM, Walsh DM. Alzheimer's disease: synaptic dysfunction and A β . *Mol Neurodegener.* 2009; 4:48–62. [PubMed: 19930651]
 53. Bitan G, Kirkitadze MD, Lomakin A, Vollers SS, Benedek GB, Teplow DB. Amyloid beta -protein (A β) assembly: A β 40 and A β 42 oligomerize through distinct pathways. *Proc Natl Acad Sci U S A.* 2003; 100:330–335. [PubMed: 12506200]
 54. Haass C, Schlossmacher MG, Hung AY, Vigo-Pelfrey C, Mellon A, Ostaszewski BL, Lieberburg I, Koo EH, Schenk D, Teplow DB. Amyloid beta-peptide is produced by cultured cells during normal metabolism. *Nature.* 1992; 359:322–325. [PubMed: 1383826]
 55. Seubert P, Vigo-Pelfrey C, Esch F, Lee M, Dovey H, Davis D, Sinha S, Schlossmacher M, Whaley J, Swindlehurst C. Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature.* 1992; 359:325–327. [PubMed: 1406936]
 56. Shoji M, Golde TE, Ghiso J, Cheung TT, Estus S, Shaffer LM, Cai XD, McKay DM, Tintner R, Frangione B. Production of the Alzheimer amyloid beta protein by normal proteolytic processing. *Science.* 1992; 258:126–129. [PubMed: 1439760]
 57. Piccini A, Russo C, Gliozzi A, Relini A, Vitali A, Borghi R, Giliberto L, Armirotti A, D'Arrigo C, Bachi A, Cattaneo A, Canale C, Torrassa S, Saido TC, Markesbery W, Gambetti P, Tabaton M. Beta-amyloid is different in normal aging and in Alzheimer disease. *J Biol Chem.* 2005; 280:34186–34192. [PubMed: 16103127]
 58. Stefani M, Dobson CM. Protein aggregation and aggregate toxicity: new insights into protein folding, misfolding diseases and biological evolution. *J Mol Med.* 2003; 81:678–699. [PubMed: 12942175]
 59. Scheuner D, Eckman C, Jensen M, Song X, Citron M, Suzuki N, Bird TD, Hardy J, Hutton M, Kukull W, Larson E, Levy-Lahad E, Viitanen M, Peskind E, Poorkaj P, Schellenberg G, Tanzi R, Wasco W, Lannfelt L, Selkoe D, Younkin S. Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nat Med.* 1996; 2:864–870. [PubMed: 8705854]

60. Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ. Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature*. 2002; 416:535–539. [PubMed: 11932745]
61. Kawarabayashi T, Younkin LH, Saido TC, Shoji M, Ashe KH, Younkin SG. Age-dependent changes in brain, CSF, and plasma amyloid (beta) protein in the Tg2576 transgenic mouse model of Alzheimer's disease. *J Neurosci*. 2001; 21:372–381. [PubMed: 11160418]
62. Selkoe DJ. Clearing the brain's amyloid cobwebs. *Neuron*. 2001; 32:177–180. [PubMed: 11683988]
63. Caccamo A, Oddo S, Billings LM, Green KN, Martinez-Coria H, Fisher A, LaFerla FM. M1 receptors play a central role in modulating AD-like pathology in transgenic mice. *Neuron*. 2006; 49:671–682. [PubMed: 16504943]
64. Nitsch RM, Deng M, Tennis M, Schoenfeld D, Growdon JH. The selective muscarinic M1 agonist AF102B decreases levels of total Abeta in cerebrospinal fluid of patients with Alzheimer's disease. *Ann Neurol*. 2000; 48:913–918. [PubMed: 11117548]
65. Lesne S, Ali C, Gabriel C, Croci N, MacKenzie ET, Glabe CG, Plotkine M, Marchand-Verrecchia C, Vivien D, Buisson A. NMDA receptor activation inhibits alpha-secretase and promotes neuronal amyloid-beta production. *J Neurosci*. 2005; 25:9367–9377. [PubMed: 16221845]
66. Hooper NM, Turner AJ. The search for alpha-secretase and its potential as a therapeutic approach to Alzheimer's disease. *Curr Med Chem*. 2002; 9:1107–1119. [PubMed: 12052175]
67. Lichtenthaler SF, Haass C. Amyloid at the cutting edge: activation of alpha-secretase prevents amyloidogenesis in an Alzheimer disease mouse model. *J Clin Invest*. 2004; 113:1384–1387. [PubMed: 15146234]
68. Carroll RC, Beattie EC, von Zastrow M, Malenka RC. Role of AMPA receptor endocytosis in synaptic plasticity. *Nat Rev Neurosci*. 2001; 2:315–324. [PubMed: 11331915]
69. Malinow R, Mainen ZF, Hayashi Y. LTP mechanisms: from silence to four-lane traffic. *Curr Opin Neurobiol*. 2000; 10:352–357. [PubMed: 10851179]
70. Koo EH, Squazzo SL, Selkoe DJ, Koo CH. Trafficking of cell-surface amyloid beta-protein precursor. I. Secretion, endocytosis and recycling as detected by labeled monoclonal antibody. *J Cell Sci*. 1996; 109:991–998. [PubMed: 8743946]
71. Carey RM, Balcz BA, Lopez-Coviella I, Slack BE. Inhibition of dynamin-dependent endocytosis increases shedding of the amyloid precursor protein ectodomain and reduces generation of amyloid beta protein. *BMC Cell Biol*. 6:30–40. [PubMed: 16095541]
72. Buckner RL, Snyder AZ, Shannon BJ, LaRossa G, Sachs R, Fotenos AF, Sheline YI, Klunk WE, Mathis CA, Morris JC, Mintun MA. Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *J Neurosci*. 2005; 25:7709–7717. [PubMed: 16120771]
73. Deshpande A, Kawai H, Metherate R, Glabe CG, Busciglio J. A role for synaptic zinc in activity-dependent Abeta oligomer formation and accumulation at excitatory synapses. *J Neurosci*. 2009; 29:4004–4015. [PubMed: 19339596]
74. Lacor PN, Buniel MC, Furlow PW, Clemente AS, Velasco PT, Wood M, Viola KL, Klein WL. Abeta oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci*. 2007; 27:796–807. [PubMed: 17251419]
75. Snyder EM, Nong Y, Almeida CG, Paul S, Moran T, Choi EY, Nairn AC, Salter MW, Lombroso PJ, Gouras GK, Greengard P. Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci*. 2005; 8:1051–1058. [PubMed: 16025111]
76. Shankar GM, Bloodgood BL, Townsend M, Walsh DM, Selkoe DJ, Sabatini BL. Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J Neurosci*. 2007; 27:2866–2875. [PubMed: 17360908]
77. Zoghbi HY, Orr HT. Glutamine repeats and neurodegeneration. *Annu Rev Neurosci*. 2000; 23:217–247. [PubMed: 10845064]
78. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL. Diffusible, nonfibrillar

- ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A*. 1998; 95:6448–6453. [PubMed: 9600986]
79. Walsh DM, Hartley DM, Kusumoto Y, Fezoui Y, Condron MM, Lomakin A, Benedek GB, Selkoe DJ, Teplow DB. Amyloid beta-protein fibrillogenesis. Structure and biological activity of protofibrillar intermediates. *J Biol Chem*. 1999; 274:25945–25952. [PubMed: 10464339]
 80. Pearson HA, Peers C. Physiological roles for amyloid beta peptides. *J Physiol*. 2006; 575:5–10. [PubMed: 16809372]
 81. Sheng M, Kim MJ. Postsynaptic signaling and plasticity mechanisms. *Science*. 2002; 298:776–780. [PubMed: 12399578]
 82. Rodrigues SM, Schafe GE, LeDoux JE. Molecular mechanisms underlying emotional learning and memory in the lateral amygdala. *Neuron*. 2004; 44:75–91. [PubMed: 15450161]
 83. Puzzo D, Privitera L, Leznik E, Fa M, Staniszewski A, Palmeri A, Arancio O. Picomolar amyloid-beta positively modulates synaptic plasticity and memory in hippocampus. *J Neurosci*. 2008; 28:14537–14545. [PubMed: 19118188]
 84. Cirrito JR, May PC, O'Dell MA, Taylor JW, Parsadanian M, Cramer JW, Audia JE, Nissen JS, Bales KR, Paul SM, DeMattos RB, Holtzman DM. In vivo assessment of brain interstitial fluid with microdialysis reveals plaque-associated changes in amyloid-beta metabolism and half-life. *J Neurosci*. 2003; 23:8844–8853. [PubMed: 14523085]
 85. Dawson GR, Seabrook GR, Zheng H, Smith DW, Graham S, O'Dowd G, Bowery BJ, Boyce S, Trumbauer ME, Chen HY, Van der Ploeg LH, Sirinathsinghji DJ. Age-related cognitive deficits, impaired long-term potentiation and reduction in synaptic marker density in mice lacking the beta-amyloid precursor protein. *Neuroscience*. 1999; 90:1–13. [PubMed: 10188929]
 86. Laird FM, Cai H, Savonenko AV, Farah MH, He K, Melnikova T, Wen H, Chiang HC, Xu G, Koliatsos VE, Borchelt DR, Price DL, Lee HK, Wong PC. BACE1, a major determinant of selective vulnerability of the brain to amyloid-beta amyloidogenesis, is essential for cognitive, emotional, and synaptic functions. *J Neurosci*. 2005; 25:11693–11709. [PubMed: 16354928]
 87. Saura CA, Choi SY, Beglopoulos V, Malkani S, Zhang D, Shankaranarayana Rao BS, Chattarji S, Kelleher RJ III, Kandel ER, Duff K, Kirkwood A, Shen J. Loss of presenilin function causes impairments of memory and synaptic plasticity followed by age-dependent neurodegeneration. *Neuron*. 2004; 42:23–36. [PubMed: 15066262]
 88. Rozmahel R, Huang J, Chen F, Liang Y, Nguyen V, Ikeda M, Levesque G, Yu G, Nishimura M, Mathews P, Schmidt SD, Mercken M, Bergeron C, Westaway D, George-Hyslop P. Normal brain development in PS1 hypomorphic mice with markedly reduced gamma-secretase cleavage of betaAPP. *Neurobiol Aging*. 2002; 23:187–194. [PubMed: 11804702]
 89. Phinney AL, Drisaldi B, Schmidt SD, Lugowski S, Coronado V, Liang Y, Horne P, Yang J, Sekoulidis J, Coomaraswamy J, Chishti MA, Cox DW, Mathews PM, Nixon RA, Carlson GA, George-Hyslop P, Westaway D. In vivo reduction of amyloid-beta by a mutant copper transporter. *Proc Natl Acad Sci U S A*. 2003; 100:14193–14198. [PubMed: 14617772]
 90. Pawlik M, Sastre M, Calero M, Mathews PM, Schmidt SD, Nixon RA, Levy E. Overexpression of human cystatin C in transgenic mice does not affect levels of endogenous brain amyloid Beta Peptide. *J Mol Neurosci*. 2004; 22:13–18. [PubMed: 14742906]
 91. Mastrangelo P, Mathews PM, Chishti MA, Schmidt SD, Gu Y, Yang J, Mazzella MJ, Coomaraswamy J, Horne P, Strome B, Pelly H, Levesque G, Ebeling C, Jiang Y, Nixon RA, Rozmahel R, Fraser PE, George-Hyslop P, Carlson GA, Westaway D. Dissociated phenotypes in presenilin transgenic mice define functionally distinct gamma-secretases. *Proc Natl Acad Sci U S A*. 2005; 102:8972–8977. [PubMed: 15951428]
 92. Wasling P, Daborg J, Riebe I, Andersson M, Portelius E, Blennow K, Hanse E, Zetterberg H. Synaptic retrogenesis and amyloid-beta in Alzheimer's disease. *J Alzheimers Dis*. 2009; 16:1–14. [PubMed: 19158416]
 93. Lopez-Toledano MA, Shelanski ML. Neurogenic effect of beta-amyloid peptide in the development of neural stem cells. *J Neurosci*. 2004; 24:5439–5444. [PubMed: 15190117]
 94. Zou K, Gong JS, Yanagisawa K, Michikawa M. A novel function of monomeric amyloid beta-protein serving as an antioxidant molecule against metal-induced oxidative damage. *J Neurosci*. 2002; 22:4833–4841. [PubMed: 12077180]

95. Whitson JS, Selkoe DJ, Cotman CW. Amyloid beta protein enhances the survival of hippocampal neurons in vitro. *Science*. 1989; 243:1488–1490. [PubMed: 2928783]
96. Kim J, Onstead L, Randle S, Price R, Smithson L, Zwizinski C, Dickson DW, Golde T, McGowan E. Abeta40 inhibits amyloid deposition in vivo. *J Neurosci*. 2007; 27:627–633. [PubMed: 17234594]
97. Giuffrida ML, Caraci F, Pignataro B, Cataldo S, De Bona P, Bruno V, Molinaro G, Pappalardo G, Messina A, Palmigiano A, Garozzo D, Nicoletti F, Rizzarelli E, Copani A. Beta-amyloid monomers are neuroprotective. *J Neurosci*. 2009; 29:10582–10587. [PubMed: 19710311]
98. Yankner BA, Duffy LK, Kirschner DA. Neurotrophic and neurotoxic effects of amyloid beta protein: reversal by tachykinin neuropeptides. *Science*. 1990; 250:279–282. [PubMed: 2218531]
99. Lee CC, Huang CC, Wu MY, Hsu KS. Insulin stimulates postsynaptic density-95 protein translation via the phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling pathway. *J Biol Chem*. 2005; 280:18543–18550. [PubMed: 15755733]
100. Ding Q, Vaynman S, Akhavan M, Ying Z, Gomez-Pinilla. Insulin-like growth factor I interfaces with brain-derived neurotrophic factor-mediated synaptic plasticity to modulate aspects of exercise-induced cognitive function. *Neuroscience*. 2006; 140:823–833. [PubMed: 16650607]
101. Sugano T, Yanagita T, Yokoo H, Satoh S, Kobayashi H, Wada A. Enhancement of insulin-induced PI3K/Akt/GSK-3beta and ERK signaling by neuronal nicotinic receptor/PKC-alpha/ERK pathway: up-regulation of IRS-1/-2 mRNA and protein in adrenal chromaffin cells. *J Neurochem*. 2006; 98:20–33. [PubMed: 16805793]
102. Hoyer S. Causes and consequences of disturbances of cerebral glucose metabolism in sporadic Alzheimer disease: therapeutic implications. *Adv Exp Med Biol*. 2004; 541:135–152. [PubMed: 14977212]
103. Bondy CA, Cheng CM. Signaling by insulin-like growth factor 1 in brain. *Eur J Pharmacol*. 2004; 490:25–31. [PubMed: 15094071]
104. Lee HK, Kumar P, Fu Q, Rosen KM, Querfurth HW. The insulin/Akt signaling pathway is targeted by intracellular beta-amyloid. *Mol Biol Cell*. 2009; 20:1533–1544. [PubMed: 19144826]
105. Horwood JM, Dufour F, Laroche S, Davis S. Signalling mechanisms mediated by the phosphoinositide 3-kinase/Akt cascade in synaptic plasticity and memory in the rat. *Eur J Neurosci*. 2006; 23:3375–3384. [PubMed: 16820027]
106. Park CR, Seeley RJ, Craft S, Woods SC. Intracerebroventricular insulin enhances memory in a passive-avoidance task. *Physiol Behav*. 2000; 68:509–514. [PubMed: 10713291]
107. Craft S, Watson GS. Insulin and neurodegenerative disease: shared and specific mechanisms. *Lancet Neurol*. 2004; 3:169–178. [PubMed: 14980532]
108. Wei W, Wang X, Kusiak JW. Signaling events in amyloid beta-peptide-induced neuronal death and insulin-like growth factor I protection. *J Biol Chem*. 2002; 277:17649–17656. [PubMed: 11882652]
109. Xie L, Helmerhorst E, Taddei K, Plewright B, Van Bronswijk W, Martins R. Alzheimer's beta-amyloid peptides compete for insulin binding to the insulin receptor. *J Neurosci*. 2002; 22:RC221. [PubMed: 12006603]
110. Townsend M, Mehta T, Selkoe DJ. Soluble Abeta inhibits specific signal transduction cascades common to the insulin receptor pathway. *J Biol Chem*. 2007; 282:33305–33312. [PubMed: 17855343]
111. Markus A, Zhong J, Snider WD. Raf and akt mediate distinct aspects of sensory axon growth. *Neuron*. 2002; 35:65–76. [PubMed: 12123609]
112. Izzo E, Martin-Fardon R, Koob GF, Weiss F, Sanna PP. Neural plasticity and addiction: PI3-kinase and cocaine behavioral sensitization. *Nat Neurosci*. 2002; 5:1263–1264. [PubMed: 12436114]
113. Chan TO, Rittenhouse SE, Tsichlis PN. AKT/PKB and other D3 phosphoinositide-regulated kinases: kinase activation by phosphoinositide-dependent phosphorylation. *Annu Rev Biochem*. 1999; 68:965–1014. [PubMed: 10872470]
114. Brunet A, Datta SR, Greenberg ME. Transcription-dependent and -independent control of neuronal survival by the PI3K-Akt signaling pathway. *Curr Opin Neurobiol*. 2001; 11:297–305. [PubMed: 11399427]

115. Asano T, Yao Y, Shin S, McCubrey J, Abbruzzese JL, Reddy SA. Insulin receptor substrate is a mediator of phosphoinositide 3-kinase activation in quiescent pancreatic cancer cells. *Cancer Res.* 2005; 65:9164–9168. [PubMed: 16230374]
116. Franke TF, Kaplan DR, Cantley LC. PI3K: downstream AKTion blocks apoptosis. *Cell.* 1997; 88:435–437. [PubMed: 9038334]
117. Baki L, Shioi J, Wen P, Shao Z, Schwarzman A, Gama-Sosa M, Neve R, Robakis NK. PS1 activates PI3K thus inhibiting GSK-3 activity and tau overphosphorylation: effects of FAD mutations. *EMBO J.* 2004; 23:2586–2596. [PubMed: 15192701]
118. Cross DA, Alessi DR, Cohen P, Andjelkovich M, Hemmings BA. Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B. *Nature.* 1995; 378:785–789. [PubMed: 8524413]
119. Zheng WH, Kar S, Quirion R. Insulin-like growth factor-1-induced phosphorylation of the forkhead family transcription factor FKHRL1 is mediated by Akt kinase in PC12 cells. *J Biol Chem.* 2000; 275:39152–39158. [PubMed: 10995739]
120. Griffin RJ, Moloney A, Kelliher M, Johnston JA, Ravid R, Dockery P, O'Connor R, O'Neill C. Activation of Akt/PKB, increased phosphorylation of Akt substrates and loss and altered distribution of Akt and PTEN are features of Alzheimer's disease pathology. *J Neurochem.* 2005; 93:105–117. [PubMed: 15773910]
121. Phiel CJ, Wilson CA, Lee VM, Klein PS. GSK-3alpha regulates production of Alzheimer's disease amyloid-beta peptides. *Nature.* 2003; 423:435–439. [PubMed: 12761548]
122. Hong M, Chen DC, Klein PS, Lee VM. Lithium reduces tau phosphorylation by inhibition of glycogen synthase kinase-3. *J Biol Chem.* 1997; 272:25326–25332. [PubMed: 9312151]
123. Pei JJ, Braak E, Braak H, Grundke-Iqbal I, Iqbal K, Winblad B, Cowburn RF. Distribution of active glycogen synthase kinase 3beta (GSK-3beta) in brains staged for Alzheimer disease neurofibrillary changes. *J Neuropathol Exp Neurol.* 1999; 58:1010–1019. [PubMed: 10499443]
124. Lucas JJ, Hernandez F, Gomez-Ramos P, Moran MA, Hen R, Avila J. Decreased nuclear beta-catenin, tau hyperphosphorylation and neurodegeneration in GSK-3beta conditional transgenic mice. *EMBO J.* 2001; 20:27–39. [PubMed: 11226152]
125. Hetman M, Cavanaugh JE, Kimelman D, Xia Z. Role of glycogen synthase kinase-3beta in neuronal apoptosis induced by trophic withdrawal. *J Neurosci.* 2000; 20:2567–2574. [PubMed: 10729337]
126. Willert K, Nusse R. Beta-catenin: a key mediator of Wnt signaling. *Curr Opin Genet Dev.* 1998; 8:95–102. [PubMed: 9529612]
127. Abe K, Hisatomi R, Misawa M. Amyloid beta peptide specifically promotes phosphorylation and nuclear translocation of the extracellular signal-regulated kinase in cultured rat cortical astrocytes. *J Pharmacol Sci.* 2003; 93:272–278. [PubMed: 14646244]
128. Lewis TS, Shapiro PS, Ahn NG. Signal transduction through MAP kinase cascades. *Adv Cancer Res.* 1998; 74:49–139. [PubMed: 9561267]
129. Seger R, Krebs EG. The MAPK signaling cascade. *FASEB J.* 1995; 9:726–735. [PubMed: 7601337]
130. Costa RM, Federov NB, Kogan JH, Murphy GG, Stern J, Ohno M, Kucherlapati R, Jacks T, Silva AJ. Mechanism for the learning deficits in a mouse model of neurofibromatosis type 1. *Nature.* 2002; 415:526–530. [PubMed: 11793011]
131. Sweatt JD. Mitogen-activated protein kinases in synaptic plasticity and memory. *Curr Opin Neurobiol.* 2004; 14:311–317. [PubMed: 15194111]
132. Agell N, Bachs O, Rocamora N, Villalonga P. Modulation of the Ras/Raf/MEK/ERK pathway by Ca⁽²⁺⁾, and calmodulin. *Cell Signal.* 2002; 14:649–654. [PubMed: 12020764]
133. Lever IJ, Pezet S, McMahon SB, Malcangio M. The signaling components of sensory fiber transmission involved in the activation of ERK MAP kinase in the mouse dorsal horn. *Mol Cell Neurosci.* 2003; 24:259–270. [PubMed: 14572451]
134. Blenis J. Signal transduction via the MAP kinases: Proceed at your own RSK. *Proc Natl Acad Sci U S A.* 1993; 90:5889–5892. [PubMed: 8392180]

135. Xing J, Ginty DD, Greenberg ME. Coupling of the RAS-MAPK pathway to gene activation by RSK2, a growth factor-regulated CREB kinase. *Science*. 1996; 273:959–963. [PubMed: 8688081]
136. Sheng M, Thompson MA, Greenberg ME. CREB: a Ca⁽²⁺⁾-regulated transcription factor phosphorylated by calmodulin-dependent kinases. *Science*. 1991; 252:1427–1430. [PubMed: 1646483]
137. McDonald DR, Bamberger ME, Combs CK, Landreth GE. beta-Amyloid fibrils activate parallel mitogen-activated protein kinase pathways in microglia and THP1 monocytes. *J Neurosci*. 1998; 18:4451–4460. [PubMed: 9614222]
138. Nishizuka Y. Intracellular signaling by hydrolysis of phospholipids and activation of protein kinase C. *Science*. 1992; 258:607–614. [PubMed: 1411571]
139. Alkon DL, Rasmussen H. A spatial-temporal model of cell activation. *Science*. 1988; 239:998–1005. [PubMed: 2830669]
140. Sweatt JD. The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory. *J Neurochem*. 2001; 76:1–10. [PubMed: 11145972]
141. Cole G, Dobkins KR, Hansen LA, Terry RD, Saitoh T. Decreased levels of protein kinase C in Alzheimer brain. *Brain Res*. 1988; 452:165–174. [PubMed: 3165303]
142. Bhagavan S, Ibarreta D, Ma D, Kozikowski AP, Etcheberrigaray R. Restoration of TEA-induced calcium responses in fibroblasts from Alzheimer's disease patients by a PKC activator. *Neurobiol Dis*. 1998; 5:177–187. [PubMed: 9848089]
143. Etcheberrigaray R, Tan M, Dewachter I, Kuiperi C, Van da I, Wera S, Qiao L, Bank B, Nelson TJ, Kozikowski AP, Van Leuven F, Alkon DL. Therapeutic effects of PKC activators in Alzheimer's disease transgenic mice. *Proc Natl Acad Sci U S A*. 2004; 101:11141–11146. [PubMed: 15263077]
144. Garrido JL, Godoy JA, Alvarez A, Bronfman M, Inestrosa NC. Protein kinase C inhibits amyloid beta peptide neurotoxicity by acting on members of the Wnt pathway. *FASEB J*. 2002; 16:1982–1984. [PubMed: 12397090]
145. Hardingham GE, Bading H. The Yin and Yang of NMDA receptor signalling. *Trends Neurosci*. 2003; 26:81–89. [PubMed: 12536131]
146. Hardingham GE, Fukunaga Y, Bading H. Extrasynaptic NMDARs oppose synaptic NMDARs by triggering CREB shut-off and cell death pathways. *Nat Neurosci*. 2002; 5:405–414. [PubMed: 11953750]
147. Papadia S, Stevenson P, Hardingham NR, Bading H, Hardingham GE. Nuclear Ca²⁺ and the cAMP response element-binding protein family mediate a late phase of activity-dependent neuroprotection. *J Neurosci*. 2005; 25:4279–4287. [PubMed: 15858054]
148. Mao Z, Bonni A, Xia F, Nadal-Vicens M, Greenberg ME. Neuronal activity-dependent cell survival mediated by transcription factor MEF2. *Science*. 1999; 286:785–790. [PubMed: 10531066]
149. Zhang SJ, Zou M, Lu L, Lau D, Ditzel DA, Delucinge-Vivier C, Aso Y, Descombes P, Bading H. Nuclear calcium signaling controls expression of a large gene pool: identification of a gene program for acquired neuroprotection induced by synaptic activity. *PLoS Genet*. 2009; 5:e1000604. [PubMed: 19680447]
150. LaFerla FM. Calcium dyshomeostasis and intracellular signalling in Alzheimer's disease. *Nat Rev Neurosci*. 2002; 3:862–872. [PubMed: 12415294]
151. Small DH, Mok SS, Bornstein JC. Alzheimer's disease and Abeta toxicity: from top to bottom. *Nat Rev Neurosci*. 2001; 2:595–598. [PubMed: 11484003]
152. Wang HY, Lee DH, Davis CB, Shank RP. Amyloid peptide Abeta(1–42) binds selectively and with picomolar affinity to alpha7 nicotinic acetylcholine receptors. *J Neurochem*. 2000; 75:1155–1161. [PubMed: 10936198]
153. Dougherty JJ, Wu J, Nichols RA. Beta-amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *J Neurosci*. 2003; 23:6740–6747. [PubMed: 12890766]
154. Zucker RS, Regehr WG. Short-term synaptic plasticity. *Annu Rev Physiol*. 2002; 64:355–405. [PubMed: 11826273]

155. Jia Y, Zhou J, Tai Y, Wang Y. TRPC channels promote cerebellar granule neuron survival. *Nat Neurosci.* 2007; 10:559–567. [PubMed: 17396124]
156. Ramsden M, Plant LD, Webster NJ, Vaughan PF, Henderson Z, Pearson HA. Differential effects of unaggregated and aggregated amyloid beta protein (1–40) on K(+) channel currents in primary cultures of rat cerebellar granule and cortical neurones. *J Neurochem.* 2001; 79:699–712. [PubMed: 11701773]
157. Shaywitz AJ, Greenberg ME. CREB: a stimulus-induced transcription factor activated by a diverse array of extracellular signals. *Annu Rev Biochem.* 1999; 68:821–861. [PubMed: 10872467]
158. Wiggin GR, Soloaga A, Foster JM, Murray-Tait V, Cohen P, Arthur JS. MSK1 and MSK2 are required for the mitogen- and stress-induced phosphorylation of CREB and ATF1 in fibroblasts. *Mol Cell Biol.* 2002; 22:2871–2881. [PubMed: 11909979]
159. Gass P, Riva MA. CREB, neurogenesis and depression. *Bioessays.* 2007; 29:957–961. [PubMed: 17876779]
160. Wood MA, Kaplan MP, Park A, Blanchard EJ, Oliveira AM, Lombardi TL, Abel T. Transgenic mice expressing a truncated form of CREB-binding protein (CBP) exhibit deficits in hippocampal synaptic plasticity and memory storage. *Learn Mem.* 2005; 12:111–119. [PubMed: 15805310]
161. Bayly R, Chuen L, Currie RA, Hyndman BD, Casselman R, Blobel GA, LeBrun DP. E2A-PBX1 interacts directly with the KIX domain of CBP/p300 in the induction of proliferation in primary hematopoietic cells. *J Biol Chem.* 2004; 279:55362–55371. [PubMed: 15507449]
162. Chrivia JC, Kwok RP, Lamb N, Hagiwara M, Montminy MR, Goodman RH. Phosphorylated CREB binds specifically to the nuclear protein CBP. *Nature.* 1993; 365:855–859. [PubMed: 8413673]
163. Oike Y, Hata A, Mamiya T, Kaname T, Noda Y, Suzuki M, Yasue H, Nabeshima T, Araki K, Yamamura K. Truncated CBP protein leads to classical Rubinstein-Taybi syndrome phenotypes in mice: implications for a dominant-negative mechanism. *Hum Mol Genet.* 1999; 8:387–396. [PubMed: 9949198]
164. Yamamoto-Sasaki M, Ozawa H, Saito T, Rosler M, Riederer P. Impaired phosphorylation of cyclic AMP response element binding protein in the hippocampus of dementia of the Alzheimer type. *Brain Res.* 1999; 824:300–303. [PubMed: 10196463]
165. Hardingham GE, Arnold FJ, Bading H. Nuclear calcium signaling controls CREB-mediated gene expression triggered by synaptic activity. *Nat Neurosci.* 2001; 4:261–267. [PubMed: 11224542]
166. Toescu EC, Verkhatsky A, Landfield PW. Ca²⁺ regulation and gene expression in normal brain aging. *Trends Neurosci.* 2004; 27:614–620. [PubMed: 15374673]
167. Yao ZX, Papadopoulos V. Function of beta-amyloid in cholesterol transport: a lead to neurotoxicity. *FASEB J.* 2002; 16:1677–1679. [PubMed: 12206998]
168. Jick H, Zornberg GL, Jick SS, Seshadri S, Drachman DA. Statins and the risk of dementia. *Lancet.* 2000; 356:1627–1631. [PubMed: 11089820]
169. Wolozin B, Kellman W, Ruosseau P, Celesia GG, Siegel G. Decreased prevalence of Alzheimer disease associated with 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors. *Arch Neurol.* 2000; 57:1439–1443. [PubMed: 11030795]
170. Fassbender K, Simons M, Bergmann C, Stroick M, Lutjohann D, Keller P, Runz H, Kuhl S, Bertsch T, von Bergmann K, Hennerici M, Beyreuther K, Hartmann T. Simvastatin strongly reduces levels of Alzheimer's disease beta-amyloid peptides Abeta 42 and Abeta 40 in vitro and in vivo. *Proc Natl Acad Sci U S A.* 2001; 98:5856–5861. [PubMed: 11296263]
171. Bodovitz S, Klein WL. Cholesterol modulates alpha-secretase cleavage of amyloid precursor protein. *J Biol Chem.* 1996; 271:4436–4440. [PubMed: 8626795]
172. Mahley RW. Apolipoprotein E: cholesterol transport protein with expanding role in cell biology. *Science.* 1988; 240:622–630. [PubMed: 3283935]
173. Hofman A, Ott A, Breteler MM, Bots ML, Slooter AJ, van Harskamp F, van Duijn CN, Van Broeckhoven C, Grobbee DE. Atherosclerosis, apolipoprotein E, and prevalence of dementia and Alzheimer's disease in the Rotterdam Study. *Lancet.* 1997; 349:151–154. [PubMed: 9111537]

174. Pfrieger FW, Barres BA. Synaptic efficacy enhanced by glial cells in vitro. *Science*. 1997; 277:1684–1687. [PubMed: 9287225]
175. Goritz C, Mauch DH, Pfrieger FW. Multiple mechanisms mediate cholesterol-induced synaptogenesis in a CNS neuron. *Mol Cell Neurosci*. 2005; 29:190–201. [PubMed: 15911344]
176. Brewer GJ, Boehler MD, Jones TT, Wheeler BC. NbActiv4 medium improvement to Neurobasal/B27 increases neuron synapse densities and network spike rates on multielectrode arrays. *J Neurosci Methods*. 2008; 170:181–187. [PubMed: 18308400]
177. Nakamura M, Shishido N, Nunomura A, Smith MA, Perry G, Hayashi Y, Nakayama K, Hayashi T. Three histidine residues of amyloid-beta peptide control the redox activity of copper and iron. *Biochemistry*. 2007; 46:12737–12743. [PubMed: 17929832]
178. Curtain CC, Ali F, Volitakis I, Cherny RA, Norton RS, Beyreuther K, Barrow CJ, Masters CL, Bush AI, Barnham KJ. Alzheimer's disease amyloid-beta binds copper and zinc to generate an allosterically ordered membrane-penetrating structure containing superoxide dismutase-like subunits. *J Biol Chem*. 2001; 276:20466–20473. [PubMed: 11274207]
179. Terry RD, Masliah E, Salmon DP, Butters N, DeTeresa R, Hill R, Hansen LA, Katzman R. Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. *Ann Neurol*. 1991; 30:572–580. [PubMed: 1789684]
180. Wei W, Nguyen LN, Kessels HW, Hagiwara H, Sisodia S, Malinow R. Amyloid beta from axons and dendrites reduces local spine number and plasticity. *Nat Neurosci*. 2010; 13:190–196. [PubMed: 20037574]
181. Yuste R, Bonhoeffer T. Genesis of dendritic spines: insights from ultrastructural and imaging studies. *Nat Rev Neurosci*. 2004; 5:24–34. [PubMed: 14708001]
182. Harris KM, Kater SB. Dendritic spines: cellular specializations imparting both stability and flexibility to synaptic function. *Annu Rev Neurosci*. 1994; 17:341–371. [PubMed: 8210179]
183. Carlisle HJ, Kennedy MB. Spine architecture and synaptic plasticity. *Trends Neurosci*. 2005; 28:182–187. [PubMed: 15808352]
184. Segal M. Dendritic spines and long-term plasticity. *Nat Rev Neurosci*. 2005; 6:277–284. [PubMed: 15803159]
185. Ferrer I, Gullotta F. Down's syndrome and Alzheimer's disease: dendritic spine counts in the hippocampus. *Acta Neuropathol*. 1990; 79:680–685. [PubMed: 2141748]
186. Moolman DL, Vitolo OV, Vonsattel JP, Shelanski ML. Dendrite and dendritic spine alterations in Alzheimer models. *J Neurocytol*. 2004; 33:377–387. [PubMed: 15475691]
187. Spires TL, Meyer-Luehmann M, Stern EA, McLean PJ, Skoch J, Nguyen PT, Bacskai BJ, Hyman BT. Dendritic spine abnormalities in amyloid precursor protein transgenic mice demonstrated by gene transfer and intravital multiphoton microscopy. *J Neurosci*. 2005; 25:7278–7287. [PubMed: 16079410]
188. Jacobsen JS, Wu CC, Redwine JM, Comery TA, Arias R, Bowlby M, Martone R, Morrison JH, Pangalos MN, Reinhart PH, Bloom FE. Early-onset behavioral and synaptic deficits in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A*. 2006; 103:5161–5166. [PubMed: 16549764]
189. Sze C, Bi H, Kleinschmidt-DeMasters BK, Filley CM, Martin LJ. N-Methyl-D-aspartate receptor subunit proteins and their phosphorylation status are altered selectively in Alzheimer's disease. *J Neurol Sci*. 2001; 182:151–159. [PubMed: 11137521]
190. Mishizen-Eberz AJ, Rissman RA, Carter TL, Ikonovic MD, Wolfe BB, Armstrong DM. Biochemical and molecular studies of NMDA receptor subunits NR1/2A/2B in hippocampal subregions throughout progression of Alzheimer's disease pathology. *Neurobiol Dis*. 2004; 15:80–92. [PubMed: 14751773]
191. Harigaya Y, Shoji M, Shirao T, Hirai S. Disappearance of actin-binding protein, drebrin, from hippocampal synapses in Alzheimer's disease. *J Neurosci Res*. 1996; 43:87–92. [PubMed: 8838578]
192. Meyer-Luehmann M, Spires-Jones TL, Prada C, Garcia-Alloza M, de Calignon A, Rozkalne A, Koenigsknecht-Talboo J, Holtzman DM, Bacskai BJ, Hyman BT. Rapid appearance and local toxicity of amyloid-beta plaques in a mouse model of Alzheimer's disease. *Nature*. 2008; 451:720–724. [PubMed: 18256671]

193. Tsai J, Grutzendler J, Duff K, Gan WB. Fibrillar amyloid deposition leads to local synaptic abnormalities and breakage of neuronal branches. *Nat Neurosci.* 2004; 7:1181–1183. [PubMed: 15475950]
194. Moreno H, Yu E, Pigo G, Hernandez AI, Kim N, Moreira JE, Sugimori M, Llinas RR. Synaptic transmission block by presynaptic injection of oligomeric amyloid beta. *Proc Natl Acad Sci U S A.* 2009; 106:5901–5906. [PubMed: 19304802]
195. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL. Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A.* 1998; 95:6448–6453. [PubMed: 9600986]
196. Selkoe DJ. Soluble oligomers of the amyloid beta-protein impair synaptic plasticity and behavior. *Behav Brain Res.* 2008; 192:106–113. [PubMed: 18359102]
197. Bhatt DH, Zhang S, Gan WB. Dendritic spine dynamics. *Annu Rev Physiol.* 2009; 71:261–282. [PubMed: 19575680]
198. Hua JY, Smith SJ. Neural activity and the dynamics of central nervous system development. *Nat Neurosci.* 2004; 7:327–332. [PubMed: 15048120]
199. Varghese K, Molnar P, Das M, Bhargava N, Lambert S, Kindy MS, Hickman JJ. A new target for amyloid beta toxicity validated by standard and high-throughput electrophysiology. *PLoS One.* 2010; 5:e8643. [PubMed: 20062810]
200. Takahashi RH, Almeida CG, Kearney PF, Yu F, Lin MT, Milner TA, Gouras GK. Oligomerization of Alzheimer's beta-amyloid within processes and synapses of cultured neurons and brain. *J Neurosci.* 2004; 24:3592–3599. [PubMed: 15071107]
201. Koffie RM, Meyer-Luehmann M, Hashimoto T, Adams KW, Mielke ML, Garcia-Alloza M, Micheva KD, Smith SJ, Kim ML, Lee VM, Hyman BT, Spires-Jones TL. Oligomeric amyloid beta associates with postsynaptic densities and correlates with excitatory synapse loss near senile plaques. *Proc Natl Acad Sci U S A.* 2009; 106:4012–4017. [PubMed: 19228947]
202. Mokin M, Lindahl JS, Keifer J. Immediate-early gene-encoded protein Arc is associated with synaptic delivery of GluR4-containing AMPA receptors during in vitro classical conditioning. *J Neurophysiol.* 2006; 95:215–224. [PubMed: 16339507]
203. Vitolo OV, Sant'Angelo A, Costanzo V, Battaglia F, Arancio O, Shelanski M. Amyloid beta - peptide inhibition of the PKA/CREB pathway and long-term potentiation: reversibility by drugs that enhance cAMP signaling. *Proc Natl Acad Sci U S A.* 2002; 99:13217–13221. [PubMed: 12244210]
204. Nimchinsky EA, Sabatini BL, Svoboda K. Structure and function of dendritic spines. *Annu Rev Physiol.* 2002; 64:313–353. [PubMed: 11826272]
205. Smith DL, Pozueta J, Gong B, Arancio O, Shelanski M. Reversal of long-term dendritic spine alterations in Alzheimer disease models. *Proc Natl Acad Sci U S A.* 2009; 106:16877–16882. [PubMed: 19805389]
206. Nowak G, Clifton GL, Godwin ML, Bakajsova D. Activation of ERK1/2 pathway mediates oxidant-induced decreases in mitochondrial function in renal cells. *Am J Physiol Renal Physiol.* 2006; 291:F840–F855. [PubMed: 16705147]
207. Kaye R, Sokolov Y, Edmonds B, McIntire TM, Milton SC, Hall JE, Glabe CG. Permeabilization of lipid bilayers is a common conformation-dependent activity of soluble amyloid oligomers in protein misfolding diseases. *J Biol Chem.* 2004; 279:46363–46366. [PubMed: 15385542]
208. Glabe CG, Kaye R. Common structure and toxic function of amyloid oligomers implies a common mechanism of pathogenesis. *Neurology.* 2006; 66:S74–S78. [PubMed: 16432151]
209. Snow AD, Kinsella MG, Parks E, Sekiguchi RT, Miller JD, Kimata K, Wight TN. Differential binding of vascular cell-derived proteoglycans (perlecan, biglycan, decorin, and versican) to the beta-amyloid protein of Alzheimer's disease. *Arch Biochem Biophys.* 1995; 320:84–95. [PubMed: 7793988]
210. Baude A, Nusser Z, Roberts JD, Mulvihill E, McIlhinney RA, Somogyi P. The metabotropic glutamate receptor (mGluR1 alpha) is concentrated at perisynaptic membrane of neuronal subpopulations as detected by immunogold reaction. *Neuron.* 1993; 11:771–787. [PubMed: 8104433]

211. Takumi Y, Ramirez-Leon V, Laake P, Rinvik E, Ottersen OP. Different modes of expression of AMPA and NMDA receptors in hippocampal synapses. *Nat Neurosci.* 1999; 2:618–624. [PubMed: 10409387]
212. Gu Z, Liu W, Yan Z. β -Amyloid impairs AMPA receptor trafficking and function by reducing Ca^{2+} /calmodulin-dependent protein kinase II synaptic distribution. *J Biol Chem.* 2009; 284:10639–10649. [PubMed: 19240035]
213. Hu NW, Klyubin I, Anwyl R, Rowan MJ. GluN2B subunit-containing NMDA receptor antagonists prevent $\text{A}\beta$ -mediated synaptic plasticity disruption in vivo. *Proc Natl Acad Sci U S A.* 2009; 106:20504–20509. [PubMed: 19918059]
214. Raina P, Santaguida P, Ismaila A, Patterson C, Cowan D, Levine M, Booker L, Oremus M. Effectiveness of cholinesterase inhibitors and memantine for treating dementia: evidence review for a clinical practice guideline. *Ann Intern Med.* 2008; 148:379–397. [PubMed: 18316756]
215. Lipton SA. Pathologically-activated therapeutics for neuroprotection: mechanism of NMDA receptor block by memantine and S-nitrosylation. *Curr Drug Targets.* 2007; 8:621–632. [PubMed: 17504105]
216. Parsons CG, Stoffler A, Danysz W. Memantine: a NMDA receptor antagonist that improves memory by restoration of homeostasis in the glutamatergic system--too little activation is bad, too much is even worse. *Neuropharmacology.* 2007; 53:699–723. [PubMed: 17904591]
217. Haass C, Selkoe DJ. Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid β -peptide. *Nat Rev Mol Cell Biol.* 2007; 8:101–112. [PubMed: 17245412]
218. Roselli F, Tirard M, Lu J, Hutzler P, Lamberti P, Livrea P, Morabito M, Almeida OF. Soluble β -amyloid1-40 induces NMDA-dependent degradation of postsynaptic density-95 at glutamatergic synapses. *J Neurosci.* 2005; 25:11061–11070. [PubMed: 16319306]
219. De Felice FG, Vieira MN, Bomfim TR, Decker H, Velasco PT, Lambert MP, Viola KL, Zhao WQ, Ferreira ST, Klein WL. Protection of synapses against Alzheimer's-linked toxins: insulin signaling prevents the pathogenic binding of $\text{A}\beta$ oligomers. *Proc Natl Acad Sci U S A.* 2009; 106:1971–1976. [PubMed: 19188609]
220. Domingues A, Almeida S, Da Cruz e Silva EF, Oliveira CR, Rego AC. Toxicity of β -amyloid in HEK293 cells expressing NR1/NR2A or NR1/NR2B N-methyl-D-aspartate receptor subunits. *Neurochem Int.* 2007; 50:872–880. [PubMed: 17403555]
221. Kelly BL, Ferreira A. β -Amyloid-induced dynamin 1 degradation is mediated by N-methyl-D-aspartate receptors in hippocampal neurons. *J Biol Chem.* 2006; 281:28079–28089. [PubMed: 16864575]
222. De Felice FG, Velasco PT, Lambert MP, Viola K, Fernandez SJ, Ferreira ST, Klein WL. $\text{A}\beta$ oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine. *J Biol Chem.* 2007; 282:11590–11601. [PubMed: 17308309]
223. Engert F, Bonhoeffer T. Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature.* 1999; 399:66–70. [PubMed: 10331391]
224. Hayashi Y, Shi SH, Esteban JA, Piccini A, Poncer JC, Malinow R. Driving AMPA receptors into synapses by LTP and CaMKII: requirement for GluR1 and PDZ domain interaction. *Science.* 2000; 287:2262–2267. [PubMed: 10731148]
225. Kopec CD, Li B, Wei W, Boehm J, Malinow R. Glutamate receptor exocytosis and spine enlargement during chemically induced long-term potentiation. *J Neurosci.* 2006; 26:2000–2009. [PubMed: 16481433]
226. Kawamoto EM, Lepsch LB, Boaventura MF, Munhoz CD, Lima LS, Yshii LM, Avellar MC, Curi R, Mattson MP, Scavone C. Amyloid β -peptide activates nuclear factor- κ B through an N-methyl-D-aspartate signaling pathway in cultured cerebellar cells. *J Neurosci Res.* 2008; 86:845–860. [PubMed: 17969100]
227. Nakamura S, Murayama N, Noshita T, Katsuragi R, Ohno T. Cognitive dysfunction induced by sequential injection of amyloid- β and ibotenate into the bilateral hippocampus; protection by memantine and MK-801. *Eur J Pharmacol.* 2006; 548:115–122. [PubMed: 16952349]

228. Coan EJ, Irving AJ, Collingridge GL. Low-frequency activation of the NMDA receptor system can prevent the induction of LTP. *Neurosci Lett*. 1989; 105:205–210. [PubMed: 2577224]
229. Zorumski CF, Izumi Y. Modulation of LTP induction by NMDA receptor activation and nitric oxide release. *Prog Brain Res*. 1998; 118:173–182. [PubMed: 9932441]
230. Carroll RC, Lissin DV, von Zastrow M, Nicoll RA, Malenka RC. Rapid redistribution of glutamate receptors contributes to long-term depression in hippocampal cultures. *Nat Neurosci*. 1999; 2:454–460. [PubMed: 10321250]
231. Snyder EM, Philpot BD, Huber KM, Dong X, Fallon JR, Bear MF. Internalization of ionotropic glutamate receptors in response to mGluR activation. *Nat Neurosci*. 2001; 4:1079–1085. [PubMed: 11687813]
232. Xiao MY, Zhou Q, Nicoll RA. Metabotropic glutamate receptor activation causes a rapid redistribution of AMPA receptors. *Neuropharmacology*. 2001; 41:664–671. [PubMed: 11640920]
233. Zhou Q, Homma KJ, Poo MM. Shrinkage of dendritic spines associated with long-term depression of hippocampal synapses. *Neuron*. 2004; 44:749–757. [PubMed: 15572107]
234. Kaminsky YG, Marlatt MW, Smith MA, Kosenko EA. Subcellular and metabolic examination of amyloid-beta peptides in Alzheimer disease pathogenesis: evidence for Abeta (25–35). *Exp Neurol*. 2010; 221:26–37. [PubMed: 19751725]
235. Inestrosa NC, Varela-Nallar L, Grabowski CP, Colombres M. Synaptotoxicity in Alzheimer's disease: the Wnt signaling pathway as a molecular target. *IUBMB Life*. 2007; 59:316–321. [PubMed: 17505971]
236. Magdesian MH, Carvalho MM, Mendes FA, Saraiva LM, Juliano MA, Juliano L, Garcia-Abreu J, Ferreira ST. Amyloid-beta binds to the extracellular cysteine-rich domain of Frizzled and inhibits Wnt/beta-catenin signaling. *J Biol Chem*. 2008; 283:9359–9368. [PubMed: 18234671]
237. Levin ED, Simon BB. Nicotinic acetylcholine involvement in cognitive function in animals. *Psychopharmacology (Berl)*. 1998; 138:217–230. [PubMed: 9725745]
238. Jones S, Sudweeks S, Yakel JL. Nicotinic receptors in the brain: correlating physiology with function. *Trends Neurosci*. 1999; 22:555–561. [PubMed: 10542436]
239. Gray R, Rajan AS, Radcliffe KA, Yakehiro M, Dani JA. Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature*. 1996; 383:713–716. [PubMed: 8878480]
240. Genzen JR, McGehee DS. Short- and long-term enhancement of excitatory transmission in the spinal cord dorsal horn by nicotinic acetylcholine receptors. *Proc Natl Acad Sci U S A*. 2003; 100:6807–6812. [PubMed: 12748382]
241. Girod R, Barazangi N, McGehee D, Role LW. Facilitation of glutamatergic neurotransmission by presynaptic nicotinic acetylcholine receptors. *Neuropharmacology*. 2000; 39:2715–2725. [PubMed: 11044742]
242. Hu M, Waring JF, Gopalakrishnan M, Li J. Role of GSK-3beta activation and alpha7 nAChRs in Abeta(1-42)-induced tau phosphorylation in PC12 cells. *J Neurochem*. 2008; 106:1371–1377. [PubMed: 18485099]
243. Dineley KT, Bell KA, Bui D, Sweatt JD. beta -Amyloid peptide activates alpha 7 nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *J Biol Chem*. 2002; 277:25056–25061. [PubMed: 11983690]
244. Dougherty JJ, Wu J, Nichols RA. Beta-amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *J Neurosci*. 2003; 23:6740–6747. [PubMed: 12890766]
245. Pettit DL, Shao Z, Yakel JL. beta-Amyloid(1-42) peptide directly modulates nicotinic receptors in the rat hippocampal slice. *J Neurosci*. 2001; 21(1–5):RC120. [PubMed: 11150356]
246. Wu J, Kuo YP, George AA, Xu L, Hu J, Lukas RJ. beta-Amyloid directly inhibits human alpha4beta2-nicotinic acetylcholine receptors heterologously expressed in human SH-EP1 cells. *J Biol Chem*. 2004; 279:37842–37851. [PubMed: 15234980]
247. Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, Reitz AB. beta-Amyloid(1-42) binds to alpha7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem*. 2000; 275:5626–5632. [PubMed: 10681545]
248. Garcia-Osta A, Alberini CM. Amyloid beta mediates memory formation. *Learn Mem*. 2009; 16:267–272. [PubMed: 19318468]

249. Small DH, Maksel D, Kerr ML, Ng J, Hou X, Chu C, Mehrani H, Unabia S, Azari MF, Loiacono R, Aguilar MI, Chebib M. The beta-amyloid protein of Alzheimer's disease binds to membrane lipids but does not bind to the alpha7 nicotinic acetylcholine receptor. *J Neurochem*. 2007; 101:1527–1538. [PubMed: 17286584]
250. Liu Q, Kawai H, Berg DK. beta -Amyloid peptide blocks the response of alpha 7-containing nicotinic receptors on hippocampal neurons. *Proc Natl Acad Sci U S A*. 2001; 98:4734–4739. [PubMed: 11274373]
251. Seguela P, Wadiche J, Dineley-Miller K, Dani JA, Patrick JW. Molecular cloning, functional properties, and distribution of rat brain alpha 7: a nicotinic cation channel highly permeable to calcium. *J Neurosci*. 1993; 13:596–604. [PubMed: 7678857]
252. Chwang WB, O'Riordan KJ, Levenson JM, Sweatt JD. ERK/MAPK regulates hippocampal histone phosphorylation following contextual fear conditioning. *Learn Mem*. 2006; 13:322–328. [PubMed: 16741283]
253. Davies P, Maloney AJ. Selective loss of central cholinergic neurons in Alzheimer's disease. *Lancet*. 1976; 2:1403. [PubMed: 63862]
254. Perry EK, Perry RH, Blessed G, Tomlinson BE. Necropsy evidence of central cholinergic deficits in senile dementia. *Lancet*. 1977; 1:189. [PubMed: 64712]
255. Caspersen C, Wang N, Yao J, Sosunov A, Chen X, Lustbader JW, Xu HW, Stern D, McKhann G, Yan SD. Mitochondrial Abeta: a potential focal point for neuronal metabolic dysfunction in Alzheimer's disease. *FASEB J*. 2005; 19:2040–2041. [PubMed: 16210396]
256. Muirhead KE, Borger E, Aitken L, Conway SJ, Gunn-Moore FJ. The consequences of mitochondrial amyloid beta-peptide in Alzheimer's disease. *Biochem J*. 2010; 426:255–270. [PubMed: 20175748]
257. Du H, Guo L, Fang F, Chen D, Sosunov AA, McKhann GM, Yan Y, Wang C, Zhang H, Molkentin JD, Gunn-Moore FJ, Vonsattel JP, Arancio O, Chen JX, Yan SD. Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med*. 2008; 14:1097–1105. [PubMed: 18806802]
258. Du H, Yan SS. Mitochondrial permeability transition pore in Alzheimer's disease: cyclophilin D and amyloid beta. *Biochim Biophys Acta*. 2010; 1802:198–204. [PubMed: 19616093]
259. Singh P, Suman S, Chandna S, Das TK. Possible role of amyloid-beta, adenine nucleotide translocase and cyclophilin-D interaction in mitochondrial dysfunction of Alzheimer's disease. *Bioinformation*. 2009; 3:440–445. [PubMed: 19759867]
260. Schubert A, Grimm S. Cyclophilin D, a component of the permeability transition-pore, is an apoptosis repressor. *Cancer Res*. 2004; 64:85–93. [PubMed: 14729611]
261. Takuma K, Fang F, Zhang W, Yan S, Fukuzaki E, Du H, Sosunov A, McKhann G, Funatsu Y, Nakamichi N, Nagai T, Mizoguchi H, Ibi D, Hori O, Ogawa S, Stern DM, Yamada K, Yan SS. RAGE-mediated signaling contributes to intraneuronal transport of amyloid-beta and neuronal dysfunction. *Proc Natl Acad Sci U S A*. 2009; 106:20021–20026. [PubMed: 19901339]
262. Ren Y, Xu HW, Davey F, Taylor M, Aiton J, Coote P, Fang F, Yao J, Chen D, Chen JX, Yan SD, Gunn-Moore FJ. Endophilin I expression is increased in the brains of Alzheimer disease patients. *J Biol Chem*. 2008; 283:5685–5691. [PubMed: 18167351]
263. Lustbader JW, Cirilli M, Lin C, Xu HW, Takuma K, Wang N, Caspersen C, Chen X, Pollak S, Chaney M, Trinchese F, Liu S, Gunn-Moore F, Lue LF, Walker DG, Kuppusamy P, Zewier ZL, Arancio O, Stern D, Yan SS, Wu H. ABAD directly links Abeta to mitochondrial toxicity in Alzheimer's disease. *Science*. 2004; 304:448–452. [PubMed: 15087549]
264. Yao J, Irwin RW, Zhao L, Nilsen J, Hamilton RT, Brinton RD. Mitochondrial bioenergetic deficit precedes Alzheimer's pathology in female mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A*. 2009; 106:14670–14675. [PubMed: 19667196]
265. Alikhani N, Ankarcona M, Glaser E. Mitochondria and Alzheimer's disease: amyloid-beta peptide uptake and degradation by the presequence protease, hPreP. *J Bioenerg Biomembr*. 2009; 41:447–451. [PubMed: 19798557]
266. Takuma K, Yao J, Huang J, Xu H, Chen X, Luddy J, Trillat AC, Stern DM, Arancio O, Yan SS. ABAD enhances Abeta-induced cell stress via mitochondrial dysfunction. *FASEB J*. 2005; 19:597–598. [PubMed: 15665036]

267. Hansson Petersen CA, Alikhani N, Behbahani H, Wiehager B, Pavlov PF, Alafuzoff I, Leinonen V, Ito A, Winblad B, Glaser E, Ankarcrona M. The amyloid beta-peptide is imported into mitochondria via the TOM import machinery and localized to mitochondrial cristae. *Proc Natl Acad Sci U S A*. 2008; 105:13145–13150. [PubMed: 18757748]
268. Yao J, Taylor M, Davey F, Ren Y, Aiton J, Coote P, Fang F, Chen JX, Yan SD, Gunn-Moore FJ. Interaction of amyloid binding alcohol dehydrogenase/Abeta mediates up-regulation of peroxiredoxin II in the brains of Alzheimer's disease patients and a transgenic Alzheimer's disease mouse model. *Mol Cell Neurosci*. 2007; 35:377–382. [PubMed: 17490890]
269. Rizzuto R. Calcium mobilization from mitochondria in synaptic transmitter release. *J Cell Biol*. 2003; 163:441–443. [PubMed: 14610050]
270. Brown MR, Sullivan PG, Geddes JW. Synaptic mitochondria are more susceptible to Ca²⁺-overload than nonsynaptic mitochondria. *J Biol Chem*. 2006; 281:11658–11668. [PubMed: 16517608]
271. Chang DT, Honick AS, Reynolds IJ. Mitochondrial trafficking to synapses in cultured primary cortical neurons. *J Neurosci*. 2006; 26:7035–7045. [PubMed: 16807333]
272. Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH. Mitochondria are a direct site of A beta accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression. *Hum Mol Genet*. 2006; 15:1437–1449. [PubMed: 16551656]
273. Anandatheerthavarada HK, Biswas G, Robin MA, Avadhani NG. Mitochondrial targeting and a novel transmembrane arrest of Alzheimer's amyloid precursor protein impairs mitochondrial function in neuronal cells. *J Cell Biol*. 2003; 161:41–54. [PubMed: 12695498]
274. Mungarro-Menchaca X, Ferrera P, Moran J, Arias C. beta-Amyloid peptide induces ultrastructural changes in synaptosomes and potentiates mitochondrial dysfunction in the presence of ryanodine. *J Neurosci Res*. 2002; 68:89–96. [PubMed: 11933053]
275. Reddy PH, Beal MF. Amyloid beta, mitochondrial dysfunction and synaptic damage: implications for cognitive decline in aging and Alzheimer's disease. *Trends Mol Med*. 2008; 14:45–53. [PubMed: 18218341]
276. Serrano F, Klann E. Reactive oxygen species and synaptic plasticity in the aging hippocampus. *Ageing Res Rev*. 2004; 3:431–443. [PubMed: 15541710]
277. Thiels E, Klann E. Hippocampal memory and plasticity in superoxide dismutase mutant mice. *Physiol Behav*. 2002; 77:601–605. [PubMed: 12527006]
278. Parihar MS, Brewer GJ. Simultaneous age-related depolarization of mitochondrial membrane potential and increased mitochondrial reactive oxygen species production correlate with age-related glutamate excitotoxicity in rat hippocampal neurons. *J Neurosci Res*. 2007; 85:1018–1032. [PubMed: 17335078]
279. Parihar MS, Kunz EA, Brewer GJ. Age-related decreases in NAD(P)H and glutathione cause redox declines before ATP loss during glutamate treatment of hippocampal neurons. *J Neurosci Res*. 2008; 86:2339–2352. [PubMed: 18438923]
280. Hirai K, Aliev G, Nunomura A, Fujioka H, Russell RL, Atwood CS, Johnson AB, Kress Y, Vinters HV, Tabaton M, Shimohama S, Cash AD, Siedlak SL, Harris PL, Jones PK, Petersen RB, Perry G, Smith MA. Mitochondrial abnormalities in Alzheimer's disease. *J Neurosci*. 2001; 21:3017–3023. [PubMed: 11312286]
281. Bubber P, Haroutunian V, Fisch G, Blass JP, Gibson GE. Mitochondrial abnormalities in Alzheimer brain: mechanistic implications. *Ann Neurol*. 2005; 57:695–703. [PubMed: 15852400]
282. Parker WD Jr, Filley CM, Parks JK. Cytochrome oxidase deficiency in Alzheimer's disease. *Neurology*. 1990; 40:1302–1303. [PubMed: 2166249]
283. Butterfield DA. Amyloid beta-peptide [1-42]-associated free radical-induced oxidative stress and neurodegeneration in Alzheimer's disease brain: mechanisms and consequences. *Curr Med Chem*. 2003; 10:2651–2659. [PubMed: 14529455]
284. Mattson MP. Metal-catalyzed disruption of membrane protein and lipid signaling in the pathogenesis of neurodegenerative disorders. *Ann N Y Acad Sci*. 2004; 1012:37–50. [PubMed: 15105254]

285. Drake J, Link CD, Butterfield DA. Oxidative stress precedes fibrillar deposition of Alzheimer's disease amyloid beta-peptide (1-42) in a transgenic *Caenorhabditis elegans* model. *Neurobiol Aging*. 2003; 24:415–420. [PubMed: 12600717]
286. Cardoso SM, Santos S, Swerdlow RH, Oliveira CR. Functional mitochondria are required for amyloid beta-mediated neurotoxicity. *FASEB J*. 2001; 15:1439–1441. [PubMed: 11387250]
287. Brewer GJ. Epigenetic oxidative redox shift (EORS) theory of aging unifies the free radical and insulin signaling theories. *Exp Gerontol*. 2010; 45:173–179. [PubMed: 19945522]
288. Abramov AY, Duchen MR. Mechanisms underlying the loss of mitochondrial membrane potential in glutamate excitotoxicity. *Biochim Biophys Acta*. 2008; 1777:953–964. [PubMed: 18471431]
289. Jurgensen S, Ferreira ST. Nicotinic receptors, amyloid-beta, and synaptic failure in Alzheimer's disease. *J Mol Neurosci*. 2010; 40:221–229. [PubMed: 19690986]
290. Palop JJ, Chin J, Mucke L. A network dysfunction perspective on neurodegenerative diseases. *Nature*. 2006; 443:768–773. [PubMed: 17051202]




Fig. 1. Model of picomolar A β -induced insulin-PI3K-Akt-ERK signalling plus mitochondrial targets of intracellular A β

Extracellular A β at picomolar concentration binds to the insulin receptor (IR) and activates PKB/Akt via PDK-1. PKB/Akt translocates into the nucleus and phosphorylates CREB. Activation of the lipid kinase PI3K is critical for the activation of PKB by PDK. PDK1 phosphorylates the activation loop of a number of protein serine/threonine kinases of the AGC kinase superfamily, including protein kinase B (PKB α ; also called Akt1). Akt may also maintain the integrity of the mitochondria by a unknown mechanism or by a specific mechanism of Bad phosphorylation. Akt can also inhibit apoptosis by phosphorylation and inactivation of caspase-9. ERK1/2 are activated by upstream MAPKK, such as MEK1/2, and MAPKKK, such as c-Raf. MEK1/2 induce ERK1/2 activation via dual phosphorylation on threonine 202 and tyrosine 204 residues. Phosphorylation of ERK leads to the activation of a number of transcription factors, important in controlling differentiation, neuronal survival, learning and memory plasticity. For example, ERK activates pro-survival transcription factor CREB, by activating both p90RSK and MSK1/2.

Picomolar extracellular A β also binds nAChR, glutamate receptors (NMDAR) and Ca $^{2+}$ ion channels (e.g. VDCCs, TRPC) and causes Ca $^{2+}$ influx at controlled rates into the cytoplasm and mitochondria. Increased cytosolic calcium concentrations initiate the activation of several kinase-dependent signalling cascades including activation of PKC leading to CREB activation and phosphorylation at Ser133, a process critical for protein synthesis-dependent synaptic plasticity and LTP. PKC- α also activates ERK by interacting with Ras or Raf-1. Mitochondria are critical targets of intracellular A β . A β interacts with CypD, a protein component of the membrane permeability transition pore (MPTP). The interaction of CypD with A β causes functional modification of this protein leading to MPTP opening. A β also binds with another mitochondrial protein, ABAD to distort the enzyme's structure, rendering it inactive. This causes an increase in reactive oxygen species and oxidative stress leading to initiation of apoptosis.

Table 1

Neuron survival and memory functions of A β peptides.

Peptide	Form	Source	Concentration (nM)	Function	Reference
A β 1-40	monomer	Synthetic	10 μ M	Inhibits formation of A β 1-42 β -sheets	94
A β 1-40	monomer	Synthetic	0.1-10	Neurotrophic for E18 rat neurons	98
A β 1-40 & A β 1-42	Monomer or mixed	Synthetic	0.01-1	Increases rat cortical neuronal cell viability	28
A β 1-40 & A β 1-42	Mono + oligomers; anti-A β antibody	Synthetic	0.1	Rat memory formation & consolidation	248
A β 1-40 & A β 1-42	monomer	Synthetic	1-10; 1-1000	Increased survival (PI-3K, AKT) and protection from NMDA toxicity	97
A β 1-42	monomer	Synthetic	0.2	Enhanced LTP in mouse hippocampal slices + improves mouse memory by ICV injection	83
A β 1-42	Monomers +oligomers	Synthetic	0.01-1	Inhibited nicotine- evoked increases of Ca ²⁺ in rat synaptosomes	153