

Growth Temperature-Dependent Contributions of Response Regulators, σ^B , PrfA, and Motility Factors to *Listeria monocytogenes* Invasion of Caco-2 Cells

Reid A. Ivy, Yvonne C. Chan, Barbara M. Bowen, Kathryn J. Boor, and Martin Wiedmann

Abstract

Foodborne pathogens encounter rapidly changing environmental conditions during transmission, including exposure to temperatures below 37°C. The goal of this study was to develop a better understanding of the effects of growth temperatures and temperature shifts on regulation of invasion phenotypes and invasion-associated genes in *Listeria monocytogenes*. We specifically characterized the effects of *L. monocytogenes* growth at different temperatures (30°C vs. 37°C) on (i) the contributions to Caco-2 invasion of different regulators (including σ^B , PrfA, and 14 response regulators [RRs]) and invasion proteins (i.e., InlA and FlaA), and on (ii) *gadA*, *plcA*, *inlA*, and *flaA* transcript levels and their regulation. Overall, Caco-2 invasion efficiency was higher for *L. monocytogenes* grown at 30°C than for bacteria grown at 37°C ($p=0.0051$ for the effect of temperature on invasion efficiency; analysis of variance); the increased invasion efficiency of the parent strain 10403S (serotype 1/2a) observed after growth at 30°C persisted for 2.5 h exposure to 37°C. For *L. monocytogenes* grown at 30°C, the motility RRs DegU and CheY and σ^B , but not PrfA, significantly contributed to Caco-2 invasion efficiency. For *L. monocytogenes* grown at 37°C, none of the 14 RRs tested significantly contributed to Caco-2 invasion, whereas σ^B and PrfA contributed synergistically to invasion efficiency. At both growth temperatures there was significant synergism between the contributions to invasion of FlaA and InlA; this synergism was more pronounced after growth at 30°C than at 37°C. Our data show that growth temperature affects invasion efficiency and regulation of virulence-associated genes in *L. monocytogenes*. These data support increasing evidence that a number of environmental conditions can modulate virulence-associated phenotypes of foodborne bacterial pathogens, including *L. monocytogenes*.

Introduction

LISTERIA MONOCYTOGENES is a Gram-positive, non-spore-forming rod that is capable of causing disease in humans and animals. *L. monocytogenes* can enter intestinal epithelia via an internalization process initiated by the interaction of InlA, expressed on the surface of invading *L. monocytogenes*, and E-cadherin, expressed on the epithelial cell surface. Although the organism is widespread in nature, 99% of *L. monocytogenes* infections are foodborne (Mead *et al.*, 1999). *L. monocytogenes* therefore encounter a variety of different environments and associated stress conditions during transmission from the environment through foods to humans, including a wide range of temperature, pH, and osmotic stress conditions.

The alternative sigma factor, σ^B , the pleiotropic transcriptional regulator PrfA, and two-component regulatory systems (TCS) have all been shown to regulate key processes important for *L. monocytogenes* stress response and virulence

(Kallipolitis and Ingmer, 2001; Kazmierczak *et al.*, 2003; Williams *et al.*, 2005a; Scotti *et al.*, 2007). In addition to a large stress response regulon, σ^B specifically regulates the transcription of genes involved in responses to stresses encountered during passage through the gastrointestinal system such as osmotic stress and acid stress. σ^B also regulates transcription of genes encoding different internalins that are involved in entry into host cells (Raffelsbauer *et al.*, 1998; Kazmierczak *et al.*, 2003; Kim *et al.*, 2005; McGann *et al.*, 2007b). PrfA regulates the *L. monocytogenes* core virulence genes *plcA*, *plcB*, *hly*, *mpl*, and *actA*, which are important for escape from the vacuole and cell-to-cell spread (Scotti *et al.*, 2007) and also co-regulates, with σ^B , other virulence genes, including *inlA* (McGann *et al.*, 2008). Among the 16 putative or confirmed TCS in *L. monocytogenes*, a number of them have been shown to regulate bacterial adaptation (Autret *et al.*, 2003; Brondsted *et al.*, 2003; Kallipolitis *et al.*, 2003; Dons *et al.*, 2004; Mandin *et al.*, 2005; Sleator and Hill, 2005; Williams *et al.*, 2005b; Larsen

et al., 2006), including VirR, DegU and CheY, which have been implicated in regulating mechanisms contributing to host cell invasion (Dons *et al.*, 2004; Knudsen *et al.*, 2004; Mandin *et al.*, 2005).

Growth temperature has been shown to have a profound effect on activities of key *L. monocytogenes* regulators (Leimeister-Wachter *et al.*, 1992; Liu *et al.*, 2002; Dons *et al.*, 2004; Chan *et al.*, 2007b; McGann *et al.*, 2007a; van der Veen *et al.*, 2007). For example, expression levels of the genes comprising the PrfA-regulated *Listeria* pathogenicity island I, which are important in escape from the host cell vacuole and cell-to-cell spread, are maximal at 37°C due to a *prfA* mRNA thermosensor that represses PrfA translation at lower temperatures (Johansson *et al.*, 2002). Also, certain σ^B -dependent genes have been shown to be involved in adaptation to cold temperatures (Becker *et al.*, 2000; Chan *et al.*, 2007a), and σ^B -dependent internalin genes, including *inlC2* and *inlD*, have been shown to be expressed at higher levels at temperatures $\leq 30^\circ\text{C}$ than at 37°C (McGann *et al.*, 2007a). Finally, *L. monocytogenes* motility, which has been shown to contribute to host cell invasion (Dons *et al.*, 2004) and increased virulence in a mouse model (O'Neil and Marquis, 2006), is temperature dependent. At 37°C, the transcription of *flaA* (the gene encoding the flagellin structural protein) and other motility-associated genes is repressed by MogR (Grundling *et al.*, 2004; Shen and Higgins, 2006), though the stringency of this repression may vary among different strains of *L. monocytogenes*. For example, repression of motility genes and associated motility phenotypes has been shown to be less stringent in *L. monocytogenes* 10403S than in other strains (Grundling *et al.*, 2004; Way *et al.*, 2004).

Although *L. monocytogenes* will have likely been adapted to a temperature below 37°C before ingestion, many studies on *L. monocytogenes* host cell invasion have been done only on *L. monocytogenes* cells grown at 37°C and have only determined the effects of single gene mutations on invasion (Cotter *et al.*, 1999; Kallipolitis and Ingmer, 2001; Autret *et al.*, 2003; Williams *et al.*, 2005a). Therefore, a more comprehensive evaluation of the effect of growth temperature on the contributions of response regulators (RRs), transcriptional regulators, and motility genes as well as functional synergisms between select genes to host cell invasion is necessary to better understand how adaptation to environments outside a mammalian host affects *L. monocytogenes* virulence. The objective of this study was, thus, to characterize the effects of growth temperature (30°C vs. 37°C) on (i) the contributions of σ^B and PrfA, 14 RRs, and invasion proteins InlA and FlaA to Caco-2 invasion, and on (ii) transcript levels of *inlA*, *flaA*, *gadA*, and *plcA*. Growth temperatures of 30°C and 37°C were chosen because *L. monocytogenes* grown at 30°C and 37°C show very similar lag phase durations and growth rates, whereas *L. monocytogenes* grown at lower temperatures (e.g. 22.5°C; [Pal *et al.*, 2008]) show considerably longer lag phase durations and slower growth rates. As Caco-2 invasion efficiency has been shown to vary with bacterial growth phase (Garner *et al.*, 2006a), it is critical to use bacteria grown with similar growth parameters and synchronized to similar growth phases to ensure that differences in invasion efficiency represent a temperature effect rather than differences in growth phase. In addition, *L. monocytogenes* grown at 30°C have previously been shown to display differential expression of key virulence-associated characteristics, including (i)

downregulation of PrfA activity (Johansson *et al.*, 2002) and (ii) upregulation of motility (O'Neil and Marquis, 2006). Consistent with our choice of 30°C as a growth temperature that allows for appropriate comparison between *L. monocytogenes* grown at typical mammalian body temperature and bacteria grown under environmental conditions, others (Toledo-Arana *et al.*, 2009) have recently used 30°C as a growth temperature representing the saprophytic stage of the life of *L. monocytogenes*. Despite these key reasons for using growth temperatures of 30°C and 37°C to study temperature-dependent phenotypes of *L. monocytogenes*, we appreciate that these temperatures represent an experimental model that does not necessarily reflect natural transmission where foodborne bacteria are more likely exposed to refrigeration or room temperatures or heat shock conditions before ingestion.

Materials and Methods

Bacterial strains and growth conditions

Bacterial strains used in this study are listed in Table 1. For invasion assays, *L. monocytogenes* 10403S (Bishop and Hinrichs, 1987) and associated mutant strains were grown to early stationary phase as previously described (McGann *et al.*, 2007a). Briefly, a 12 to 18 h culture grown at 37°C with aeration (i.e., shaking at 220 rpm) in brain heart infusion broth (BHI) was diluted 1:100 into 5 mL of fresh BHI and grown at 37°C with aeration to OD₆₀₀ = 0.4. This culture was diluted 1:100 into another 5 mL of fresh BHI and grown at 30°C or 37°C with aeration to early stationary phase (defined as growth to OD₆₀₀ = 1.0, followed by an additional 3 h incubation). For temperature shift experiments, a 1 mL aliquot of the early stationary phase culture was centrifuged at 12,000 rpm for 10 min, resuspended in phosphate-buffered saline (PBS; 0.14 M NaCl, 2.7 mM KCl, 10 mM Na₂PO₄, and 1.8 mM KH₂PO₄; pH 7.2), and incubated statically at 30°C or 37°C for 0, 2.5, or 5 h.

Mutant construction

Internal, in-frame, nonpolar deletion mutant alleles in 14 RR genes (Table 1) were created using splicing overlap extension polymerase chain reaction (PCR) (see Supplemental Table S1, available online at www.liebertonline.com for primers; all supplemental material is located at <http://author.cals.cornell.edu/cals/foodsci/research/labs/wiedmann/links/ivy2010-1.cfm>) and cloned into the temperature-sensitive suicide shuttle vector pKSV7 (Smith and Youngman, 1992) as previously detailed by our group (Wiedmann *et al.*, 1998). Mutant alleles were introduced into *L. monocytogenes* 10403S using standard allelic exchange mutagenesis procedures (Camilli *et al.*, 1993). Mutant strains were confirmed with sequencing. *lmo0287* is likely to be essential, as null mutants in this gene could not be constructed here, consistent with similar studies by others (Kallipolitis and Ingmer, 2001; Williams *et al.*, 2005a).

Caco-2 invasion assays

The Caco-2 cell-line (ATCC HTB-37) was maintained at 37°C (4%–6% CO₂ and 85% humidity) in a Caco-2 medium, which was Dulbecco's minimal essential medium with Earle's salts, 1% sodium pyruvate, 20% fetal bovine serum, 1.0%

TABLE 1. STRAINS USED IN THIS STUDY

Strain designation ^a	Genotype	Description of gene product	Reference ^b
X1-001	Parent strain	Strain 10403S; serotype 1/2a	Bishop and Hinrichs (1987)
K4-006	$\Delta inlA$	Internalin A, required for internalization into selected host cells	Bakardjiev <i>et al.</i> (2004)
H6-199	$\Delta flaA$	Flagellum structural protein	O'Neil and Marquis (2006)
B4-007	$\Delta lisRK^c$	LisRK, contributes to log-phase acid resistance	This study
B2-078	$\Delta agrA$	Response regulator (RR), contributes to protein secretion	This study
B2-080	$\Delta resD$	RR, contributes to virulence gene repression in presence of select carbohydrates	This study
B2-086	$\Delta lmo1022$	Putative RR, no known role	This study
B2-096	$\Delta lmo1060$	Putative RR, no known role	This study
C5-017	$\Delta lmo2010$	Putative RR, no known role	This study
C5-019	$\Delta lmo2583$	Putative RR, no known role	This study
B2-100	$\Delta phoP$	RR, similar to <i>Bacillus subtilis</i> PhoP	This study
C5-041	$\Delta virR$	Novel RR involved in invasion of Caco-2 cells	This study
C5-036	$\Delta lmo1507$	Putative RR, no known role	This study
C5-033	$\Delta degU$	RR, regulates expression of motility genes	This study
B2-104	$\Delta kdpE$	RR, contributes to growth at high osmolarity and low temperature	This study
B2-105	$\Delta cheY$	RR, contributes to chemotaxis	This study
B2-102	$\Delta cesR$	RR, contributes to ethanol and β -lactam tolerance	This study
A1-254	$\Delta sigB$	Alternative sigma factor σ^B , regulates general stress response	Wiedmann <i>et al.</i> (1998)
B2-046	$\Delta prfA$	PrfA, regulation of virulence genes	Cheng and Portnoy (2003)
B2-068	$\Delta sigB\Delta prfA$		McGann <i>et al.</i> (2007b)
I1-001	$\Delta inlA\Delta flaA$		Marquis Lab

^aAll strain designations carry the prefix FSL.

^bReferences where a given mutant was previously described.

^cThis mutant includes an internal deletion of the *lisR* gene, which also removed a portion of the ribosome binding site of *lisK* sensory kinase gene.

RR, response regulator.

nonessential amino acids, 1.5 g/L sodium bicarbonate, and, when appropriate, penicillin G and streptomycin (each at 100 μ g/mL) (all reagents were obtained from Gibco, a subsidiary of Invitrogen). All invasion assays were performed at 37°C as previously described (Garner *et al.*, 2006b) with minor modifications. Briefly, 48 h before the assay, Caco-2 cells were seeded (from a culture passaged no >60 times) into 24-well tissue culture plates (Corning Inc.) at a density of 5×10^4 cells/well in the Caco-2 medium without antibiotics. For infection, $\sim 2 \times 10^7$ CFU *L. monocytogenes* were added to each well (representing a multiplicity of infection of ~ 200). All inocula were enumerated on BHI agar plates. Thirty minutes postinfection, the Caco-2 monolayers were washed three times with PBS to remove any unassociated *L. monocytogenes*, and the medium was replaced with a fresh Caco-2 medium. Forty-five minutes after infection, the medium was replaced with the Caco-2 medium plus 150 μ g/mL gentamycin to kill any extracellular *L. monocytogenes*. At 90 min postinfection, Caco-2 cells were washed three times with PBS and lysed with ice-cold distilled water. Intracellular *L. monocytogenes* were enumerated by plating the appropriate dilutions of the Caco-2 lysate on BHI agar, using a spiral plater (Spiral Biotech). At least three independent trials of the invasion assays were performed with duplicate wells tested for each treatment in each replicate.

Quantitative reverse transcriptase-PCR (qRT-PCR)

Transcript levels of *inlA*, *flaA*, *plcA*, *gadA*, *rpoB*, *sigB*, and *prfA* were quantified for select strains grown to early stationary phase at 30°C or 37°C using TaqMan probes and primers and the ABI Prism 7000 Sequence Detection System as previously described (Sue *et al.*, 2003; Chaturongakul and Boor, 2006; McGann *et al.*, 2007a) with one exception: copy numbers for each gene were normalized to *rpoB* levels. Primers and probes for *inlA*, *plcA*, *gadA*, *sigB*, *rpoB*, and *prfA* have been previously described (Sue *et al.*, 2004; Kim *et al.*, 2005; Kazmierczak *et al.*, 2006). *flaA* Taqman primers (*flaA*-F: 5'-TCGTAAAATAACGAAGGCATGAC-3'; *flaA*-R: AGA ACTGTTAATACGTTTACCAGATGCT-3') and the *flaA* MGB probe (FAM-5'-CAAGCGCAAGAAC-3'NFQ) were designed using Primer Express 1.0 (Applied Biosystems).

Statistical analyses

All statistical analyses were performed in JMP 7.0 (SAS Institute Inc.). Invasion efficiencies were initially analyzed using a one-way analysis of variance (ANOVA; $\alpha = 0.05$). Strain and, where appropriate, date of experiment were included as variables in the model. For each ANOVA, data were log-transformed to ensure that the data set satisfied ANOVA assumptions of normality of residuals and equality

of variances. To determine whether a particular mutant strain differed from the parent strain (e.g., in invasion efficiency), a *post hoc* Dunnett's many to one test was used (Shun *et al.*, 2003). To compare invasion efficiencies for a given strain exposed to multiple conditions (i.e., as shown in Table 2) or among strains (in cases where a strain with a double mutation was included in the comparison), a *post hoc* Tukey honestly significantly different (HSD) test was used. For qRT-PCR, normalized log copy numbers for a given gene were compared among strains by using one-way ANOVA, followed by Tukey HSD. To measure whether two gene deletions showed an effect on invasion efficiency or transcript level that is more than additive (which indicates synergism), the parent strain, single mutants, and double mutants were assigned unique allelic states by coding dummy variables (e.g., "gene1" and "gene2") for each allelic state with 1 meaning the gene is present and 0 meaning the gene is absent. A two-way ANOVA was performed to determine the effect of each allelic state on invasion efficiency or transcript level in these analyses. "Date of experiment," "gene1," "gene2," and "gene1*gene2" were included as effects in the two-way ANOVA model. A significant effect of gene1*gene2 in the model (i.e., $p < 0.05$) indicates a more than additive effect of the double mutation (i.e., deletion of both *gene1* and *gene2* in the same genetic background has a greater effect than the sum of the effects of deleting *gene1* alone and *gene2* alone).

Results

Contributions of RRs to Caco-2 invasion

Overall ANOVA analysis showed a highly significant effect of temperature on invasion efficiency ($p = 0.0051$), and all RR mutants (except $\Delta degU$ and $\Delta cheY$) and 10403S had numerically higher mean Caco-2 invasion efficiencies after growth to early stationary phase in BHI at 30°C than growth at 37°C. For several strains, including 10403S, $\Delta resD$, $\Delta lmo1022$, $\Delta phoP$, $\Delta cesR$, and $\Delta kdpE$, this difference was statistically significant ($p < 0.05$; *t*-test; Fig. 1).

For the *L. monocytogenes* parent strain and the 14 RR mutants grown at 30°C, ANOVA showed a significant effect of strain on Caco-2 invasion efficiency ($p < 0.0001$). Specifically, $\Delta degU$ and $\Delta cheY$ both had ~50-fold lower mean invasion efficiencies than the parent strain ($p < 0.0001$ for each; Dunnett's test) when grown at 30°C (Fig. 1). These results confirm the findings of other studies that have shown a role for CheY in

host cell invasion (Dons *et al.*, 2004). Unlike all other strains tested, the $\Delta cheY$ and $\Delta degU$ strains showed numerically higher invasion efficiency after growth at 37°C than at 30°C (Fig. 1).

For *L. monocytogenes* parent strain 10403S and the 14 RR mutants grown to early stationary phase at 37°C, the effect of the factor "strain" on Caco-2 invasion efficiencies was not statistically significant ($p = 0.3900$; ANOVA). Therefore, the appropriate conservative statistical test (i.e., ANOVA) found no statistically significant reduction in Caco-2 invasion for any of the 14 RR mutants grown at 37°C.

As *L. monocytogenes* motility has been reported in other *L. monocytogenes* strains to be regulated by DegU and CheY at temperatures $\leq 30^\circ\text{C}$ (Dons *et al.*, 2004; Knudsen *et al.*, 2004; Williams *et al.*, 2005a; Mauder *et al.*, 2008), we conducted swarming assays to evaluate motility of the *L. monocytogenes* 10403S, $\Delta degU$, $\Delta cheY$, and $\Delta flaA$ strains. While 10403S was clearly motile when grown at 30°C, $\Delta degU$, $\Delta cheY$, and $\Delta flaA$ showed reduced swarming (6%, 6%, and 1%, respectively, of the swarming area for 10403S grown at 30°C; see Supplemental Fig. S1, available online at www.liebertonline.com). When grown at 37°C, the parent strain showed limited and considerably reduced swarming compared to 30°C (Supplemental Fig. S1), and the three mutant strains showed no detectable swarming at this temperature. The finding that 10403S exhibited some swarming at 37°C is consistent with previous reports that *L. monocytogenes* 10403S (the parent strain used here) shows limited *flaA* expression even at 37°C (Grundling *et al.*, 2004; Way *et al.*, 2004).

Effects of temperature shifts and holds on growth temperature-dependent phenotypes

On the basis of the data reported above, we evaluated whether the increased invasion efficiency of 10403S at 30°C is maintained after a shift to 37°C. While *L. monocytogenes* 10403S grown at 30°C and held (in PBS) at 37°C for 2.5 h (30°C–37°C) showed numerically (about twofold) reduced invasion efficiency compared to bacteria either before the shift or bacteria shifted to 30°C for 2.5 h (Table 2), the invasion efficiency of the "30°C–37°C" treatment group was not significantly different from the control groups grown and held at 30°C (30°C–30°C in Table 2) ($p > 0.05$; Tukey HSD) but was significantly higher than control groups grown and held at 37°C (37°C–37°C) ($p < 0.05$; Tukey HSD). These results suggest that increased invasion efficiency of *L. monocytogenes*

TABLE 2. CACO-2 INVASION EFFICIENCIES OF *LISTERIA MONOCYTOGENES* 10403S AND SELECT MUTANTS AFTER EXPOSURE TO VARYING TEMPERATURE TREATMENTS

Growth temperature	Temperature of PBS hold (2.5 h)	Caco-2 invasion efficiency (calculated as [CFU recovered/CFU infected]×100) (SD) for ^a				
		10403S	$\Delta cheY$	$\Delta degU$	$\Delta flaA$	$\Delta inlA$
30°C	None	0.67 (0.17) ^A	0.018 (0.007) ^{B,*}	0.015 (0.008) ^{B,*}	0.014 (0.010) ^{B,*}	0.024 (0.010) ^{A,*}
30°C	30°C	0.64 (0.19) ^A	0.013 (0.003) ^{B,*}	0.010 (0.005) ^{B,*}	0.010 (0.006) ^{B,*}	0.023 (0.007) ^{A,*}
30°C	37°C	0.36 (0.10) ^A	0.006 (0.002) ^{B,*}	0.007 (0.002) ^{B,*}	0.007 (0.004) ^{B,*}	0.014 (0.005) ^{A,*}
37°C	None	0.14 (0.03) ^B	0.14 (0.06) ^A	0.20 (0.05) ^A	0.14 (0.06) ^A	0.0008 (0.0003) ^{B,*}
37°C	37°C	0.11 (0.04) ^B	0.12 (0.05) ^A	0.14 (0.10) ^A	0.09 (0.06) ^A	0.0007 (0.0007) ^{B,*}
37°C	30°C	0.11 (0.03) ^B	0.13 (0.05) ^A	0.17 (0.08) ^A	0.14 (0.07) ^A	0.0007 (0.0005) ^{B,*}

^aWithin a given column values with identical letters (A or B) are not significantly different ($p \geq 0.05$; *post hoc* Tukey HSD test). Within a given row, invasion efficiencies that are lower for a given mutant than the parent strain exposed to the same condition ($p \leq 0.05$; *post hoc* Dunnett's test) are marked with an asterisk (*). Data represent the mean and standard deviation of three biological replicates.

PBS, phosphate-buffered saline; SD, standard deviation; ANOVA, analysis of variance; HSD, honestly significantly different.




FIG. 1. Caco-2 invasion efficiencies of *Listeria monocytogenes* 10403S and 14 mutant strains with deletions in genes encoding RRs. Bacteria were grown to early stationary phase (growth to $OD_{600} = 1.0$, followed by 3 h of incubation) with aeration (i.e., shaking at 220 rpm) at 30°C or 37°C. Invasion assays were performed at 37°C. Data represent the mean of at least three biological replicates. Error bars represent SD. A $\Delta inlA$ strain was included as a control. Overall ANOVA analyses of the 14 RR mutants and the parent strain showed a significant effect of temperature ($p = 0.0051$); all strains (except $\Delta cheY$ and $\Delta degU$) showed numerically higher invasion efficiencies when grown at 30°C. Strains showing statistically higher invasion after growth at 30°C than at 37°C ($p < 0.05$; t -test) are denoted by ^. For 10403S and RR mutants, ANOVA showed a significant effect of strain on invasion efficiency for bacteria grown at 30°C ($p < 0.0001$). For bacteria grown at 37°C this effect was not significant ($p = 0.3900$). Among RR mutants grown at 30°C, $\Delta cheY$ and $\Delta degU$ showed significantly lower invasion efficiencies than the parent strain ($p < 0.0001$; *post hoc* Dunnett's; denoted by *). RR, response regulator; ANOVA, analysis of variance; SD, standard deviation.

grown at 30°C is largely maintained during a 2.5 h hold at 37°C in PBS. When *L. monocytogenes* 10403S was grown at 30°C, and shifted to and held at 37°C for 5 h (in PBS), the invasion efficiency was reduced (0.15% invasion efficiency) and virtually identical to that of *L. monocytogenes* grown at 37°C (0.14% invasion efficiency) with no significant difference ($p > 0.05$; Tukey HSD). This indicates that prolonged exposure of *L. monocytogenes* grown at 30°C to 37°C (in PBS) reduces invasion efficiencies to values typical for bacteria grown at 37°C. Growth and *de novo* protein synthesis (which are unlikely to occur in PBS) thus do not seem to be required for reduced invasiveness observed at 37°C.

When the *L. monocytogenes* parent strain grown at 37°C was switched to 30°C and held at this temperature for 2.5 h, bacteria showed no changes in invasion efficiency and maintained lower invasion efficiency typical for bacteria grown at 37°C ($p < 0.05$; Tukey HSD; Table 2), suggesting that the increased invasion phenotype cannot be induced by simply switching the organism to 30°C but requires growth at 30°C. Controls included in these experiments showed that exposure to PBS for 2.5 h had no effect on invasion efficiency; bacteria grown at 30°C and exposed to PBS at 30°C for 2.5 h as well as bacteria grown at 37°C and exposed to PBS at 37°C for 2.5 h did not differ in their invasion efficiencies from bacteria grown at 30°C or 37°C, respectively, without subsequent exposure to PBS ($p > 0.05$; Tukey HSD; Table 2).

As *L. monocytogenes* $\Delta degU$, $\Delta cheY$, and $\Delta flaA$ showed lower invasion than the parent strain after growth to early stationary phase at 30°C, we also tested whether this invasion defect was maintained after a shift from 30°C to 37°C, followed by a 2.5 h hold at this temperature (in PBS). All three strains ($\Delta degU$, $\Delta cheY$, and $\Delta flaA$) maintained lower invasion efficiencies than the parent strain even after bacteria grown at 30°C were held for 2.5 h at 37°C (Table 2). Invasion efficiencies for these strains grown at 30°C and shifted to 37°C for 2.5 h were not significantly

different ($p > 0.05$; Tukey HSD; Table 2) from invasion efficiencies for (i) strains grown at 30°C (before shift) or (ii) strains grown at 30°C and subsequently held at 30°C in PBS for 2.5 h (Table 2). The invasion efficiencies for $\Delta degU$, $\Delta cheY$, and $\Delta flaA$ grown at 30°C and held for 2.5 h at 37°C also continued to be significantly lower than the invasion efficiencies for the same strains grown at 37°C ($p < 0.05$; Tukey HSD); invasion efficiencies for these three strains in the "30°C–37°C" treatment group were ~20 to 30-fold lower than the same strains grown at 37°C (Table 2). These three mutant strains thus maintained their invasion deficient phenotype even after 2.5 h exposure to 37°C. To test the effect of prolonged exposure to 37°C on the reduced invasion efficiency of motility-deficient mutant strains, invasion efficiency of the $\Delta flaA$ strain was also evaluated for bacteria grown at 30°C and shifted to 37°C with a hold in PBS at 37°C for 5 h. Even after a hold at 37°C for 5 h, the $\Delta flaA$ strain maintained reduced invasion efficiency (0.006%) compared to the invasion efficiency of either the parent strain grown at 37°C (0.14%) or $\Delta flaA$ grown at 37°C (0.14%); both comparisons had $p < 0.05$; Tukey HSD), suggesting that increased invasion efficiency at 37°C, of a $\Delta flaA$ mutant, requires growth at 37°C and probably *de novo* protein synthesis (which is unlikely to occur in PBS). When the $\Delta degU$, $\Delta cheY$, and $\Delta flaA$ strains were grown at 37°C and shifted to 30°C with a hold at 30°C for 2.5 h, all three strains maintained the higher invasion efficiency as they displayed when grown at 37°C (Table 2).

Temperature-dependent contributions of *InlA* and *FlaA* to Caco-2 invasion

In initial experiments, the *L. monocytogenes* $\Delta inlA$ strain showed 4.4-fold higher invasion efficiency when grown at 30°C than at 37°C (Fig. 1). In subsequent experiments, when grown at 37°C, the $\Delta inlA$ strain showed, on average, greater than 100-fold lower invasion efficiency than either the parent

strain or the $\Delta degU$, $\Delta cheY$, or $\Delta flaA$ strains grown at 37°C (Table 2). However, when grown at 30°C, the $\Delta inlA$ strain showed only a 26-fold lower average invasion efficiency than the parent strain (Table 2). These experiments also confirmed a significantly higher invasion efficiency for the $\Delta inlA$ grown at 30°C than at 37°C ($p < 0.05$; Tukey HSD; Table 2). Temperature shifts (i.e., 30°C–37°C, 2.5 h or 37°C–30°C, 2.5 h) did not significantly affect these statistical differences (Table 2). These results indicate an InlA-independent increase in Caco-2 invasion for *L. monocytogenes* grown at 30°C (as compared to 37°C), which is maintained after a shift to 37°C.

To further investigate the contributions of InlA and flagellin to Caco-2 invasion after growth at 30°C and 37°C and to specifically determine whether there is synergism between InlA and flagellin in facilitating invasion, 10403S, $\Delta flaA$, $\Delta inlA$, and $\Delta inlA\Delta flaA$ were grown at 30°C or 37°C and tested for their Caco-2 invasion efficiencies (Fig. 2). After growth at 30°C, $\Delta flaA$ showed ~30-fold lower invasion efficiency than the parent strain (Fig. 2). At both 30°C and 37°C, the invasion efficiency of the $\Delta inlA\Delta flaA$ strain was lower ($p < 0.05$; Tukey HSD) than that of the parent strain (0.007% vs. 0.680% for 37°C and 0.003% vs. 2.064% for 30°C, respectively). Further, two-way ANOVA analysis showed a significant “flaA*inlA” interaction effect on Caco-2 invasion for *L. monocytogenes* grown at both 30°C ($p = 0.0015$) and 37°C ($p = 0.0269$). Interestingly, the F-ratio of the interaction effect for bacteria grown at 30°C ($F = 20.15$) was higher than that of bacteria grown at 37°C ($F = 6.97$), suggesting a greater contribution of the interaction effect to the observed variance at 30°C than at 37°C. These statistical findings indicate that the effects of deleting *inlA* and *flaA* on Caco-2 invasion are more than additive, indicating a synergism between InlA and FlaA in facilitating Caco-2 invasion for bacteria grown at either 30°C or 37°C.

Temperature-dependent contributions of transcriptional regulators σ^B and PrfA to Caco-2 invasion

To determine growth temperature effects on the contributions of σ^B and PrfA to Caco-2 invasion, $\Delta prfA$, $\Delta sigB$, and $\Delta sigB\Delta prfA$ strains grown to early stationary phase at 30°C or


37°C were used for Caco-2 invasion assays (performed at 37°C). After growth at 30°C, the invasion efficiency of $\Delta prfA$ was 2.737%, which was not significantly different ($p > 0.05$; Tukey HSD) than the parent strain (2.064% invasion efficiency); the $\Delta sigB$ (0.376% invasion) and $\Delta sigB\Delta prfA$ (0.231% invasion) strains were significantly less invasive ($p < 0.05$; Tukey HSD) than the parent strain (Fig. 3). These results indicate that only σ^B , not PrfA, is involved in invasion when *L. monocytogenes* is grown at 30°C. When the bacteria were grown at 37°C, $\Delta prfA$ (0.322%), $\Delta sigB$ (0.060%), and $\Delta sigB\Delta prfA$ (0.018%) had significantly lower invasion efficiencies ($p < 0.05$; Tukey HSD) than the parent strain (0.680%) (Fig. 3); the $\Delta sigB\Delta prfA$ strain also showed lower invasion efficiency than the parent strain and the $\Delta prfA$ strain ($p < 0.05$; Tukey HSD). A two-way ANOVA also showed a significant “sigB*prfA” interaction effect, indicating a synergism between σ^B and PrfA in regulating Caco-2 invasion for *L. monocytogenes* grown at 37°C (Table 3).

Temperature-dependent regulation of genes involved in Caco-2 invasion

To determine the effects of growth temperature on σ^B and PrfA-dependent regulation of *L. monocytogenes* genes involved in invasion, we determined transcript levels for *inlA*, *flaA*, *plcA*, *gadA*, *sigB*, *prfA*, and *rpoB* in 10403S, and $\Delta sigB$, $\Delta prfA$, and $\Delta sigB\Delta prfA$ grown to early stationary phase at 30°C or 37°C. Neither *prfA* nor *sigB* transcript levels differed between the parent strain grown at 30°C and 37°C ($p > 0.05$, *t*-test). The transcript levels for the σ^B -dependent *gadA* and the PrfA-dependent *plcA* (measured as indicators of σ^B and PrfA activity, respectively) (Fig. 4) also did not differ significantly between bacteria grown at 30°C and 37°C ($p > 0.05$; *t*-test), suggesting no differences in PrfA and σ^B activity between *L. monocytogenes* 10403S grown to early stationary phase in BHI at these two temperatures.

For bacteria grown to early stationary phase at 30°C, *inlA* transcript levels were 10 times lower in the $\Delta sigB$ strain than the parent strain ($p < 0.05$; Tukey HSD; Fig. 4), whereas there was no significant effect of the *prfA* deletion on *inlA* transcript levels ($p = 0.1282$; two-way ANOVA) (Supplemental Table S2,

FIG. 2. Caco-2 invasion efficiencies of *L. monocytogenes* 10403S, $\Delta inlA$, $\Delta flaA$, and $\Delta inlA\Delta flaA$ strains grown to early stationary phase with aeration (i.e., shaking at 220 rpm) at either 30°C or 37°C. Data represent the mean of four biological replicates. Error bars represent SDs. Strains with different letters had significantly different invasion efficiencies at a given growth temperature ($p < 0.05$; *post hoc* Tukey HSD); lower invasion for the $\Delta flaA$ strain than the parent strain (for growth at 37°C) was only borderline significant ($p = 0.0279$; Tukey HSD). Two-way ANOVA was performed within each temperature to determine effects of single and double mutations on invasion efficiency (see Table 3 for *p*-values). HSD, honestly significantly different.






FIG. 3. Caco-2 invasion efficiencies of *L. monocytogenes* 10403S, $\Delta prfA$, $\Delta sigB$, and $\Delta sigB\Delta prfA$ strains grown to early stationary phase with aeration (i.e., shaking at 220 rpm) at either 30°C or 37°C. Data for the parent strain (10403S) are the same as those shown in Figure 2. Data represent the mean of four biological replicates. Error bars represent SD. Strains with different letters have significantly different invasion efficiencies at a given growth temperature ($p < 0.05$; *post hoc* Tukey HSD). Two-way ANOVA was performed within each temperature to determine effects of single and double mutations on invasion efficiency (see Table 3 for p -values).

available online at www.liebertonline.com). These data indicate that *inlA* transcription is σ^B dependent, but PrfA independent under these conditions. For *L. monocytogenes* grown at 37°C, *inlA* transcript levels were significantly lower in $\Delta sigB$ than in the parent strain ($p < 0.05$; ANOVA; Tukey HSD; Fig. 4), whereas *inlA* transcript levels were not significantly different in $\Delta prfA$ compared to the parent strain ($p > 0.05$; Tukey HSD; Fig. 4). However, the $sigB^*prfA$ interaction effect on *inlA* transcript levels was borderline significant ($p = 0.0799$; two-way ANOVA; Supplemental Table S2), suggesting contributions of both σ^B and PrfA to regulation of *inlA* transcription in *L. monocytogenes* grown at 37°C.

In the *L. monocytogenes* parent strain, *flaA* transcript levels were higher after growth at 30°C (1.26; standard deviation = 0.07) than at 37°C (1.104; standard deviation = 0.05) ($p = 0.0014$, *t*-test; data shown in Fig. 4). Although the differ-

ence is small, these results indicate at least some temperature-dependent regulation of *flaA* transcription in 10403S. *flaA* transcript levels were not significantly different between $\Delta prfA$ and the parent strain for either growth temperature ($p > 0.05$; Tukey HSD; Fig. 4). In *L. monocytogenes* grown at 37°C, *flaA* transcript levels were not different between the $\Delta sigB$ strain and the parent strain ($p > 0.05$; Tukey HSD; Fig. 4). However, at 30°C *flaA* transcription was significantly higher in $\Delta sigB$ and $\Delta sigB\Delta prfA$ than in the parent strain ($p < 0.05$; ANOVA, Tukey HSD; Fig. 4), and two-way ANOVA showed a significant “*sigB* effect” on *flaA* transcript levels ($p = 0.0003$; Supplemental Table S2). These results are consistent with other studies that suggest a role of σ^B in down-regulating chemotaxis genes (Raengpradub *et al.*, 2008; Toledo-Arana *et al.*, 2009).

Discussion

L. monocytogenes grown at 30°C shows a higher Caco-2 invasion efficiency than bacteria grown at 37°C

Our data showed that, with the exception of mutants with deletions of motility-related genes, all *L. monocytogenes* strains showed higher Caco-2 invasion efficiency when grown at 30°C than at 37°C. This finding supports previous work (Dons *et al.*, 2004) reporting that a different *L. monocytogenes* strain (12067) showed about 10-fold increased association with Caco-2 cells and twofold increased invasion of Caco-2 cells for bacteria grown at 24°C (where 12067 was motile) compared to 37°C (where this strain was not motile); this previous study did not include a statistical evaluation of these differences in invasion efficiencies (Dons *et al.*, 2004). As most *L. monocytogenes*, including strain 10403S, show increased flagellar motility when grown at 30°C, a trait that contributes to Caco-2 cell invasion (Dons *et al.*, 2004) and intestinal colonization in mice (O’Neil and Marquis, 2006), expression of motility genes at $\leq 30^\circ\text{C}$ may explain the increased Caco-2 invasion of *L. monocytogenes* grown at $\leq 30^\circ\text{C}$ compared to bacteria grown at 37°C.

Interestingly, other bacterial pathogens have also been shown to differ in their invasiveness and virulence depending

TABLE 3. EFFECTS OF VARIOUS GENE DELETIONS ON *LISTERIA MONOCYTOGENES* CACO-2 INVASION EFFICIENCY

Deletion variable ^a	<i>p</i> -Values from two-way ANOVA for <i>L. monocytogenes</i> grown at	
	30°C	37°C
<i>sigB</i>	0.0001 ^b	<0.0001 ^b
<i>prfA</i>	0.4401	0.0014 ^c
<i>sigB</i> * <i>prfA</i>	0.226	0.0057 ^c
<i>inlA</i>	0.001 ^b	<0.0001 ^b
<i>flaA</i>	0.0009 ^b	0.0256 ^d
<i>inlA</i> * <i>flaA</i>	0.0015 ^c	0.0269 ^d

^aThe variables listed in this column represent either single-gene deletions (e.g., “*sigB*”) or interactions between two gene deletions (e.g., “*sigB***prfA*”). The p -values for the single-gene deletions measure the individual effect of deleting each respective gene. The “gene*gene” variable measures synergistic deletion effects by comparing the effect of deleting both genes to the effect of deleting either one gene or the other; significant values are marked with ^b $p \leq 0.001$, ^c $p \leq 0.01$, or ^d $p \leq 0.05$. The actual data used for these analyses are presented in Figure 2 (*inlA* and *flaA* data) and Figure 3 (*sigB* and *prfA* data).




FIG. 4. Normalized transcript levels of four genes in *L. monocytogenes* 10403S, $\Delta sigB$, $\Delta prfA$, and $\Delta sigB\Delta prfA$ grown to early stationary phase with aeration (i.e., shaking at 220 rpm) at 30°C or 37°C. Data presented as log₁₀ (target gene mRNA level/rpoB mRNA level). Bars represent the average of four biological replicates and error bars represent SD. Strains with different letters (e.g., A, B) had significantly different transcript levels ($p < 0.05$; *post hoc* Tukey HSD). Two-way ANOVA was performed for each gene at each temperature to determine effects of single and double mutations on transcript levels (see Supplemental Table S2 for p -value summary). For each gene, a t -test was used to compare transcript levels in 10403S between 30°C and 37°C. Only *flaA* levels were significantly different between temperatures ($p = 0.0014$; t -test; indicated by *).

on growth temperature (Konkel and Tilly, 2000). While in a number of bacterial pathogens virulence genes have been found to be more highly expressed at 37°C than at lower temperatures (Maurelli, 1989), some pathogens have been shown to express invasion factors at higher levels when grown at temperatures below 37°C. For example, *Yersinia pseudotuberculosis invA*, which encodes an invasin required for host cell invasion, appears to be expressed at higher levels in bacteria grown at 28°C than at 37°C (based on visual examination of Western blot data) (Isberg *et al.*, 1988). *Yersinia enterocolitica* has also been shown to express a motility phenotype when grown below 37°C, whereas motility, which may play a role in initiation of host cell invasion, is down-regulated in bacteria grown at 37°C (Young *et al.*, 2000). Therefore, adaptation to environments outside the host may increase the virulence potential of *L. monocytogenes* and other bacterial pathogens.

Enteric pathogens transmitted to humans from food and environmental sources often experience a sudden change in environmental temperature when they are ingested. Our results show that invasion phenotypes of the parent strain and

the motility mutants grown at 30°C persisted for 2.5 h after the bacteria were switched to 37°C. As 70% to 90% of human stomach contents are emptied after 2 h (Bennink *et al.*, 1999), our findings, in combination with other studies (Dons *et al.*, 2004; O'Neil and Marquis, 2006), suggest that *L. monocytogenes* grown at temperatures that permit motility [i.e., 12°C–30°C (Di Bonaventura *et al.*, 2008)] may have increased invasion potential that could be maintained during gastric passage. As regulation of motility appears to differ considerably between *L. monocytogenes* strains (Grundling *et al.*, 2004; Way *et al.*, 2004), further studies will need to use different strains to validate our findings on growth temperature dependence of Caco-2 invasion efficiency.

In addition to growth temperature, other studies (Garner *et al.*, 2006a; Andersen *et al.*, 2007) have shown that exposure to other environmental conditions (e.g., organic acids, anaerobic conditions) appear to affect the virulence potential of *L. monocytogenes*. For example *L. monocytogenes* grown in an oxygen-limited environment displayed 100-fold increased invasion efficiency in Caco-2 cells as well as increased virulence in a guinea pig model, including 10 to

100-fold higher *L. monocytogenes* fecal shedding levels, compared to numbers for bacteria grown under aerobic conditions (Andersen *et al.*, 2007). Further, *L. monocytogenes* grown in the presence of sodium lactate or NaCl showed an about 10-fold higher invasion efficiency than bacteria grown without these compounds (Garner *et al.*, 2006a). Overall, growth temperatures as well as other preinvasion environmental conditions (e.g., anaerobiosis) thus appear to affect the regulation of genes with roles in host attachment and invasion across different environmentally transmitted pathogens, including *L. monocytogenes*.

CheY and *DegU* significantly contribute to invasion in *L. monocytogenes* grown $\leq 30^\circ\text{C}$

Our data, in conjunction with other reports (e.g., Dons *et al.*, 2004), clearly show that *DegU* and *CheY* play a temperature-dependent role in enhancing invasion of host cells across different *L. monocytogenes* strain backgrounds. For example, $\Delta cheA$, $\Delta cheY$, and $\Delta cheYA$ mutants in an *L. monocytogenes* 12067 background all showed about 100-fold reduced Caco-2 cell invasion as well as reduced motility as compared to their parent strain after growth at 24°C (Dons *et al.*, 2004), whereas Williams *et al.* (2005a) found that *L. monocytogenes* EGD $\Delta degU$ and $\Delta cheY$ strains, grown at 37°C , did not show reduced invasion efficiency of Cos-1 fibroblast cells. Previous studies have shown increased swarming and flagellar motility of *L. monocytogenes* grown at 24°C compared to *L. monocytogenes* grown at 37°C (Dons *et al.*, 2004; Knudsen *et al.*, 2004; Shen and Higgins, 2006), due to repression of *L. monocytogenes* motility genes at 37°C (Grundling *et al.*, 2004; Shen and Higgins, 2006). Therefore, contributions of motility genes to virulence phenotypes are generally apparent only when *L. monocytogenes* is grown at temperatures where the organism is typically motile (i.e., $\leq 30^\circ\text{C}$) (Dons *et al.*, 2004; Knudsen *et al.*, 2004; Williams *et al.*, 2005b; O'Neil and Marquis, 2006). While *DegU* has also been shown to contribute to motility in *L. monocytogenes* grown at 24°C and to virulence in a mouse model (Williams *et al.*, 2005b), we are not aware of any previous studies reporting a specific effect of a *degU* null mutation on invasion of human intestinal epithelial cells.

While we found no contributions of other RRs, besides *DegU* and *CheY*, to invasion of Caco-2 cells, some studies have shown contributions of other RRs to virulence phenotypes in *L. monocytogenes* strains other than 10403S and in other invasion models. For example, Williams *et al.* (2005a) reported that RRs Lmo1507 and LisR are involved in invasion of Cos-1 cells. A deletion of *lisK* (encoding the LisR-associated sensory kinase) in *L. monocytogenes* LO28 was also found to affect virulence in mice (Cotter *et al.*, 1999), further supporting contributions of LisRK to *L. monocytogenes* virulence in some models. Similarly, *L. monocytogenes* EGD with a deletion of the RR VirR, grown at 37°C , has also been reported to be deficient in Caco-2 cell invasion (Mandin *et al.*, 2005), even though the *L. monocytogenes* 10403S $\Delta virR$ strain used in our current study did not show evidence for reduced invasion, again possibly reflecting strain differences. Overall, different TCS thus appears to contribute to *L. monocytogenes* virulence and virulence-associated characteristics, even though contributions appear to differ based on growth conditions, strain backgrounds, and assays used.

While InlA and FlaA show significant synergism in their contributions to Caco-2 invasion, synergism is less pronounced after growth at 37°C

While flagellin has been identified as a critical part of the motility machinery involved in Caco-2 invasion in *L. monocytogenes* grown at temperatures that allow motility gene expression (i.e., $\leq 30^\circ\text{C}$) (Dons *et al.*, 2004; O'Neil and Marquis, 2006), we found flagellin and *InlA* contribute synergistically to Caco-2 cell invasion in *L. monocytogenes* 10403S grown at 30°C and 37°C . While studies in other *L. monocytogenes* strains (e.g., 12067, EDGe) have shown that contributions of motility factors to host cell invasion typically are only apparent in *L. monocytogenes* grown at 30°C or less (Dons *et al.*, 2004; Shen and Higgins, 2006), some studies (Grundling *et al.*, 2004; Way *et al.*, 2004) have shown increased flagellar motility at 37°C for strain 10403S as compared to other strains. For example, *flaA* repression by MogR was shown to be less stringent in 10403S than in EDGe (Grundling *et al.*, 2004), and 10403S activated a flagellum-dependent innate immune response even after growth at 37°C (Way *et al.*, 2004). Critical contributions of *InlA* to invasion of human intestinal epithelial cells (Dramsai *et al.*, 1993; Lingnau *et al.*, 1995) and virulence after oral infection (Lecuit *et al.*, 1999; Garner *et al.*, 2006b) have been well established for host species carrying the E-cadherin allotype that allows for *InlA* binding (e.g., humans and guinea pigs). Although we observed a synergism between the contributions of *FlaA* and *InlA* to Caco-2 invasion in *L. monocytogenes* grown at both 30°C and 37°C , we also found that the $\Delta inlA$ strain shows increased invasion efficiency when grown at 30°C compared to 37°C , suggesting *InlA*-independent contributions of flagellar motility to Caco-2 cell invasion. This observation further supports the importance of flagellar motility in *L. monocytogenes* virulence.

σ^B , but not *PrfA*, contributes to invasion of bacteria grown at 30°C , whereas σ^B and *PrfA* show synergistic contributions to invasion if bacteria are grown at 37°C

Our data showed that (i) σ^B contributes to Caco-2 invasion for *L. monocytogenes* grown at both 30°C and 37°C and (ii) σ^B positively regulates *inlA* transcription at both of these temperatures. The observation that σ^B contributes to Caco-2 invasion in bacteria grown at both temperatures is consistent with a number of studies that have shown that σ^B is critical for invasion of intestinal cells *in vitro* (Kim *et al.*, 2004; Garner *et al.*, 2006b) and in a guinea pig model of listeriosis (Garner *et al.*, 2006b). While McGann *et al.* (2007a) found higher transcript levels in *L. monocytogenes* grown at 30°C than bacteria grown at 37°C , for certain σ^B -dependent internalin genes (i.e., *inlC2*, *inlD*, *lmo331*, and *lmo0610*) and *opuCA*, which has been shown to be regulated by both σ^A and σ^B (Cetin *et al.*, 2004; Chan *et al.*, 2007a), the same study also found that *inlA* transcript levels were similar in *L. monocytogenes* grown at 30°C and 37°C (McGann *et al.*, 2007a). Another study also reported that genes found to be σ^B dependent in the intestine were generally not differentially expressed at 30°C and 37°C (Toledo-Arana *et al.*, 2009). Overall, these data suggest that σ^B activity and σ^B -dependent regulation of invasion is similar in *L. monocytogenes* grown at 30°C and 37°C , even though some σ^B -dependent genes (e.g., *opuCA* and *inlC2D*) may be regulated by additional temperature-dependent mechanisms (McGann *et al.*, 2007a; Chan *et al.*, 2007a). We did find though that σ^B -dependent

negative regulation of *flaA* transcript levels, which was previously described by two studies (Raengpradub *et al.*, 2008; Toledo-Arana *et al.*, 2009), was only apparent here in bacteria grown at 30°C, even though this regulation has previously been reported in bacteria exposed to salt stress at 37°C (Raengpradub *et al.*, 2008). Temperature effects on σ^B -dependent regulation of some genes may thus be dependent on other environmental conditions (e.g., osmotic stress).

Contributions of PrfA to Caco-2 cell invasion in *L. monocytogenes* grown at 37°C, but not in bacteria grown at 30°C, are consistent with previous findings of low PrfA activity in *L. monocytogenes* grown at temperatures <37°C (Leimeister-Wachter *et al.*, 1992) as well as data showing that PrfA-dependent *inlA* expression is significantly higher at 37°C than at 25°C (Dramsi *et al.*, 1993). Interestingly, in our study here, *prfA* transcript levels and PrfA activity (as measured by *plcA* transcription levels) were not significantly different in the parent strain between 30°C and 37°C growth conditions; this observation may reflect low baseline PrfA activity in *L. monocytogenes* grown at 37°C in BHI (i.e., the conditions used here) as low levels of easily catabolized sugar and/or presence of other compounds (e.g., charcoal) seem to be required to induce PrfA activity at 37°C (Ripio *et al.*, 1996; Milenbachs *et al.*, 1997; Gilbreth *et al.*, 2004). While PrfA activity and PrfA-dependent phenotypes are thus clearly temperature dependent, with maximum PrfA activity in bacteria grown under certain conditions at 37°C, it is increasingly clear that transcriptional patterns and phenotypic characteristics of *L. monocytogenes* are governed by complex, environmental-condition-dependent interactions between multiple regulators.

Synergisms between PrfA and σ^B were confirmed here through formal statistical analyses that showed (i) a statistically significant interaction effect between *sigB* and *prfA* deletions on Caco-2 invasion in bacteria grown at 37°C, but not in bacteria grown at 30°C and (ii) a borderline significant interaction effect between *sigB* and *prfA* deletions on *inlA* transcript levels in bacteria grown at 37°C, but not at 30°C. While this type of temperature-dependent synergism has not previously been described, contributions of both σ^B and PrfA to Caco-2 cell invasion and co-regulation of *L. monocytogenes* virulence genes, including *inlA*, have been reported previously (Lingnau *et al.*, 1995; Kazmierczak *et al.*, 2003; Sue *et al.*, 2004; Kim *et al.*, 2005; McGann *et al.*, 2007b). The observation that *flaA* also appears to be downregulated by σ^B , possibly with an antisense RNA type mechanism (Toledo-Arana *et al.*, 2009), further supports a temperature-dependent regulatory network involving PrfA and σ^B that affects multiple effector proteins contributing to *L. monocytogenes* invasion and virulence, even though possible (temperature-dependent) contributions of PrfA itself to transcription of *flaA* and other motility regulated genes (Michel *et al.*, 1998; Milohanic *et al.*, 2003) will require further confirmation. For example, while others (Ripio *et al.*, 1997) previously reported (based on visual examination of an RNA slot blot) negative regulation of *flaA* transcription by PrfA* (i.e., a PrfA protein that is constitutively active) in *L. monocytogenes* grown at 20°C, we did not find any effect of the *prfA* deletion on *flaA* transcript levels.

Conclusions

Overall, our data show that *L. monocytogenes* use a number of regulatory mechanisms to modulate virulence gene ex-

pression, particularly expression of genes important for invasion, under different temperatures. Specifically, modulation of gene expression in *L. monocytogenes* grown at temperatures <37°C (mimicking environmental conditions before host infection) appears to increase the invasiveness of this pathogen, priming it for subsequent infection of a mammalian host. Importantly, our data also suggest an initial specific model for regulation of key invasion-associated genes at transition of *L. monocytogenes* from environment to host. During growth at temperatures below mammalian body temperatures, CheY and DegU activity induce a motility phenotype, and although PrfA-dependent virulence gene expression is minimal, *InlA* expression is assured by σ^B -dependent *inlA* transcription. After introduction into the host environment, passage through the gastrointestinal system can further activate σ^B (e.g., through acid and osmotic stress), priming the cell for intestinal cell invasion through increased *inlA* transcription (Sue *et al.*, 2004). Once *L. monocytogenes* enter the intracellular environments, PrfA-dependent gene expression becomes critical for intracellular survival and spread (Freitag *et al.*, 1993) with σ^B taking on a modulating role by downregulating expression of genes encoding cytolysins, which may cause excessive host cell damage (Ollinger *et al.*, 2008). Concurrently, motility appears to be downregulated in *L. monocytogenes* grown at 37°C, including through σ^B -dependent mechanisms (Raengpradub *et al.*, 2008; Toledo-Arana *et al.*, 2009), possibly facilitating evasion of Toll-like-receptor-mediated host responses (Hayashi *et al.*, 2001; Torres *et al.*, 2004).

Acknowledgments

We would like to acknowledge Dr. H el ene Marquis for donation of the Δ *flaA* and Δ *inlA Δ *flaA* strains, and thank Sara Milillo, Courtney Lucas Stelling, Karlyn Beer, and Wan-Lin Su for help with plasmid construction, as well as the members of the Wiedmann and Boor laboratories at Cornell University for their support and suggestions. This work was funded by USDA Special Research Grants 2003-34459-12999 and 2004-34459-14296 (to M.W.) and NIH-NIAID (R01 AI052151 to K.J.B.). Reid A. Ivy was supported by the USDA-CSREES Food and Agricultural Sciences National Needs Graduate Fellowship Grants Program (2005-38420-15776).*

Disclosure Statement

No competing financial interests exist.

References

- Andersen JB, Roldgaard B, Christensen B, and Licht T. Oxygen restriction increases the infective potential of *Listeria monocytogenes* *in vitro* in Caco-2 cells and *in vivo* in guinea pigs. *BMC Microbiol* 2007;7:55.
- Autret N, Raynaud C, Dubail I, Berche P, and Charbit A. Identification of the *agr* locus of *Listeria monocytogenes*: role in bacterial virulence. *Infect Immun* 2003;71:4463–4471.
- Bakardjiev AI, Stacy BA, Fisher SJ, and Portnoy DA. Listeriosis in the pregnant guinea pig: a model of vertical transmission. *Infect Immun* 2004;72:489–497.
- Becker LA, Evans SN, Hutkins RW, and Benson AK. Role of Sigma B in adaptation of *Listeria monocytogenes* to growth at low temperature. *J Bacteriol* 2000;182:7083–7087.

- Bennink R, Peeters M, Van den Maegdenbergh V, Geypens B, Rutgeerts P, De Roo M, and Mortelmans L. Evaluation of small-bowel transit for solid and liquid test meal in healthy men and women. *Eur J Nucl Med Mol Imaging* 1999;26:1560–1566.
- Bishop DK and Hinrichs DJ. Adoptive transfer of immunity to *Listeria monocytogenes*. The influence of *in vitro* stimulation on lymphocyte subset requirements. *J Immunol* 1987;139:2005–2009.
- Brondsted L, Kallipolitis BH, Ingmer H, and Knochel S. *kdpE* and a putative RsbQ homologue contribute to growth of *Listeria monocytogenes* at high osmolarity and low temperature. *FEMS Microbiol Lett* 2003;219:233–239.
- Camilli A, Tilney LG, and Portnoy DA. Dual roles of *plcA* in *Listeria monocytogenes* pathogenesis. *Mol Microbiol* 1993;8:143–157.
- Cetin MS, Zhang C, Hutkins RW, and Benson AK. Regulation of transcription of compatible solute transporters by the general stress sigma factor, Sigma B, in *Listeria monocytogenes*. *J Bacteriol* 2004;186:794–802.
- Chan YC, Boor KJ, and Wiedmann M. Sigma B-dependent and -independent mechanisms contribute to transcription of *Listeria monocytogenes* cold stress genes during cold shock and cold growth. *Appl Environ Microbiol* 2007a;73:6019–6029.
- Chan YC, Raengpradub S, Boor KJ, and Wiedmann M. Microarray-based characterization of the *Listeria monocytogenes* cold regulon in log- and stationary-phase cells. *Appl Environ Microbiol* 2007b;73:6484–6498.
- Chaturongakul S and Boor KJ. Sigma B activation under environmental and energy stress conditions in *Listeria monocytogenes*. *Appl Environ Microbiol* 2006;72:5197–5203.
- Cheng LW and Portnoy DA. *Drosophila* S2 cells: an alternative infection model for *Listeria monocytogenes*. *Cell Microbiol* 2003;5:875–885.
- Cotter PD, Emerson N, Gahan CGM, and Hill C. Identification and disruption of *lisRK*, a genetic locus encoding a two-component signal transduction system involved in stress tolerance and virulence in *Listeria monocytogenes*. *J Bacteriol* 1999;181:6840–6843.
- Di Bonaventura G, Piccolomini R, Paludi D, D'Orio V, Vergara A, Conter M, and Ianieri A. Influence of temperature on biofilm formation by *Listeria monocytogenes* on various food-contact surfaces: relationship with motility and cell surface hydrophobicity. *J Appl Microbiol* 2008;104:1552–1561.
- Dons L, Eriksson E, Jin Y, Rottenberg ME, Kristensson K, Larsen CN, Bresciani J, and Olsen JE. Role of flagellin and the two-component CheA/CheY system of *Listeria monocytogenes* in host cell invasion and virulence. *Infect Immun* 2004;72:3237–3244.
- Dramsi S, Kocks C, Forestier C, and Cossart P. Internalin-mediated invasion of epithelial cells by *Listeria monocytogenes* is regulated by the bacterial growth state, temperature and the pleiotropic activator PrfA. *Mol Microbiol* 1993;9:931–941.
- Freitag NE, Rong L, and Portnoy DA. Regulation of the PrfA transcriptional activator of *Listeria monocytogenes*: multiple promoter elements contribute to intracellular growth and cell-to-cell spread. *Infect Immun* 1993;61:2537–2544.
- Garner MR, James KE, Callahan MC, Wiedmann M, and Boor KJ. Exposure to salt and organic acids increases the ability of *Listeria monocytogenes* to invade Caco-2 cells but decreases its ability to survive gastric stress. *Appl Environ Microbiol* 2006a;72:5384–5395.
- Garner MR, Njaa BL, Wiedmann M, and Boor KJ. Sigma B contributes to *Listeria monocytogenes* gastrointestinal infection but not to systemic spread in the guinea pig infection model. *Infect Immun* 2006b;74:876–886.
- Gilbreth SE, Benson AK, and Hutkins RW. Catabolite repression and virulence gene expression in *Listeria monocytogenes*. *Curr Microbiol* 2004;49:95–98.
- Grundling A, Burrack LS, Bouwer HGA, and Higgins DE. From the cover: *Listeria monocytogenes* regulates flagellar motility gene expression through MogR, a transcriptional repressor required for virulence. *PNAS* 2004;101:12318–12323.
- Hayashi F, Smith KD, Ozinsky A, Hawn TR, Yi EC, Goodlett DR, Eng JK, Akira S, Underhill DM, and Aderem A. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature* 2001;410:1099–1103.
- Isberg RR, Swain A, and Falkow S. Analysis of expression and thermoregulation of the *Yersinia pseudotuberculosis* *inv* gene with hybrid proteins. *Infect Immun* 1988;56:2133–2138.
- Johansson J, Mandin P, Renzoni A, Chiaruttini C, Springer M, and Cossart P. An RNA thermosensor controls expression of virulence genes in *Listeria monocytogenes*. *Cell* 2002;110:551–561.
- Kallipolitis BH and Ingmer H. *Listeria monocytogenes* response regulators important for stress tolerance and pathogenesis. *FEMS Microbiol Lett* 2001;204:111–115.
- Kallipolitis BH, Ingmer H, Gahan CG, Hill C, and Sogaard-Andersen L. CesRK, a two-component signal transduction system in *Listeria monocytogenes*, responds to the presence of cell wall-acting antibiotics and affects [beta]-lactam resistance. *Antimicrob Agents Chemother* 2003;47:3421–3429.
- Kazmierczak MJ, Mithoe SC, Boor KJ, and Wiedmann M. *Listeria monocytogenes* Sigma B regulates stress response and virulence functions. *J Bacteriol* 2003;185:5722–5734.
- Kazmierczak MJ, Wiedmann M, and Boor KJ. Contributions of *Listeria monocytogenes* Sigma B and PrfA to expression of virulence and stress response genes during extra- and intracellular growth. *Microbiology* 2006;152:1827–1838.
- Kim H, Boor KJ, and Marquis H. *Listeria monocytogenes* Sigma B contributes to invasion of human intestinal epithelial cells. *Infect Immun* 2004;72:7374–7378.
- Kim H, Marquis H, and Boor KJ. Sigma B contributes to *Listeria monocytogenes* invasion by controlling expression of *inlA* and *inlB*. *Microbiology* 2005;151:3215–3222.
- Knudsen GM, Olsen JE, and Dons L. Characterization of DegU, a response regulator in *Listeria monocytogenes*, involved in regulation of motility and contributes to virulence. *FEMS Microbiol Lett* 2004;240:171–179.
- Konkel ME and Tilly K. Temperature-regulated expression of bacterial virulence genes. *Microb Infect* 2000;2:157–166.
- Larsen MH, Kallipolitis BH, Christiansen JK, Olsen JE, and Ingmer H. The response regulator ResD modulates virulence gene expression in response to carbohydrates in *Listeria monocytogenes*. *Mol Microbiol* 2006;61:1622–1635.
- Lecuit M, Dramsi S, Gottardi C, Fedor-Chaikin M, Gumbiner B, and Cossart P. A single amino acid in E-cadherin responsible for host specificity towards the human pathogen *Listeria monocytogenes*. *EMBO J* 1999;18:3956–3963.
- Leimeister-Wachter M, Domann E, and Chakraborty T. The expression of virulence genes in *Listeria monocytogenes* is thermoregulated. *J Bacteriol* 1992;174:947–952.
- Lingnau A, Domann E, Hudel M, Bock M, Nichterlein T, Wehland J, and Chakraborty T. Expression of the *Listeria monocytogenes* EGD *inlA* and *inlB* genes, whose products mediate bacterial entry into tissue culture cell lines, by PrfA-dependent and -independent mechanisms. *Infect Immun* 1995;63:3896–3903.

- Liu S, Graham JE, Bigelow L, Morse PD II, and Wilkinson BJ. Identification of *Listeria monocytogenes* genes expressed in response to growth at low temperature. *Appl Environ Microbiol* 2002;68:1697–1705.
- Mandin P, Fsihi H, Dussurget O, Vergassola M, Milohanic E, Toledo-Arana A, Lasa I, Johansson J, and Cossart P. VirR, a response regulator critical for *Listeria monocytogenes* virulence. *Mol Microbiol* 2005;57:1367–1380.
- Mauder N, Williams T, Fritsch F, Kuhn M, and Beier D. Response regulator DegU of *Listeria monocytogenes* controls temperature-responsive flagellar gene expression in its unphosphorylated state. *J Bacteriol* 2008;190:4777–4781.
- Maurelli AT. Temperature regulation of virulence genes in pathogenic bacteria: a general strategy for human pathogens? *Microb Pathog* 1989;7:1–10.
- McCann P, Ivanek R, Wiedmann M, and Boor KJ. Temperature-dependent expression of *Listeria monocytogenes* internalin and internalin-like genes suggests functional diversity of these proteins among the *Listeriae*. *Appl Environ Microbiol* 2007a;73:2806–2814.
- McCann P, Raengpradub S, Ivanek R, Wiedmann M, and Boor KJ. Differential regulation of *Listeria monocytogenes* internalin and internalin-like genes by Sigma B and PrfA as revealed by subgenomic microarray analyses. *Foodborne Pathog Dis* 2008;5:417–435.
- McCann P, Wiedmann M, and Boor KJ. The alternative sigma factor Sigma B and the virulence gene regulator PrfA both regulate transcription of *Listeria monocytogenes* internalins. *Appl Environ Microbiol* 2007b;73:2919–2930.
- Mead PS, Slutsker L, Dietz V, McCaig LF, Bresee JS, Shapiro C, Griffin PM, and Tauxe RV. Food-related illness and death in the United States. *Emerg Infect Dis* 1999;5:607–625.
- Michel E, Mengaud J, Galsworthy S, and Cossart P. Characterization of a large motility gene cluster containing the *cheR*, *motAB* genes of *Listeria monocytogenes* and evidence that PrfA downregulates motility genes. *FEMS Microbiol Lett* 1998;169:341–347.
- Milenbachs AA, Brown DP, Moors M, and Youngman P. Carbon-source regulation of virulence gene expression in *Listeria monocytogenes*. *Mol Microbiol* 1997;23:1075–1085.
- Milohanic E, Glaser P, Coppee J-Y, Frangeul L, Vega Y, Vazquez-Boland JA, Kunst F, Cossart P, and Buchrieser C. Transcriptome analysis of *Listeria monocytogenes* identifies three groups of genes differently regulated by PrfA. *Mol Microbiol* 2003;47:1613–1625.
- Ollinger J, Wiedmann M, and Boor KJ. SigB- and PrfA-dependent transcription of genes previously classified as putative constituents of the *Listeria monocytogenes* PrfA Regulon. *Foodborne Pathog Dis* 2008;5:281–293.
- O'Neil HS and Marquis H. *Listeria monocytogenes* flagella are used for motility, not as adhesins, to increase host cell invasion. *Infect Immun* 2006;74:6675–6681.
- Pal A, Labuza TP, and Diez-Gonzalez F. Comparison of primary predictive models to study the growth of *Listeria monocytogenes* at low temperatures in liquid cultures and selection of fastest growing ribotypes in meat and turkey product slurries. *Food Microbiol* 2008;25:460–470.
- Raengpradub S, Wiedmann M, and Boor KJ. Comparative analysis of the Sigma B-dependent stress responses in *Listeria monocytogenes* and *Listeria innocua* strains exposed to selected stress conditions. *Appl Environ Microbiol* 2008;74:158–171.
- Raffelsbauer D, Bubert A, Engelbrecht F, Scheinpflug J, Simm A, Hess J, Kaufmann SH, and Goebel W. The gene cluster *inIC2DE* of *Listeria monocytogenes* contains additional new internalin genes and is important for virulence in mice. *Mol Gen Genet* 1998;260:144–158.
- Ripio MT, Brehm K, Lara M, Suarez M, and Vazquez-Boland JA. Glucose-1-phosphate utilization by *Listeria monocytogenes* is PrfA dependent and coordinately expressed with virulence factors. *J Bacteriol* 1997;179:7174–7180.
- Ripio MT, DomÍnguez-Bernal G, Suárez M, Brehm K, Berche P, and Vázquez-Boland JA. Transcriptional activation of virulence genes in wild-type strains of *Listeria monocytogenes* in response to a change in the extracellular medium composition. *Res Microbiol* 1996;147:371–384.
- Scotti M, Monzo HJ, Lacharme-Lora L, Lewis DA, and Vazquez-Boland JA. The PrfA virulence regulon. *Microb Infect* 2007;9:1196–1207.
- Shen A and Higgins DE. The MogR transcriptional repressor regulates nonhierarchical expression of flagellar motility genes and virulence in *Listeria monocytogenes*. *PLoS Pathog* 2006;2:0283–0295.
- Shun Z, Silverberg A, Chang C, and Ouyang P. Dunnett's many-to-one test and least square means. *J Biopharm Stat* 2003;13:17–28.
- Sleator RD and Hill C. A novel role for the LisRK two-component regulatory system in listerial osmotolerance. *Clin Microbiol Infect* 2005;11:599–601.
- Smith K and Youngman P. Use of a new integrational vector to investigate compartment-specific expression of the *Bacillus subtilis* *spolM* gene. *Biochimie* 1992;74:705–711.
- Sue D, Boor KJ, and Wiedmann M. Sigma B-dependent expression patterns of compatible solute transporter genes *opuCA* and *lmo1421* and the conjugated bile salt hydrolase gene *bsh* in *Listeria monocytogenes*. *Microbiology* 2003;149:3247–3256.
- Sue D, Fink D, Wiedmann M, and Boor KJ. Sigma B-dependent gene induction and expression in *Listeria monocytogenes* during osmotic and acid stress conditions simulating the intestinal environment. *Microbiology* 2004;150:3843–3855.
- Toledo-Arana A, Dussurget O, Nikitas G, Sesto N, Guet-Revillet H, Balestrino D, Loh E, Gripenland J, Tiensuu T, Vaitkevicius K, Barthelemy M, Vergassola M, Nahori M-A, Soubigou G, Regnault B, Coppee J-Y, Lecuit M, Johansson J, and Cossart P. The *Listeria* transcriptional landscape from saprophytism to virulence. *Nature* 2009;459:950–956.
- Torres D, Barrier M, Bihl F, Quesniaux VJF, Maillat I, Akira S, Ryffel B, and Erard F. Toll-like Receptor 2 is required for optimal control of *Listeria monocytogenes* infection. *Infect Immun* 2004;72:2131–2139.
- van der Veen S, Hain T, Wouters JA, Hossain H, de Vos WM, Abee T, Chakraborty T, and Wells-Bennik MHJ. The heat-shock response of *Listeria monocytogenes* comprises genes involved in heat shock, cell division, cell wall synthesis, and the SOS response. *Microbiology* 2007;153:3593–3607.
- Way SS, Thompson LJ, Lopes JE, Hajjar AM, Kollmann TR, Freitag NE, and Wilson CB. Characterization of flagellin expression and its role in *Listeria monocytogenes* infection and immunity. *Cell Microbiol* 2004;6:235–242.
- Wiedmann M, Arvik TJ, Hurley RJ, and Boor KJ. General stress transcription factor Sigma B and its role in acid tolerance and virulence of *Listeria monocytogenes*. *J Bacteriol* 1998;180:3650–3656.
- Williams T, Bauer S, Beier D, and Kuhn M. Construction and characterization of *Listeria monocytogenes* mutants with in-

frame deletions in the response regulator genes identified in the genome sequence. *Infect Immun* 2005a;73:3152–3159.

Williams T, Joseph B, Beier D, Goebel W, and Kuhn M. Response regulator DegU of *Listeria monocytogenes* regulates the expression of flagella-specific genes. *FEMS Microbiol Lett* 2005b;252:287–298.

Young GM, Badger JL, and Miller VL. Motility is required to initiate host cell invasion by *Yersinia enterocolitica*. *Infect Immun* 2000;68:4323–4326.

Address correspondence to:
Martin Wiedmann, Dr. med. vet., Ph.D.
Department of Food Science
Cornell University
412 Stocking Hall
Ithaca, NY 14853

E-mail: mw16@cornell.edu

