

LETTER TO THE EDITOR

ARF1 Localizes to the Golgi and the *Trans*-Golgi Network

The recruitment of coat proteins for transport vesicles (COPI-, COPII-, and clathrin-coated) is mediated by the small GTPases of the ADP-ribosylation factor (ARF) family of which there are three *SAR*, 12 *ARF*, and six *ARL* genes in the *Arabidopsis thaliana* genome (Vernoud et al., 2003). These GTPases are themselves recruited by guanidine exchange factors located at the donor compartments, which convert them into active GTP-bound forms (Anders and Jürgens, 2008). The well-established inhibitor of ARF guanidine exchange factors is the fungal toxin brefeldin A (Robinson et al., 2008). Normally, the ARF-GTPases are located on membranes where transport vesicles are formed (Kawasaki et al., 2005; D'Souza-Schorey and Chavrier, 2006). According to the wealth of data from various eukaryotic organisms, these are the endoplasmic reticulum (ER) for COPII vesicles, Golgi apparatus for COPI vesicles, and the *trans*-Golgi network (TGN)/early endosome (EE) and the plasma membrane for clathrin-coated vesicles. However, a recent article in *The Plant Cell* (Böhlenius et al., 2010) reported that a barley (*Hordeum vulgare*) green fluorescent protein (GFP)-tagged ARF (Hv-ARFA1b/1c) localizes to a multivesicular body (MVB) when transiently expressed in onion epidermal cells. We take issue with this conclusion, in particular because no electron microscopy evidence was presented to confirm that the structure labeled was indeed multivesiculate. Here, we present evidence that *Arabidopsis* ARF1 localizes to the TGN and Golgi stack. We argue that multiple lines of evidence are required to make an accurate determination of protein localization, especially with respect to the endomembrane system, which has high potential for overexpression and tagging-induced artifacts (Millar et al., 2009; Moore and Murphy, 2009).

In all other higher plants so far investigated, principally *Arabidopsis* and tobacco (*Nicotiana tabacum*), the very similar At-ARFA1c (differing from the

barley ARF by only four amino acids) localizes to the Golgi apparatus/TGN. A Golgi localization was first demonstrated using an antibody generated against a full-length At-ARFA1c-glutathione S-transferase fusion protein (Pimpl et al., 2000) by both immunogold electron microscopy (Pimpl et al., 2000; Stierhof and El Kasmi, 2010) (Figures 1A and 1B) and immunofluorescence (Ritzenthaler et al., 2002) (Figures 1D, 1F, and 1G). The same result was obtained when ARF1-(X)FP constructs are expressed (Xu and Scheres, 2005; Stefano et al., 2006) (Figures 1D and 1E). Functional confirmation of an ARF1 localization at the Golgi apparatus comes additionally from several expression studies on tobacco cells with GDP-(T31N) or GTP-(Q71L) restricted ARF1 mutants, which lead to an abrogation of ER-to-Golgi transport and a redistribution of Golgi enzymes into the ER (Lee et al., 2002; Takeuchi et al., 2002; Stefano et al., 2006). The expression of these mutants (leading to an inhibition of COPI vesicle formation) is equivalent to treatment with brefeldin A, which in tobacco results in a fusion of Golgi membranes with the ER (Langhans et al., 2011). However, there is good evidence that ARF1 also localizes to the TGN. This was originally suggested by Pimpl et al. (2003) who showed that a functional ARF1 was also required for post-Golgi sorting of soluble vacuolar proteins, implying (as in mammals) that ARF1 was required for clathrin-coated vesicle formation at the TGN. Visualization of a potential ARF1 localization at the TGN was provided later by immunofluorescence in *Arabidopsis* using lines expressing SYP61–cyan fluorescent protein and VHA-a1-GFP as TGN markers (Paciorek et al., 2005; Tanaka et al., 2009) and demonstration that a FM4-64 positive compartment, distinct from the Golgi apparatus, was labeled by ARF1-GFP (Xu and Scheres, 2005; Tanaka et al., 2009). A similar, non-Golgi ARF1-positive compartment was also identified by Stefano et al. (2006) and later confirmed as the TGN by immunogold electron

microscopy by Stierhof and El-Kasmi (2010) (Figure 1A).

Böhlenius et al. (2010) based their conclusion that Hv-ARF1 localizes to MVBs on the observation of ARF1 colocalization with the *Arabidopsis* Rab GTPase ARA7/RabF2 in a transient expression assay in onion epidermal cells. We cannot confirm this observation with At-ARF1 in *Arabidopsis* roots. At-ARF1 antibodies that specifically colocalize with stably expressed ARF1-GFP (Figure 1D) did not label structures which carry stably expressed ARA7-GFP (Figure 1G). On the other hand, Xu and Scheres (2005) localized a small proportion of the ARF1-GFP signal in *Arabidopsis* root cells to an ARA7-positive organelle. This may reflect localization of a portion of ARA7 to the TGN/EE since immunogold electron microscopy of plastic sections cut from high-pressure frozen *Arabidopsis* roots has confirmed the localization of ARA7 to the boundary membrane of MVBs (Haas et al., 2007). ARA7 also has been detected at the TGN in *Arabidopsis* roots (Stierhof and El Kasmi, 2010) and in a FM4-64-positive compartment, distinct from the Golgi apparatus (Dhonukshe et al., 2006). Taken together, all of the localization data so far available using different markers as well immunogold electron microscopy indicate that the reported colocalizations of ARF1 and ARA7 reflect the presence of both of these regulators at the TGN/EE rather than at the MVB.

Is there experimental support for the speculation made by Böhlenius et al. (2010) that ARF1 at the MVB “mediates formation of vesicles with another destination, potentially in retrograde trafficking?” More precisely stated: Is there any evidence for (1) COPI- or clathrin-coated vesicle formation at the MVB and (2) recycling from MVB/late endosomes (LEs)? In the mammalian literature, there are numerous papers implicating the role of various coat-omer subunits in endocytosis (Whitney et al., 1995; Razi et al., 2009). However, COPI seems mainly to be involved in early trafficking events (e.g., effecting transferrin internalization and recycling from EE; Daro




Figure 1. ARF1 Localization, MVBs, and PVC Maturation.

(A) and **(B)** Immunogold electron microscopy with ARF1 antibodies. Bars = 500 nm in **(A)** and 100 nm in **(B)**.

(A) Tokuyasu cryosection of high-pressure frozen/freeze-substituted *Arabidopsis* roots. Thawed sections were stained with 1-nm gold particles and silver enhanced Nanogold according to Stierhof and El-Kasmi (2010). Label is present over both the Golgi stack and TGN. (Micrograph courtesy of York-Dieter Stierhof, Tübingen.)

(B) Conventional cryosection of a maize (*Zea mays*) root cap cell (for details, see Pimpl et al., 2000). Gold particles (5 nm) are seen on vesicles (arrows) at both the *cis*- (*c*) and *trans*- (*t*) faces of the Golgi stack.

(C) Ultrastructure of an MVB. Section prepared from a high-pressure frozen/freeze-substituted/plastic-embedded sample of an *Arabidopsis* root. Note the almost perfect circular profile lacking vesicle budding profiles, as well as an electron-dense putative clathrin plaque (arrowheads) at the surface. Bar = 100 nm. (Micrograph courtesy of York-Dieter Stierhof, Tübingen.)

(D) to **(G)** Double immunofluorescence with antibodies directed against proteins indicated in each panel in 5-d-old root cells of the *Arabidopsis* seedlings. In all images, the GFP fusion proteins are shown in green and the endogenous ARF1 (**[D]**, **[F]**, and **[G]**) or COPI (**[E]**) proteins in red; merged signals are yellowish. The rabbit polyclonal anti-ARF1 antibody and anti- γ -COP antibody were diluted 1:800, and mouse monoclonal anti-GFP antibody was diluted 1:600. The secondary antibodies anti-rabbit CY3 (Sigma-Aldrich) and anti-mouse Alexa 488 (Invitrogen) were diluted 1:600. Bars = 4 μ m.

(D) Endogenous ARF1 (detected by ARF1 antibody) colocalizes with signal from GFP antibody in a line expressing At-ARFA1c fused to enhanced GFP (ARF1-GFP).

(E) ARF1-GFP (detected by GFP antibody) colocalizes with COPI (detected by antibody against the Golgi marker γ -COP; Pimpl et al., 2000).

(F) ARF1 (detected by ARF1 antibody) also colocalizes with an anti-GFP signal in a line expressing the TGN marker VHA-a1-GFP.

(G) No significant colocalization was observed between the PVC/MVB marker GFP-ARA7 (detected by GFP antibody) and endogenous ARF1.

(H) to **(J)** Transient expression of fluorescently tagged TGN (YFP-AtSYP61) and PVC/MVB (AtVSR2-RFP) markers in tobacco mesophyll protoplasts. After short expression periods (6 h), both markers show complete colocalization. However, the two markers becoming increasingly separate with increasing length of expression. This can be interpreted as demonstrating the formation of the PVC/MVB at the TGN and its gradual maturation away from the TGN. Bars = 5 μ m.

et al., 1997) or in anterograde trafficking between EE and LE (Aniento et al., 1996). Although there is evidence for the participation of COPI in transport from the LE to vacuole in yeast (Gabriely et al., 2007), in no eukaryotic organism, including plants, have COPI proteins actually been demonstrated at the surface of MVB/LEs.

With respect to clathrin, to our knowledge, there are no reports for clathrin-coated vesicle formation on LEs in either mammalian or yeast cells. The same is true for plant MVBs, which in high-pressure frozen/plastic-embedded preparations have a circular profile normally without vesicle budding profiles (Figure 1C). However, plant MVBs have a densely labeled tuft at their surface, and this is probably the double-layered clathrin plaque related to the ESCRT-0 complex (Sachse et al., 2002; Wollert and Hurley, 2010). In addition, it should be noted that receptor recycling from endosomes is facilitated by the pentameric retromer complex (Bonifacino and Hurley, 2008). Retromer binds mainly to the tubular protrusions of the EE in mammalian cells (Mari et al., 2008), but in both yeast and plant cells, the situation is controversial, with evidence pointing to retromer at both the EE (Strochlic et al., 2007; Niemes et al., 2010) and MVB/LE (Jaillais et al., 2008; Balderhaar et al., 2010). However, it has recently been shown that retromer is recruited by the Rab GTPase Rab7 and not by an ARF-type GTPase (Seaman et al., 2009). Therefore, a role for ARF1 in retrograde trafficking from the MVB remains questionable.

The question now is, might such a colocalization reflect stable resident populations or a transitory, dynamic situation? Mammalian cell biologists usually regard multivesicular late endosomes to be the result of a gradual transformation (maturation) from pleiomorphic early endosomes (van Weering et al., 2010). Thus, when performing transient expression experiments as Böhlenius et al. (2010) have done, it cannot be ruled out that overlapping signals for ARF1 and ARA7 merely reflect an early stage in MVB formation from the TGN/EE. This can in fact be demonstrated in tobacco mesophyll protoplasts, which show a colocalization of TGN/EE and MVB/LE markers early in the expression period but a clear separation at a later time point (Figures 1H to 1J). The mechanism underlying this long delay in separation of

markers is not clear. Although performed with plasmid concentrations that do not interfere with the appearance of Golgi stack markers, it cannot be ruled out that the overexpression of the TGN and MVB markers temporarily perturbs their trafficking.

For reasons indicated above, attempting to localize a protein solely on the basis of transient expression of an (X)FP-tagged fusion construct can be misleading. Not only are such studies open to question because of their transient nature, but in such experiments, the proteins in question are sometimes highly overexpressed relative to their endogenous counterparts. This could be particularly troublesome when dealing with a regulatory protein like a GTPase whose overexpression could well perturb membrane flux leading to a temporary mislocalization of marker proteins. Ideally, the organelle/compartments to which a transiently expressed construct locates should be the same as that where a stably expressed construct resides and even better in a genetic background where the function of the tagged protein can be assessed. Therefore, to rule out problems of overexpression, the localization of the endogenous protein should also be determined by immunofluorescence and/or immunogold electron microscopy with a specific antibody.

David G. Robinson and David Scheuring
Department of Plant Cell Biology
Centre for Organismal Studies
University of Heidelberg
D-69120 Heidelberg, Germany

Satoshi Naramoto and Jiri Friml
Department of Plant Systems Biology
VIB and Department of Molecular
Genetics, Ghent University
B-9052 Ghent, Belgium

REFERENCES

- Anders, N., and Jürgens, G.** (2008). Large ARF guanine nucleotide exchange factors in membrane trafficking. *Cell. Mol. Life Sci.* **65**: 3433–3445.
- Aniento, F., Gu, F., Parton, R.G., and Gruenberg, J.** (1996). An endosomal beta

COP is involved in the pH-dependent formation of transport vesicles destined for late endosomes. *J. Cell Biol.* **133**: 29–41.

- Balderhaar, H.J., Arit, H., Ostrowicz, C., Bröcker, C., Sündermann, F., Brandt, R., Babst, M., and Ungermann, C.** (2010). The Rab GTPase Ypt7 is linked to retromer-mediated receptor recycling and fusion at the yeast late endosome. *J. Cell Sci.* **123**: 4085–4094.
- Böhlenius, H., Mørch, S.M., Godfrey, D., Nielsen, M.E., and Thordal-Christensen, H.** (2010). The multivesicular body-localized GTPase ARFA1b/1c is important for callose deposition and ROR2 syntaxin-dependent preinvasive basal defense in barley. *Plant Cell* **22**: 3831–3844.
- Bonifacino, J.S., and Hurley, J.H.** (2008). Retromer. *Curr. Opin. Cell Biol.* **20**: 427–436.
- Daro, E., Sheff, D., Gomez, M., Kreis, T., and Mellman, I.** (1997). Inhibition of endosome function in CHO cells bearing a temperature-sensitive defect in the coatamer (COPI) component epsilon-COP. *J. Cell Biol.* **139**: 1747–1759.
- Dhonukshe, P., Baluska, F., Schlicht, M., Hlavacka, A., Samaj, J., Friml, J., and Gadella, T.W., Jr.** (2006). Endocytosis of cell surface material mediates cell plate formation during plant cytokinesis. *Dev. Cell* **10**: 137–150.
- D'Souza-Schorey, C., and Chavrier, P.** (2006). ARF proteins: Roles in membrane traffic and beyond. *Nat. Rev. Mol. Cell Biol.* **7**: 347–358.
- Gabriely, G., Kama, R., and Gerst, J.E.** (2007). Involvement of specific COPI subunits in protein sorting from the late endosome to the vacuole in yeast. *Mol. Cell. Biol.* **27**: 526–540.
- Haas, T.J., Sliwinski, M.K., Martínez, D.E., Preuss, M., Ebine, K., Ueda, T., Nielsen, E., Odorizzi, G., and Otegui, M.S.** (2007). The *Arabidopsis* AAA ATPase SKD1 is involved in multivesicular endosome function and interacts with its positive regulator LYST-INTERACTING PROTEIN5. *Plant Cell* **19**: 1295–1312.
- Jaillais, Y., Fobis-Loisy, I., Miège, C., and Gaude, T.** (2008). Evidence for a sorting endosome in *Arabidopsis* root cells. *Plant J.* **53**: 237–247.
- Kawasaki, M., Nakayama, K., and Wakatsuki, S.** (2005). Membrane recruitment of effector proteins by Arf and Rab GTPases. *Curr. Opin. Struct. Biol.* **15**: 681–689.
- Langhans, M., Förster, S., Helmchen, G., and Robinson, D.G.** (February 28, 2011). Differential effects of the brefeldin A analogue (6R)-hydroxy-BFA in tobacco and *Arabidopsis*. *J. Exp. Bot.* <http://dx.doi.org/10.1093/jxb/err007>.
- Lee, M.H., Min, M.K., Lee, Y.J., Jin, J.B., Shin, D.H., Kim, D.H., Lee, K.H., and Hwang, I.**

- (2002). ADP-ribosylation factor 1 of Arabidopsis plays a critical role in intracellular trafficking and maintenance of endoplasmic reticulum morphology in Arabidopsis. *Plant Physiol.* **129**: 1507–1520.
- Mari, M., Bujny, M.V., Zeuschner, D., Geerts, W.J., Griffith, J., Petersen, C.M., Cullen, P.J., Klumperman, J., and Geuze, H.J.** (2008). SNX1 defines an early endosomal recycling exit for sortilin and mannose 6-phosphate receptors. *Traffic* **9**: 380–393.
- Millar, A.H., Carrie, C., Pogson, B., and Whelan, J.** (2009). Exploring the function-location nexus: using multiple lines of evidence in defining the subcellular location of plant proteins. *Plant Cell* **21**: 1625–1631.
- Moore, I., and Murphy, A.** (2009). Validating the location of fluorescent protein fusions in the endomembrane system. *Plant Cell* **21**: 1632–1636.
- Niemes, S., Langhans, M., Viotti, C., Scheuring, D., San Wan Yan, M., Jiang, L., Hillmer, S., Robinson, D.G., and Pimpl, P.** (2010). Retromer recycles vacuolar sorting receptors from the trans-Golgi network. *Plant J.* **61**: 107–121.
- Paciorek, T., Zazimalová, E., Ruthardt, N., Petrášek, J., Stierhof, Y.D., Kleine-Vehn, J., Morris, D.A., Emans, N., Jürgens, G., Geldner, N., and Friml, J.** (2005). Auxin inhibits endocytosis and promotes its own efflux from cells. *Nature* **435**: 1251–1256.
- Pimpl, P., Hanton, S.L., Taylor, J.P., Pinto-daSilva, L.L., and Denecke, J.** (2003). The GTPase ARF1p controls the sequence-specific vacuolar sorting route to the lytic vacuole. *Plant Cell* **15**: 1242–1256.
- Pimpl, P., Movafeghi, A., Coughlan, S., Denecke, J., Hillmer, S., and Robinson, D.G.** (2000). In situ localization and in vitro induction of plant COPI-coated vesicles. *Plant Cell* **12**: 2219–2236.
- Razi, M., Chan, E.Y., and Tooze, S.A.** (2009). Early endosomes and endosomal coatome are required for autophagy. *J. Cell Biol.* **185**: 305–321.
- Ritzenthaler, C., Nebenführ, A., Movafeghi, A., Stussi-Garaud, C., Behnia, L., Pimpl, P., Staehelin, L.A., and Robinson, D.G.** (2002). Reevaluation of the effects of brefeldin A on plant cells using tobacco Bright Yellow 2 cells expressing Golgi-targeted green fluorescent protein and COPI antisera. *Plant Cell* **14**: 237–261.
- Robinson, D.G., Langhans, M., Saint-Jore-Dupas, C., and Hawes, C.** (2008). BFA effects are tissue and not just plant specific. *Trends Plant Sci.* **13**: 405–408.
- Sachse, M., Urbé, S., Oorschot, V., Strous, G.J., and Klumperman, J.** (2002). Bilayered clathrin coats on endosomal vacuoles are involved in protein sorting toward lysosomes. *Mol. Biol. Cell* **13**: 1313–1328.
- Seaman, M.N., Harbour, M.E., Tattersall, D., Read, E., and Bright, N.** (2009). Membrane recruitment of the cargo-selective retromer subcomplex is catalysed by the small GTPase Rab7 and inhibited by the Rab-GAP TBC1D5. *J. Cell Sci.* **122**: 2371–2382.
- Stefano, G., Renna, L., Chatre, L., Hanton, S.L., Moreau, P., Hawes, C., and Brandizzi, F.** (2006). In tobacco leaf epidermal cells, the integrity of protein export from the endoplasmic reticulum and of ER export sites depends on active COPI machinery. *Plant J.* **46**: 95–110.
- Stierhof, Y.D., and El Kasm, F.** (2010). Strategies to improve the antigenicity, ultrastructure preservation and visibility of trafficking compartments in Arabidopsis tissue. *Eur. J. Cell Biol.* **89**: 285–297.
- Strochlic, T.I., Setty, T.G., Sitaram, A., and Burd, C.G.** (2007). Grd19/Snx3p functions as a cargo-specific adapter for retromer-dependent endocytic recycling. *J. Cell Biol.* **177**: 115–125.
- Takeuchi, M., Ueda, T., Yahara, N., and Nakano, A.** (2002). Arf1 GTPase plays roles in the protein traffic between the endoplasmic reticulum and the Golgi apparatus in tobacco and Arabidopsis cultured cells. *Plant J.* **31**: 499–515.
- Tanaka, H., Kitakura, S., De Rycke, R., De Groodt, R., and Friml, J.** (2009). Fluorescence imaging-based screen identifies ARF GEF component of early endosomal trafficking. *Curr. Biol.* **19**: 391–397.
- van Weering, J.R., Verkade, P., and Cullen, P.J.** (2010). SNX-BAR proteins in phosphoinositide-mediated, tubular-based endosomal sorting. *Semin. Cell Dev. Biol.* **21**: 371–380.
- Vernoud, V., Horton, A.C., Yang, Z., and Nielsen, E.** (2003). Analysis of the small GTPase gene superfamily of Arabidopsis. *Plant Physiol.* **131**: 1191–1208.
- Whitney, J.A., Gomez, M., Sheff, D., Kreis, T.E., and Mellman, I.** (1995). Cytoplasmic coat proteins involved in endosome function. *Cell* **83**: 703–713.
- Wollert, T., and Hurley, J.H.** (2010). Molecular mechanism of multivesicular body biogenesis by ESCRT complexes. *Nature* **464**: 864–869.
- Xu, J., and Scheres, B.** (2005). Dissection of Arabidopsis ADP-RIBOSYLATION FACTOR 1 function in epidermal cell polarity. *Plant Cell* **17**: 525–536.

Reply: On ARF1 Localizes to the Golgi and the *Trans*-Golgi Network: Future Challenge in Plant Multivesicular Body Studies

Vesicle trafficking is a large and complex research area, still in its infancy in plants. The Letter to the Editor in *The Plant Cell* by Robinson et al. (2011) is written by long-time experts in this area and makes a sincere and honest effort to clarify that “ARF1 localizes to the Golgi and the TGN.” The letter is a comment on our recent article “The Multivesicular Body-Localized GTPase ARFA1b/1c Is Important for Callose Deposition and ROR2 Syntaxin-Dependent Preinvasive Basal Defense

in Barley” published in *The Plant Cell* (Böhlenius et al., 2010), which questions the subcellular location of ARF1. Therefore, the letter is well justified.

Robinson et al. suggest that our ARF GTPase localization to multivesicular bodies (MVBs) is erroneous, perhaps due to overexpression from a strong promoter, and that no ARF GTPases have been associated with MVBs, since no coat-mediated vesicle budding occurs on this cellular compartment. This explanation is certainly possible. However, we remain open to the possibility that our result could

be due to differences between dicots and monocots, or, perhaps more likely, that differences in both location and function may exist within the ARFA subfamily. Robinson et al. present an example that illustrates how colocalization can be misleading. However, the example involves two integral membrane proteins (syntaxin, SYP61, and vacuolar sorting receptor [VSR2]), both of which pass through the ER-Golgi pathway before reaching their final separate destinations. Therefore, it is not surprising that they colocalize for a period of time after transformation. This