

Letter to Editor

Normal olfaction range of Rasht residents using a new test designed for the region

JRMS 2010; 15(6): 371-372

Disorders of sensing smell and taste can present diagnostic and therapeutic challenges for the otolaryngologists,¹ especially for forensic problems and especially in Islamic countries-the importance of olfactory sensation in Islamic laws is as high as whole body of a human being.

There are many methods for qualitative and quantitative evaluation of olfactory sensation in the world.² Discrimination and detecting of odors is largely cultural and therefore learnt³; hence, it is not surprising if the smell tests which had been designed and applied in other cultures and countries would have some restrictions in being applied in another country.³⁻⁵

Unfortunately there isn't any locally designed test currently used in Iran. Also, the current tests are difficult to interpret and are usually expensive to run.

In order to develop a standard smell test for Iranians and determination of the normal range of olfactory sense in this region, a research composed of two stages was conducted; to find the most popular odorous items in the region, fifty six well known materials were presented to two hundred, 15-60 years old,

normal residents of Rasht, Guilan province, Iran. Sixteen materials with the highest scores were selected in this stage: gasoline, alcohol, tea, rice, soap, cinnamon, origan, garlic, onion, washing powder, mint, rose water, lemon juice, olive oil, vinegar, and menthol. Fifteen of these were the olfactory nerve stimulators, and one (menthol salicylate) was the trigeminal nerve stimulator; this is unlike other tests that use ammonia as the trigeminal nerve stimulator, because ammonia has tang and unpleasant smell and in high doses may damage the olfactory system.

Then, these 16 materials were presented in identical containers to 150 normal citizens (77 women, 73 men) at 2 centimeter distance from their noses, for 3 seconds. Each nostril was tested separately; therefore the total score for two nostril and 16 materials was 32 for each person. The subjects would choose an answer in a 4 choice questionnaire; so, malingering individuals can be identified (someone who had less than 25% of total score). Thirty of the subjects were re-tested with the designed sniff box for evaluation of reproducibility of the test.

Table 1. Mean of olfactory scores in different age and sex groups

Age groups (years)	Male		Female	
	Mean	SD	Mean	SD
15-30	30.72	1.42	31.15	0.85
31-45	30.91	0.86	31.26	1.42
46-60	30.00	1.72	30.79	1.00

SD: Standard Deviation

*Hooshang Gerami^a, Shadman Nemati^{*a}, Rahmatollah Banan^a, Rezvan Rouhi^b*

^a Otolaryngology Head and Neck Surgery Department and Research Center, Guilan University of Medical Sciences, Faculty of Medicine, Rasht, Iran.

^b Resident of Otolaryngology Head and Neck Surgery, Guilan University of Medical Sciences, Faculty of Medicine, Rasht, Iran.

* Corresponding Author

E-mail: nemati@gums.ac.ir

Considering 97 percentile, the normal range of olfactory score for 15-60 years old residents of Rasht was 28-32. The mean scores in different age and sex groups are presented in table 1. Test re test reliability was more than 96%.

In each age group, the mean olfactory scores of women were higher than men. With increasing age, the scores were decreased.

The designed test is cheap and available and significantly will reduce our need to expensive

foreign kits. Perhaps it would be better saying this is a large scale pilot study that must be tested furthermore in other studies and trials.

Acknowledgements

Our special thanks to Dr. Shahryar Dadgari, resident of otolaryngology who contributed in examination of the cases, and Dr. Kambiz Forghanparast who analyzed the data.

Conflict of Interests

Authors have no conflict of interests.

References

1. Wrobel BB, Leopold DA. Clinical assessment of patients with smell and taste disorders. *Otolaryngol Clin North Am* 2004;37(6):1127-42.
2. Rubin DB, Cleland TA. Dynamical mechanisms of odor processing in olfactory bulb mitral cells. *J Neurophysiol* 2006;96(2):555-68.
3. Drake-Lee A. Physiology of the nose and paranasal sinuses. In: Gleeson MJ, Browning GG, Burton MJ, editors. *Scott-Brown's otorhinolaryngology, head and neck surgery*. 7th ed. London: Hodder Arnold; 2008. p. 1355-69.
4. Hvastja L, Zanuttini L. Odour memory and odour hedonics in children. *Perception* 1989;18(3):391-6.
5. Dorland AL, Eric H. Physiology of olfaction. In: Cummings C, Haughey B, Thomas JR, Harker L, Robbins K, Flint P, et al, editors. *Cummings otolaryngology head and neck surgery*. 4th ed. Philadelphia: Mosby; 2005. p. 865-98.