

# Are Hill's criteria for causality satisfied for vitamin D and periodontal disease?

William B. Grant<sup>1,\*</sup> and Barbara J. Boucher<sup>2</sup>

<sup>1</sup>Sunlight, Nutrition; and Health Research Center (SUNARC); San Francisco, CA USA; <sup>2</sup>Centre for Diabetes; Bart's and The London School of Medicine and Dentistry; Queen Mary University of London; Blizard Institute of Cell and Molecular Science; London, UK

**Key words:** vitamin D, periodontal disease, criteria, causality, Bradford-Hill

There is mounting evidence that periodontal disease (PD) is linked to low serum 25-hydroxyvitamin D [25(OH)D] concentrations in addition to recognized risk factors like diet and smoking. This paper reviews this evidence using Hill's criteria for causality in a biological system. Evidence for strength of association, consistency, cohesion and 'dose-effects' [biological 'gradients'] include strong inverse correlations between serum 25(OH)D and PD cross-sectionally and that PD is consistently more prevalent in darker vs. lighter skinned people and increases at higher latitudes with analogy for gingivitis and for disorders associated with PD whose risks also increase with hypovitaminosis D. Evidence for plausibility includes that vitamin D increases calcium absorption and protects bone strength; induces formation of cathelicidin and other defensins that combat bacterial infection; reduces tissue production of destructive matrix metalloproteinases actively associated with PD and that prevalence of PD varies with common vitamin D receptor polymorphisms. Experimental evidence from limited supplementation studies [using calcium and vitamin D] shows that supplementation reduces tooth loss. Thus, existing evidence for hypovitaminosis D as a risk factor for PD to date meets Hill's criteria for causality in a biological system. Further experimental evidence for effectiveness and temporality, preferably from randomized controlled trials of vitamin D supplementation [adjusting for other PD risk factors including diet and smoking to reduce confounding] are necessary to confirm causality. If confirmed, dentists and periodontists could perform a valuable service to their patients by discussing the importance of adequate vitamin D status and how to avoid deficiency.

## Introduction

Periodontal disease (PD) is a chronic condition where bacterial biofilms adhere to the gingiva (periodontal tissues) leading to host responses within periodontal tissues, and inducing inflammatory damage resulting in breakdown of the connective tissue that anchors teeth to alveolar bone.<sup>1</sup> If left unchecked, PD leads to tooth loss. The proportion of the US population with PD in 1999–2004

according to the National Health and Nutrition Examination Survey (NHANES) increased from 3.8% for those aged 20–34 years to 10.4% for those aged 35–49 years and to 11.9% for those aged 50–64 years.<sup>2</sup> PD rates also varied with ethnicity at 5.8% in Whites, 16.8% in Blacks and 13.8% in Mexican-Americans.<sup>2</sup> However, another study, also based on NHANES, found 3.6% in Whites, 7.2% in Blacks and 4.4% for Mexican-Americans.<sup>3</sup> Overall, periodontitis affects almost 40% of the entire adult population in the UK.<sup>4</sup>

Since oral bacteria are important risk factors for PD,<sup>5</sup> and since activated vitamin D [1,25-dihydroxyvitamin D], through induction of cathelicidin (LL-37) and other defensins, effectively fights bacterial infections,<sup>6</sup> it seems reasonable to expect that maintenance of adequate repletion with vitamin D could reduce the risk of PD. The most convincing evidence would be provided by well-conducted randomized controlled trials (RCTs). However, there has been only one to date, and it provided evidence of a moderate benefit with respect to periodontal health for combined vitamin D and calcium supplementation.<sup>7</sup> A cross-sectional study in the United States found a significant inverse correlation between PD and serum 25-hydroxyvitamin D [25(OH)D] concentrations.<sup>8</sup> While useful, cross-sectional studies alone cannot establish a causal connection between low serum 25(OH)D concentrations and risk of PD.

Another way to examine causality in a biological system was proposed by A. Bradford Hill.<sup>9</sup> Hill designated nine criteria that could be evaluated: strength of association, consistency, specificity, temporality, biological gradient, plausibility, coherence, experiment and analogy. Other researchers added the ruling out of confounding factors and bias.<sup>10</sup> Not all criteria need be satisfied, but the more that are, the stronger the case for causality. Some examples of applying Hill's criteria to examine causal links include dietary factors in the risk of Alzheimer disease,<sup>11</sup> the first paper to make such a link, and vitamin D as a risk reduction factor for several types of cancer nearly thirty years after the ultraviolet-B (UVB)-vitamin D-cancer hypothesis was proposed.<sup>12</sup>

This paper reviews the journal literature in order to address the question of whether the evidence supports a role for better supplies of vitamin D in reducing the risk of PD, using Hill's criteria for causality.

## Results

**Strength of association.** Data from the National Health and Nutrition Examination Survey (NHANES) III survey, conducted

\*Correspondence to: William B. Grant; Email: wbgrant@infonline.net  
Submitted: 04/19/10; Revised: 05/22/10; Accepted: 05/25/10  
Previously published online:  
[www.landesbioscience.com/journals/dermatoendocrinology/article/12488](http://www.landesbioscience.com/journals/dermatoendocrinology/article/12488)

in two phases between 1988 and 1994 were analyzed in terms of periodontal attachment loss as a function of serum 25(OH)D concentrations for males and females for two age groups, 20–49 years and >50 years. Associations were adjusted for age, race or ethnicity, smoking, diabetes, calcium intake, BMI, poverty income ratio, gingival bleeding, survey phase and dental examiner. The trends for those aged 20–49 years were not significant. For those >50 years, the standardized  $\beta$  coefficients varied from 0.39 (95% CI, 0.17–0.68) for 25(OH)D <40.2 nmol/L to 1.0 (reference) for 25(OH)D >85.6 nmol/L ( $p_{\text{for trend}}$ , 0.001) for males and 0.26 (95% CI, 0.09–0.43) ( $p_{\text{for trend}}$ , 0.001) for females.<sup>8</sup> Since these results were adjusted for many factors that affect risk of PD other than diet and vitamin D, they suggest a strong association of PD with poor vitamin D status.

Chronic marginal gingivitis, a chronic inflammation of the gingival tissues that is induced by bacterial dental plaque, is often associated with PD. In a related study using NHANES III data, Dietrich et al.<sup>15</sup> found a significant trend for prevalence of bleeding across quintiles of 25(OH)D; for the highest (32.4 nmol/L) vs. the lowest (99.6 nmol/L) the OR was 0.90 (0.86–0.95) after adjustment for the same factors as for PD but with the addition of vitamin C, hormone replacement therapy among women, missing teeth, full crown coverage and frequency of dental visits.

**Consistency.** While there are few studies of PD with respect to serum 25(OH)D concentrations, there are other studies that can be used to indicate the effect of vitamin D on PD risk. One such set of studies compares PD prevalence among African-Americans and White-Americans.<sup>3,16–21</sup> In the first of these studies, Blacks had an average of 78% of their sites with attachment loss and the average level of loss in those sites was approximately 4 mm, as compared to 65% and 3.1 mm for Whites.<sup>16</sup> In a recent study in Brazil, lighter-skinned Black people (Pardos) and dark-skinned Black people (Pretos) presented higher levels of PD when compared with white people [odds ratio (OR) = 1.5; 95% CI 1.2; 1.8 and OR = 1.6; 95% CI 1.2; 2.1, respectively] even after controlling for age, gender, schooling, per capita income and geographic region.<sup>21</sup> While such studies should be examined to determine whether factors other than serum 25(OH)D concentration could explain the findings, the fact that the mean serum 25(OH)D values for African-Americans was 16 ng/mL [64 nmol/L] while that for White-Americans was 26 ng/mL [104 nmol/L]<sup>22</sup> also supports the assumption that differences in vitamin D status [serum 25(OH)D concentration] could account for much of the disparity. A recent review concluded that differences in serum 25(OH)D levels between Black and White Americans could account for a large fraction of the health disparities observed.<sup>23</sup>

**Temporality.** This condition requires that beneficial effects should follow, not precede, the agent. This is generally the case for vitamin D. However, several papers have raised the possibility that low serum 25(OH)D concentrations measured for those who already have a disease or condition could reflect a changed lifestyle with less time spent out of doors, and, conversely, higher levels could be associated with increased outdoor activity, better health or the use of vitamin D supplementation because of its known or suspected health benefits. Two cross-sectional studies, one on PD,<sup>8</sup> the other on gingivitis,<sup>15</sup> are part of a much larger

set of studies using the NHANES dataset. Nearly all of the 357 NHANES papers mentioning associations of disease state with serum 25(OH)D concentrations have been supported by other studies. In addition, PD does not generally cause sufferers to stay indoors as might be the case with cancer or cardiovascular disease. There have been no prospective studies, but the one RCT of calcium and vitamin D supplementation on the effect of tooth loss in the elderly found a modest protective effect.<sup>7</sup>

**Mechanisms.** The first mechanism that is proposed to explain the role of vitamin D in reducing the risk of PD and gingivitis is induction of LL-37 and defensins by 1,25-dihydroxyvitamin D. This mechanism has been found to explain how solar UVB and vitamin D reduce the risk of tuberculosis.<sup>24,25</sup> Markers of higher vitamin D repletion have also been found to be associated with reduced risk of influenza,<sup>26</sup> pneumonia<sup>27</sup> and septicemia.<sup>28,29</sup> There are several good reviews of this mechanism.<sup>6,30–33</sup> A recent study found that saliva from orally healthy individuals is capable of protecting LL-37 from proteolysis by *Porphyromonas gingivalis*, a bacterium associated with periodontitis. This protective activity was found to be necessary to enable the direct bactericidal activity of LL-37 (as tested on *E. coli*) in the presence of virulence-associated proteases of *P. gingivalis*.<sup>34</sup>

**Matrix metalloproteinases (MMPs).** MMPs are enzymes that assist in modulation of interstitial tissue by digestion of its supportive matrix, especially collagen and MMP-9, in particular, is known to be present in large amounts in active PD,<sup>35</sup> as it is in active atheromatous plaque and this enzyme is inhibited by tissue inhibitor of matrix metalloproteinase-1 (TIMP-1). Higher plasma MMP-3, MMP-8 and MMP-9 concentrations have been found in PD patients compared with healthy controls (all  $p < 0.05$ ).<sup>36</sup> Furthermore, a study of British Bangladeshi adults, free of known diabetes or major illness, showed that a year of modest vitamin D supplementation led to a decrease in plasma circulating matrix metalloproteinase 9 (MMP-9) by -69% and of CRP by -23%.<sup>37</sup> Activated vitamin D has also been shown to suppress the increases in MMP9 formation induced in white cells by tubercle bacilli<sup>38</sup> as well as of MMP-3.<sup>39</sup> MMP-3 is associated with acute coronary syndrome<sup>40</sup> and unstable coronary disease.<sup>41</sup> Increases in circulating MMP-9 are seen in acute coronary events and associated with increases in cardiovascular disease risk factors<sup>42</sup> though this was not confirmed by Kaplan et al.<sup>43</sup> Since vitamin D modulates MMP9 in bone, and bone formation,<sup>44</sup> this second mechanism provides a possible link through which vitamin D status could influence both gum disease, tooth attachment and preservation of alveolar bone as well as cardiovascular disease and general bone health.

**Vitamin D receptors.** Vitamin D receptors (VDRs) are found in nearly every cell of the body. The hormonal metabolite of vitamin D, 1,25-hydroxyvitamin D, is the ligand that activates them by fitting into a specific cleft in the VDR. Ligand-bound VDRs heterodimerize with the retinoid X receptor to induce many of the effects of vitamin D through binding of the VDR; RXR complex to vitamin D response elements in the promoter regions of target genes, leading to modulation of expression of many of the 1,000 or more target genes. In about two-thirds of cases, they activate or enhance gene expression, and in the other third, they reduce

gene expression. Much of the action of vitamin D is achieved through the VDRs. There are many VDR polymorphisms, *BsmI*, *ApaI*, *TaqI* and *FokI* being the best known. VDR polymorphisms appear to be associated with variation in expression of various other genes even though polymorphisms such as *ApaI*, *BsmI*, *FokI* and *TaqI* are in non expressed portions of the VDR gene, possibly due to variation in the induction of tertiary structure of the VDR.<sup>45</sup> Investigations of disease associations with VDR polymorphisms can give an indication of whether vitamin D status may affect a given health or disease outcome. For PD, the VDRs that have received the most attention are *ApaI*, *BsmI*, *FokI* and *TaqI* (reviewed in ref. 46–52).

The question arises whether these polymorphisms could be related to effects on alveolar bone through calcium metabolism or by the ability to combat bacterial infection. To address that question, Pubmed was searched for papers on VDR, osteoporosis and infectious diseases. Papers were found for bones,<sup>53,54</sup> osteoporosis,<sup>55</sup> *Mycobacterium malmoense* pulmonary disease,<sup>56</sup> tuberculosis,<sup>57,58</sup> acute lower respiratory tract infection,<sup>59</sup> insulin secretory capacity,<sup>60</sup> type 1 diabetes,<sup>61,62</sup> lepromatous leprosy,<sup>63</sup> *Staphylococcus aureus* infection<sup>64</sup> and advanced prostate cancer.<sup>65,66</sup> For bones, the TT allele frequency was higher for those with higher mean forearm and spine bone mass density.<sup>53</sup> For tuberculosis, the Tt allele was found associated with better response to treatment,<sup>56,57,67</sup> as was the AA genotype.<sup>68</sup> Thus, these studies suggest that these polymorphisms are important for both functions but do not allow any distinction to be made between the roles of calcium metabolism and bactericidal efficacy.

**Supplementation studies.** There has been one small RCT of vitamin D and calcium supplementation and risk of PD. In this study, 11 of the 82 subjects (13%) taking supplements and 17 of the 63 subjects (27%) taking placebo lost one or more teeth (OR = 0.4; 95% CI: 0.2 to 0.9). During the 2-year follow-up period, 31 of the 77 subjects (40%) with total calcium intake of at least 1,000 mg per day lost one or more teeth compared with 40 of the 68 subjects (59%) who consumed less (OR = 0.5; 95% CI: 0.2 to 0.9).<sup>7</sup> Since both vitamin D and calcium were used, it is difficult to distinguish between the effects of the two supplements.

In a related cross-sectional study of chronic periodontitis in relation to calcium and vitamin D supplementation in Illinois and Missouri, a comparison was made of 23 subjects who had been taking vitamin D ( $\geq 400$  IU/day) and calcium ( $\geq 1,000$  mg/day) supplementation for >18 months (takers) and 28 subjects who were taking neither vitamin D nor calcium supplementation and had dietary intakes of calcium <1,000 mg/day and of vitamin D <400 IU/day (non-takers). The mean values of vitamin D and calcium intakes for the takers were 1,049 (781–1,317) IU/d and 1,769 (1,606–1,933) mg/day and 156 (117–195) IU/d and 642 (505–779) mg/d for the non-takers. Measures of periodontitis were consistently higher in non-takers than in takers [ $p < 0.05$ ], despite the small numbers of subjects<sup>69</sup> and despite the fact that participants had been enrolled in periodontal maintenance programs for >6 months before supplementation, which may have reduced the potential for benefits from vitamin D and calcium.

**Table 1.** Regression results for tooth loss for data in Dunning (1953)<sup>71</sup>

Number of states	UVB ( $\beta$ , p)	Urban ( $\beta$ , p)	T July ( $\beta$ , p)	Adj. R <sup>2</sup> , F, p
48	-0.72*	0.39*		0.64, 42*
47	-0.55*		-0.25, 0.08	0.52, 26*
47		0.37, 0.001	-0.60	0.48, 22*

\* $p < 0.001$ .

**Analogies with dental caries in the U.S.** Dental caries is analogous to PD in that it is caused by oral bacteria.<sup>5</sup> There are two studies on the effect of sunlight on risk of dental caries on adolescents and young men from the early-to-mid twentieth century. In the first, it was reported that adolescent males between the ages of 12 and 14 years had half as many carious lesions in the sunny west (3,000 hours of sunlight/year) than those in the much less sunny northeast (<2,200 hours of sunlight/year) whilst the rates in the portions of the country with intermediate annual hours of sunlight fell between those for the extremes of sun exposure.<sup>70</sup> In the second study, dental condition rankings were developed for young men entering the armed forces for World War I and World War II.<sup>71</sup> The northeast had the worst ranking, while states near Texas had the best ranking. A simple ecological analysis of the rankings using summertime UVB<sup>72</sup> and urban residence for 1,960 found an adjusted R<sup>2</sup> = 0.64 standardized coefficient beta for UVB = -0.72,  $p < 0.001$ , and for urban residence the adjusted R<sup>2</sup> = 0.39,  $p < 0.001$  (See Table 1).

**Confounding factors: smoking, diet and calcium intake.**

The most important recognized factors for PD are age (greater than 40 years), smoking,<sup>73</sup> diet<sup>74,75</sup> and calcium intake.<sup>76</sup> However, even when these factors are included in studies examining links between PD and systemic and infectious diseases, there is often a large portion of causation unaccounted for.

Bone mineral density (BMD) is linked to calcium, diet and smoking, as well as other factors. Many studies have found PD and osteoporosis to be comorbid.<sup>77–80</sup> However, osteoporosis may not be directly linked to dietary calcium intake because national diets high in calcium are also high in animal sources of protein,<sup>81</sup> which lowers the pH of the digestive tract, reducing calcium absorption and thus increasing calcium removal from bone.<sup>82</sup> Studies of correlations of changes in BMD with changes in PD status may provide some insight into the relative roles of the various factors. In a study in Pennsylvania, “absolute BMD and percentage change in BMD were similar in dentate and edentulous women. We found no difference in BMD or in absolute or percentage change in BMD between women with or without periodontal disease” and the authors concluded. “Little evidence exists for an association between edentulousness, periodontal disease and longitudinal changes in BMD.”<sup>83</sup> “In a cohort of 1,347 (137 edentulous) older men followed for an average of 2.7 year after recruitment from the Osteoporotic Fractures in Men Study, random half-mouth dental measures included clinical attachment loss (CAL), pocket depth (PD), calculus, plaque and bleeding. BMD was measured at the hip, spine and whole-body, by dual-energy X-ray absorptiometry,

and at the heel by ultrasound. After adjustment for age, smoking, race, education, body mass index and calculus, there was no association between number of teeth, periodontitis, periodontal disease progression, and either BMD or annualized rate of BMD change.<sup>84</sup> However, both of these studies were short-term in nature. Longer-term studies are probably required to find links. For example, in studies of measured and imputed maternal serum 25(OH)D and bone mineral density in the offspring aged nine years, both the estimated exposure to UVB radiation during late pregnancy and the maternal use of vitamin D supplements contributed to maternal 25(OH)-vitamin D concentration ( $p < 0.0001$  and  $p = 0.01$ , respectively) which in turn contributed to childhood bone mass ( $p = 0.03$ ).<sup>85,86</sup> In adults, supplementation has led to lesser reductions in BMD over time<sup>87</sup> and to reductions in fracture rates with 5 years of supplementation at ~820 IU/day.<sup>88</sup>

**Correlation with other diseases.** PD has itself been linked to many systemic and infectious diseases,<sup>89-92</sup> including many types of cancer,<sup>93,94</sup> carotid atheroma,<sup>95</sup> cardiovascular disease,<sup>96,97</sup> diabetes mellitus,<sup>98,99</sup> pneumonia<sup>100</sup> and rheumatoid arthritis.<sup>101</sup> Lower serum 25(OH)D concentrations have been found associated with risk of many types of chronic and infectious diseases in the past few years.<sup>102</sup> Such diseases include many types of cancer,<sup>12,103-105</sup> cardiovascular disease,<sup>106</sup> diabetes mellitus<sup>106</sup> and pneumonia.<sup>107</sup> Thus, the hypothesis that PD was a risk factor for the diseases associated with it was proposed. A number of other hypotheses have been postulated, including common susceptibility, systemic inflammation with increased circulating cytokines and mediators, local infection and cross-reactivity or molecular mimicry between bacterial antigens and self-antigens.<sup>90</sup>

In many association studies, commonly accepted risk factors for PD, such as smoking, were included in the analysis but did not explain the association. There is some evidence from intervention trials that periodontal therapy, which decreases the intraoral bacterial bioburden and reduces periodontal inflammation, can have a significant impact on systemic inflammatory status.<sup>108</sup> However, it has not been established that treating PD reduces the risk of any of the associated diseases.<sup>109</sup>

Another hypothesis is that a high carbohydrate diet increases both the risk of PD and of systemic diseases such as diabetes and cardiovascular disease, thereby explaining the link between the two.<sup>74</sup> Indeed, there is evidence that added sugar intake is a risk factor for cardiovascular disease.<sup>110-113</sup>

An alternative hypothesis, proposed here, is that low serum 25(OH)D concentrations increase the risk of both PD and of diseases associated with PD. **Table 2** includes examples of studies of PD with other diseases known to be associated with low serum 25(OH)D, namely cardiovascular disease, coronary heart disease, cognitive impairment, pneumonia, and many types of cancer. If this is the case, vitamin D supplementation for those with PD may prove to reduce the risk of the associated diseases.

Recently it was proposed that signaling through the receptor for advanced glycation end products (RAGE) might explain the periodontal-systemic disease link. This receptor, which is frequently associated with proinflammatory responses, has been shown to be activated by various ligands such as high mobility

group box-1 (HMGB1/amphoterin), amyloid fibrils, trans-tyreine, Mac-1 (Integrin Mac-1), as well as advanced glycation end products (AGEs), and recent studies indicate that signaling through RAGE is implicated as an underlying condition in diverse pathologies. It was recently demonstrated that hormonal vitamin D can blunt the deleterious impact of AGEs on endothelial cells, which provides further support for the vitamin D hypothesis.<sup>116-117</sup>

## Discussion

This paper reviews the evidence for the association of PD with low serum 25(OH)D concentrations. The more important evidence includes:- strong inverse correlation between serum 25(OH)D and both PD and gingivitis cross-sectionally; consistent data showing higher prevalence of PD in darker as compared to lighter-skinned people living in the same geographical locations. Several feasible mechanisms for this association are presented; that vitamin D increases calcium absorption and upregulates intracellular calcium; induces production of cathelicidin and other defensins with bacteriocidal effects on oral bacteria and reduces production of MMPs, especially of the MMP9 found in high concentrations in active PD. In addition, PD severity varies with two common VDR polymorphisms. Limited studies of vitamin D and calcium supplementation report reduced tooth loss with combined supplementation but could not distinguish between the effects of vitamin D and those of calcium. Analogy with dental caries include; that higher solar UVB exposure correlated inversely with dental caries in young men in the United States in the early-to-mid twentieth century; that there is co-morbidity of caries with systemic and infectious diseases for which low serum 25(OH)D concentration is a risk factor. The more commonly recognized risk factors for PD, such as age, diet and smoking, can confound studies related to serum 25(OH)D concentration but studies of co-morbid diseases with PD not examining serum 25(OH)D, suggest that these associations are not fully accounted for by the classical risk factors.

Thus, the evidence for hypovitaminosis D as a risk factor for PD meets the criteria for causality in a biological system proposed by Hill<sup>9</sup> and discussed by Weed<sup>10</sup> reasonably well even though some of his criteria are better satisfied than others. The cross-sectional evidence for this association and for variation with vitamin D polymorphisms supports the suggestion of causality as do the results of the single randomized controlled trial of vitamin D supplementation for PD though the benefits could not be distinguished from those of the calcium given. The fact that there are recognized mechanisms by which hypovitaminosis D tissue can worsen the risks of gingival inflammation, worsen tissue destruction, increase bone loss and reduce the efficiency of defensive mechanisms after bacterial infection, also supports the likelihood of causality.

Thus, there is consistent evidence using Hills Criteria, that hypovitaminosis D is a risk factor for PD but further evidence, either from randomized controlled trials of vitamin D supplementation or careful prospective studies, is required in order to determine whether hypovitaminosis D is causal for PD.

**Table 2.** Associations of periodontal disease with chronic and infectious diseases themselves known to be associated with hypovitaminosis D

Disease or health outcome	Study parameters	Findings:- Risk/Odds/Hazard Ratio [95% confidence interval]	Reference
Cardiovascular disease	215 epidemiological studies, 47 were observational 29 of these suitable for meta-analysis. 16,720 subjects in cross-sectional studies, 1,464 cases, 1,726 controls and 147,821 in cohort studies.	Pooled OR calculated from 22 case-control & cross-sectional studies 2.35 [1.87; 2.96; p < 0.0001]. RR for incident cardiovascular disease in the 7 cohort studies = 1.34 [1.27; 1.42; p < 0.0001].	96
Coronary heart disease	Meta-analysis of risk of PD with elevated markers of systemic bacterial exposure Total N of subjects = 12,074	OR = 1.75 [1.32; 2.34; p < 0.001]	114
Cognitive function	NHANES 2001 to 2002. 803 dentate participants aged 60 or older, completing periodontal and cognitive examination.	OR = 0.69 [0.51; 0.94] per/standard deviation (SD) increase in the DSST score	115
Pneumonia	Cohort study of 277 men and 420 women aged 80 years old examined in Japan in 1997.	OR for death from pneumonia (adjusted) = 3.9 in those with 10 or more teeth and with pockets >4 mm deep vs. those with 10 or more teeth but without pockets.	100
Cancers	Cohort study with 48,375 men and 18 years of follow-up were available (1986 to January 31, 2004; median follow-up 17.7 yrs), during which time 5,720 incident cancer cases were documented (excluding nonmelanoma skin cancer and nonaggressive prostate cancer).	By cancer site, significant associations for those with periodontal disease were found for lung (1.36 [1.15–1.60]), kidney (1.49 [1.12–1.97]), pancreas (1.54 [1.16–2.04]) and haematological cancers (1.30 [1.11–1.53]). Fewer teeth at baseline (0–16) marked OR for lung cancer of 1.70 [1.37–2.11], for those with 0–16 teeth vs. those with 25–32 teeth. In never-smokers, periodontal disease was associated with increases in total (1.21 [1.06–1.39]) and haematological cancers (1.35 [1.01–1.81]). But not for lung cancer (0.96 [0.46–1.98]).  The most consistent risk increases were seen for oral and esophageal cancers in periodontal disease. Gastric and pancreatic cancers were associated in some but not all studies. Lung, prostate, hematologic and other cancers were not consistently associated or there were too few studies for analysis.	93  94
	Review of approximately 40 studies		

## Materials and Methods

This paper is a review of the peer reviewed medical literature. The primary data source was www.pubmed.gov. The search terms were used with ‘periodontal disease’ or ‘periodontitis’ in searching this database. Since some of the evidence relates to vitamin D receptor (VDR) polymorphisms the search was extended to additional diseases linked to variation in VDR polymorphisms. The *BsmI*, *Apal*, *TaqI* and *FokI* VDR polymorphisms have received the greatest attention with respect to PD and other diseases,<sup>13,14</sup> so they were the ones included in searching for links to other

diseases. In addition, a limited search was made for reports of large studies or meta-analyses discussing links between PD and systemic and infectious diseases.

## Disclosure

W.B.G. receives funding from the UV Foundation (McLean, VA), the Sunlight Research Forum (Veldhoven), Bio-Tech-Pharmaceutical (Fayetteville, AR) and the Vitamin D Society (San Luis Obispo, CA), and has previously received funding from the Vitamin D Society (Canada).

## References

- Graves D. Cytokines that promote periodontal tissue destruction. *J Periodontol* 2008; 79:1585-91.
- Dye BA, Tan S, Smith V, Lewis BG, Barker LK, Thornton-Evans G, et al. Trends in oral health status: United States, 1988–1994 and 1999–2004. *Vital Health Stat* 11 2007; 248:1-92.
- Borrell LN, Crawford ND. Social disparities in periodontitis among United States adults 1999–2004. *Community Dent Oral Epidemiol* 2008; 36:383-91.
- D’Aiuto F, Graziani F, Tetè S, Gabriele M, Tonetti MS. Periodontitis: from local infection to systemic diseases. *Int J Immunopathol Pharmacol* 2005; 18:1-11.
- Law V, Seow WK, Townsend G. Factors influencing oral colonization of mutans streptococci in young children. *Aust Dent J* 2007; 52:93-100; quiz 59.
- Kamen DL, Tangpricha V. Vitamin D and molecular actions on the immune system: modulation of innate and autoimmunity. *J Mol Med* 2010; 88:441-50.
- Krall EA, Wehler C, Garcia RI, Harris SS, Dawson-Hughes B. Calcium and vitamin D supplements reduce tooth loss in the elderly. *Am J Med* 2001; 111:452-6.
- Dietrich T, Joshipura KJ, Dawson-Hughes B, Bischoff-Ferrari HA. Association between serum concentrations of 25-hydroxyvitamin D3 and periodontal disease in the US population. *Am J Clin Nutr* 2004; 80:108-13.
- Hill AB. The Environment and Disease: Association or Causation? *Proc R Soc Med* 1965; 58:295-300.
- Weed DL, Gorelic LS. The practice of causal inference in cancer epidemiology. *Cancer Epidemiol Biomarkers Prev* 1996; 5:303-11.
- Grant WB. Dietary links to Alzheimer’s disease. *Alz Dis Rev* 1997; 2:42-55.
- Grant WB. How strong is the evidence that solar ultraviolet B and vitamin D reduce the risk of cancer? An examination using Hill’s criteria for causality. *Dermato-Endocrinology* 2009; 1:14-21.
- Uitterlinden AG, Fang Y, van Meurs JB, van Leeuwen H, Pols HA. Vitamin D receptor gene polymorphisms in relation to Vitamin D related disease states. *J Steroid Biochem Mol Biol* 2004; 89:187-93.
- Kostner K, Denzer N, Muller CS, Klein R, Tilgen W, Reichrath J. The relevance of vitamin D receptor (VDR) gene polymorphisms for cancer: a review of the literature. *Anticancer Res* 2009; 29:3511-36.
- Dietrich T, Nunn M, Dawson-Hughes B, Bischoff-Ferrari HA. Association between serum concentrations of 25-hydroxyvitamin D and gingival inflammation. *Am J Clin Nutr* 2005; 82:575-80.
- Beck JD, Koch GG, Rozier RG, Tudor GE. Prevalence and risk indicators for periodontal attachment loss in a population of older community-dwelling blacks and whites. *J Periodontol* 1990; 61:521-8.
- Brown LF, Beck JD, Rozier RG. Incidence of attachment loss in community-dwelling older adults. *J Periodontol* 1994; 65:316-23.

18. Umeda M, Chen C, Bakker I, Contreras A, Morrison JL, Slots J. Risk indicators for harboring periodontal pathogens. *J Periodontol* 1998; 69:1111-8.
19. Craig RG, Boylan R, Yip J, Bamgboye P, Koutsoukos J, Mijares D, et al. Prevalence and risk indicators for destructive periodontal diseases in 3 urban American minority populations. *J Clin Periodontol* 2001; 28:524-35.
20. Borrell LN, Burt BA, Taylor GW. Prevalence and trends in periodontitis in the USA: from the NHANESIII to the NHANES, 1988 to 2000. *J Dent Res* 2005; 84:924-30.
21. Peres MA, Antunes JL, Boing AF, Peres KG, Bastos JL. Skin colour is associated with periodontal disease in Brazilian adults: a population-based oral health survey. *J Clin Periodontol* 2007; 34:196-201.
22. Ginde AA, Liu MC, Camargo CA Jr. Demographic differences and trends of vitamin D insufficiency in the US population, 1988–2004. *Arch Intern Med* 2009; 169:626-32.
23. Grant WB, Peiris AN. Possible role of serum 25-hydroxyvitamin D in Black-White health disparities in the United States. *J Am Med Directors Assoc.* in press.
24. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006; 311:1770-3.
25. Liu PT, Stenger S, Tang DH, Modlin RL. Cutting edge: vitamin D-mediated human antimicrobial activity against *Mycobacterium tuberculosis* is dependent on the induction of cathelicidin. *J Immunol* 2007; 179:2060-3.
26. Cannell JJ, Vieth R, Umhau JC, Holick MF, Grant WB, Madronich S, et al. Epidemic influenza and vitamin D. *Epidemiol Infect* 2006; 134:1129-40.
27. Grant WB, Giovannucci E. The possible roles of solar ultraviolet-B radiation and vitamin D in reducing case-fatality rates from the 1918–1919 influenza pandemic in the United States. *Dermato-Endocrinology* 2009; 1:215-9.
28. Grant WB. Solar ultraviolet-B irradiance and vitamin D may reduce the risk of septicemia. *Dermato-Endocrinology* 2009; 1:25-30.
29. Jeng L, Yamshchikov AV, Judd SE, Blumberg HM, Martin GS, Ziegler TR, et al. Alterations in vitamin D status and anti-microbial peptide levels in patients in the intensive care unit with sepsis. *J Transl Med* 2009; 7:28.
30. Zasloff M. Inducing endogenous antimicrobial peptides to battle infections. *Proc Natl Acad Sci USA* 2006; 103:8913-4.
31. Bikle DD. Vitamin D and the immune system: role in protection against bacterial infection. *Curr Opin Nephrol Hypertens* 2008; 17:348-52.
32. Adams JS, Ren S, Liu PT, Chun RF, Lagishetty V, Gombart AF, et al. Vitamin D-directed rheostatic regulation of monocyte antibacterial responses. *J Immunol* 2009; 182:4289-95.
33. Gombart AF. The vitamin D-antimicrobial peptide pathway and its role in protection against infection. *Future Microbiol* 2009; 4:1151-65.
34. Gutner M, Chaushu S, Balter D, Bachrach G. Saliva enables the antimicrobial activity of LL-37 in the presence of proteases of *Porphyromonas gingivalis*. *Infect Immun* 2009; 77:5558-63.
35. Lorencini M, Silva JA, de la Hoza CL, Carvalho HF, Stach-Machado DR. Changes in MMPs and inflammatory cells in experimental gingivitis. *Histol Histopathol* 2009; 24:157-66.
36. Marcaccini AM, Novaes AB Jr, Meschiari CA, Souza SL, Palioto DB, Sorgi CA, et al. Circulating matrix metalloproteinase-8 (MMP-8) and MMP-9 are increased in chronic periodontal disease and decrease after non-surgical periodontal therapy. *Clin Chim Acta* 2009; 409:117-22.
37. Timms PM, Mannan N, Hitman GA, Noonan K, Mills PG, Syndercombe-Court D, et al. Circulating MMP9, vitamin D and variation in the TIMP-1 response with VDR genotype: mechanisms for inflammatory damage in chronic disorders? *QJM* 2002; 95:787-96.
38. Coussens A, Timms PM, Boucher BJ, Venton TR, Ashcroft AT, Skolimowska KH, et al. 1alpha,25-dihydroxyvitamin D3 inhibits matrix metalloproteinases induced by *Mycobacterium tuberculosis* infection. *Immunology* 2009; 127:539-48.
39. Kobayashi H, Asano K, Kanai K, Suzuki H. Suppressive activity of vitamin D3 on matrix metalloproteinase production from cholesteatoma keratinocytes in vitro. *Mediators Inflamm* 2005; 2005:210-5.
40. Liu PY, Li YH, Chan SH, Lin LJ, Wu HL, Shi GY, et al. Genotype-phenotype association of matrix metalloproteinase-3 polymorphism and its synergistic effect with smoking on the occurrence of acute coronary syndrome. *Am J Cardiol* 2006; 98:1012-7.
41. White AJ, Duffy SJ, Walton AS, Ng JF, Rice GE, Mukherjee S, et al. Matrix metalloproteinase-3 and coronary remodelling: implications for unstable coronary disease. *Cardiovasc Res* 2007; 75:813-20.
42. Garvin P, Nilsson L, Carstensen J, Jonasson L, Kristenson M. Circulating matrix metalloproteinase-9 is associated with cardiovascular risk factors in a middle-aged normal population. *PLoS One* 2008; 3:1774.
43. Kaplan RC, Smith NL, Zucker S, Heckbert SR, Rice K, Psaty BM. Matrix metalloproteinase-3 (MMP3) and MMP9 genes and risk of myocardial infarction, ischemic stroke and hemorrhagic stroke. *Atherosclerosis* 2008; 201:130-7.
44. Lips P. Vitamin D physiology. *Prog Biophys Mol Biol* 2006; 92:4-8.
45. Norman AW, Ishizuka S, Okamura WH. Ligands for the vitamin D endocrine system: different shapes function as agonists and antagonists for genomic and rapid response receptors or as a ligand for the plasma vitamin D binding protein. *J Steroid Biochem Mol Biol* 2001; 76:49-59.
46. Inagaki K, Krall EA, Fleet JC, Garcia RI. Vitamin D receptor alleles, periodontal disease progression and tooth loss in the VA dental longitudinal study. *J Periodontol* 2003; 74:161-7.
47. de Brito Junior RB, Scarel-Caminaga RM, Trevisatto PC, de Souza AP, Barros SP. Polymorphisms in the vitamin D receptor gene are associated with periodontal disease. *J Periodontol* 2004; 75:1090-5.
48. Park KS, Nam JH, Choi J. The short vitamin D receptor is associated with increased risk for generalized aggressive periodontitis. *J Clin Periodontol* 2006; 33:524-8.
49. Nibali L, Parkar M, D'Aiuto F, Suvan JE, Brett PM, Griffiths GS, et al. Vitamin D receptor polymorphism (-1056 Taq-I) interacts with smoking for the presence and progression of periodontitis. *J Clin Periodontol* 2008; 35:561-7.
50. Wang C, Zhao H, Xiao L, Xie C, Fan W, Sun S, et al. Association between vitamin D receptor gene polymorphisms and severe chronic periodontitis in a Chinese population. *J Periodontol* 2009; 80:603-8.
51. Amano Y, Komiyama K, Makishima M. Vitamin D and periodontal disease. *J Oral Sci* 2009; 51:11-20.
52. Zhang L, Meng HX, Zhao HS, Li QY, Xu L, Chen ZB, et al. [Correlation study on polymorphisms of vitamin D receptor gene in patients with periodontitis]. *Beijing Da Xue Xue Bao* 2010; 42:37-40.
53. Vupputuri MR, Goswami R, Gupta N, Ray D, Tandon N, Kumar N. Prevalence and functional significance of 25-hydroxyvitamin D deficiency and vitamin D receptor gene polymorphisms in Asian Indians. *Am J Clin Nutr* 2006; 83:1411-9.
54. Arabi A, Zahed L, Mahfoud Z, El-Onsi L, Nabulsi M, Maalouf J, et al. Vitamin D receptor gene polymorphisms modulate the skeletal response to vitamin D supplementation in healthy girls. *Bone* 2009; 45:1091-7.
55. Seremak-Mrozikiewicz A, Drews K, Mrozikiewicz PM, Bartkowiak-Wieczorek J, Marcinkowska M, Wawrzyniak A, et al. Correlation of vitamin D receptor gene (VDR) polymorphism with osteoporotic changes in Polish postmenopausal women. *Neuro Endocrinol Lett* 2009; 30:540-6.
56. Gelder CM, Hart KW, Williams OM, Lyons E, Welsh KI, Campbell IA, et al. Vitamin D receptor gene polymorphisms and susceptibility to *Mycobacterium mageritense* pulmonary disease. *J Infect Dis* 2000; 181:2099-102.
57. Roth DE, Soto G, Arenas F, Bautista CT, Ortiz J, Rodriguez R, et al. Association between vitamin D receptor gene polymorphisms and response to treatment of pulmonary tuberculosis. *J Infect Dis* 2004; 190:920-7.
58. Gao L, Tao Y, Zhang L, Jin Q. Vitamin D receptor genetic polymorphisms and tuberculosis: updated systematic review and meta-analysis. *Int J Tuberc Lung Dis* 2010; 14:15-23.
59. Roth DE, Jones AB, Prosser C, Robinson JL, Vohra S. Vitamin D receptor polymorphisms and the risk of acute lower respiratory tract infection in early childhood. *J Infect Dis* 2008; 197:676-80.
60. Ogunkolade BW, Boucher BJ, Prah J, Bustin SA, Burren JM, Noonan K, et al. Vitamin D receptor (VDR) mRNA and VDR protein levels in relation to vitamin D status, insulin secretory capacity and VDR genotype in Bangladeshi Asians. *Diabetes* 2002; 51:2294-300.
61. Ponsoy AL, Pezic A, Ellis J, Morley R, Cameron F, Carlin J, et al. Variation in associations between allelic variants of the vitamin D receptor gene and onset of type 1 diabetes mellitus by ambient winter ultraviolet radiation levels: a meta-regression analysis. *Am J Epidemiol* 2008; 168:358-65.
62. Panierakis C, Goulielmos G, Mamoulakis D, Petraki E, Papavasiliou E, Galanakis E. Vitamin D receptor gene polymorphisms and susceptibility to type 1 diabetes in Crete, Greece. *Clin Immunol* 2009; 133:276-81.
63. Velarde Félix JS, Cázarez Salazar SG, Castro Velázquez R, Rendón Maldonado JG, Rangel Villalobos H. [Association between the TaqI polymorphism of the vitamin D receptor gene and lepromatous leprosy in a Mexican population sample]. *Salud Publica Mex* 2009; 51:59-61. Spanish.
64. Panierakis C, Goulielmos G, Mamoulakis D, Maraki S, Papavasiliou E, Galanakis E. *Staphylococcus aureus* nasal carriage might be associated with vitamin D receptor polymorphisms in type 1 diabetes. *Int J Infect Dis* 2009; 13:437-43.
65. John EM, Schwartz GG, Koo J, Van Den Berg D, Ingles SA. Sun exposure, vitamin D receptor gene polymorphisms and risk of advanced prostate cancer. *Cancer Res* 2005; 65:5470-9.
66. Chen L, Davey Smith G, Evans DM, Cox A, Lawlor DA, Donovan J, et al. Genetic variants in the vitamin D receptor are associated with advanced prostate cancer at diagnosis: findings from the prostate testing for cancer and treatment study and a systematic review. *Cancer Epidemiol Biomarkers Prev* 2009; 18:2874-81.
67. Wilbur AK, Kubatko LS, Hurtado AM, Hill KR, Stone AC. Vitamin D receptor gene polymorphisms and susceptibility *M. tuberculosis* in native Paraguayans. *Tuberculosis (Edinb)* 2007; 87:329-37.
68. Babb C, van der Merwe L, Beyers N, Pheiffer C, Walz G, Duncan K, et al. Vitamin D receptor gene polymorphisms and sputum conversion time in pulmonary tuberculosis patients. *Tuberculosis (Edinb)* 2007; 87:295-302.
69. Miley DD, Garcia MN, Hildebolt CF, Shannon WD, Couture RA, Anderson Speare CL, et al. Cross-sectional study of vitamin D and calcium supplementation effects on chronic periodontitis. *J Periodontol* 2009; 80:1433-9.
70. East BR. Mean annual hours of sunshine and the incidence of dental caries. *Am J Public Health Nations Health* 1939; 29:777-80.

71. Dunning JM. The influence of latitude and distance from seacoast on dental disease. *J Dent Res* 1953; 32:811-29.
72. Leffell DJ, Brash DE. Sunlight and skin cancer. *Sci Am* 1996; 275:52-3.
73. Zee KY. Smoking and periodontal disease. *Aust Dent J* 2009; 54:44-50.
74. Huijoe P. Dietary carbohydrates and dental-systemic diseases. *J Dent Res* 2009; 88:490-502.
75. Palacios C, Josphipura K, Willett W. Nutrition and health: guidelines for dental practitioners. *Oral Dis* 2009; 15:369-81.
76. Nishida M, Grossi SG, Dunford RG, Ho AW, Trevisan M, Genco RJ. Calcium and the risk for periodontal disease. *J Periodontol* 2000; 71:1057-66.
77. Gomes-Filho IS, Passos Jde S, Cruz SS, Vianna MI, Cerqueira Ede M, Oliveira DC, et al. The association between postmenopausal osteoporosis and periodontal disease. *J Periodontol* 2007; 78:1731-40.
78. Swoboda JR, Kiyak HA, Darveau R, Persson GR. Correlates of periodontal decline and biologic markers in older adults. *J Periodontol* 2008; 79:1920-6.
79. Erdogan O, Incki KK, Benlidayi ME, Seydaoglu G, Kelekci S. Dental and radiographic findings as predictors of osteoporosis in postmenopausal women. *Geriatr Gerontol Int* 2009; 9:155-64.
80. Moedano DE, Irigoyen ME, Borges-Yáñez A, Flores-Sánchez I, Rotter RC. Osteoporosis, the risk of vertebral fracture and periodontal disease in an elderly group in Mexico City. *Gerodontology* 2009; In press.
81. Meyer HE, Pedersen JL, Løken EB, Tverdal A. Dietary factors and the incidence of hip fracture in middle-aged Norwegians. A prospective study. *Am J Epidemiol* 1997; 145:117-23.
82. Jajoo R, Song L, Rasmussen H, Harris SS, Dawson-Hughes B. Dietary acid-base balance, bone resorption, and calcium excretion. *J Am Coll Nutr* 2006; 25:224-30.
83. Famili P, Cauley J, Suzuki JB, Weyant R. Longitudinal study of periodontal disease and edentulism with rates of bone loss in older women. *J Periodontol* 2005; 76:11-5.
84. Phipps KR, Chan BK, Madden TE, Geurs NC, Reddy MS, Lewis CE, Orwoll ES. Longitudinal study of bone density and periodontal disease in men. *J Dent Res* 2007; 86:1110-4.
85. Javaid MK, Crozier SR, Harvey NC, Gale CR, Dennison EM, Boucher BJ, et al. Princess Anne Hospital Study Group. Maternal vitamin D status during pregnancy and childhood bone mass at age 9 years: a longitudinal study. *Lancet* 2006; 367:36-43.
86. Sayers A, Tilling K, Boucher BJ, Noonan K, Tobias JH. Predicting ambient ultraviolet from routine meteorological data; its potential use as an instrumental variable for vitamin D status in pregnancy in a longitudinal birth cohort in the UK. *Int J Epidemiol* 2009; 38:1681-8.
87. Tang BM, Eslick GD, Nowson C, Smith C, Bensoussan A. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet* 2007; 370:657-66.
88. Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D3 (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomised double blind controlled trial. *BMJ* 2003; 326:469.
89. Price WA. Nutrition and Physical Degeneration: A Comparison of Primitive and Modern Diets and Their Effects. Paul B. Hoeber, Inc., Medical Book Department of Harper & Brothers, New York London, 1939.
90. Seymour GJ, Ford PJ, Cullinan MP, Leishman S, Yamazaki K. Relationship between periodontal infections and systemic disease. *Clin Microbiol Infect* 2007; 13:3-10.
91. Kuo LC, Polson AM, Kang T. Associations between periodontal diseases and systemic diseases: a review of the inter-relationships and interactions with diabetes, respiratory diseases, cardiovascular diseases and osteoporosis. *Public Health* 2008; 122:417-33.
92. Williams RC, Barnett AH, Claffey N, Davis M, Gadsby R, Kelleth M, et al. The potential impact of periodontal disease on general health: a consensus view. *Curr Med Res Opin* 2008; 24:1635-43.
93. Michaud DS, Liu Y, Meyer M, Giovannucci E, Josphipura K. Periodontal disease, tooth loss and cancer risk in male health professionals: a prospective cohort study. *Lancet Oncol* 2008; 9:550-8.
94. Fitzpatrick SG, Katz J. The association between periodontal disease and cancer: A review of the literature. *J Dent* 2010; 38:83-95.
95. Tzorbatozoglou ID, Sfyroeras GS, Giannoukas AD. Periodontitis and carotid atheroma: is there a causal relationship? *Int Angiol* 2010; 29:27-9.
96. Blaizot A, Vergnes JN, Nuwwareh S, Amar J, Sixou M. Periodontal diseases and cardiovascular events: meta-analysis of observational studies. *Int Dent J* 2009; 59:197-209.
97. Cronin A. Periodontal disease is a risk marker for coronary heart disease? *Evid Based Dent* 2009; 10:22.
98. Dunning T. Periodontal disease—the overlooked diabetes complication. *Nephrol Nurs J* 2009; 36:489-95.
99. Santacroce L, Carlaio RG, Bottalico L. Does it make sense that diabetes is reciprocally associated with periodontal disease? *Endocr Metab Immune Disord Drug Targets* 2010; 10:57-70.
100. Awano S, Ansai T, Takata Y, Soh I, Akifusa S, Hamasaki T, et al. Oral health and mortality risk from pneumonia in the elderly. *J Dent Res* 2008; 87:334-9.
101. Smolik I, Robinson D, El-Gabalawy HS. Periodontitis and rheumatoid arthritis: epidemiologic, clinical and immunologic associations. *Compend Contin Educ Dent* 2009; 30:188-90.
102. Holick MF. Vitamin D deficiency. *N Engl J Med* 2007; 357:266-81.
103. Garland CF, Garland FC, Gorham ED, Lipkin M, Newmark H, Mohr SB, et al. The role of vitamin D in cancer prevention. *Am J Public Health* 2006; 96:252-61.
104. Grant WB, Garland CF. The association of solar ultraviolet B (UVB) with reducing risk of cancer: multifactorial ecologic analysis of geographic variation in age-adjusted cancer mortality rates. *Anticancer Res* 2006; 26:2687-99.
105. Garland CF, Gorham ED, Mohr SB, Garland FC. Vitamin D for cancer prevention: global perspective. *Ann Epi* 2009; 19:468-83.
106. Parker J, Hashmi O, Dutton D, Mavrodaris A, Stranges S, Kandala NB, et al. Levels of vitamin D and cardiometabolic disorders: systematic review and meta-analysis. *Maturitas* 2010; 65:225-36.
107. Grant WB, Giovannucci D. The possible roles of solar ultraviolet-B radiation and vitamin D in reducing case-fatality rates from the 1918–1919 influenza pandemic in the United States. *Dermato-Endocrinology* 2009; 1:215-9.
108. Mealey BL, Rose LF. Diabetes mellitus and inflammatory periodontal diseases. *Compend Contin Educ Dent* 2008; 29:402-8.
109. Scannapieco FA, Dasanayake AP, Chhun N. “Does periodontal therapy reduce the risk for systemic diseases?” *Dent Clin North Am* 2010; 54:163-81.
110. Yudkin J. Sugar and ischaemic heart disease. *Practitioner* 1967; 198:680-3.
111. Grant WB. Reassessing the role of sugar in the etiology of heart disease. *J Orthomolec Med*, 1998; 13:95-104; 1998; 13:241-3.
112. Grant WB. Milk and other dietary influences on coronary heart disease. *Altern Med Rev* 1998; 3:281-94.
113. Schaefer EJ, Gleason JA, Dansinger ML. The effects of low-fat, high-carbohydrate diets on plasma lipoproteins, weight loss and heart disease risk reduction. *Curr Atheroscler Rep* 2005; 7:421-7.
114. Mustapha IZ, Debrey S, Oladubu M, Ugarte R. Markers of systemic bacterial exposure in periodontal disease and cardiovascular disease risk: a systematic review and meta-analysis. *J Periodontol* 2007; 78:2289-302.
115. Yu YH, Kuo HK. Association between cognitive function and periodontal disease in older adults. *J Am Geriatr Soc* 2008; 56:1693-7.
116. Katz J, Wallet S, Cha S. Periodontal disease and the oral-systemic connection: “is it all the RAGE?”. *Quintessence Int* 2010; 41:229-37.
117. Talmor Y, Golan E, Bencheitrit S, Bernheim J, Klein O, Green J, Rashid G. Calcitriol blunts the deleterious impact of advanced glycation end products on endothelial cells. *Am J Physiol Renal Physiol* 2008; 294:1059-64.