

Diaqua(1,4,8,11-tetraazacyclotetradecane- $\kappa^4 N^1, N^4, N^8, N^{11}$)copper(II) didodecanoate dihydrate

Nur Syamimi Ahmad Tajidi, Norbani Abdullah* and Zainudin Arifin

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: nursyamimi@crackberry.com


Received 31 March 2011; accepted 6 April 2011

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.047; wR factor = 0.129; data-to-parameter ratio = 21.5.

The title compound, $[Cu(C_{10}H_{24}N_4)(H_2O)_2][CH_3(CH_2)_{10}CO_2]_2 \cdot 2H_2O$, consists of one cationic copper(II) complex, two dodecanoate anions and two water solvent molecules. The Cu^{II} atom is located on an inversion center and is chelated by the four aza N atoms of the neutral 1,4,8,11-tetraazacyclotetradecane (cyclam) ligand and by two water molecules in axial positions, giving an octahedral coordination geometry, distorted as a consequence of the Jahn–Teller effect. The uncoordinated water molecules link the complex cations and the dodecanoate counter-ions through $O-H \cdots O$ hydrogen bonding, forming a layer structure parallel to (001). Intermolecular $N-H \cdots O$ interactions also occur.

Related literature

For the complexation of cyclam with transition metals, see: Ahmad Tajidi *et al.* (2010*a,b,c,d*); Lindoy *et al.* (2003); Holanda *et al.* (2007); Sreedaran *et al.* (2008); Zgolli *et al.* (2010).



Experimental

Crystal data

$[Cu(C_{10}H_{24}N_4)(H_2O)_2] \cdot (C_{12}H_{23}O_2)_2 \cdot 2H_2O$
 $M_r = 734.54$
Triclinic, $P\bar{1}$
 $a = 6.9972$ (4) Å
 $b = 8.8164$ (5) Å
 $c = 17.1495$ (10) Å
 $\alpha = 96.218$ (3)°

$\beta = 99.137$ (3)°
 $\gamma = 98.329$ (3)°
 $V = 1024.13$ (10) Å³
 $Z = 1$
Mo $K\alpha$ radiation
 $\mu = 0.58$ mm⁻¹
 $T = 150$ K
 $0.41 \times 0.41 \times 0.08$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 1997)
 $T_{min} = 0.796$, $T_{max} = 0.955$
7085 measured reflections
4623 independent reflections
4138 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.045$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.129$
 $S = 1.07$
4623 reflections

215 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 0.54$ e Å⁻³
 $\Delta\rho_{min} = -0.52$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O1W-H1WB \cdots O2$	0.90	1.91	2.774 (2)	160
$O1W-H1WA \cdots O2^i$	0.90	1.81	2.694 (2)	168
$O2W-H2WB \cdots O1W$	0.90	1.93	2.8037 (19)	164
$O2W-H2WA \cdots O1^{ii}$	0.90	1.89	2.777 (2)	168
$N2-H2 \cdots O1^i$	0.93	2.25	3.030 (2)	141
$N1-H1 \cdots O1W^{iii}$	0.93	2.12	2.982 (2)	153

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x, y - 1, z$; (iii) $x + 1, y, z$.

Data collection: SMART (Bruker, 1997); cell refinement: SAINT (Bruker, 1997); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics:

ORTEP-III (Burnett & Johnson, 1996), *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *pubCIF* (Westrip, 2010).

This project was financed by the University of Malaya (grant No. A-50101-DA000-B21519). The authors thank Mr Harry Adams for his support and cooperation.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2672).

References

- Ahmad Tajidi, N. S., Abdullah, N., Arifin, Z., Tan, K. W. & Ng, S. W. (2010a). *Acta Cryst.* **E66**, m887.
- Ahmad Tajidi, N. S., Abdullah, N., Arifin, Z., Tan, K. W. & Ng, S. W. (2010b). *Acta Cryst.* **E66**, m888.
- Ahmad Tajidi, N. S., Abdullah, N., Arifin, Z., Tan, K. W. & Ng, S. W. (2010c). *Acta Cryst.* **E66**, m889.
- Ahmad Tajidi, N. S., Abdullah, N., Arifin, Z., Tan, K. W. & Ng, S. W. (2010d). *Acta Cryst.* **E66**, m890.
- Bruker (1997). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEP-III*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Holanda, A. K. M., da Silva, F. O. N., Carvalho, I. M. M., Batista, A. A., Ellena, J., Castellano, E. E., Moreira, I. S. & Lopes, L. G. F. (2007). *Polyhedron*, **26**, 4653–4658.
- Lindoy, L. F., Mahinay, M. S., Skelton, B. W. & White, A. H. (2003). *J. Coord. Chem.* **56**, 1203–1213.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sreedaran, S., Bharathi, K. S., Rahiman, A. K., Rajesh, K., Nirmala, G. & Narayanan, V. (2008). *J. Coord. Chem.* **22**, 3594–3609.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zgolli, D. Z., Boughzala, H. & Driss, A. (2010). *Acta Cryst.* **E66**, m265–m266.

supplementary materials

Acta Cryst. (2011). E67, m588-m589 [doi:10.1107/S1600536811012773]

Diaqua(1,4,8,11-tetraazacyclotetradecane- $\kappa^4N^1,N^4,N^8,N^{11}$)copper(II) didodecanoate dihydrate

N. S. Ahmad Tajidi, N. Abdullah and Z. Arifin

Comment

Copper(II) cyclam complexes are potential functional materials in the field of molecular electronic, photonics and spintronics whose properties may be tuned by steric and electronic effects. The present complex represents our attempt to synthesize a functional material that possesses metallomesogenic properties for such applications. Several cyclam complexes with copper(II) (Ahmad Tajidi *et al.*, 2010*a,b,c,d*) and other transition metals (Lindoy *et al.*, 2003; Holanda *et al.*, 2007; Sreedaran *et al.*, 2008; Zgolli *et al.*, 2010) have been reported.

In the complex, the Cu^{II} atom, located on an inversion center, is coordinated to the 1,4,8,11-tetraazacyclotetradecane through the four aza-N atoms forming the basal plane of a distorted octahedra whose apices are occupied by two water molecules. Two solvate water molecules link anion and cations through O-H \cdots O hydrogen bondings (Fig. 1, Table 1). The relatively long Cu-O(water) distance, 2.455 (1)Å, is a consequence of the Jahn-Teller effect resulting in the distorted octahedron coordination geometry.

O-H \cdots O and N-H \cdots O Hydrogen bonds involving the coordinated and non coordinated water molecules, the carboxylate O atoms as well as the N atoms of the cyclam build up a two dimensional network forming a layer parallel to the (0 0 1) plane (Table 1, Fig. 2).

Experimental

An ethanolic solution of cyclam (2.50 mmol, 50 ml) was added to a warm ethanolic solution of dimeric copper(II) didodecanoate (1.25 mmol, 100 ml), forming a clear purple solution. The solution was then gently heated for 2 h. Purple plates formed upon cooling to room temperature. The yield was 60%.

Refinement

All H atoms attached to C atoms and N atom were fixed geometrically and treated as riding on their parent atoms with C—H = 0.98 Å (methyl) or 0.99 Å (methylene) and N—H = 0.93 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C or N})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{Cmethyl})$. H atoms of water molecule were located in difference Fourier maps and included in the subsequent refinement using restraints (O—H = 0.89 (1)Å and H \cdots H = 1.42 (2)Å) with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. In the last cycles of refinement they were treated as riding on their parent O atoms.

Figures




Fig. 1. View of the title compound with the atom labeling scheme. Ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii. Hydrogen bonds are shown as dashed lines. [Symmetry code: (i) $-x+2, -y, -z+2$]




Fig. 2. Partial packing view showing the formation of layer through O-H...O and N-H...O hydrogen bonds. H atoms not involved in hydrogen bondings have been omitted for clarity. Hydrogen bonds are shown as dashed lines. [Symmetry codes: (ii) $-x+1, -y+1, -z+2$; (iii) $x, y-1, z$; (iv) $x+1, y, z$]

Diaqua(1,4,8,11-tetraazacyclotetradecane- $\kappa^4N^1, N^4, N^8, N^{11}$)copper(II) didodecanoate dihydrate

Crystal data

$[\text{Cu}(\text{C}_{10}\text{H}_{24}\text{N}_4)(\text{H}_2\text{O})_2](\text{C}_{12}\text{H}_{23}\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$

$M_r = 734.54$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.9972\ (4)\ \text{\AA}$

$b = 8.8164\ (5)\ \text{\AA}$

$c = 17.1495\ (10)\ \text{\AA}$

$\alpha = 96.218\ (3)^\circ$

$\beta = 99.137\ (3)^\circ$

$\gamma = 98.329\ (3)^\circ$

$V = 1024.13\ (10)\ \text{\AA}^3$

$Z = 1$

$F(000) = 403$

$D_x = 1.191\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2586 reflections

$\theta = 2.8\text{--}27.6^\circ$

$\mu = 0.58\ \text{mm}^{-1}$

$T = 150\ \text{K}$

Plate, violet

$0.41 \times 0.41 \times 0.08\ \text{mm}$

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Sheldrick, 2008)

$T_{\min} = 0.796, T_{\max} = 0.955$

7085 measured reflections

4623 independent reflections

4138 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.045$

$\theta_{\max} = 27.6^\circ, \theta_{\min} = 2.8^\circ$

$h = -9 \rightarrow 8$

$k = -11 \rightarrow 11$

$l = 0 \rightarrow 22$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.047$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.129$	H-atom parameters constrained
$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0591P)^2 + 0.109P]$
4623 reflections	where $P = (F_o^2 + 2F_c^2)/3$
215 parameters	$(\Delta/\sigma)_{\max} = 0.001$
0 restraints	$\Delta\rho_{\max} = 0.54 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.52 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	1.0000	0.0000	1.0000	0.01254 (12)
O2W	0.6641 (2)	0.00995 (16)	0.93529 (9)	0.0230 (3)
H2WA	0.5631	-0.0637	0.9110	0.034*
H2WB	0.6086	0.0954	0.9319	0.034*
N1	1.1080 (2)	0.18383 (18)	0.95170 (10)	0.0151 (3)
H1	1.2439	0.1913	0.9617	0.018*
N2	0.9722 (2)	0.15474 (18)	1.09264 (9)	0.0159 (3)
H2	0.8416	0.1686	1.0853	0.019*
C1	1.1016 (3)	0.0404 (3)	0.81813 (12)	0.0234 (5)
H1A	1.0855	0.0541	0.7610	0.028*
H1B	1.2416	0.0352	0.8367	0.028*
C2	1.0491 (3)	0.1814 (2)	0.86470 (12)	0.0210 (4)
H2A	1.1157	0.2767	0.8484	0.025*
H2B	0.9057	0.1799	0.8518	0.025*
C3	1.0621 (3)	0.3232 (2)	0.99684 (13)	0.0203 (4)
H3A	0.9248	0.3356	0.9778	0.024*
H3B	1.1501	0.4166	0.9888	0.024*
C4	1.0903 (3)	0.3035 (2)	1.08367 (13)	0.0207 (4)

supplementary materials

H4A	1.2308	0.3037	1.1043	0.025*
H4B	1.0478	0.3902	1.1145	0.025*
C5	1.0207 (3)	0.1110 (3)	1.17381 (12)	0.0221 (4)
H5A	0.9975	0.1934	1.2135	0.027*
H5B	1.1615	0.1019	1.1851	0.027*
O1	0.3901 (2)	0.76965 (15)	0.84556 (8)	0.0201 (3)
O2	0.3977 (3)	0.54859 (17)	0.89662 (9)	0.0277 (4)
C6	1.5268 (4)	0.8928 (3)	0.31756 (16)	0.0379 (6)
H6A	1.6020	0.9753	0.3585	0.057*
H6B	1.5357	0.9224	0.2646	0.057*
H6C	1.5804	0.7970	0.3229	0.057*
C7	1.3129 (3)	0.8673 (3)	0.32783 (13)	0.0302 (5)
H7A	1.2590	0.9636	0.3204	0.036*
H7B	1.2380	0.7852	0.2856	0.036*
C8	1.2814 (3)	0.8219 (3)	0.40846 (12)	0.0216 (4)
H8A	1.3601	0.9020	0.4509	0.026*
H8B	1.3301	0.7233	0.4151	0.026*
C9	1.0669 (3)	0.8029 (3)	0.41888 (12)	0.0217 (4)
H9A	1.0194	0.9024	0.4136	0.026*
H9B	0.9879	0.7250	0.3755	0.026*
C10	1.0328 (3)	0.7534 (3)	0.49849 (12)	0.0227 (4)
H10A	1.1154	0.8293	0.5420	0.027*
H10B	1.0755	0.6520	0.5030	0.027*
C11	0.8192 (3)	0.7402 (3)	0.50970 (12)	0.0223 (4)
H11A	0.7362	0.6671	0.4652	0.027*
H11B	0.7779	0.8426	0.5069	0.027*
C12	0.7825 (3)	0.6855 (3)	0.58845 (13)	0.0241 (5)
H12A	0.8202	0.5819	0.5907	0.029*
H12B	0.8676	0.7572	0.6330	0.029*
C13	0.5699 (3)	0.6768 (3)	0.59969 (12)	0.0225 (4)
H13A	0.4846	0.6099	0.5534	0.027*
H13B	0.5348	0.7817	0.6002	0.027*
C14	0.5274 (3)	0.6143 (2)	0.67581 (12)	0.0214 (4)
H14A	0.6137	0.6802	0.7222	0.026*
H14B	0.5598	0.5086	0.6750	0.026*
C15	0.3140 (3)	0.6090 (2)	0.68655 (12)	0.0202 (4)
H15A	0.2270	0.5519	0.6379	0.024*
H15B	0.2853	0.7160	0.6928	0.024*
C16	0.2673 (3)	0.5319 (2)	0.75860 (11)	0.0192 (4)
H16A	0.3117	0.4302	0.7555	0.023*
H16B	0.1231	0.5128	0.7554	0.023*
C17	0.3612 (3)	0.6252 (2)	0.83941 (11)	0.0154 (4)
O1W	0.5329 (2)	0.29510 (16)	0.95510 (9)	0.0208 (3)
H1WA	0.5554	0.3337	1.0070	0.031*
H1WB	0.4965	0.3679	0.9255	0.031*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.01428 (19)	0.01012 (17)	0.01465 (18)	0.00220 (12)	0.00653 (13)	0.00193 (12)
O2W	0.0123 (7)	0.0189 (7)	0.0358 (9)	0.0048 (6)	0.0002 (6)	-0.0017 (6)
N1	0.0121 (8)	0.0136 (8)	0.0210 (8)	0.0032 (6)	0.0051 (6)	0.0044 (6)
N2	0.0122 (8)	0.0170 (8)	0.0185 (8)	0.0028 (6)	0.0044 (6)	-0.0010 (6)
C1	0.0183 (11)	0.0389 (12)	0.0173 (10)	0.0080 (9)	0.0088 (8)	0.0097 (9)
C2	0.0174 (10)	0.0254 (10)	0.0231 (10)	0.0038 (8)	0.0059 (8)	0.0122 (8)
C3	0.0161 (10)	0.0101 (8)	0.0367 (12)	0.0022 (7)	0.0100 (9)	0.0036 (8)
C4	0.0175 (10)	0.0136 (9)	0.0298 (11)	-0.0002 (8)	0.0080 (8)	-0.0042 (8)
C5	0.0207 (11)	0.0306 (11)	0.0149 (9)	0.0069 (9)	0.0037 (8)	-0.0024 (8)
O1	0.0231 (8)	0.0157 (7)	0.0213 (7)	0.0048 (6)	0.0033 (6)	0.0001 (5)
O2	0.0426 (10)	0.0209 (8)	0.0188 (7)	0.0039 (7)	0.0029 (7)	0.0052 (6)
C6	0.0286 (14)	0.0546 (17)	0.0364 (14)	0.0053 (12)	0.0196 (11)	0.0141 (12)
C7	0.0229 (12)	0.0478 (14)	0.0213 (11)	0.0013 (10)	0.0098 (9)	0.0090 (10)
C8	0.0182 (11)	0.0290 (11)	0.0185 (10)	0.0018 (8)	0.0077 (8)	0.0037 (8)
C9	0.0189 (11)	0.0281 (11)	0.0170 (10)	-0.0017 (8)	0.0059 (8)	0.0020 (8)
C10	0.0204 (11)	0.0292 (11)	0.0196 (10)	0.0018 (9)	0.0086 (8)	0.0030 (8)
C11	0.0203 (11)	0.0272 (11)	0.0191 (10)	-0.0008 (8)	0.0089 (8)	0.0000 (8)
C12	0.0216 (11)	0.0313 (12)	0.0215 (11)	0.0041 (9)	0.0100 (9)	0.0044 (9)
C13	0.0225 (11)	0.0275 (11)	0.0190 (10)	0.0017 (9)	0.0104 (8)	0.0025 (8)
C14	0.0201 (11)	0.0271 (11)	0.0182 (10)	0.0040 (8)	0.0072 (8)	0.0028 (8)
C15	0.0214 (11)	0.0253 (10)	0.0137 (9)	0.0019 (8)	0.0065 (8)	-0.0002 (8)
C16	0.0220 (11)	0.0189 (10)	0.0160 (9)	-0.0008 (8)	0.0063 (8)	0.0006 (7)
C17	0.0127 (9)	0.0180 (9)	0.0178 (9)	0.0046 (7)	0.0076 (7)	0.0019 (7)
O1W	0.0201 (8)	0.0195 (7)	0.0234 (7)	0.0054 (6)	0.0033 (6)	0.0025 (6)

Geometric parameters (\AA , $^\circ$)

Cu1—N1 ⁱ	2.0048 (16)	C7—C8	1.521 (3)
Cu1—N1	2.0048 (16)	C7—H7A	0.9900
Cu1—N2 ⁱ	2.0319 (15)	C7—H7B	0.9900
Cu1—N2	2.0319 (15)	C8—C9	1.527 (3)
O2W—H2WA	0.8998	C8—H8A	0.9900
O2W—H2WB	0.8994	C8—H8B	0.9900
N1—C2	1.481 (2)	C9—C10	1.521 (3)
N1—C3	1.483 (2)	C9—H9A	0.9900
N1—H1	0.9300	C9—H9B	0.9900
N2—C4	1.482 (3)	C10—C11	1.527 (3)
N2—C5	1.485 (2)	C10—H10A	0.9900
N2—H2	0.9300	C10—H10B	0.9900
C1—C5 ⁱ	1.510 (3)	C11—C12	1.529 (3)
C1—C2	1.526 (3)	C11—H11A	0.9900
C1—H1A	0.9900	C11—H11B	0.9900
C1—H1B	0.9900	C12—C13	1.523 (3)
C2—H2A	0.9900	C12—H12A	0.9900

supplementary materials

C2—H2B	0.9900	C12—H12B	0.9900
C3—C4	1.503 (3)	C13—C14	1.526 (3)
C3—H3A	0.9900	C13—H13A	0.9900
C3—H3B	0.9900	C13—H13B	0.9900
C4—H4A	0.9900	C14—C15	1.528 (3)
C4—H4B	0.9900	C14—H14A	0.9900
C5—C1 ⁱ	1.510 (3)	C14—H14B	0.9900
C5—H5A	0.9900	C15—C16	1.530 (3)
C5—H5B	0.9900	C15—H15A	0.9900
O1—C17	1.250 (2)	C15—H15B	0.9900
O2—C17	1.264 (2)	C16—C17	1.529 (3)
C6—C7	1.522 (3)	C16—H16A	0.9900
C6—H6A	0.9800	C16—H16B	0.9900
C6—H6B	0.9800	O1W—H1WA	0.8986
C6—H6C	0.9800	O1W—H1WB	0.9006
N1 ⁱ —Cu1—N1	180.0	H7A—C7—H7B	107.6
N1 ⁱ —Cu1—N2 ⁱ	86.18 (7)	C7—C8—C9	113.37 (18)
N1—Cu1—N2 ⁱ	93.82 (7)	C7—C8—H8A	108.9
N1 ⁱ —Cu1—N2	93.82 (7)	C9—C8—H8A	108.9
N1—Cu1—N2	86.18 (7)	C7—C8—H8B	108.9
N2 ⁱ —Cu1—N2	180.000 (1)	C9—C8—H8B	108.9
H2WA—O2W—H2WB	100.8	H8A—C8—H8B	107.7
C2—N1—C3	111.39 (15)	C10—C9—C8	113.87 (17)
C2—N1—Cu1	117.57 (13)	C10—C9—H9A	108.8
C3—N1—Cu1	107.35 (12)	C8—C9—H9A	108.8
C2—N1—H1	106.6	C10—C9—H9B	108.8
C3—N1—H1	106.6	C8—C9—H9B	108.8
Cu1—N1—H1	106.6	H9A—C9—H9B	107.7
C4—N2—C5	112.33 (16)	C9—C10—C11	113.57 (17)
C4—N2—Cu1	106.22 (12)	C9—C10—H10A	108.9
C5—N2—Cu1	116.78 (12)	C11—C10—H10A	108.9
C4—N2—H2	107.0	C9—C10—H10B	108.9
C5—N2—H2	107.0	C11—C10—H10B	108.9
Cu1—N2—H2	107.0	H10A—C10—H10B	107.7
C5 ⁱ —C1—C2	114.02 (17)	C10—C11—C12	113.90 (18)
C5 ⁱ —C1—H1A	108.7	C10—C11—H11A	108.8
C2—C1—H1A	108.7	C12—C11—H11A	108.8
C5 ⁱ —C1—H1B	108.7	C10—C11—H11B	108.8
C2—C1—H1B	108.7	C12—C11—H11B	108.8
H1A—C1—H1B	107.6	H11A—C11—H11B	107.7
N1—C2—C1	111.41 (15)	C13—C12—C11	113.29 (18)
N1—C2—H2A	109.3	C13—C12—H12A	108.9
C1—C2—H2A	109.3	C11—C12—H12A	108.9
N1—C2—H2B	109.3	C13—C12—H12B	108.9
C1—C2—H2B	109.3	C11—C12—H12B	108.9
H2A—C2—H2B	108.0	H12A—C12—H12B	107.7
N1—C3—C4	108.34 (15)	C12—C13—C14	114.26 (18)

N1—C3—H3A	110.0	C12—C13—H13A	108.7
C4—C3—H3A	110.0	C14—C13—H13A	108.7
N1—C3—H3B	110.0	C12—C13—H13B	108.7
C4—C3—H3B	110.0	C14—C13—H13B	108.7
H3A—C3—H3B	108.4	H13A—C13—H13B	107.6
N2—C4—C3	108.71 (16)	C13—C14—C15	113.37 (17)
N2—C4—H4A	109.9	C13—C14—H14A	108.9
C3—C4—H4A	109.9	C15—C14—H14A	108.9
N2—C4—H4B	109.9	C13—C14—H14B	108.9
C3—C4—H4B	109.9	C15—C14—H14B	108.9
H4A—C4—H4B	108.3	H14A—C14—H14B	107.7
N2—C5—C1 ⁱ	111.54 (17)	C14—C15—C16	113.17 (17)
N2—C5—H5A	109.3	C14—C15—H15A	108.9
C1 ⁱ —C5—H5A	109.3	C16—C15—H15A	108.9
N2—C5—H5B	109.3	C14—C15—H15B	108.9
C1 ⁱ —C5—H5B	109.3	C16—C15—H15B	108.9
H5A—C5—H5B	108.0	H15A—C15—H15B	107.8
C7—C6—H6A	109.5	C15—C16—C17	114.68 (17)
C7—C6—H6B	109.5	C15—C16—H16A	108.6
H6A—C6—H6B	109.5	C17—C16—H16A	108.6
C7—C6—H6C	109.5	C15—C16—H16B	108.6
H6A—C6—H6C	109.5	C17—C16—H16B	108.6
H6B—C6—H6C	109.5	H16A—C16—H16B	107.6
C8—C7—C6	114.1 (2)	O1—C17—O2	124.53 (19)
C8—C7—H7A	108.7	O1—C17—C16	118.96 (17)
C6—C7—H7A	108.7	O2—C17—C16	116.46 (17)
C8—C7—H7B	108.7	H1WA—O1W—H1WB	109.5
C6—C7—H7B	108.7		
C3—N1—C2—C1	178.39 (16)	C7—C8—C9—C10	178.49 (19)
Cu1—N1—C2—C1	-57.1 (2)	C8—C9—C10—C11	177.93 (18)
C5 ⁱ —C1—C2—N1	70.0 (2)	C9—C10—C11—C12	178.14 (18)
C2—N1—C3—C4	169.59 (16)	C10—C11—C12—C13	178.57 (17)
Cu1—N1—C3—C4	39.57 (18)	C11—C12—C13—C14	176.93 (18)
C5—N2—C4—C3	168.83 (15)	C12—C13—C14—C15	179.07 (18)
Cu1—N2—C4—C3	40.02 (17)	C13—C14—C15—C16	174.42 (17)
N1—C3—C4—N2	-53.9 (2)	C14—C15—C16—C17	70.0 (2)
C4—N2—C5—C1 ⁱ	179.78 (15)	C15—C16—C17—O1	30.9 (3)
Cu1—N2—C5—C1 ⁱ	-57.2 (2)	C15—C16—C17—O2	-151.66 (19)
C6—C7—C8—C9	177.8 (2)		

Symmetry codes: (i) $-x+2, -y, -z+2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H1WB \cdots O2	0.90	1.91	2.774 (2)	160.
O1W—H1WA \cdots O2 ⁱⁱ	0.90	1.81	2.694 (2)	168.
O2W—H2WB \cdots O1W	0.90	1.93	2.8037 (19)	164.

supplementary materials

O2W—H2WA...O1 ⁱⁱⁱ	0.90	1.89	2.777 (2)	168.
N2—H2...O1 ⁱⁱ	0.93	2.25	3.030 (2)	141.
N1—H1...O1W ^{iv}	0.93	2.12	2.982 (2)	153.

Symmetry codes: (ii) $-x+1, -y+1, -z+2$; (iii) $x, y-1, z$; (iv) $x+1, y, z$.

Fig. 1




Fig. 2

