

Published in final edited form as:

Phys Chem Chem Phys. 2011 April 7; 13(13): 5926–5930. doi:10.1039/c0cp02774j.

A Direct Comparison of Azide and Nitrile Vibrational Probes

Xin Sonia Gai, Basil A. Coutifaris, Scott H. Brewer*, and Edward E. Fenlon*

Franklin & Marshall College, Department of Chemistry, Lancaster, PA 17604-3003 USA

Abstract

The synthesis of 2'-azido-5-cyano-2'-deoxyuridine, **N₃CNdU (1)**, from trityl-protected 2'-amino-2'-deoxyuridine was accomplished in four steps with a 12.5% overall yield. The IR absorption positions and profiles of the azide and nitrile group of **N₃CNdU** were investigated in 14 different solvents and water/DMSO solvent mixtures. The azide probe was superior to the nitrile probe in terms of its extinction coefficient, which is 2–4 times larger. However, the nitrile IR absorbance profile is generally less complicated by accidental Fermi resonance. The IR frequencies of both probes undergo a substantial red shift upon going from water to aprotic solvents such as THF or DMSO. DFT calculations supported the hypothesis that the molecular origin of the higher observed frequency in water is primarily due to hydrogen bonds between the probes and water molecules.

Keywords

Infrared spectroscopy; azide; nitrile; nucleosides; DNA; density functional theory

Introduction

The use of vibrational probes to study biomolecules has a long and rich history. In most of the early work the probe was a ligand bound to hemoproteins.¹ For example, four decades ago McCoy and Caughey used infrared spectroscopy to investigate the binding of ligands such as azide, cyanide, and thiocyanate to metmyoglobin and methemoglobin.² More recently, Suydam and Boxer³ examined a series of vibrational probes for proteins and concluded that the most useful ones have the following properties: (1) an IR absorption band that is narrow, intense, and in a clear region of the spectrum; (2) sensitive to changes in local environment; (3) small size to minimize structural perturbations; and (4) chemical stability. It is therefore no surprise that nitriles and azides have emerged as the most useful vibrational probes, since both meet these criteria. Within the past decade the use of nitriles to study non-hemoproteins has become prevalent, and more recently azides have also been used for this purpose.

In terms of nitriles, the unnatural amino acid (UAA) *p*-cyanophenylalanine (**pCNPhe**) has been inserted into proteins both synthetically^{4–12} and by *in vivo* nonsense suppression.^{13–15} The nitrile of **pCNPhe** served as a vibrational probe to study the hydration and electrostatic environments of several protein systems, including the MLCK peptide–calmodulin complex,⁴ the mastoparan-X peptide,^{5,9} myoglobin,¹³ and ribonuclease S;¹² the latter was

*Corresponding authors., scott.brewer@fandm.edu, Phone: (717) 358-4766, Fax: (717) 291-4343, edward.fenlon@fandm.edu, Phone: (717) 291-4201, Fax: (717) 291-4343.

Electronic supplementary information (ESI) available: Full synthetic procedures; FTIR absorbance spectra of **N₃CNdU** in 14 solvents; line shape analysis of the FTIR absorbance spectrum of **N₃CNdU** in THF and methanol; variable temperature spectra of **N₃CNdU** in water; DFT geometry-optimized structures.

also studied with the UAAs *m*-cyanophenylalanine and *S*-cyanohomocysteine. The chemical conversion of cysteine thiols into thiocyanates or aryl nitriles has been utilized to introduce a vibrational probe into numerous peptides and proteins.^{16–18} The vibrational Stark effect of a nitrile-containing enzyme inhibitor was effectively used to measure the electric field in the active site of human aldose reductase.¹⁹ The UAA 5-cyanotryptophan has also been employed as an infrared probe of protein hydration status.²⁰

In terms of azides, the UAAs azidoalanine and azidohomoalanine have been synthetically incorporated into peptide fragments of the A β peptide and the N-terminal domain of the ribosomal protein L9 (NTL9), respectively, to serve as vibrational probes of protein structure.^{21, 22} An engineered aminoacyl-tRNA synthetase/tRNA pair has been employed to incorporate the UAA *p*-azidophenylalanine (**pN₃Phe**) into the G-protein coupled receptor rhodopsin to study the helix movements during light activation via infrared spectroscopy.^{23, 24} Aromatic azides such as **pN₃Phe** have an additional property of photoreactivity. The instability of this probe is a disadvantage for vibrational experiments but it can be utilized for photocrosslinking experiments.^{25, 26} Azido-nicotinamide adenine dinucleotide was recently synthesized and shown to be a 2D IR probe of the active sites of NAD-dependent enzymes.²⁷ These same workers showed that the structurally related 3-azidopyridine was less suitable as a vibrational probe because of a complex absorption profile resulting from accidental Fermi resonance.²⁸

The use of nitrile and azide vibrational probes in nucleic acids has lagged behind the protein work. These functional groups were first incorporated into nucleic acids decades ago, but the idea to employ them as vibrational probes was first proposed in 2007 when the Stark tuning rates of a number of nitrile- and azido-nucleosides were published.²⁹ Krummel and Zanni³⁰ have demonstrated that the vibrational coupling of ¹⁴N- and ¹⁵N-labeled 5-cyano-2'-deoxyuridines (**CNdU**) can be used to measure distances and angles in a DNA oligomer, while the effects of solvent, hydrogen bonded heterodimer formation, and temperature on the nitrile and azido IR absorbance bands of **CNdU** and 2'-azido-2'-deoxyuridine (**N₃dU**), respectively, have also been reported.^{31, 32} Here, we believe for the first time, both nitrile and azido functional groups have been incorporated into the same biomolecular monomer to facilitate a direct comparison of the vibrational characteristics of these important probes. 2'-Azido-5-cyano-2'-deoxyuridine (**N₃CNdU**, **1**) was synthesized and the IR absorption profiles of the azide and nitrile groups in multiple solvents and solvent mixtures are compared. This work builds upon Cho's comparison of amino acids containing azide, nitrile, or thiocyanate groups in three different solvents.²¹ Future incorporation of **N₃CNdU** into nucleic acid oligomers will allow simultaneous monitoring of *both* the sugar backbone region and the major groove *at a specific nucleotide site* through analysis of the azide and nitrile IR absorbance bands, respectively.

Results and Discussion

Synthetic Chemistry

The synthesis of **N₃CNdU** (**1**) was accomplished in four steps with a 12.5% overall yield, Scheme 1. Trityl-protected 2'-amino-2'-deoxyuridine **2**³³ was subjected to a copper-catalyzed diazotransfer reaction using triflic azide to afford azide **3** in 70% yield. Aromatic bromination of **3** with NBS provided bromide **4** in moderate yield (54%). Cyanation with potassium cyanide followed by detritylation using dichloroacetic acid afforded a 33% yield of **1** for the two steps.

Infrared Spectroscopy

N₃CNdU in Water—The dependence of the position, shape, and complexity of the IR absorbance bands corresponding to the azide asymmetric or the nitrile symmetric stretch of N₃CNdU was explored in a number of solvents and solvent mixtures. The solvents and solvent mixtures were selected to present a wide range of local environments around the azide and nitrile groups of N₃CNdU to compare and contrast the utility of these groups as vibrational probes. Figure 1 shows the normalized FTIR absorbance spectra of N₃CNdU dissolved in water, tetrahydrofuran (THF), or methanol. The IR absorbance spectrum of N₃CNdU in water shows two well-resolved bands at 2124.1 cm⁻¹ and 2242.7 cm⁻¹ corresponding to the azide asymmetric and the nitrile symmetric stretching frequency, respectively. The fwhm of the azide IR absorbance band is ~26 cm⁻¹ compared to a fwhm of ~10 cm⁻¹ for the nitrile IR absorbance band. These experimental results are consistent with previous studies of the single-modified nucleosides: N₃dU³² and CNdU³¹. A third distinct difference between the bands is that the azide IR absorbance band is a factor of two larger than the nitrile IR absorbance band based upon the peak absorbance of each band. This difference increases to a factor of five when considering the relative integrated intensity of each band due to the significant difference in the fwhm.

N₃CNdU in Tetrahydrofuran (THF)—The nitrile IR absorbance band of N₃CNdU dissolved in the aprotic solvent THF is centered at 2233.5 cm⁻¹. This position represents a 9.2 cm⁻¹ red shift of the nitrile stretching frequency in THF relative to water. The higher frequency in water is the result of H-bonding between the nitrile group and water, which is absent in THF. This red shift is consistent with previous experimental studies of the solvent induced frequency shifts of nitrile-modified molecules such as acetonitrile, pCNPhe, thiocyanates, and CNdU.^{4, 31, 34, 35}

The impact of H-bonding with water on the nitrile stretching frequency of N₃CNdU was further explored using DFT calculations at the B3PW91/6-31++G(d,p) level. The calculations resulted in two distinct geometry-optimized H-bonding configurations between the water molecule and the nitrile group of the model 5-cyanouracil (see Supporting Information for the geometry optimized structures). The first geometry was characterized by a C≡N...H angle of 153° between the nitrile group of 5-cyanouracil and the water molecule where a hydrogen atom of the water molecule interacted with the lone pair on the nitrogen atom of the nitrile group. This σ-H-bond resulted in a 8.6 cm⁻¹ blue shift in the nitrile stretching frequency relative to isolated 5-cyanouracil. The C≡N...H angle in the second configuration was 90°, where a hydrogen atom of the water molecule interacted with the π-orbital of the nitrile group. The π-H-bond geometry resulted in a 8.4 cm⁻¹ red shift in the nitrile group. Consequently, the nitrile stretching frequency is dependent on both the presence and geometry of H-bonding interactions with water. A balance of these H-bonding interactions results in the experimentally observed blue shift between the THF and aqueous solutions. The calculations were not in quantitative agreement with the experimental values since the calculations were performed in the gas phase, they do not include effects of anharmonicity, and limitations in the basis set.³⁶⁻⁴⁵ However, these computational results are in agreement with previous theoretical studies exploring the impact of H-bonding interactions between the nitrile group of acetonitrile and water on the nitrile stretching frequency.^{34, 36}

The azide IR absorbance band of N₃CNdU in THF displays a complex absorption profile with a maximum absorbance at 2111.5 cm⁻¹. This band position represents a 12.6 cm⁻¹ red shift relative to its position in water. The azide IR absorbance band for N₃CNdU in THF, however, contains a pronounced high frequency shoulder resulting in an absorbance band comprised of at least two spectral components. The central frequency of the primary

component of the azide IR absorbance band of N_3CNdU in THF is 2109.8 cm^{-1} and the secondary component (the high frequency shoulder) is centered at 2135.2 cm^{-1} . These positions are based upon a line shape analysis of the azide IR absorbance band utilizing two line shape functions described by Eqn. 1 (see Supporting Information). This added complexity is not surprising based upon previous experimental studies of the azide IR absorbance band of aryl azides such as 3-azidopyridine and pN_3Phe .^{23, 28, 46, 47}

The observed complexity of the azide IR absorbance band of N_3CNdU in THF is likely the result of either multiple conformations of N_3CNdU in THF or accidental Fermi resonance involving the azide asymmetric stretch and a normally forbidden combination or overtone band of N_3CNdU . The presence of two distinct conformations giving rise to the complex absorption profile is unlikely since the azide IR absorbance band for N_3dU in THF does not show a pronounced high frequency shoulder (the band is only slightly asymmetric).^{32, 48} Consequently, we interpret the 2109.8 cm^{-1} component to be the azide asymmetric stretching frequency and the secondary component at 2135.2 cm^{-1} to be the normally forbidden combination or overtone band participating in the Fermi resonance. The assignment of Fermi resonance is in agreement with a previous literature study,²⁸ although 2D IR studies of N_3CNdU are needed for confirmation.^{27, 28, 48–53}

Similar to the nitrile IR absorbance band, the observed red shift in the azide IR absorbance band of N_3CNdU in THF relative to water is due to specific solute-solvent interactions (H-bonding) between the azide group and water. The sensitivity of the azide asymmetric stretch to H-bonding with water was investigated by DFT calculations at the B3PW91/6-31++G(d,p) level. The calculations resulted in two geometry-optimized H-bonding configurations. The calculations were performed with the model 2-azido-1,2-dideoxyribose and one explicit water molecule (see Supporting Information for the geometry-optimized structures). The first configuration showed a σ -H-bond between the azide group and a water molecule characterized by a $\text{N}=\text{N}\cdots\text{H}$ angle of 150° . This σ -H-bond formed by an interaction between the hydrogen atom of the water molecule and the terminal nitrogen atom of the azide group resulted in a 7.4 cm^{-1} blue shift in the azide asymmetric stretching frequency relative to isolated 2-azido-1,2-dideoxyribose. The second H-bond geometry involved a π -H-bond between the water molecule and the azide group characterized by a $\text{N}=\text{N}\cdots\text{H}$ angle of 101° . This π -H-bond resulted in a 0.4 cm^{-1} red shift in the azide asymmetric stretching frequency relative to isolated 2-azido-1,2-dideoxyribose. These calculations are in agreement with previous theoretical studies probing the impact of H-bonding between the azide group of methyl azide and water.⁵⁴ The calculations here demonstrate the sensitivity of the azide asymmetric stretch to H-bonding with water, although the calculations are not in quantitative agreement with the experimental results due, in part, to limitations in the basis set.^{41–45}

N_3CNdU in Methanol—The azide and nitrile IR absorbance bands of N_3CNdU dissolved in methanol both display complex absorption profiles. The azide IR absorbance band has an absorbance maximum at 2113.5 cm^{-1} with a high frequency shoulder, which is between the band position observed in water and THF due to the potential of H-bonding between methanol and the azide group. Similar to the azide IR absorbance band of N_3CNdU in THF, the azide IR absorbance band in methanol is comprised of at least two spectral components. The central frequency of the major component is 2111.9 cm^{-1} and the frequency of the minor component is 2135.6 cm^{-1} based upon line shape analysis of the experimental band to line shape functions described by Eqn. 1 (see Supporting Information). Once again, this complex absorption profile could be the result of different conformations of N_3CNdU but is more likely due to accidental Fermi resonance, although 2D IR experiments are needed for a definitive determination.^{27, 28, 48–53}

The nitrile IR absorbance band of N_3CNdU in methanol has an absorbance maximum at 2237.4 cm^{-1} and is comprised of two components positioned at 2236.3 cm^{-1} and 2241.1 cm^{-1} as determined by line shape analysis (see Supporting Information). The complex absorbance band is likely due to two distinct solvation states,^{35, 49, 55} although Fermi resonance has been observed previously for nitrile absorbance bands.^{34, 56, 57} Based upon literature precedence, we attribute the high frequency component to a solvation state where the nitrogen lone pair of the nitrile group is involved in a σ -H-bond with the hydroxyl hydrogen atom of methanol.^{49, 55} This assignment is supported by the similarity of the frequency with the observed position of the nitrile IR absorbance band in water. The low frequency component is due to a solvation state devoid of a σ -H-bond between the nitrile group and water. It is hypothesized that these two distinct solvation states give rise to the complex nitrile absorption profile in methanol.⁴⁹

N_3CNdU in Water/DMSO Mixtures—Figure 2A shows the dependence of the azide and nitrile IR absorbance bands on the systematic variation of the solvent system from water to DMSO in 12.5% increments. These solvent mixtures represent a wide range of local environments for the azide and nitrile groups of N_3CNdU both in terms of solvent dielectric constants and the potential for specific solute-solvent interactions (H-bonding). The FTIR spectrum of N_3CNdU in DMSO is similar to the spectrum in THF in that the azide IR absorbance band shows a complex profile consisting of at least two components while the nitrile IR absorbance profile is a single band. Again, the complexity of the azide IR absorbance band is likely due to accidental Fermi resonance.²⁸

The frequency of maximum absorbance for the azide and nitrile IR absorbance bands of N_3CNdU as a function of the solvent composition is plotted in Figures 2B and 2C, respectively. The positions of both bands decrease monotonically as the percent water is decreased in the water/DMSO solvent mixture. The position of the azide IR absorbance band decreases from 2124.4 to 2108.7 cm^{-1} , while the nitrile IR absorbance band decreases from 2242.7 to 2230.2 cm^{-1} upon going from water to DMSO. The 15.7 cm^{-1} and 12.5 cm^{-1} red shifts are due to changes in H-bond interactions between the azide and nitrile groups with water, respectively. The azide IR absorbance band displays a larger peak extinction for each solvent mixture relative to the nitrile IR absorbance band while exhibiting a larger solvent induced frequency shift. The magnitude of the shift in response to solvent composition highlights the potential utility of azide and nitrile vibrational probes of local environments in nucleic acids.

Additional Solvent and Temperature Studies on N_3CNdU —The IR absorption profiles of N_3CNdU in ten additional solvents (2-propanol, 1-butanol, *tert*-butanol, 2,2,2-trifluoroethanol, acetone, nitromethane, formamide, *N*-methylformamide, *N,N*-dimethylformamide, and propylene carbonate) were investigated (see Supporting Information). The azide IR absorbance band is asymmetric in each of these solvents, but is less pronounced in 2,2,2-trifluoroethanol, nitromethane, and formamide. This complexity is presumably due to accidental Fermi resonance. The nitrile IR absorbance band also showed a complex absorption profile in 2-propanol, 1-butanol, *tert*-butanol, formamide, and *N*-methylformamide (see Supporting Information). In the alcoholic solvents the nitrile complexity is likely due to different solvation states corresponding to H-bonding and non-H-bonding configurations between the hydroxyl group of the solvent and the nitrile group of N_3CNdU . The solvent induced frequency shifts of the vibrational probes did not correlate with solvent parameters such as solvent dielectric constant or solvent dipole moment, similar to previous studies of other nitrile or azide modified molecules.^{21, 34} Additionally, both the azide and nitrile IR bands of N_3CNdU showed a modest dependence on temperature in aqueous solution (see Supporting Information).

Conclusions

The three-atom azide and two-atom nitrile groups are effective vibrational probes of local environments as demonstrated by the solvent dependent positions of the azide asymmetric and nitrile symmetric stretching frequencies in N_3CNdU . On the whole, the azide probe is superior to the nitrile probe because its extinction coefficient is 2–4 times larger. However, the nitrile IR absorbance profile is generally less complicated by accidental Fermi resonance although both the azide and nitrile IR absorbance profiles are effectively a single band in the biologically relevant aqueous solution. The IR frequencies of both probes exhibit a pronounced red shift upon going from water to DMSO (or THF). The frequencies of both observables were higher in water due to H-bonding with water, as confirmed by DFT calculations. As noted previously for 3-azidopyridine,²⁸ the potential complex absorption profiles for the azide and nitrile probes must be considered when using them to study biomolecular structure and dynamics. These complexities^{28, 57, 58} could result in significant difficulties in the interpretation of the infrared signature of azide and nitrile groups in biological systems. A general experimental method to address this issue is currently under investigation.

The azide and nitrile groups of N_3CNdU have the potential to simultaneously probe both the sugar backbone region and the major groove, respectively, of nucleic acids. The vibrational signatures of both groups are well resolved from each other permitting the solvent-dependent azide asymmetric and nitrile symmetric stretching frequencies to yield site-specific information about nucleic acid structure and dynamics. For instance, this minimally invasive modified nucleoside has the potential to aid in the investigation of conformation changes associated with DNA-protein interactions and RNA folding.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to Carol Strausser for assistance in the editing of the manuscript, Lisa Mertzman for obtaining materials and supplies, and Beth Buckwalter for acquiring the NMR spectra. This work was supported by an F&M Snavely Summer Research stipend and a Snavely Research Award to XSG, the William M. and Lucille M. Hackman Scholars Program to BAC, a Mellon/CPC New Tasks/New Tools grant to EEF, and the NIH (R15GM093330) to SHB/EEF.

References

1. Jung C. *J Mol Recognit.* 2000; 13:325–351. [PubMed: 11114067]
2. McCoy S, Caughey WS. *Biochemistry.* 1970; 9:2387–2393. [PubMed: 5423257]
3. Suydam IT, Boxer SG. *Biochemistry.* 2003; 42:12050–12055. [PubMed: 14556636]
4. Getahun Z, Huang CY, Wang T, De Leon B, DeGrado WF, Gai F. *J Am Chem Soc.* 2003; 125:405–411. [PubMed: 12517152]
5. Tucker MJ, Getahun Z, Nanda V, DeGrado WF, Gai F. *J Am Chem Soc.* 2004; 126:5078–5079. [PubMed: 15099085]
6. Aprilakis KN, Taskent H, Raleigh DP. *Biochemistry.* 2007; 46:12308–12313. [PubMed: 17924662]
7. Tucker MJ, Oyola R, Gai F. *J Phys Chem B.* 2005; 109:4788–4795. [PubMed: 16851563]
8. Tucker MJ, Oyola R, Gai F. *Biopolymers.* 2006; 83:571–576. [PubMed: 16917881]
9. Tang J, Signarvic RS, DeGrado WF, Gai F. *Biochemistry.* 2007; 46:13856–13863. [PubMed: 17994771]
10. Marek P, Gupta R, Raleigh DP. *Chembiochem.* 2008; 9:1372–1374. [PubMed: 18478525]

11. Glasscock JM, Zhu YJ, Chowdhury P, Tang J, Gai F. *Biochemistry*. 2008; 47:11070–11076. [PubMed: 18816063]
12. Fafarman AT, Boxer SG. *J Phys Chem B*. 2010; 114:13536–13544. [PubMed: 20883003]
13. Schultz KC, Supekova L, Ryu Y, Xie J, Perera R, Schultz PG. *J Am Chem Soc*. 2006; 128:13984–13985. [PubMed: 17061854]
14. Miyake-Stoner SJ, Miller AM, Hammill JT, Peeler JC, Hess KR, Mehl RA, Brewer SH. *Biochemistry*. 2009; 48:5953–5962. [PubMed: 19492814]
15. Taskent-Sezgin H, Chung J, Patsalo V, Miyake-Stoner SJ, Miller AM, Brewer SH, Mehl RA, Green DF, Raleigh DP, Carrico I. *Biochemistry*. 2009; 48:9040–9046. [PubMed: 19658436]
16. Fafarman AT, Webb LJ, Chuang JI, Boxer SG. *J Am Chem Soc*. 2006; 128:13356–13357. [PubMed: 17031938]
17. McMahon HA, Alfieri KN, Clark CAA, Londergan CH. *J Phys Chem Lett*. 2010; 1:850–855. [PubMed: 20228945]
18. Jo H, Culik RM, Korendovych IV, DeGrado WF, Gai F. *Biochemistry*. 2010; 49:10354–10356. [PubMed: 21077670]
19. Suydam IT, Snow CD, Pande VS, Boxer SG. *Science*. 2006; 313:200–204. [PubMed: 16840693]
20. Waegele MM, Tucker MJ, Gai F. *Chem Phys Lett*. 2009; 478:249–253. [PubMed: 20161057]
21. Oh KI, Lee JH, Joo C, Han H, Cho M. *J Phys Chem B*. 2008; 112:10352–10357. [PubMed: 18671422]
22. Taskent-Sezgin H, Chung JA, Banerjee PS, Nagarajan S, Dyer RB, Carrico I, Raleigh DP. *Angew Chem, Int Ed*. 2010; 49:7473–7475.
23. Ye SX, Huber T, Vogel R, Sakmar TP. *Nat Chem Biol*. 2009; 5:397–399. [PubMed: 19396177]
24. Ye SX, Zaitseva E, Caltabiano G, Schertler GFX, Sakmar TP, Deupi X, Vogel R. *Nature*. 2010; 464:1386–1389. [PubMed: 20383122]
25. Zhang KC, Diehl MR, Tirrell DA. *J Am Chem Soc*. 2005; 127:10136–10137. [PubMed: 16028902]
26. Carrico IS, Maskarinec SA, Heilshorn SC, Mock ML, Liu JC, Nowatzki PJ, Franck C, Ravichandran G, Tirrell DA. *J Am Chem Soc*. 2007; 129:4874–4875. [PubMed: 17397163]
27. Dutta S, Cook RJ, Houtman JCD, Kohen A, Cheatum CM. *Anal Biochem*. 2010; 407:241–246. [PubMed: 20705046]
28. Nydegger MW, Dutta S, Cheatum CM. *J Chem Phys*. 2010; 133:134506. [PubMed: 20942545]
29. Silverman LN, Pitzer ME, Ankomah PO, Boxer SG, Fenlon EE. *J Phys Chem B*. 2007; 111:11611–11613. [PubMed: 17877390]
30. Krummel AT, Zanni MT. *J Phys Chem B*. 2008; 112:1336–1338. [PubMed: 18197662]
31. Watson MD, Gai XS, Gillies AT, Brewer SH, Fenlon EE. *J Phys Chem B*. 2008; 112:13188–13192. [PubMed: 18816094]
32. Gai XS, Fenlon EE, Brewer SH. *J Phys Chem B*. 2010; 114:7958–7966. [PubMed: 20496915]
33. McGee DPC, Vaughn-Settle A, Vargeese C, Zhai YS. *J Org Chem*. 1996; 61:781–785. [PubMed: 11667006]
34. Reimers JR, Hall LE. *J Am Chem Soc*. 1999; 121:3730–3744.
35. Maienschein-Cline MG, Londergan CH. *J Phys Chem A*. 2007; 111:10020–10025. [PubMed: 17867661]
36. Choi JH, Oh KI, Lee H, Lee C, Cho M. *J Chem Phys*. 2008; 128:134506. [PubMed: 18397076]
37. Lindquist BA, Corcelli SA. *J Phys Chem B*. 2008; 112:6301–6303. [PubMed: 18438998]
38. Lindquist BA, Haws RT, Corcelli SA. *J Phys Chem B*. 2008; 112:13991–14001. [PubMed: 18855431]
39. Oh KI, Choi JH, Lee JH, Han JB, Lee H, Cho M. *J Chem Phys*. 2008; 128:154504. [PubMed: 18433232]
40. Waegele MM, Gai F. *J Phys Chem Lett*. 2010; 1:781–786. [PubMed: 20436926]
41. Boese AD, Martin JML, Klopper W. *J Phys Chem A*. 2007; 111:11122–11133. [PubMed: 17929777]
42. Lee JS. *J Chem Phys*. 2007; 127:085104. [PubMed: 17764303]

43. Riley KE, Hobza P. *J Phys Chem A*. 2007; 111:8257–8263. [PubMed: 17649987]
44. Santra B, Michaelides A, Scheffler M. *J Chem Phys*. 2007; 127:184104. [PubMed: 18020627]
45. Scott JN, Nucci NV, Vanderkooi JM. *J Phys Chem A*. 2008; 112:10939–10948. [PubMed: 18839935]
46. Lieber E, Rao CNR, Thomas AE, Oftedahl E, Minnis R, Nambury CVN. *Spectrochim Acta*. 1963; 19:1135–1144.
47. Dyall LK, Kemp JE. *Aust J Chem*. 1967; 20:1395–1402.
48. Tucker MJ, Gai XS, Fenlon EE, Brewer SH, Hochstrasser RM. *Phys Chem Chem Phys*. 2011; 13:2237–2241. [PubMed: 21116553]
49. Kim YS, Hochstrasser RM. *J Phys Chem B*. 2009; 113:8231–8251. [PubMed: 19351162]
50. Jeon J, Yang S, Choi JH, Cho M. *Acc Chem Res*. 2009; 42:1280–1289. [PubMed: 19456096]
51. Ganim Z, Chung HS, Smith AW, Deflores LP, Jones KC, Tokmakoff A. *Acc Chem Res*. 2008; 41:432–441. [PubMed: 18288813]
52. Shim SH, Gupta R, Ling YL, Strasfeld DB, Raleigh DP, Zanni MT. *Proc Natl Acad Sci USA*. 2009; 106:6614–6619. [PubMed: 19346479]
53. Maekawa H, De Poli M, Toniolo C, Ge NH. *J Am Chem Soc*. 2009; 131:2042–2043. [PubMed: 19199613]
54. Choi JH, Oh KI, Cho MH. *J Chem Phys*. 2008; 129:174512. [PubMed: 19045363]
55. Lindquist BA, Furse KE, Corcelli SA. *Phys Chem Chem Phys*. 2009; 11:8119–8132. [PubMed: 19756266]
56. Andrews SS, Boxer SG. *J Phys Chem A*. 2000; 104:11853–11863.
57. Tucker MJ, Kim YS, Hochstrasser RM. *Chem Phys Lett*. 2009; 470:80–84. [PubMed: 20160952]
58. Kinnaman CS, Cremeens ME, Romesberg FE, Corcelli SA. *J Am Chem Soc*. 2006; 128:13334–13335. [PubMed: 17031927]




Figure 1. FTIR absorbance spectra of N_3CNDU dissolved in water, tetrahydrofuran (THF), or methanol recorded at 293 K with a concentration of 50 mM. The maximum absorbance of each spectrum has been normalized to unity and the spectra have been offset for comparison.





Figure 2.

A. FTIR absorbance spectra of N_3CNDU in water/DMSO mixtures ranging from 0 to 100% water in 12.5% increments recorded at 293 K with a concentration of 50 mM. The maximum absorbance of each spectrum has been normalized to unity and the spectra have been offset for comparison. B. Dependence of the position of maximum absorbance of the azide IR absorbance band on solvent composition. C. Dependence of the nitrile IR absorbance band position on solvent composition.



Scheme 1.