

Activation of P2X Receptors Induces Apoptosis in Human Retinal Pigment Epithelium

Dongli Yang,¹ Susan G. Elner,¹ Andrea J. Clark,¹ Bret A. Hughes,^{1,2} Howard R. Petty,^{1,3} and Victor M. Elner^{1,4}

PURPOSE. The retinal pigment epithelium (RPE) is considered a primary site of pathology in age-related macular degeneration (AMD), which is the most prevalent form of irreversible blindness worldwide in the elderly population. Extracellular adenosine triphosphate (ATP) acts as a key signaling molecule in numerous cellular processes, including cell death. The purpose of this study was to determine whether extracellular ATP induces apoptosis in cultured human RPE.

METHODS. RPE apoptosis was evaluated by caspase-3 activation, Hoechst staining, and DNA fragmentation. Intracellular Ca²⁺ levels were determined by both a cell-based fluorometric Ca²⁺ assay and a ratiometric Ca²⁺ imaging technique. P2X₇ mRNA and protein expression were detected by reverse transcription-polymerase chain reaction (RT-PCR) and confocal microscopy, respectively.

RESULTS. The authors found that both the endogenous P2X₇ agonist ATP and the synthetic, selective P2X₇ agonist 2',3'-O-(4-benzoylbenzoyl)-ATP (BzATP) induced RPE apoptosis, which was significantly inhibited by P2X₇ antagonist oxidized ATP (oATP) but not by the P2 receptor antagonist suramin; both ATP and BzATP increase intracellular Ca²⁺ via extracellular Ca²⁺ influx; both ATP- and BzATP-induced Ca²⁺ responses were significantly inhibited by oATP but not by suramin; ATP-induced apoptosis was significantly inhibited or blocked by BAPTA-AM or by low or no extracellular Ca²⁺; and P2X₇ receptor mRNA and protein were expressed in RPE cells.

CONCLUSIONS. These findings suggest that P2X receptors, especially P2X₇ receptors, contribute to ATP- and BzATP-induced Ca²⁺ signaling and apoptosis in the RPE. Abnormal Ca²⁺ homeostasis through the activation of P2X receptors could cause the dysfunction and apoptosis of RPE that underlie AMD. (*Invest Ophthalmol Vis Sci.* 2011;52:1522-1530) DOI:10.1167/iov.10-6172

Extracellular adenosine triphosphate (ATP) acts as a key signaling molecule in numerous cellular processes and is viewed as an endogenous danger signal released in large quantities by cells after inflammation, oxidative stress, and cell

injury.^{1,2} It activates a class of cell-surface nucleotide receptors termed P2 receptors that are further categorized into P2Y receptors and P2X receptors.³ P2 receptors are widely expressed in excitable and nonexcitable cells, where they play important functions.⁴⁻⁸ P2Y receptor expression and function have been reported in human, rat, bovine, and rabbit retinal pigment epithelium (RPE).⁹⁻¹⁸

Little is known of P2X receptors in the RPE. Ryan et al.¹³ suggested that in addition to P2Y receptors, cultured rat RPE cells exhibited functional P2X receptors. Dutot et al.¹⁹ showed that ATP and a selective P2X₇ agonist 2',3'-O-(4-benzoylbenzoyl)-ATP (BzATP) stimulated YO-PRO-1 dye uptake and confocal immunofluorescence microscopy detected P2X₇ receptor protein in a human RPE cell line, ARPE-19 cells. Among seven P2X receptors, the P2X₇ receptor is unique because it plays a critical role in oxidative stress, cell death, and inflammation as well as in several diseases such as Alzheimer's disease and kidney diseases.^{20,21} However, the role of this receptor in the RPE is unknown. Because oxidative stress, cell death, and inflammation are implicated in age-related macular degeneration (AMD) and apoptotic RPE death underlies AMD,^{22,23} we hypothesized that ATP may induce RPE apoptosis by activation of the P2X₇ receptor. By combining molecular, functional, and pharmacologic approaches, we show that the P2X₇ receptor is expressed in native and cultured human RPE and that activation of the P2X₇ receptor induces both Ca²⁺ signaling and apoptosis in RPE cells.

METHODS

Materials

Ninety-six-well black clear-bottom plates were purchased from Fisher Scientific (Costar; Fisher Scientific, Pittsburgh, PA), and 35-mm glass bottom culture dishes were purchased from MatTek Corporation (Ashland, MA). Hoechst 33342, ATP, BzATP, brilliant blue G (BBG), KN-62, suramin, 1,2-Bis (2-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid tetrakis (acetoxymethyl ester) (BAPTA-AM), and oxidized ATP (oATP) were purchased from Sigma-Aldrich (St. Louis, MO). Hanks' balanced salt solutions (HBSS), N-(2-hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid) (HEPES), a RNA isolation reagent (Trizol), Taq DNA polymerase, AlexaFluor 555 goat anti-rabbit IgG, and Indo-1-AM (acetoxymethyl ester) were obtained from Invitrogen (Carlsbad, CA). Rabbit polyclonal anti-P2X₇ antibody was purchased from Abcam, Inc. (Cambridge, MA). Mounting medium with DAPI was purchased from Vector Laboratories (Vectashield; Vector Laboratories, Burlingame, CA). Caspase-3 assay kit was purchased from Biotium, Inc. (NucView 488; Biotium, Inc., Hayward, CA). DNase I (DNAfree) and first-strand synthesis kit (RETROscript) were purchased from Ambion (Austin, TX). Oligonucleotides were synthesized by Integrated DNA Technologies, Inc. (Coralville, IA).

Human RPE Cell Culture

Human RPE cells were isolated from donor eyes by enzymatic digestion as previously described.^{24,25} The protocol adhered to the provisions of

From the Departments of ¹Ophthalmology and Visual Sciences, ²Molecular and Integrative Physiology, ³Microbiology and Immunology, and ⁴Pathology, University of Michigan, Ann Arbor, Michigan.

Supported by National Institutes of Health Grants EY09441 (VME), EY08850 (BAH), CA74120 (HRP), and P30EY07003 (Core). VME is a recipient of a Lew R. Wasserman Merit Award from Research to Prevent Blindness.

Submitted for publication July 6, 2010; revised September 29, 2010; accepted October 8, 2010.

Disclosure: D. Yang, None; S.G. Elner, None; A.J. Clark, None; B.A. Hughes, None; H.R. Petty, None; V.M. Elner, None

Corresponding author: Victor M. Elner, W. K. Kellogg Eye Center, Department of Ophthalmology and Visual Sciences, University of Michigan, 1000 Wall Street, Ann Arbor, MI 48105-0714; velner@umich.edu.

the Declaration of Helsinki for the use of human tissue in research. In all experiments, parallel assays were performed on RPE cells between passages 3 and 6. RPE cells were seeded at the same time and density from the same parent cultures, then grown in phenol red-free complete DMEM/F12 for at least 4 days until 85% to 100% confluence. RPE cells were placed in serum-free media overnight before treatments.

Detection of Caspase-3 Activation

Caspase-3 activation was measured by caspase-3 substrate (NucView 488; Biotium), as described previously.²⁶ Fluorescence intensity of activated caspase-3 was measured by ImageJ software (developed by Wayne Rasband, National Institutes of Health, Bethesda, MD; available at <http://rsb.info.nih.gov/ij/index.html>).

Hoechst Fluorescence Staining

Nuclear staining was performed as a second marker of RPE apoptosis.²⁶ The numbers of stained RPE cells that exhibited apoptotic nuclear condensation and fragmentation were scored as apoptotic. RPE cells from at least 10 high-power microscopic fields from each group of cultures from each of three donors were counted and averaged. Data were normalized to the mean of control RPE cultures.

Double Staining with Caspase-3 Substrate and Hoechst

At the end of control and experimental incubations, RPE cells were successively stained with caspase-3 substrate (NucView 488) for 30 minutes and Hoechst 33342 for 10 minutes at room temperature.²⁶ The caspase-3 substrate is cleaved by activated caspase-3 to release a dye that stains the cell nuclei green, whereas Hoechst 33342 stains the cell nuclei of healthy cells faintly blue and those of the apoptotic cells bluish-white.

Cell Death Detection ELISA

RPE apoptosis was also evaluated by DNA fragmentation, as measured by an ELISA kit (Cell Death Detection ELISA^{PLUS}; Roche Applied Science, Indianapolis, IN) according to procedures outlined by the manufacturer.

Cell-Based Fluorometric Ca²⁺ Assay

Intracellular Ca²⁺ levels were quantitatively determined by cell-based fluorometric Ca²⁺ assay using Indo-1-AM. RPE cells grown on 96-well culture plates were incubated with Indo-1-AM (5 μ M) for 1 hour at 37°C in the dark, after which RPE cells were washed thoroughly, and control medium (HBSS/HEPES), ATP, or BzATP was added to the RPE cells. Indo-1 was excited at 355 nm, and the fluorescence emission from Indo-1 was measured at 405 nm and 485 nm with a fluorometer (FlexStation Scanning Fluorometer; Molecular Devices, Sunnyvale, CA). Fluorescence data were collected at 5-second intervals throughout the course of each experiment. Data are expressed as Indo-1 fluorescence ratios (F405/F485), which are used as a direct index of intracellular Ca²⁺ concentrations ([Ca²⁺]_i).

Ratiometric Calcium Imaging

Intracellular Ca²⁺ levels were also determined using fluorescence microscopy. RPE cells grown on 35-mm glass-bottom culture dishes were labeled as described and then mounted on the stage of an inverted fluorescence microscope (Nikon). Cells were excited at 355 nm, and fluorescence images were collected simultaneously at the dual-emission wavelengths (405 nm and 485 nm) using an imaging apparatus (Dual-View; Optical Insights, Suwanee, GA) and imaging software (MetaFluor Ratio Imaging Software; Universal Imaging Corporation, West Chester, PA). Analysis was carried out using the MetaFluor Analysis software (Universal Imaging Corporation, PA). The fluorescence intensity ratio (F405/F485) was used as a direct index of [Ca²⁺]_i.

Total RNA Isolation

Total RNA was isolated from cultured human RPE cells using a RNA isolation reagent (Trizol; Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. The concentration of total RNA was measured by ultraviolet spectrophotometry.

RT-PCR Analysis

RT-PCR was used to detect P2X₇ mRNA, as described previously.²⁷ PCR was performed with P2X₇-specific primers with the forward primer sequence 5'-GAACCAGCAGCTACTAGGGGAGAAG-3' and the reverse primer sequence 5'-TGGGCAGGTTGGCAAAGTCAGC-3'.²⁸ The house-keeping gene, *hypoxanthine guanine phosphoribosyl transferase 1* (*HPRT1*), served as a control. The forward primer for *HPRT1* was 5'-ACCGTGTGTTAGAAAAGTAAGAAG-3', and the reverse primer was 5'-AGGGAAGTCTGACAAAAGATTC-3'.²⁹ The PCR products were generated by adding *Taq* DNA polymerase and cycled 40 times for P2X₇, or *HPRT1* (1 minute at 94°C, 0.5 minute at 64°C, 0.5 minute at 72°C), followed by a 7-minute extension at 72°C. The RT-PCR products were separated by 1.5% agarose gel electrophoresis.

Confocal Microscopy

Human RPE cells were plated onto 8-well chamber slides and grown for at least 4 days, rinsed twice in PBS, and fixed for 15 minutes in 3% paraformaldehyde in PBS. Cells were then rinsed and permeabilized for 5 minutes at room temperature in 0.2% Triton X-100/PBS. Cells were blocked in 6% BSA/10% normal goat serum in PBS for 1 hour and incubated with 8 μ g/mL rabbit polyclonal anti-P2X₇ antibody diluted in 1% BSA in PBS overnight at 4°C. The specificity of the anti-P2X₇ antibody was confirmed by omitting the primary antibody. The cells were then incubated with 10 μ g/mL Alexa Fluor 555 goat anti-rabbit IgG diluted in 1% BSA in PBS for 2 hours at room temperature. Finally, the cells were washed, mounted in a mounting medium with DAPI (Vectashield; Vector Laboratories), and viewed with confocal microscope (TCS SP5; Leica, Wetzlar, Germany). Digital images were collected.

Statistical Analysis

Data are expressed as mean \pm SD and evaluated by Student's unpaired *t*-test or one-way analysis of variance (ANOVA), followed by a Student-Newman-Keul's post hoc test. *P* < 0.05 is considered statistically significant.

RESULTS

ATP and BzATP Induce RPE Apoptosis

We assessed whether ATP increases RPE apoptosis. We used three apoptotic markers—activated caspase-3, nuclear condensation, and DNA fragmentation—to evaluate the effects of extracellular ATP on the multiple biochemical processes accompanying RPE apoptosis. Figure 1A shows images of RPE cells treated with ATP in the presence or absence of oATP, a P2X₇ antagonist, for 6 hours. ATP induced caspase-3 activation and nuclear condensation, both of which were blocked by oATP. The percentages of RPE cells with apoptotic nuclei, as judged by Hoechst 33342 staining (Fig. 1B; *P* < 0.001), and of RPE cells with activated caspase-3 (Fig. 1C; *P* < 0.05) in ATP-treated RPE cells were statistically greater than those of control RPE cell cultures. The P2X₇ antagonist, oATP, significantly reduced the percentages of RPE cells with apoptotic nuclei (Fig. 1B; *P* < 0.001) and activated caspase-3 (Fig. 1C; *P* < 0.05). At 24 hours, ATP significantly increased DNA fragmentation, as measured by cell death detection ELISA; this increase was completely blocked by oATP (Fig. 1D; *P* < 0.001). Each set of experiments was repeated on RPE cells isolated from three donors.




FIGURE 1. ATP induces apoptosis in human RPE cells. (A) RPE cells were preincubated with or without oxidized ATP (oATP; 300 μ M) and then exposed to 100 μ M ATP with or without oATP for 6 hours. The number of cells with activated caspase-3 and the number of apoptotic RPE cells were determined by double labeling with caspase-3 substrate and Hoechst 33342. Scale bar, 10 μ m. (B) RPE cells were treated and stained by Hoechst 33342 as described in (A), and apoptotic RPE nuclei were determined. (C) RPE cells were treated and stained by caspase-3 substrate as described in (A), and activated caspase 3-positive RPE cells were determined. (D) RPE cells were treated as described in (A) for 24 hours. DNA fragmentation or released mono-nucleosomes and oligonucleosomes were measured by ELISA. Data are presented as mean \pm SD. * P < 0.05 and ** P < 0.001 compared with control. # P < 0.05 and ## P < 0.01, compared with ATP alone.

We next tested the effects of BzATP on RPE apoptosis. BzATP is a synthetic, selective P2X₇ receptor agonist and is widely used in other systems.³⁰⁻³² As shown in Figure 2, BzATP significantly increased RPE apoptosis, which was completely blocked or significantly inhibited by the P2X₇ antagonists BBG (P < 0.001), KN-62 (P < 0.001), and oATP (P < 0.05), whereas suramin had no effect on BzATP-induced RPE apoptosis (P > 0.05), suggesting the involvement of P2X₇ receptors in ATP- and BzATP-induced RPE apoptosis.




FIGURE 2. BzATP induces apoptosis in human RPE cells. RPE cells were treated with 1 mM benzoylbenzoyl adenosine triphosphate (BzATP) in the presence or in the absence of P2 receptor antagonists, brilliant blue G (BBG), KN-62, oxidized ATP (oATP), or suramin for 24 hours and then were stained by caspase-3 substrate. Activated caspase-3 fluorescence intensity was measured. Data are presented as mean \pm SD. *** P < 0.001 compared with control. # P < 0.05 and ### P < 0.001 compared with BzATP alone.

ATP and BzATP Increase RPE Intracellular Ca²⁺ Level

To assess the level of P2X₇ receptor activity, we monitored [Ca²⁺]_i using two different methods: a cell-based fluorometric Ca²⁺ assay and a Ca²⁺ imaging technique. As shown in Figure 3A, 100 μ M ATP induced an increase in [Ca²⁺]_i. The transient rise declined to a level that remained higher than the baseline or control [Ca²⁺]_i. Stimulation with 100 μ M BzATP also produced an increase in [Ca²⁺]_i, but the BzATP-induced signal was lower than that caused by equimolar ATP (Figs. 3A, 3B). The ATP-induced Ca²⁺ peak or sustained Ca²⁺ level at 10 minutes was significantly higher than the control level and the baseline Ca²⁺ reading (Fig. 3B; P < 0.001 or P < 0.01). The BzATP-induced Ca²⁺ peak (Fig. 3B; P < 0.001) and sustained Ca²⁺ level (Fig. 3B; P < 0.05) were significantly higher than baseline and control Ca²⁺ levels. Using ratiometric calcium imaging, we confirmed that ATP-induced Ca²⁺ peak or Ca²⁺ level at 10 minutes is significantly greater than baseline [Ca²⁺]_i (Fig. 3C). This Ca²⁺ imaging result is representative of five independent experiments.

Both ATP and BzATP Trigger Extracellular Ca²⁺ Influx

We then asked whether the ATP- and BzATP-induced increases in intracellular Ca²⁺ were caused by the release of Ca²⁺ from intracellular stores, influx from the extracellular environment, or both. To test this, additional experiments were performed in the presence or absence of extracellular Ca²⁺. In the presence of extracellular Ca²⁺, ATP or BzATP triggered a fast and sustained [Ca²⁺]_i increase (Fig. 4). In the absence of extracellular Ca²⁺, ATP induced a much smaller [Ca²⁺]_i increase in some traces (Fig. 4; RPE cells from donor 1) or no increase at all in other traces (Fig. 4; RPE cells from donors 1 and 2), suggesting that the observed ATP-induced [Ca²⁺]_i was caused mainly by extracellular Ca²⁺ influx. In the absence of extracellular Ca²⁺,




FIGURE 3. ATP and BzATP induce intracellular Ca^{2+} increase in human RPE cells. (A) Temporal plot of F405/F485 ratio changes recorded from Indo-1-AM-labeled RPE cells before and after 100 μM ATP or 100 μM BzATP application. Shown are the mean traces of six to nine traces for each condition from RPE cells derived from three different donors. Arrow: time when ATP, BzATP, or control medium (HBSS/HEPES) was added to the RPE cell cultures. (B) Summary of baseline Ca^{2+} (before ATP application), Ca^{2+} peak, and Ca^{2+} at 10 minutes in RPE cells derived from three different donors. Ca^{2+} concentrations were indicated by F405/F485 ratios. Data represent the mean \pm SD. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$, compared with its own baseline or control. ### $P < 0.001$ compared with ATP alone. (C) Differential interference contrast (DIC) image and pseudocolor ratio (F405/F485) images of a representative field of Indo-1-AM-labeled RPE cells taken before and after ATP application. Scale bar, 20 μm .

the BzATP-induced increase in $[\text{Ca}^{2+}]_i$ was completely blocked in all traces, indicating BzATP-induced $[\text{Ca}^{2+}]_i$ is only from extracellular Ca^{2+} . In addition to the experiments shown in Figure 4, similar results were obtained from five additional independent experiments.

P2 Receptor Antagonists Affect ATP- or BzATP-Induced Ca^{2+} Level

We next asked whether P2 receptor antagonists block the induced Ca^{2+} rise. To this end, we preincubated RPE cells with




FIGURE 4. ATP and BzATP trigger extracellular Ca^{2+} influx in human RPE cells. Human RPE cells (1×10^4) were plated onto a 96-well plate, cultured, and labeled with Indo-1-AM. After washes, the RPE cells were stimulated with 100 μM ATP or 100 μM BzATP in the presence (+ Ca^{2+}) or absence (- Ca^{2+}) of extracellular Ca^{2+} . The responses were measured using a fluorometer. Gray line: individual traces. Black line: mean trace of individual traces in each condition. Arrow: time when ATP or BzATP was added to the RPE cell cultures. Similar results were obtained from five additional independent experiments.

oATP (300 μ M), an irreversible inhibitor of the P2X₇ receptors,^{33,34} or with suramin (50 μ M), which was reported to inhibit the P2X₁, P2X₂, P2X₃, P2X₅, P2Y₁, and P2Y₂ receptors at similar concentrations,^{9,31,35} followed by application of ATP or BzATP. As shown in Figure 5, pretreatment of human RPE cells with oATP dramatically reduced the ATP- and BzATP-induced Ca²⁺ peak ($P < 0.001$), abolished ATP-induced sustained Ca²⁺ levels, and inhibited BzATP-induced sustained Ca²⁺ levels. In contrast, suramin did not inhibit the Ca²⁺ rise evoked by ATP or BzATP (Fig. 6).

Buffering Intracellular Ca²⁺ or Decreasing Extracellular Ca²⁺ Inhibits ATP-Induced Apoptosis

BAPTA-AM is a cell-permeable free calcium chelator that is widely used to reduce intracellular Ca²⁺ levels in many systems, including RPE cells.⁹ If ATP-induced RPE apoptosis is dependent on an increase in [Ca²⁺]_i, then pretreatment of RPE cells with BAPTA-AM would inhibit ATP-induced RPE apoptosis. As expected, BAPTA-AM pretreatment dose dependently inhibited ATP-induced RPE apoptosis, as measured by DNA fragmentation (Fig. 7; $P < 0.01$ and $P < 0.001$ for 5 μ M and 10 μ M BAPTA-AM, respectively). If the extracellular Ca²⁺ influx contributes to the ATP-induced rise in [Ca²⁺]_i, then removing extracellular Ca²⁺ or reducing extracellular Ca²⁺ would be expected to block ATP-induced RPE apoptosis. As shown in Figure 7, removing extracellular Ca²⁺ ($P < 0.01$) or reducing extracellular Ca²⁺ ($P < 0.001$) from normal (1.8 mM) to low (0.3 mM) concentrations significantly inhibited or blocked ATP-induced apoptosis. These results were obtained from RPE cells derived from three different donors.

RPE Expresses P2X₇ receptor

Our data suggest that functionally active P2X₇ receptors are expressed by human RPE cells. To obtain molecular evidence

for this, we performed RT-PCR experiments and confocal immunofluorescence microscopy. Figure 8A shows the results of RT-PCR of RNA extracted from native human RPE cells, cultured human RPE cells derived from three donors, and ARPE-19 cells. RT-PCR of all RPE cells generated single bands with the expected size of 476 bp, demonstrating the presence of P2X₇ receptor mRNA in human RPE cells. To demonstrate that this message yielded P2X₇ receptor protein, confocal immunofluorescence microscopy was performed. As shown in Figure 8B, P2X₇ receptor protein expression was confirmed.

DISCUSSION

This study provides the first evidence that P2X₇ receptors are expressed in native and cultured human RPE cells and that activation of P2X receptors induces both Ca²⁺ signaling and apoptosis. Extracellular ATP can induce apoptosis through the ligation of P2X and P2Y receptors.³⁶ The P2X receptors, particularly P2X₇ receptor, have been shown to play a more important role in the induction of apoptosis than the P2Y receptors.^{37,38}

P2Y Receptors in the RPE

P2Y receptors have been implicated in RPE function, and P2Y mRNA and protein have been identified in cultured human RPE cells, rabbit, rat, and monkey RPE cells.^{12,16-18, 39-41} Activation of P2Y₂ receptors increases [Ca²⁺]_i, RPE fluid transport, and retinal reattachment in rat and rabbit models of retinal detachment.^{9,11,12,14,15,42} In addition to P2Y₂ receptors, P2Y₁ and P2Y₆ receptors were reported to regulate Ca²⁺ levels in cultured human RPE cells,¹⁷ and stimulation of ARPE-19 cells with extracellular nucleotides induced IL-8 gene expression and protein secretion, possibly through P2Y₂ and P2Y₆ receptors.¹⁸ Among the three functional receptors (P2Y₁, P2Y₂, and P2Y₆) identified in human RPE, P2Y₂ receptor can be activated




FIGURE 5. Effects of oxidized ATP on ATP- or BzATP-induced Ca²⁺ in human RPE cells. RPE cells were treated with or without P2X₇ receptor antagonist or oxidized ATP (oATP; 300 μ M) and then were exposed to 100 μ M ATP (A, B) or BzATP (C, D). Histograms show the effects of oATP on the ATP- or BzATP-induced Ca²⁺ peak and the Ca²⁺ at 10 minutes. Data represent the mean \pm SD. * $P < 0.05$ and *** $P < 0.001$, compared with baseline. ### $P < 0.001$ compared with ATP or BzATP alone.




FIGURE 6. Effects of suramin on ATP- or BzATP-induced Ca²⁺ in human RPE cells. The RPE cells were pretreated with or without P2 receptor antagonist, suramin (50 μM), and then exposed to 100 μM ATP or BzATP in the presence (suramin-ATP or suramin-BzATP) or in the absence (ATP or BzATP) of suramin. The responses were measured using a fluorometer. *Gray line*: individual traces. *Black line*: mean trace of individual traces in each condition. *Arrow*: time when ATP or BzATP was added to the RPE cell cultures. Similar results were obtained from RPE cells derived from another donor.

by ATP. Suramin is known to block P2Y₂ receptors in different cell types, including RPE cells.^{9,31,35} However, we found that pretreatment of RPE cells with suramin did not significantly inhibit the ATP-induced increase in intracellular Ca²⁺ levels (Fig. 6). P2X receptors are ATP-gated ion channels that are permeable to Ca²⁺, and the influx of extracellular Ca²⁺ contributes to the initial Ca²⁺ peak when P2X receptors are activated.^{20,43} P2Y receptors are G-protein-coupled receptors, and Ca²⁺ released from intracellular stores contributes to the initial Ca²⁺ increase when P2Y receptors are activated.¹⁷ Therefore, removal of extracellular Ca²⁺ can help determine whether ATP acts on P2X receptors, P2Y receptors, or both. We show here that, in the absence of extracellular Ca²⁺, ATP-induced Ca²⁺ responses were almost completely blocked (Fig. 4), suggesting that P2X receptors rather than P2Y receptors contribute to the ATP-induced Ca²⁺ responses in human RPE cells under our experimental conditions.

Harada et al.³⁷ reported that stimulation of P2Y₂ or P2Y₄ receptors, or both, induced cell proliferation, whereas stimulation of P2X₇ receptors induced cell apoptosis in glomerular mesangial cells. Based on this, we suggest that the relative expression of P2X and P2Y receptors by RPE cells could determine cell fate: proliferation or apoptosis in response to extracellular ATP. Furthermore, local concentrations of extra-

cellular ATP are important in determining cell death. We observed that 100 μM ATP increased human RPE apoptosis as judged by three different approaches (Fig. 1). Consistent with our results, Sugiyama et al.³² showed that ATP at 100 μM, but not 30 μM, significantly increased rat retinal neuron death. Thus, ATP may activate P2Y receptors and P2X₁₋₆ receptors at lower concentrations (<100 μM) and P2X₇ receptors at higher concentrations (≥100 μM) in RPE cells.

P2X Receptors in the RPE

P2X receptors have two transmembrane domains and can form trimeric channels by polymerization of their subunits homomERICALLY or heteromERICALLY. Functional P2X receptors include six homomeric channels and seven heteromeric channels.^{35,44} The ability of P2X receptors to act as direct conduits for Ca²⁺ influx or indirect activators of voltage-gated Ca²⁺ channels underlies their important roles in Ca²⁺-based signaling responses.⁴⁴

ATP was found to be released by RPE cells^{10,16,45-47} and by neural retina^{48,49} and to be capable of acting on P2X receptors in the RPE cells in an autocrine or a paracrine manner. The study by Sullivan et al.¹² supports the presence of functional P2X receptors in the RPE cells. Sullivan et al.¹² applied 100 μM ATP to cultured human RPE cells and found that ATP induced an initial Ca²⁺ peak and sustained rise in [Ca²⁺]_i. In the absence of extracellular Ca²⁺, the ATP-induced Ca²⁺ peak was reduced, and the sustained [Ca²⁺]_i increase was abolished.¹² Ryan et al.¹³ presented evidence that cultured rat RPE cells exhibited functional P2Y and P2X receptors and showed that ATP-induced increases in [Ca²⁺]_i did not depend on extracellular Ca²⁺ influx. The discrepancies between studies could be explained by cell strains or cell lines, cell sensitivity, cell proliferation state, ATP concentration, and exposure time to ATP. Dutot et al.¹⁹ showed that YO-PRO-1 dye uptake was increased in ATP- and BzATP-stimulated ARPE-19 cells and P2X₇ receptor protein was detected in ARPE-19 cells. In this




FIGURE 7. Effects of BAPTA-AM and decreasing extracellular Ca²⁺ on ATP-induced apoptosis in human RPE cells. The RPE cells were pretreated with or without a cell-permeable intracellular Ca²⁺ chelator, BAPTA-AM (5 μM, 10 μM) for 30 minutes and then were exposed to 100 μM ATP for 24 hours in the absence and presence of BAPTA-AM in normal extracellular Ca²⁺ (1.8 mM). RPE cells were also exposed to 100 μM ATP for 24 hours in low extracellular Ca²⁺ (0.3 mM) or in the nominal absence of extracellular Ca²⁺. DNA fragmentation or released mononucleosomes and oligonucleosomes were measured by ELISA. Data are presented as mean ± SD. ****P* < 0.001 compared with control (unstimulated cells in normal extracellular Ca²⁺). ##*P* < 0.01 and ###*P* < 0.001 compared with ATP-stimulated cells in normal extracellular Ca²⁺.




FIGURE 8. P2X₇ receptor expression in human RPE cells. (A) RT-PCR analysis. Total RNAs isolated from native human RPE (lane 1), cultured human RPE cells derived from three donors (lanes 2–4), and ARPE-19 cells (lane 5) were treated with DNase I and reverse transcribed. PCR was performed using a primer set specific for P2X₇ receptor or *HPRT1*, which served as an endogenous control. (B) Immunofluorescence labeling of P2X₇ in human RPE cells. Immunofluorescence labeling omitting primary antibody serves as a negative control. Scale bar, 40 μ m.

study, we detected not only P2X₇ receptor protein but also P2X₇ receptor mRNA in human RPE cells. Our functional data indicate that P2X receptors contribute to both ATP- and BzATP-induced Ca²⁺ increases and apoptosis in human RPE cells because the BzATP- and ATP-triggered Ca²⁺ responses were abolished or largely blocked after the removal of extracellular Ca²⁺. Our findings that the reduction or removal of extracellular Ca²⁺ or the buffering of intracellular Ca²⁺ with BAPTA-AM significantly inhibited or blocked ATP-induced apoptosis suggest that RPE apoptosis is triggered by the ATP-induced rise in [Ca²⁺]_i (Fig. 4). However, ATP-induced apoptosis seems to be lower at low extracellular Ca²⁺ than in the nominal absence of extracellular Ca²⁺ (Fig. 7), indicating that some extracellular Ca²⁺ may be required for RPE cell survival. Our pharmacologic data further support the notion that the P2X₇ receptors contribute to the responses of RPE cells to ATP because ATP-induced Ca²⁺ influx and apoptosis were blocked by oATP, and the selective P2X₇ agonist BzATP induced a Ca²⁺ influx that was significantly inhibited by oATP. Furthermore, BzATP-induced RPE apoptosis was blocked or significantly inhibited by P2X₇ receptor antagonists BBG, KN-62, and oATP. However, Ca²⁺ influx evoked by ATP was higher than that by equimolar BzATP, indicating that in addition to P2X₇, other P2X receptors might be present because BzATP is known as a much better P2X₇ agonist than ATP.⁴³ Both ATP and BzATP-triggered Ca²⁺ influx were insensitive to suramin, suggesting that P2X₄ and P2X₇ receptors may contribute to the induced Ca²⁺ influx given that P2X₁, P2X₂, P2X₃, and P2X₅ receptors, but not P2X₄ and P2X₇ receptors, were found to be sensitive to suramin,^{35,50} and the P2X₆ receptor cannot form a homomeric channel by itself.^{35,44} Further studies are needed to test whether P2X₄ homomeric channels, P2X₄/P2X₇ heteromeric channels, or both are expressed in the RPE. This is important because P2X₄ and P2X₇ receptors are coexpressed in immune cells and epithelial cells, and heteromerization can change both the functional and pharmacological properties of P2X receptors.^{51,52} Future studies to determine whether knockdown of P2X₇ reduces ATP-induced RPE Ca²⁺ responses and apoptosis may further support our findings. Taken together, our results support the idea that P2X, especially P2X₇, receptors mediate ATP-induced Ca²⁺ signaling and apoptosis in human RPE.

Pathophysiological Implications

P2X₇ receptor requires a relatively high ATP concentration to be activated, with a 50% effective concentration (EC₅₀) of approximately 0.1 to 1 mM compared with other P2X receptors with EC₅₀ of approximately 1 to 10 μ M.^{35,53} Given that extracellular ATP is normally in the low micromolar range, the

activation of P2X₇ may be not favored under physiological conditions.

We show here that activation of the P2X receptors, especially the P2X₇ receptor, increases both fast and sustained Ca²⁺ levels within RPE cells and induces RPE apoptosis. We have demonstrated previously that proinflammatory cytokines induce reactive oxygen species in human RPE cells²⁵ and that oxidative stress increases mononuclear phagocyte-induced mouse RPE apoptosis, especially when mononuclear phagocytes were activated by IFN- γ and superoxide dismutase 2 (SOD2) expression was reduced by partial knockout of the *SOD2* gene.²⁶ Proinflammatory cytokines and ATP can be released at sites of inflammation and can upregulate P2X₇ receptor expression in human monocytes, astrocytes, and epithelial cells.^{34,54–57} Thus, it is possible that proinflammatory cytokines and ATP released during pathologic conditions may increase P2X₇ expression in the RPE and thus increase the vulnerability of RPE cells to extracellular ATP-induced apoptosis.

P2X₇ receptor is also involved in inflammation and oxidative stress in many cell types.^{20,58–60} Cell death, inflammation, and oxidative stress are implicated in AMD.^{22,23,61,62} Therefore, defining the role of P2X₇ receptor in the RPE under physiological and pathophysiological conditions could have important implications for the pathogenesis of AMD. Selectively interfering with P2X receptor expression and activation could generate new preventions and therapies for AMD.

In summary, the present study provides the first evidence of functional P2X₇ receptor expressed in human RPE and demonstrates that activation of P2X receptors, especially P2X₇ receptor, induces Ca²⁺ signaling and RPE apoptosis. It is tempting to speculate that the P2X₇ receptor identified here could impact RPE function physiologically and pathologically.

Acknowledgments

The authors thank Donald G. Puro for his constructive and helpful suggestions.

References

- Schwiebert EM, Zsembery A. Extracellular ATP as a signaling molecule for epithelial cells. *Biochim Biophys Acta*. 2003;1615:7–32.
- Wang X, Arcuino G, Takano T, et al. P2X₇ receptor inhibition improves recovery after spinal cord injury. *Nat Med*. 2004;10:821–827.
- Abbracchio MP, Burnstock G. Purinoceptors: are there families of P2X and P2Y purinoceptors? *Pharmacol Ther*. 1994;64:445–475.
- Surprenant A, Rassendren F, Kawashima E, North RA, Buell G. The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X₇). *Science*. 1996;272:735–738.

5. North RA, Barnard EA. Nucleotide receptors. *Curr Opin Neurobiol.* 1997;7:346-357.
6. Rassendren F, Buell GN, Virginio C, Collo G, North RA, Surprenant A. The permeabilizing ATP receptor, P2X₇: cloning and expression of a human cDNA. *J Biol Chem.* 1997;272:5482-5486.
7. MacKenzie AB, Surprenant A, North RA. Functional and molecular diversity of purinergic ion channel receptors. *Ann N Y Acad Sci.* 1999;868:716-729.
8. Garcia-Marcos M, Pochet S, Marino A, Dehaye JP. P2X₇ and phospholipid signalling: the search of the "missing link" in epithelial cells. *Cell Signal.* 2006;18:2098-2104.
9. Peterson WM, Meggyesy C, Yu K, Miller SS. Extracellular ATP activates calcium signaling, ion, and fluid transport in retinal pigment epithelium. *J Neurosci.* 1997;17:2324-2337.
10. Pearson RA, Dale N, Llaudet E, Mobbs P. ATP released via gap junction hemichannels from the pigment epithelium regulates neural retinal progenitor proliferation. *Neuron.* 2005;46:731-744.
11. Stalmans P, Himpens B. Confocal imaging of Ca²⁺ signaling in cultured rat retinal pigment epithelial cells during mechanical and pharmacologic stimulation. *Invest Ophthalmol Vis Sci.* 1997;38:176-187.
12. Sullivan DM, Erb L, Anglade E, Weisman GA, Turner JT, Csaky KG. Identification and characterization of P2Y₂ nucleotide receptors in human retinal pigment epithelial cells. *J Neurosci Res.* 1997;49:43-52.
13. Ryan JS, Baldrige WH, Kelly MEM. Purinergic regulation of cation conductances and intracellular Ca²⁺ in cultured rat retinal pigment epithelial cells. *J Physiol.* 1999;520:745-759.
14. Maminishkis A, Jalickee S, Blaug SA, et al. The P2Y₂ receptor agonist INS37217 stimulates RPE fluid transport in vitro and retinal reattachment in rat. *Invest Ophthalmol Vis Sci.* 2002;43:3555-3566.
15. Takahashi J, Hikichi T, Mori F, Kawahara A, Yoshida A, Peterson WM. Effect of nucleotide P2Y₂ receptor agonists on outward active transport of fluorescein across normal blood-retina barrier in rabbit. *Exp Eye Res.* 2004;78:103-108.
16. Reigada D, Lu W, Zhang X, et al. Degradation of extracellular ATP by the retinal pigment epithelium. *Am J Physiol Cell Physiol.* 2005;289:C617-C624.
17. Tovell VE, Sanderson J. Distinct P2Y receptor subtypes regulate calcium signaling in human retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci.* 2008;49:350-358.
18. Relvas LJ, Bouffloux C, Marcet B, et al. Extracellular nucleotides and interleukin-8 production by ARPE cells: potential role of danger signals in blood-retinal barrier activation. *Invest Ophthalmol Vis Sci.* 2009;50:1241-1246.
19. Dutot M, Liang H, Pauloin T, et al. Effects of toxic cellular stresses and divalent cations on the human P2X₇ cell death receptor. *Mol Vis.* 2008;14:889-897.
20. Khakh BS, North RA. P2X receptors as cell-surface ATP sensors in health and disease. *Nature.* 2006;442:527-532.
21. Turner CM, Elliott JJ, Tam FW. P2 receptors in renal pathophysiology. *Purinergic Signal.* 2009;5:513-520.
22. Zarbin MA. Current concepts in the pathogenesis of age-related macular degeneration. *Arch Ophthalmol.* 2004;122:598-614.
23. Gehrs KM, Anderson DH, Johnson LV, Hageman GS. Age-related macular degeneration—emerging pathogenetic and therapeutic concepts. *Ann Med.* 2006;38:450-471.
24. Elnor SG, Strieter RM, Elnor VM, Rollins BJ, Del Monte MA, Kunkel SL. Monocyte chemotactic protein gene expression by cytokine-treated human retinal pigment epithelial cells. *Lab Invest.* 1991;64:819-825.
25. Yang D, Elnor SG, Bian ZM, Till GO, Petty HR, Elnor VM. Pro-inflammatory cytokines increase reactive oxygen species through mitochondria and NADPH oxidase in cultured RPE cells. *Exp Eye Res.* 2007;85:462-472.
26. Yang D, Elnor SG, Lin LR, Reddy VN, Petty HR, Elnor VM. Association of superoxide anions with retinal pigment epithelial cell apoptosis induced by mononuclear phagocytes. *Invest Ophthalmol Vis Sci.* 2009;50:4998-5005.
27. Yang D, Swaminathan A, Zhang X, Hughes BA. Expression of Kir7.1 and a novel Kir7.1 splice variant in native human retinal pigment epithelium. *Exp Eye Res.* 2008;86:81-91.
28. Cheewatrakoolpong B, Gilchrest H, Anthes JC, Greenfeder S. Identification and characterization of splice variants of the human P2X₇ ATP channel. *Biochem Biophys Res Commun.* 2005;332:17-27.
29. Hageman GS, Anderson DH, Johnson LV, et al. A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci U S A.* 2005;102:7227-7232.
30. Sugiyama T, Kawamura H, Yamanishi S, Kobayashi M, Katsumura K, Puro DG. Regulation of P2X₇-induced pore formation and cell death in pericyte-containing retinal microvessels. *Am J Physiol Cell Physiol.* 2005;288:C568-C576.
31. Zhang X, Zhang M, Laties AM, Mitchell CH. Stimulation of P2X₇ receptors elevates Ca²⁺ and kills retinal ganglion cells. *Invest Ophthalmol Vis Sci.* 2005;46:2183-2191.
32. Sugiyama T, Oku H, Shibata M, Fukuhara M, Yoshida H, Ikeda T. Involvement of P2X₇ receptors in the hypoxia-induced death of rat retinal neurons. *Invest Ophthalmol Vis Sci.* 2010;51:3236-3243.
33. Murgia M, Hanau S, Pizzo P, Ripa M, Di Virgilio F. Oxidized ATP: an irreversible inhibitor of the macrophage purinergic P2Z receptor. *J Biol Chem.* 1993;268:8199-8203.
34. Humphreys BD, Dubyak GR. Induction of the P2z/P2X₇ nucleotide receptor and associated phospholipase D activity by lipopolysaccharide and IFN-gamma in the human THP-1 monocytic cell line. *J Immunol.* 1996;157:5627-5637.
35. North RA, Surprenant A. Pharmacology of cloned P2X receptors. *Annu Rev Pharmacol Toxicol.* 2000;40:563-580.
36. Coutinho-Silva R, Stahl L, Cheung KK, et al. P2X and P2Y purinergic receptors on human intestinal epithelial carcinoma cells: effects of extracellular nucleotides on apoptosis and cell proliferation. *Am J Physiol Gastrointest Liver Physiol.* 2005;288:G1024-G1035.
37. Harada H, Chan CM, Loesch A, Unwin R, Burnstock G. Induction of proliferation and apoptotic cell death via P2Y and P2X receptors, respectively, in rat glomerular mesangial cells. *Kidney Int.* 2000;57:949-958.
38. Sylte MJ, Kuckleburg CJ, Inzana TJ, Bertics PJ, Czuprynski CJ. Stimulation of P2X receptors enhances lipooligosaccharide-mediated apoptosis of endothelial cells. *J Leukoc Biol.* 2005;77:958-965.
39. Cowlen MS, Zhang VZ, Warnock L, Moyer CF, Peterson WM, Yerxa BR. Localization of ocular P2Y₂ receptor gene expression by in situ hybridization. *Exp Eye Res.* 2003;77:77-84.
40. Pintor J, Sánchez-Nogueiro J, Irazu M, Mediero A, Peláez T, Peral A. Immunolocalisation of P2Y receptors in the rat eye. *Purinergic Signal.* 2004;1:83-90.
41. Fries JE, Wheeler-Schilling TH, Guenther E, Kohler K. Expression of P2Y₁, P2Y₂, P2Y₄, and P2Y₆ receptor subtypes in the rat retina. *Invest Ophthalmol Vis Sci.* 2004;45:3410-3417.
42. Meyer CH, Hotta K, Peterson WM, Toth CA, Jaffe GJ. Effect of INS37217, a P2Y₂ receptor agonist, on experimental retinal detachment and electroretinogram in adult rabbits. *Invest Ophthalmol Vis Sci.* 2002;43:3567-3574.
43. North RA. Molecular physiology of P2X receptors. *Physiol Rev.* 2002;82:1013-1067.
44. Dubyak GR. Go it alone no more—P2X₇ joins the society of heteromeric ATP-gated receptor channels. *Mol Pharmacol.* 2007;72:1402-1405.
45. Mitchell CH. Release of ATP by a human retinal pigment epithelial cell line: potential for autocrine stimulation through subretinal space. *J Physiol.* 2001;534:193-202.
46. Eldred JA, Sanderson J, Wormstone M, Reddan JR, Duncan G. Stress-induced ATP release from and growth modulation of human lens and retinal pigment epithelial cells. *Biochem Soc Trans.* 2003;31:1213-1215.
47. Reigada D, Mitchell CH. Release of ATP from RPE cells involves both CFTR and vesicular transport. *Am J Physiol Cell Physiol.* 2005;288:C132-C140.
48. Neal M, Cunningham J. Modulation by endogenous ATP of the light-evoked release of ACh from retinal cholinergic neurones. *Br J Pharmacol.* 1994;113:1085-1087.
49. Newman EA. Glial cell inhibition of neurons by release of ATP. *J Neurosci.* 2003;23:1659-1666.

50. Jones CA, Chessell IP, Simon J, et al. Functional characterization of the P2X₄ receptor orthologues. *Br J Pharmacol*. 2000;129:388-394.
51. Ma W, Korngreen A, Weil S, et al. Pore properties and pharmacological features of the P2X receptor channel in airway ciliated cells. *J Physiol*. 2006;571:503-517.
52. Guo C, Masin M, Qureshi OS, Murrell-Lagnado RD. Evidence for functional P2X₄/P2X₇ heteromeric receptors. *Mol Pharmacol*. 2007;72:1447-1456.
53. Nihei OK, Savino W, Alves LA. Procedures to characterize and study P2Z/P2X₇ purinoceptor: flow cytometry as a promising practical, reliable tool. *Mem Inst Oswaldo Cruz*. 2000;95:415-428.
54. Narcisse L, Scemes E, Zhao Y, Lee SC, Brosnan CF. The cytokine IL-1beta transiently enhances P2X₇ receptor expression and function in human astrocytes. *Glia*. 2005;49:245-258.
55. Turner CM, Tam FW, Lai PC, et al. Increased expression of the pro-apoptotic ATP-sensitive P2X₇ receptor in experimental and human glomerulonephritis. *Nephrol Dial Transplant*. 2007;22:386-395.
56. Welter-Stahl L, da Silva CM, Schachter J, et al. Expression of purinergic receptors and modulation of P2X₇ function by the inflammatory cytokine IFNgamma in human epithelial cells. *Biochim Biophys Acta*. 2009;1788:1176-1187.
57. Al-Shukaili A, Al-Kaabi J, Hassan B. A comparative study of interleukin-1beta production and P2X₇ expression after ATP stimulation by peripheral blood mononuclear cells isolated from rheumatoid arthritis patients and normal healthy controls. *Inflammation*. 2008;31:84-90.
58. Parvathenani LK, Tertyshnikova S, Greco CR, Roberts SB, Robertson B, Posmantur R. P2X₇ mediates superoxide production in primary microglia and is up-regulated in a transgenic mouse model of Alzheimer's disease. *J Biol Chem*. 2003;278:13309-13317.
59. Hewinson J, Mackenzie AB. P2X₇ receptor-mediated reactive oxygen and nitrogen species formation: from receptor to generators. *Biochem Soc Trans*. 2007;35:1168-1170.
60. Hewinson J, Moore SF, Glover C, Watts AG, MacKenzie AB. A key role for redox signaling in rapid P2X₇ receptor-induced IL-1 beta processing in human monocytes. *J Immunol*. 2008;180:8410-8420.
61. Gehrs KM, Jackson JR, Brown EN, Allikmets R, Hageman GS. Complement, age-related macular degeneration and a vision of the future. *Arch Ophthalmol*. 2010;128:349-358.
62. Detrick B, Hooks JJ. Immune regulation in the retina. *Immunol Res*. 2010;47:153-161.