

EDITORIAL

Breast cancer genome heterogeneity: a challenge to personalised medicine?

Charles Swanton^{1,2*}, Rebecca A Burrell¹ and P Andrew Futreal³

Abstract

Implementation of high-throughput genomics sequencing approaches into routine laboratory practice has raised the potential for the identification of multiple breast cancer targets suitable for future therapeutic intervention in order to improve cancer outcomes. Results from these studies have revealed bewildering breast cancer genome complexity with very few aberrations occurring in common between breast cancers. In addition, such complexity is compounded by evidence of genomic heterogeneity occurring within individual breast cancers. Such inter-tumoural and intra-tumoural heterogeneity is likely to present a challenge to personalised therapeutic approaches that might be circumvented through the definition of genome instability mechanisms governing such diversity and their exploitation using synthetic lethal approaches.

Next-generation sequencing approaches have enabled the sequencing of the human cancer genome at unprecedented speed, resolution and cost. Several such studies have recently been reported in both oestrogen receptor-positive and oestrogen receptor-negative breast cancer [1-3]. Results of these cancer-genome sequencing studies have highlighted the tremendous complexity and heterogeneity between cancer genomes from different patients with the same breast cancer histopathological phenotype (inter-tumoural heterogeneity). For example, none of the novel fusion genes identified by Stephens and colleagues were present more than once in any of the 24 cancers studied, and three expressed in-frame fusion genes selected for follow-up were not present in an additional 288 breast cancers studied [2]. In a further

twist to breast cancer complexity, Navin and colleagues have recently described profound heterogeneity within individual breast tumours (intra-tumoural heterogeneity), where multiple tumour subpopulations have been identified, each with distinct genomic profiles [4].

Both patterns of heterogeneity present challenges from a therapeutic perspective. Heterogeneity within an individual tumour raises the likelihood that if driver mutations can be identified and subsequently targeted, resistance to therapy may develop rapidly due to the genomic variation from one cancer cell clone to the next, as has recently been reported in non-small cell lung cancer [5]. Inter-tumoural heterogeneity implies that potentially different driver mutations may be responsible for cancer cell survival and growth from one patient to the next.

Given the cost (approaching \$1 billion [6]) and lead time (10 to 15 years) in drug development, it is economically challenging to develop the next generation of anticancer drugs against each target, suitable for only a small cohort of patients in an individualised approach. Furthermore, the prohibitive costs and challenges imposed by both industry and regulators for combining targeted therapeutics may mitigate against the development of rational drug combinations to target intra-tumoural heterogeneity to limit the acquisition of drug resistance.

Such genomic heterogeneity both between and within individual tumours presents an economically intractable problem requiring a change in drug development strategic approaches. Cancer cell heterogeneity and the continued genomic diversity acquired from one cancer cell division to another may promote cancer cell stress or dependence on alternative cellular pathways that are potentially targetable, as witnessed by success with poly(ADP-ribose) polymerase inhibition in patients who harbour germline BRCA1/2 mutations [7,8].

Recent observations clearly indicate that other patterns of genome instability leading to tumour heterogeneity, initiated by specific defects in the mismatch repair apparatus [9] or chromosome mis-segregation, may also be targetable. Unequal segregation of whole chromosomes at mitosis generates heterogeneity that is

*Correspondence: charles.swanton@cancer.org.uk

¹Translational Cancer Therapeutics Laboratory, Cancer Research UK London Research Institute, 44 Lincoln's Inn Fields, London WC2A 3PX, UK
Full list of author information is available at the end of the article

associated with poor prognosis in solid tumours [10] and early tumour relapse in animal models [11]. Studies in model eukaryotic organisms have identified that aneuploidy is associated with vulnerability to inhibitors of protein folding and synthesis [12]. Finally, evidence is emerging that cancer cell heterogeneity can be a reversible epigenetic event contributing to drug tolerance in cancer cell models that can be attenuated through insulin-like growth factor-1 receptor pathway inhibition [13].

Next-generation sequencing studies have revealed new patterns of genomic instability. Stephens and colleagues identified tandem duplications occurring in large numbers in oestrogen receptor-negative–progesterone receptor-negative breast cancers, and speculate that this pattern of genomic instability may be attributable to an underlying defective DNA maintenance process [2]. Defining the underlying mechanisms responsible for these tandem duplications and potential strategies to exploit them is clearly important.

The identification of common targets upon which tumours rely to sustain and develop heterogeneity is now an experimentally tractable problem in cancer medicine. Inactivation of key cancer cell survival specific to these processes might enhance the efficacy of anticancer drug treatment. Since normal cells may not routinely require such survival pathways due to their genetic identity from cell to cell, the development of anticancer drugs that inactivate genome-instability survival pathways might have an enhanced therapeutic window. Importantly, such an approach may present a more economically viable solution compared with the current strategy of targeting diverse driver mutations in molecularly heterogeneous tumours.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

CS and RB are funded by the Medical Research Council and Cancer Research UK. PAF is funded by the Wellcome trust.

Author details

¹Translational Cancer Therapeutics Laboratory, Cancer Research UK London Research Institute, 44 Lincoln's Inn Fields, London WC2A 3PX, UK. ²Royal Marsden Hospital, Department of Medicine, Breast and Drug Development Units, Downs Road, Sutton SM2 5PT, UK. ³Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SA, UK.

Published: 1 February 2011

References

1. Shah SP, Morin RD, Khattra J, Prentice L, Pugh T, Burleigh A, Delaney A, Gelmon K, Guliany R, Senz J, Steidl C, Holt RA, Jones S, Sun M, Leung G, Moore

- R, Severson T, Taylor GA, Teschendorff AE, Tse K, Turashvili G, Varhol R, Warren RL, Watson P, Zhao Y, Caldas C, Huntsman D, Hirst M, Marra MA, Aparicio S: **Mutational evolution in a lobular breast tumour profiled at single nucleotide resolution.** *Nature* 2009, **461**:809-813.
2. Stephens PJ, McBride DJ, Lin ML, Varela I, Pleasance ED, Simpson JT, Stebbings LA, Leroy C, Edkins S, Mudie LJ, Greenman CD, Jia M, Latimer C, Teague JW, Lau KW, Burton J, Quail MA, Swerdlow H, Churcher C, Natrajan R, Sieuwerts AM, Martens JW, Silver DP, Langerod A, Russnes HE, Foekens JA, Reis-Filho JS, van 't Veer L, Richardson AL, Borresen-Dale AL, *et al.*: **Complex landscapes of somatic rearrangement in human breast cancer genomes.** *Nature* 2009, **462**:1005-1010.
3. Ding L, Ellis MJ, Li S, Larson DE, Chen K, Wallis JW, Harris CC, McLellan MD, Fulton RS, Fulton LL, Abbott RM, Hoog J, Dooling DJ, Koboldt DC, Schmidt H, Kalicki J, Zhang Q, Chen L, Lin L, Wendl MC, McMichael JF, Magrini VJ, Cook L, McGrath SD, Vickery TL, Appelbaum E, Deschryver K, Davies S, Guintoli T, Crowder R, *et al.*: **Genome remodelling in a basal-like breast cancer metastasis and xenograft.** *Nature* 2010, **464**:999-1005.
4. Navin N, Krasnitz A, Rodgers L, Cook K, Meth J, Kendall J, Riggs M, Eberling Y, Troge J, Grubor V, Levy D, Lundin P, Maner S, Zetterberg A, Hicks J, Wigler M: **Inferring tumor progression from genomic heterogeneity.** *Genome Res* 2010, **20**:68-80.
5. Turke AB, Zejnullahu K, Wu YL, Song Y, Dias-Santagata D, Lifshits E, Toschi L, Rogers A, Mok T, Sequist L, Lindeman NI, Murphy C, Akhavanfard S, Yeap BY, Xiao Y, Capelletti M, Iafrate AJ, Lee C, Christensen JG, Engelman JA, Janne PA: **Preexistence and clonal selection of MET amplification in EGFR mutant NSCLC.** *Cancer Cell* 2010, **17**:77-88.
6. DiMasi JA, Hansen RW, Grabowski HG: **The price of innovation: new estimates of drug development costs.** *J Health Econ* 2003, **22**:151-185.
7. Fong PC, Boss DS, Yap TA, Tutt A, Wu P, Mergui-Roelvink M, Mortimer P, Swaisland H, Lau A, O'Connor MJ, Ashworth A, Carmichael J, Kaye SB, Schellens JH, de Bono JS: **Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers.** *N Engl J Med* 2009, **361**:123-134.
8. Tutt A, Robson M, Garber JE, Domchek SM, Audeh MW, Weitzel JN, Friedlander M, Arun B, Loman N, Schmutzler RK, Wardley A, Mitchell G, Earl H, Wickens M, Carmichael J: **Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and advanced breast cancer: a proof-of-concept trial.** *Lancet* 2010, **376**:235-244.
9. Martin SA, McCabe N, Mullarkey M, Cummins R, Burgess DJ, Nakabeppu Y, Oka S, Kay E, Lord CJ, Ashworth A: **DNA polymerases as potential therapeutic targets for cancers deficient in the DNA mismatch repair proteins MSH2 or MLH1.** *Cancer Cell* 2010, **17**:235-248.
10. Walthers A, Houlston R, Tomlinson I: **Association between chromosomal instability and prognosis in colorectal cancer: a meta-analysis.** *Gut* 2008, **57**:941-950.
11. Sotillo R, Schwartzman JM, Socci ND, Benezra R: **Mad2-induced chromosome instability leads to lung tumour relapse after oncogene withdrawal.** *Nature* 2010, **464**:436-440.
12. Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, Amon A: **Effects of aneuploidy on cellular physiology and cell division in haploid yeast.** *Science* 2007, **317**:916-924.
13. Sharma SV, Lee DY, Li B, Quinlan MP, Takahashi F, Maheswaran S, McDermott U, Azizian N, Zou L, Fischbach MA, Wong KK, Brandstetter K, Wittner B, Ramaswamy S, Classon M, Settleman J: **A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations.** *Cell* 2010, **141**:69-80.

doi:10.1186/bcr2807

Cite this article as: Swanton C, *et al.*: Breast cancer genome heterogeneity: a challenge to personalised medicine? *Breast Cancer Research* 2011, **13**:104.