

Published in final edited form as:

*Schizophr Res.* 2011 July ; 129(2-3): 128–132. doi:10.1016/j.schres.2011.03.021.

## Paternal age and psychiatric disorders: findings from a Dutch population registry

Jacobine E Buizer-Voskamp<sup>a,b</sup>, Wijnand Laan<sup>c</sup>, Wouter G Staal<sup>d</sup>, Eric AM Hennekam<sup>e</sup>, Maartje F Aukes<sup>a,c</sup>, Fabian Termorshuizen<sup>c</sup>, René S Kahn<sup>a</sup> for GROUP Consortium, Marco PM Boks<sup>a,c</sup>, and Roel A Ophoff<sup>a,b,f</sup>

Jacobine E Buizer-Voskamp: J.E.Buizer@umcutrecht.nl; Wijnand Laan: Wijnand.Laan@gmail.com; Wouter G Staal: W.Staal@umcutrecht.nl; Eric AM Hennekam: F.A.M.Hennekam@umcutrecht.nl; Maartje F Aukes: M.Aukes@umcutrecht.nl; Fabian Termorshuizen: F.Termorshuizen@umcutrecht.nl; René S Kahn: R.Kahn@umcutrecht.nl; Marco PM Boks: M.P.M.Boks@umcutrecht.nl

<sup>a</sup>Rudolf Magnus Institute of Neuroscience, Department of Psychiatry, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands <sup>b</sup>Department of Medical Genetics, University Medical Center Utrecht, Universiteitsweg 100, 3584 CG Utrecht, The Netherlands <sup>c</sup>Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Universiteitsweg 100, 3584 CG Utrecht, The Netherlands <sup>d</sup>Department of Child and Adolescent Psychiatry, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands <sup>e</sup>Department of Clinical Genetics, University Medical Center Utrecht, Postbus 85090, 3508 AB Utrecht, The Netherlands <sup>f</sup>Center for Neurobehavioral Genetics, University of California, 695 Charles E. Young Drive South, Los Angeles, CA 90095-1761, USA

### Abstract

**Background**—We measured the association between paternal age and schizophrenia (SCZ), autism spectrum disorders (ASD), major depressive disorder (MDD), and bipolar disorder (BPD) in the Dutch population.

**Methods**—In total, 14 231 patients and 56 924 matched controls were collected and analysed for an association with paternal age by logistic regression.

**Results**—ASD is significantly associated with increased paternal age: Older fathers >40 years of age have a 3.3 times increased odds of having a child with ASD compared to young fathers <20 years of age. SCZ has significant associations for fathers aged >35 years (OR=1.27, 95% Confidence Interval: 1.05, 1.53). For MDD, both younger and older fathers have increased odds. No association was found for BPD.

**Conclusions**—The effects of paternal age as a risk factor are different for ASD and SCZ on one hand, and the affective disorders on the other hand. Different types of association might indicate different biological or psychosocial mechanisms. Late paternity (associated with predispositions to psychiatric disorders) seems the most probable explanation for the association with paternal age.

© 2011 Elsevier B.V. All rights reserved.

Corresponding author: Roel A Ophoff, Center for Neurobehavioral Genetics, University of California, Gonda Center #4357C, Box 951761, 695 Charles E. Young Drive South, Los Angeles, CA 90095-1761, USA; Tel: +1 310 794 9602; Fax: +1 310 794 9613; ROphoff@mednet.ucla.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Keywords

Bipolar Disorder; Child Development Disorders; Pervasive; Depressive Disorder; Major; Paternal Age; Schizophrenia

---

## 1.1 Introduction

Psychiatric disorders show substantial similarities in symptoms. Also, it is increasingly clear that they share genetic and environmental risk factors (Hyman, 2007), which likely operate through mediating characteristics that alter risk for a number of different outcomes (Dick *et al.*, 2010). One of those possible shared vulnerability factors is increased paternal age. Accumulating evidence from epidemiological studies has suggested an association between increased paternal age and complex disorders, such as autism, schizophrenia, and bipolar disorder (Dalman and Allebeck, 2002; Frans *et al.*, 2008; Malaspina *et al.*, 2001; Shelton *et al.*, 2010; Sipos *et al.*, 2004). Different associations have been found between paternal age and psychiatric disorders in the offspring. So, the question remains whether paternal age has different risk effects for major psychiatric disorders such as autism spectrum disorders (ASD), schizophrenia (SCZ), major depressive disorder (MDD), and bipolar disorder (BPD) in the offspring.

In total, 17 cohort studies and 10 case-control studies investigated the association between increased paternal age and SCZ, ASD, MDD and/or BPD in the offspring (Brown *et al.*, 2002; Byrne *et al.*, 2003; Croen *et al.*, 2007; Dalman and Allebeck, 2002; Durkin *et al.*, 2008; Ekeus *et al.*, 2006; Frans *et al.*, 2008; Gillberg, 1982; Glasson *et al.*, 2004; Grether *et al.*, 2009; King *et al.*, 2009; Lauritsen *et al.*, 2005; Laursen *et al.*, 2007; Lopez-Castroman *et al.*, 2009; Malaspina *et al.*, 2002; Malaspina *et al.*, 2001; Menezes *et al.*, 2010; Petersen *et al.*, 2011; Rasmussen, 2006; Reichenberg *et al.*, 2006; Sasanfar *et al.*, 2010; Shelton *et al.*, 2010; Sipos *et al.*, 2004; Torrey *et al.*, 2009; Tsuchiya *et al.*, 2008; Tsuchiya *et al.*, 2005; Zammit *et al.*, 2003). Several studies with up to 13 297 patients (Miller *et al.*, 2010) found that higher paternal age doubles or triples the risk for SCZ in the offspring of men above 40 years of age (Brown *et al.*, 2002; Byrne *et al.*, 2003; Dalman and Allebeck, 2002; Laursen *et al.*, 2007; Malaspina *et al.*, 2002; Malaspina *et al.*, 2001; Rasmussen, 2006; Sipos *et al.*, 2004; Torrey *et al.*, 2009; Tsuchiya *et al.*, 2005; Zammit *et al.*, 2003). ASD was reported to be associated with both increased maternal and paternal age, with a recent emphasis for an increased paternal age (Croen *et al.*, 2007; Durkin *et al.*, 2008; Glasson *et al.*, 2004; Grether *et al.*, 2009; King *et al.*, 2009; Lauritsen *et al.*, 2005; Reichenberg *et al.*, 2006; Sasanfar *et al.*, 2010; Shelton *et al.*, 2010; Tsuchiya *et al.*, 2008). The number of ASD patients ranged from a few hundred for case-control studies to thousands of patients within birth cohort studies. However, for both SCZ and ASD the results remain inconsistent. Results include reports that the age of both parents is associated, that only one is associated (maternal or paternal), and that there is no association of SCZ and ASD with parental age. In contrast to the large number of studies investigating the association between paternal age and SCZ and ASD, data on the association with BPD or MDD are rare (Menezes *et al.*, 2010). Two studies reported an association between increased paternal age and the risk of BPD in the offspring (Frans *et al.* 2008 (Frans *et al.*, 2008): 13 428 BPD patients and Menezes *et al.* 2010 (Menezes *et al.*, 2010): 493 BPD patients), but we are not aware of studies that investigate the relation between paternal age and MDD.

A systematic analysis of paternal age is necessary to assess its relevance as risk factor for SCZ, ASD, MDD, and BPD in the offspring. On one hand, lack of specificity in the association between increased paternal age and psychiatric diagnoses suggests that paternal age is related to phenotypes shared among disorders or associated with common – perhaps

genetic – etiology (Smith *et al.*, 2009). On the other hand, diverse associations between paternal age and psychiatric diagnoses may reflect different risk effects and therefore other biological or psychosocial mechanisms. Few studies examined the effect of paternal age across different diagnostic categories simultaneously (Ekeus *et al.*, 2006; Gillberg, 1982; Lopez-Castroman *et al.*, 2009), but none of these included different DSM-IV-TR diagnoses. In this study we examine the association between paternal age and SCZ, ASD, MDD, and BPD in a single large Dutch population-based sample. As far as we are aware, this is the first large-scale population-based study to compare the four main DSM-IV-TR psychiatric diagnoses on paternal age in a single homogeneous cohort.

## 1.2 Materials and Methods

### 1.2.1 Subjects

Patients were collected through the Psychiatric Case Registry Middle Netherlands (PCR-MN), a case registry from the central part of the Netherlands. The PCR-MN contains data on psychiatric in- and outpatient admissions and diagnoses. Currently over 175 000 cases are registered in the PCR-MN. Patients diagnosed with SCZ, ASD, MDD, or BPD between January 1999 and December 2008 were identified and included in the study. Each diagnosis was defined by the corresponding codes from DSM-IV-TR. Disorders were categorized as schizophrenia spectrum disorder if they were given a DSM-IV code of 295.1 - 295.4, 295.6, 295.7, 295.9, 298.8, and 298.9; as autism spectrum disorder if they received a DSM-IV code of 299.00, 299.10, or 299.80; as major depressive disorder if they were given a DSM-IV code of 296.20 – 296.26, 296.30 – 296.36 or 311.00; and as bipolar disorder if they received a DSM-IV code of 296.00 – 296.06, 296.40 – 296.46, 296.50 – 296.56, 296.60 – 296.66, 296.70, 296.80, and 296.89. Patients were linked to a record in the civil registry of Statistics Netherlands (CBS) based upon gender, date of birth, and postal code. CBS makes individual, but anonymous data of the Dutch population available for scientific research. Among these characteristics is the parent-children registry. The likelihood to identify both parents of a patient within the civil registry decreases from 90% for those born after 1987 to about zero percent for those born before 1947. Therefore we only included those who were born after 1947. Twins were excluded because it is not possible to differentiate between these two siblings if they live in the same postal-code area. A fourfold number of matched control subjects was collected from the CBS database for each diagnostic group, matched for year of birth, place of birth, and gender. The data available in the PCR-MN is processed in accordance with its security and confidentiality policy. The use of data linked to the civil registry is allowed by Dutch CBS-law article 41 (Statistics-Netherlands, 2004). The use of the civil registry is approved by the Dutch law for the protection of privacy and is monitored by the Central Commission for Statistics of the CBS.

### 1.2.2 Parental Age Determination and Statistical Analyses

Parental age at time of birth of the proband was calculated in months. We investigated paternal age as a categorical measure by logistic regression (using Statistical Package for the Social Sciences (SPSS), 14<sup>th</sup> edition). The level of significance was set to 5% and the test was performed two-tailed. The reference group was set at age 25-29 years (in accordance to a recent meta-analysis by Miller *et al.*, 2010 (Miller *et al.*, 2010)).

Social economic status and ethnic background were considered as confounders. The average income of the residential area at time of birth of the patient was taken as a proxy for social economic status. The ethnic background was categorized into three groups: native Dutch, non-Dutch Western European, or non-Western European.

We investigated the possible influence of maternal age by performing different analyses for mothers aged <30 years and >30 years. We also performed separate analyses for male and female offspring, as it has been suggested that paternal age has a sexually dimorphic effect for psychiatric traits (Miller *et al.*, 2010).

### 1.3 Results

Of all patients with a SCZ, ASD, MDD or BPD diagnosis identified in the PCR-MN database, 79.6% could be uniquely linked to a record in the CBS database. Of the 20.4% that could not be linked 11.9% gave more than one unique match and 8.5% gave no match to a record in the civil registry. Of those who were linked to a unique record we could identify both parents for 69.7% of cases. This resulted in a final sample of 2564 SCZ patients, 2262 ASD patients, 8284 patients with MDD and 1121 patients with BPD. A total of 56 924 matched controls were included through the CBS database for the different diagnostic groups (Table 1).

Several studies report age of the other parent as a strong confounder. However, paternal and maternal ages are suggested to be highly related (Fokkema *et al.*, 2008; Olsen and Zhu, 2009). We examined this by computing the Pearson's correlation coefficient. Paternal and maternal ages were highly correlated for all diagnostic groups, with all coefficients between the 0.7 and 0.8 (fathers are 2-3 years older than mothers). Because of this strong relatedness it is not possible to investigate both age effects in the continuous model. We therefore focused on paternal age effects only. The strong correlation makes it also conceivable that paternal age effects can be caused by factors associated with an increased age difference between father and mother instead (Durkin *et al.*, 2008). We therefore also adjusted for the age difference between parents (together with social economic status and ethnic background). This would cover both the confounders of age difference and age of the other parent.

Logistic regression shows a significant association between increased paternal age and ASD risk in the children: Older fathers at >40 years of age have a 3.3 times increased odds ratios (OR) of having a child with ASD compared to young fathers <20 years of age. Also the prevalence of SCZ in the offspring is significantly associated with increased paternal age. Older fathers of 35 years of age and above show significant odds ratios compared to fathers at 25-29 years of age. MDD shows a different pattern of association compared to SCZ and ASD. There is not a linear association with paternal age, but a U-shaped relationship: both the youngest and oldest paternal age categories show significantly increased odds ratios of MDD in their offspring. Our results provide no evidence for any association between BPD and paternal age (Table 2). Maternal age shows no additional association effects for any diagnostic category for the separate analyses of younger (<30 years of age) and older (>30 years of age) mothers. Also gender has no impact on the rate of any psychiatric disorder in the model (data not shown).

It is possible that these paternal age effects in patients from the case registry are not generalizable to differently collected samples (Berkson, 1946; Parkes *et al.*, 2006). We investigated possible differences between these population-based patients and those collected in hospitals with the use of two available clinical samples for SCZ and ASD. We found both clinic-based samples to differ significantly in paternal age from our patients from the population-based registry, showing that the way in which subjects are collected can be a source of bias (see Supplemental Information for methods and results).

## 1.4 Discussion

In this study we investigated 14 231 patient and 56 924 controls from the Dutch population on the relevance of paternal age as a risk factor for psychiatric disorders in the offspring. We find significant associations between paternal age and autism spectrum disorders (ASD), schizophrenia (SCZ), and major depressive disorder (MDD), but not for bipolar disorder (BPD). Interestingly, population demographics in the Western world show a clear increase in father's mean age at childbirth. Dutch fathers have one of the highest ages at birth in Europe: 34.2 years of age in 2006 (Fokkema *et al.*, 2008). Our analyses show that the prevalence of ASD in the offspring is significantly associated with an increased paternal age, both in the continuous analysis and in the analysis using age categories. Older fathers (>40 years of age) have a 3.3 times increased odds for having a child with ASD compared to young fathers (<20 years of age). A similar trend is seen for SCZ: fathers above 35 years of age have significantly higher odds ratios of having a child with the disorder. The relationship between MDD in the offspring and paternal age is U-shaped, meaning that both younger and older fathers are associated with a higher prevalence of MDD in the offspring. This is a markedly different pattern from that observed for ASD and SCZ. It is noteworthy that we find no evidence that BPD is associated with paternal age.

The linear association between increased paternal age and ASD and SCZ in the offspring is most commonly found in literature (Miller *et al.*, 2010). For the affective disorders the evidence for paternal age as a risk factor is less pronounced. We found only one study to describe a U-shaped relationship as we found for MDD (Krishnaswamy *et al.*, 2009). However, this study describes the relation between younger and older fathers with increased rate of common mental disorders. Our negative findings for BPD contrast with two previous studies (Frans *et al.*, 2008; Menezes *et al.*, 2010) reporting increased paternal age to be associated with BPD in the offspring.

Our clearly different effects for paternal age as a risk factor for ASD and SCZ on one hand and the affective disorders on the other hand is striking. Gillberg *et al.* (Gillberg, 1982) described already in 1982 that increased parental age might have different effects on the risk of psychiatric disorders in the offspring. Our results for ASD and SCZ contrasting MDD and BPD are another indication that paternal age might have different risk effects for the major psychiatric disorders.

Different types of association could reflect different biological or psychosocial mechanisms. A molecular mechanism for the associations could be the occurrence of *de novo* mutations or epigenetic changes in the germline (Crow, 2003; McClellan *et al.*, 2007). Diminished DNA repair mechanisms, declining testosterone levels and other circumstances during mitotic divisions associated with late paternal age could lead to new mutations arising in sperm (Bassett *et al.*, 2010; Lopez-Castroman *et al.*, 2009). The fact that neuropsychiatric disorders are influenced by a large number of individually rare deleterious mutations, many of which have occurred in the present or recent generations, may favor this hypothesis (Carroll and Owen, 2009; McClellan *et al.*, 2007). However, it is not clear how a biological explanation would only apply to ASD, SCZ (and possibly MDD), but not to BPD.

Another important explanation could be confounding by genetic vulnerability. Older fathers may have personality traits (such as schizotypal or autistiform traits), which would result in difficulty establishing relationships and thereby an increased age of parenthood. Because of the high heritability, the presence of these personality traits in the fathers could result in an increased risk of developing a psychiatric disorder in the offspring (Zammit *et al.*, 2003). The finding that ASD, SCZ, and MDD, but not BPD are associated with increased paternal age supports this hypothesis, since recent findings show that patients in the premorbid phase



of BPD have improved social and cognitive functions. Also, in contrast to ASD and SCZ, BPD is not associated with reduced fecundity (MacCabe *et al.*, 2010). These findings suggest that fathers with bipolar sub clinical symptoms may not suffer from diminished social functioning explaining the lack of an increased paternal age of patients with BPD.

The hypothesis of an association between late paternity and predisposition to psychiatric disorders is supported by a recent study by Petersen et al. (Petersen et al., 2011). They confirm the association between paternal age and schizophrenia. However, they also show that the risk of SCZ did not depend on paternal age at birth of the proband, but rather on the age at which the father had his first child. The findings of Petersen et al. are the opposite of what would be predicted if de novo mutations or epigenetic changes were a major reason for the effect of paternal age.

It is possible that multiple exposures may increase the risk of psychiatric disorders through shared pathways (i.e. genetics, epigenetics, hormonal alterations, environmental factors). This will be represented as a generalized increased risk with paternal age (Shelton *et al.*, 2010). In accordance to recent findings, our results suggest that the association with increased paternal age is due to factors related to higher paternal age at first fatherhood.

An important strength of our study is that we investigated the four major psychiatric disorders simultaneously, with large sample sizes for each diagnostic group. We used data from a psychiatric case registry, which records all psychiatric diagnosis of the complete population living in the study area. Therefore there is little risk of selection bias, which may occur in clinically selected samples. However, some residual bias may still be present. As can be seen in table 1, the ASD subjects are from a younger generation than the other diagnostic groups, which could bias our results. These different generations reflect the rising of father's mean age at childbirth during the past decades, with older fathers in the ASD group compared to the other groups. However, the age of fathers from the ASD matched controls is also higher compared to the other matched control groups. This generation effect for ASD should therefore have no effect on the outcome of the analyses. Even with the recently highlighted under-diagnosis of ASD there is no reason to assume that this bias is systematic towards children with younger parents, although it would be very interesting to investigate this in more detail. Secondly, diagnoses within the PCR-MN are not based on standardized interviews that are ordinarily used for research purposes. However, because the diagnoses are made in a clinical context by experienced psychiatrists trained to use DSM-IV diagnostic criteria, we do not expect diagnoses to be much different when based on standardized questionnaires and interviews and we particularly do not expect a systematic bias in diagnosis related to paternal age. Finally, the absence of information on non-paternity is a limitation. However, we expect this to be equal for cases and controls.

In conclusion, in this large-scale population-based study we report an association between paternal age and the prevalence of ASD, SCZ, and MDD, but not BPD in the offspring. Whatever the cause behind paternal age as risk factor for psychiatric traits in the children, our results show clear differences between ASD and SCZ compared to the affective disorders. This could reflect different biological or psychosocial mechanisms. Most likely the results can be explained by social selection as pointed out by Petersen et al. (Petersen et al., 2011) who reported that not age at birth of the schizophrenia patient but rather paternal age at birth of the first child explains the association. Our differential effects of paternal age in SCZ, ASD, and affective disorders are in line with this finding.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## References

- Bassett A, Scherer W, Brzustowicz L. Copy number variations in schizophrenia: critical review and new perspectives on concepts of genetics and disease. *Am J Psychiatry*. 2010; 167(8):899–914. [PubMed: 20439386]
- Berkson J. Limitations of the application of fourfold table analysis to hospital data. *Biometrics Bull*. 1946; 2(3):47–53.
- Brown AS, Schaefer CA, Wyatt RJ, Begg MD, Goetz R, Bresnahan MA, Harkavy-Friedman J, Gorman JM, Malaspina D, Susser ES. Paternal age and risk of schizophrenia in adult offspring. *Am J Psychiatry*. 2002; 159(9):1528–1533. [PubMed: 12202273]
- Byrne M, Agerbo E, Ewald H, Eaton WW, Mortensen PB. Parental age and risk of schizophrenia. *Arch Gen Psychiatry*. 2003; 60(7):673–678. [PubMed: 12860771]
- Carroll L, Owen M. Genetic overlap between autism, schizophrenia and bipolar disorder. *Genome Med*. 2009; 1(10):102. [PubMed: 19886976]
- Croen LA, Najjar DV, Fireman B, Grether JK. Maternal and paternal age and risk of autism spectrum disorders. *Arch Pediatr Adolesc Med*. 2007; 161(4):334–340. [PubMed: 17404129]
- Crow J. Development: there's something curious about paternal-age effects. *Science*. 2003; 301(5633):606–607. [PubMed: 12893932]
- Dalman C, Allebeck P. Paternal age and schizophrenia: Further support for an association. *Am J Psychiatry*. 2002; 159(9):1591–1592. [PubMed: 12202282]
- Dick D, Riley B, Kendler K. Nature and nurture in neuropsychiatric genetics: where do we stand? *Dialogues Clin Neurosci*. 2010; 12(1):7–23. [PubMed: 20373663]
- Durkin M, Maenner M, Newschaffer C, Lee L, Cunniff C, Daniels J, Kirby R, Leavitt L, Miller L, Zahorodny W, Schieve L. Advanced parental age and the risk of autism spectrum disorder. *Am J Epidemiol*. 2008; 168(11):1268–1276. [PubMed: 18945690]
- Ekeus C, Olausson PO, Hjern A. Psychiatric morbidity is related to parental age: a national cohort study. *Psychol Med*. 2006; 36(2):269–276. [PubMed: 16420715]
- Fokkema T, De Valk H, De Beer J, Van Duin C. The Netherlands: childbearing within the context of a “poldermodel” society. *Demographic Res*. 2008; 19(21):743–794.
- Frans EM, Sandin S, Reichenberg A, Lichtenstein P, Langström N, Hultman CM. Advancing paternal age and bipolar disorder. *Arch Gen Psychiatry*. 2008; 65(9):1034–1040. [PubMed: 18762589]
- Gillberg C. Parental age in child psychiatric clinic attenders. *Acta Psychiatr Scand*. 1982; 66(6):471–478. [PubMed: 7180566]
- Glasson EJ, Bower C, Petterson B, De Klerk N, Chaney G, Hallmayer JF. Perinatal factors and the development of autism: a population study. *Arch Gen Psychiatry*. 2004; 61(6):618–627. [PubMed: 15184241]
- Grether J, Anderson M, Croen L, Smith D, Windham G. Risk of autism and increasing maternal and paternal age in a large North American population. *Am J Epidemiol*. 2009; 170(9):1118–1126. [PubMed: 19783586]
- Hyman S. Can neuroscience be integrated into the DSM-V? *Nat Rev Neurosci*. 2007; 8(9):725–732. [PubMed: 17704814]
- King M, Fountain C, Dakhllallah D, Bearman P. Estimated autism risk and older reproductive age. *Am J Public Health*. 2009; 99(9):1673–1679. [PubMed: 19608957]
- Krishnaswamy S, Subramaniam K, Indran H, Ramachandran P, Indran T, Indran R, Aziz J. Parental age and common mental disorders. *World J Biol Psychiatry*. 2009; 10(4 Pt 2):518–523. [PubMed: 19191074]
- Lauritsen MB, Pedersen CB, Mortensen PB. Effects of familial risk factors and place of birth on the risk of autism: a nationwide register-based study. *J Child Psychol Psychiatry*. 2005; 46(9):963–971. [PubMed: 16108999]
- Laursen T, Munk-Olsen T, Nordentoft M, Bo Mortensen P. A comparison of selected risk factors for unipolar depressive disorder, bipolar affective disorder, schizoaffective disorder, and schizophrenia from a Danish population-based cohort. *J Clin Psychiatry*. 2007; 68(7):1673–1681. [PubMed: 18052560]

- Lopez-Castroman J, Gomez D, Carballo Belloso J, Fernandez-Navarro P, Perez-Rodriguez M, Villamor I, Navarrete F, Ginestar C, Currier D, Torres M, Navio-Acosta M, Saiz-Ruiz J, Jimenez-Arriero M, Baca-Garcia E. Differences in maternal and paternal age between schizophrenia and other psychiatric disorders. *Schizophr Res*. 2009; 116(2-3):184–190. [PubMed: 19945257]
- Maccabe J, Lambe M, Cnattingius S, Sham P, David A, Reichenberg A, Murray R, Hultman C. Excellent school performance at age 16 and risk of adult bipolar disorder: national cohort study. *Br J Psychiatry*. 2010; 196(2):109–115. [PubMed: 20118454]
- Malaspina D, Corcoran C, Fahim C, Berman A, Harkavy-Friedman J, Yale S, Goetz D, Goetz R, Harlap S, Gorman J. Paternal age and sporadic schizophrenia: evidence for de novo mutations. *Am J Med Genet Part B*. 2002; 114(3):299–303.
- Malaspina D, Harlap S, Fennig S, Heiman D, Nahon D, Feldman D, Susser ES. Advancing paternal age and the risk of schizophrenia. *Arch Gen Psychiatry*. 2001; 58(4):361–367. [PubMed: 11296097]
- Mcclellan J, Susser E, King M. Schizophrenia: a common disease caused by multiple rare alleles. *Br J Psychiatry*. 2007; 190:194–199. [PubMed: 17329737]
- Menezes P, Lewis G, Rasmussen F, Zammit S, Sips A, Harrison G, Tynelius P, Gunnell D. Paternal and maternal ages at conception and risk of bipolar affective disorder in their offspring. *Psychol Med*. 2010; 40(3):477–185. [PubMed: 19627644]
- Miller B, Messias E, Miettunen J, Alaraisanen A, Jarvelin J, Koponen H, Rasanen P, Isohanni M, Kirkpatrick B. Meta-analysis of paternal age and schizophrenia risk in male versus female offspring. *Schizophr Bull*. 2010; 1093/schbul/sbq011
- Olsen J, Zhu JL. Re: “Advanced parental age and the risk of autism spectrum disorder”. *Am J Epidemiol*. 2009; 169(11):1406. [PubMed: 19395694]
- Parkes J, Kerr C, McDowell B, Cosgrove A. Recruitment bias in a population-based study of children with cerebral palsy. *Pediatrics*. 2006; 118(4):1616–1622. [PubMed: 17015554]
- Petersen L, Mortensen P, Pedersen C. Paternal age at birth of first child and risk of schizophrenia. *Am J Psychiatry*. 2011; 168(1):82. [PubMed: 20952457]
- Rasmussen F. Paternal age, size at birth, and size in young adulthood - risk factors for schizophrenia. *Eur J Endocrinol*. 2006; 155:S65–S69.
- Reichenberg A, Gross R, Weiser M, Bresnahan M, Silverman J, Harlap S, Rabinowitz J, Shulman C, Malaspina D, Lubin G, Knobler HY, Davidson M, Susser E. Advancing paternal age and autism. *Arch Gen Psychiatry*. 2006; 63(9):1026–1032. [PubMed: 16953005]
- Sasanfar R, Haddad S, Tolouei A, Ghadami M, Yu D, Santangelo S. Paternal age increases the risk for autism in an Iranian population sample. *Mol Autism*. 2010; 1186/2040-2392-1-2
- Shelton J, Tancredi D, Herz-Picciotto I. Independent and dependent contributions of advanced maternal and paternal ages to autism risk. *Autism Res*. 2010; 3(1):1–10. [PubMed: 20020537]
- Sips A, Rasmussen F, Harrison G, Tynelius P, Lewis G, Leon DA, Gunnell D. Paternal age and schizophrenia: A population based cohort study. *BMJ*. 2004; 329(7474):1070–1073. [PubMed: 15501901]
- Smith R, Kember R, Mill J, Fernandes C, Schalkwyk L, Buxbaum J, Reichenberg A. Advancing paternal age is associated with deficits in social and exploratory behaviors in the offspring: a mouse model. *PLoS One*. 2009; 4(12):e8456. [PubMed: 20041141]
- Statistics-Netherlands. Wet op het Centraal bureau voor de statistiek. 2004
- Torrey E, Buka S, Cannon T, Goldstein J, Seidman L, Liu T, Hadley T, Rosso I, Bearden C, Yolken R. Paternal age as a risk factor for schizophrenia: how important is it? *Schizophr Res*. 2009; 114(1-3):1–5. [PubMed: 19683417]
- Tsuchiya KJ, Matsumoto K, Miyachi T, Tsujii M, Nakamura K, Takagai S, Kawai M, Yagi A, Iwaki K, Suda S, Sugihara G, Iwata Y, Matsuzaki H, Sekine Y, Suzuki K, Sugiyama T, Mori N, Takei N. Paternal age at birth and high-functioning autistic-spectrum disorder in offspring. *Br J Psychiatry*. 2008; 193(4):316–321. [PubMed: 18827294]
- Tsuchiya KJ, Takagai S, Kawai M, Matsumoto H, Nakamura K, Minabe Y, Mori N, Takei N. Advanced paternal age associated with an elevated risk for schizophrenia in offspring in a Japanese population. *Schizophr Res*. 2005; 76(2-3):337–342. [PubMed: 15949666]



Zammit S, Allebeck P, Dalman C, Lundberg I, Hemmingson T, Owen MJ, Lewis G. Paternal age and risk for schizophrenia. *Br J Psychiatry*. 2003; 183:405–408. [PubMed: 14594914]

**Table 1**

Demographic characteristics of patients with schizophrenia (SCZ), autism spectrum disorders (ASD), major depression disorder (MDD), or bipolar disorder (BPD) and matched controls.

		Cases	Controls
<b>SCZ</b>	N	2564	10 256
	Mean age (SD)	36.66 (9.69)	36.74 (9.66)
	Male / Female ratio	64.8% / 35.2%	64.9% / 35.1%
	Ethnicity:		
	<i>Native Dutch (%)</i>	1737 (67.7)	8316 (81.1)
	<i>Non-Dutch Western European (%)</i>	52 (2.0)	192 (1.9)
	<i>Turkey (%)</i>	82 (3.2)	247 (2.4)
	<i>Morocco (%)</i>	225 (8.8)	426 (4.2)
	<i>Surinam/Netherlands Antilles/Aruba (%)</i>	147 (5.7)	278 (2.7)
	<i>Other (%)</i>	321 (12.5)	797 (7.8)
Mean paternal age (SD)	30.84 (5.84)	30.56 (5.47)	
<b>ASD</b>	N	2262	9048
	Mean age (SD)	19.38 (11.28)	19.33 (11.23)
	Male / Female ratio	80.9% / 19.1%	80.7% / 19.3%
	Ethnicity:		
	<i>Native Dutch (%)</i>	1911 (84.5)	7080 (78.2)
	<i>Non-Dutch Western European (%)</i>	48 (2.1)	176 (1.9)
	<i>Turkey (%)</i>	25 (1.1)	227 (2.5)
	<i>Morocco (%)</i>	35 (1.5)	497 (5.5)
	<i>Surinam/Netherlands Antilles/Aruba (%)</i>	51 (2.3)	252 (2.8)
	<i>Other (%)</i>	192 (8.5)	816 (9.0)
Mean paternal age (SD)	32.96 (5.53)	32.47 (5.51)	
<b>MDD</b>	N	8284	33 136
	Mean age (SD)	37.23 (10.54)	37.32 (10.52)
	Male / Female ratio	34.2% / 65.8%	34.3% / 65.7%
	Ethnicity:		
	<i>Native Dutch (%)</i>	6429 (77.6)	27 566 (83.2)
	<i>Non-Dutch Western European (%)</i>	149 (1.8)	618 (1.9)
	<i>Turkey (%)</i>	316 (3.8)	651 (2.0)
	<i>Morocco (%)</i>	615 (7.4)	1130 (3.4)
	<i>Surinam/Netherlands Antilles/Aruba (%)</i>	197 (2.4)	735 (2.2)
	<i>Other (%)</i>	578 (7.0)	2436 (7.4)
Mean paternal age (SD)	30.54 (5.76)	30.48 (5.39)	
<b>BPD</b>	N	1121	4484
	Mean age (SD)	40.71 (10.03)	40.71 (10.02)
	Male / Female ratio	46.1% / 53.9%	46.1% / 53.9%
	Ethnicity:		
	<i>Native Dutch (%)</i>	937 (83.6)	3749 (83.6)

	Cases	Controls
<i>Non-Dutch Western European (%)</i>	26 (2.3)	80 (1.8)
<i>Turkey (%)</i>	15 (1.3)	76 (1.7)
<i>Morocco (%)</i>	24 (2.1)	147 (3.3)
<i>Surinam/Netherlands Antilles/Aruba (%)</i>	27 (2.4)	105 (2.3)
<i>Other (%)</i>	92 (8.2)	327 (7.3)
Mean paternal age (SD)	30.75 (5.41)	30.54 (5.50)

N: Number; SD: Standard Deviation

**Table 2**

Multivariate Logistic Regression for paternal age as a categorical measurement

Paternal age	<20 yr	20 – 24 yr	25 – 29 yr	30 – 34 yr	35 – 39 yr	≥40 yr
<b>SCZ</b>	<b>OR</b>	<b>95% CI</b>	<b>OR</b>	<b>95% CI</b>	<b>OR</b>	<b>95% CI</b>
<i>N</i> Cases	36	324	874	751	380	199
<i>N</i> Controls	87	1271	3840	3146	1293	619
<i>OR-crude</i>	1.82	1.22, 2.70	Ref	1.05	1.29	1.41
<i>OR-adjusted</i>	1.47	0.98, 2.21	Ref	1.04	1.24	1.27
						<b>1.05, 1.53</b>
<b>ASD</b>		*				*
<i>N</i> Cases	3	108	576	846	505	224
<i>N</i> Controls	40	623	2345	3375	1886	779
<i>OR-crude</i>	0.31	0.09, 0.99	Ref	1.02	1.09	1.17
<i>OR-adjusted</i>	0.37	0.11, 1.20	Ref	1.00	1.07	1.23
		<b>0.60, 0.95</b>		0.91, 1.15	0.95, 1.25	0.98, 1.39
				0.89, 1.13	0.93, 1.23	<b>1.01, 1.50</b>
<b>MDD</b>		*			*	*
<i>N</i> Cases	125	1148	2871	2454	1136	550
<i>N</i> Controls	299	4160	12 382	10 151	4364	1780
<i>OR-crude</i>	1.80	1.46, 2.23	Ref	1.04	1.12	1.33
<i>OR-adjusted</i>	1.65	1.33, 2.05	Ref	1.04	1.09	1.22
		<b>1.07, 1.24</b>		0.98, 1.11	<b>1.01, 1.18</b>	<b>1.09, 1.36</b>
<b>BPD</b>						
<i>N</i> Cases	17	117	406	364	149	68
<i>N</i> Controls	43	583	1645	1328	630	255
<i>OR-crude</i>	1.60	0.90, 2.84	Ref	1.11	0.96	1.08
<i>OR-adjusted</i>	1.68	0.94, 3.01	Ref	1.12	0.99	1.14
		0.65, 1.02		0.95, 1.30	0.78, 1.18	0.81, 1.44
		0.65, 1.03		0.96, 1.32	0.80, 1.22	0.84, 1.55

OR: Odds-ratio; N: Number; Yr: Years; Odds-ratios are presented with corresponding 95% confidence intervals. Adjusted OR were corrected for average income of the neighborhood, difference in age between the father and the mother and the ethnic background. Bold values indicate significant adjusted odds-ratios (\*)