



Published in final edited form as:

*Pharmacol Biochem Behav.* 2011 August ; 99(2): 190–210. doi:10.1016/j.pbb.2011.02.002.

## Alzheimer's Disease and Age-Related Memory Decline (Preclinical)

**Alvin V. Terry Jr, Patrick M. Callahan, Brandon Hall, and Scott J. Webster**

Department of Pharmacology and Toxicology and Small Animal Behavior Core, Medical College of Georgia, Augusta, Georgia, 30912

### Abstract

An unfortunate result of the rapid rise in geriatric populations worldwide is the increasing prevalence of age-related cognitive disorders such as Alzheimer's disease (AD). AD is a devastating neurodegenerative illness that is characterized by a profound impairment of cognitive function, marked physical disability, and an enormous economic burden on the afflicted individual, caregivers, and society in general. The rise in elderly populations is also resulting in an increase in individuals with related (potentially treatable) conditions such as "Mild Cognitive Impairment" (MCI) which is characterized by a less severe (but abnormal) level of cognitive impairment and a high-risk for developing dementia. Even in the absence of a diagnosable disorder of cognition (e.g., AD, MCI), the perception of increased forgetfulness and declining mental function is a clear source of apprehension in the elderly. This is a valid concern given that even a modest impairment of cognitive function is likely to be associated with significant disability in a rapidly evolving, technology-based society. Unfortunately, the currently available therapies designed to improve cognition (i.e., for AD and other forms of dementia) are limited by modest efficacy, adverse side effects, and their effects on cognitive function are not sustained over time. Accordingly, it is incumbent on the scientific community to develop safer and more effective therapies that improve and/or sustain cognitive function in the elderly allowing them to remain mentally active and productive for as long as possible. As diagnostic criteria for memory disorders evolve, the demand for pro-cognitive therapeutic agents is likely to surpass AD and dementia to include MCI and potentially even less severe forms of memory decline. The purpose of this review is to provide an overview of the contemporary therapeutic targets and preclinical pharmacologic approaches (with representative drug examples) designed to enhance memory function.

### Keywords

cognition; drug development; pro-cognitive; neurotransmitter receptors; Mild Cognitive Impairment; dementia

## 1. Introduction

According to the United Nations (UN, 2007) and the United States Centers for Disease Control and Prevention (CDC, 2003) a combination of declining fertility and increasing life

© 2010 Elsevier Inc. All rights reserved.

**Corresponding Author:** Alvin V. Terry Jr., Ph.D., Professor of Pharmacology and Toxicology, CB-3618, Medical College of Georgia, 1120 Fifteenth Street, Augusta, Georgia 30912-2450, Phone 706-721-9462, Fax 706-721-2347, aterry@mcg.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

expectancies is a phenomenon that is resulting in an unprecedented growth of elderly populations worldwide. An unfortunate result of this trend is the concomitant rise in the number of people suffering from age-related forms of dementia such as Alzheimer's disease (AD). In the United States alone the number of individuals with AD could be considered epidemic at approximately 5.3 million (Herbert et al., 2003), and unfortunately, this number is expected to increase dramatically in the near future due to the number of aging baby boomers (CDC, 2007;Kinsella et al., 2009). This issue has many dimensions at both the individual level and on societies as a whole. On an individual level, AD significantly shortens life expectancy and it is one of the major causes of physical disability, institutionalization, and decreased quality of life among the elderly (reviewed Qiu et al., 2009). On a societal level, the marked rise in the number of dementia patients is likely to create an extraordinary (potentially unsustainable) economic burden and demand on social systems and health care-related institutions. It is also important to note that the population estimates (for AD) highlighted above do not include individuals with related (potentially treatable) conditions such as "Mild Cognitive Impairment" (MCI) which is characterized by a less severe (but abnormal) level of cognitive impairment (reviewed Petersen et al., 2004) and a high-risk for developing dementia (reviewed, Luck et al., 2010). Even in the absence of a diagnosable disorder of cognition (e.g., AD, MCI), the perception of increased forgetfulness and declining memory function is a clear source of distress, embarrassment, and low self esteem in the elderly (Imhof et al., 2006;Ohman et al., 2008, see also review, Ballard, 2010). Undoubtedly these factors have (at least in part) driven sales of over the counter nutritional supplements (e.g., ginko biloba, phosphatidyl serine) that promise improved memory function (now a billion dollar industry in the United States alone), despite the lack of any clear evidence of their effectiveness (Kennedy, 2004, see also review, Lanni et al., 2008).

Collectively, the information discussed above indicates that it is incumbent on the scientific community to develop safer and more effective pro-cognitive agents that could allow even the oldest in our societies to remain socially active and productive. The purpose of this review is to provide a brief overview of the contemporary therapeutic targets and preclinical pharmacologic approaches (with representative drug examples) designed to enhance memory function. The pharmacologic agents reviewed are not all-inclusive but represent examples from the more actively pursued (i.e., current) areas of investigation. The review does not address the wide variety of potentially exciting disease modifying (mechanism-based) treatments that are in development for AD based on the amyloid cascade hypothesis (vaccines,  $\beta$  and  $\gamma$  secretase inhibitors), or therapies targeting tau and neurofibrillary tangle formation, neuroinflammation, etc.

## 2. Current Treatments

Since the FDA approval of the acetylcholinesterase inhibitor (AChEI) tacrine in 1993, the primary therapeutic approach to the cognitive loss associated with AD (and other forms of dementia) has been that of a cholinergic replacement strategy. This approach is based on studies spanning more than 35 years in both humans and lower animals indicating that acetylcholine (cholinergic) pathways in the brain modulate a number of important processes including attention, working memory, and other aspects of cognition (reviewed, Terry and Buccafusco, 2003). Furthermore, the brains of cognitively impaired elderly patients and those who suffered from AD commonly exhibit damage or abnormalities in cholinergic pathways. Both postmortem and antemortem studies in aged humans and AD patients, as well as animal experiments (i.e., in aged animals and AD-related animal models) show decreased high affinity choline uptake; decreased acetylcholine synthesis and release, and decreased numbers of nicotinic and muscarinic acetylcholine receptors (Auld, et al., 2002). Collectively such findings have lead to the so called "Cholinergic Hypothesis of AD"

(reviewed, Bartus, 2000). With the exception of the relatively recent FDA approval of the NMDA antagonist, memantine (see below) in the United States, only the clinical data derived from studies based on the cholinergic hypothesis (more specifically, with AChEIs) have provided convincing evidence of an adequate level of efficacy and reliability in AD balanced with an acceptable burden of side effects.

## 2.1 AChEIs

FDA-approved therapy with donepezil, rivastigmine, and galantamine is based on several large clinical studies in patients diagnosed with mild to moderate AD (Boada-Rovira et al., 2004; Mintzer and Kershaw, 2003; Raskind et al., 2000; Winblad et al., 2001). The compounds produced subtle but statistically significant cognitive improvements in the AD study patients. Further, there are some (albeit controversial) retrospective data to suggest that these agents may enhance activities of daily living and improve behavioral disturbances in AD (reviewed, Standridge, 2004). However, there are a number of limitations associated with AChEI's. This drug class can be associated with a number of peripheral side effects (e.g., nausea, vomiting, diarrhea, tremors) and it is only effective in a limited number of patients. In addition, few would argue that the compounds in this class have potent neuroprotective activity, the ability to preserve cognition over significant periods of time, or to prevent the inevitable outcome of the illness.

## 2.2 Memantine

Memantine, the other FDA-approved therapeutic agent for the cognitive symptoms of AD is a low to moderate affinity uncompetitive (open-channel) NMDA receptor antagonist which binds preferentially to the NMDA receptor-operated cation channels. These actions are believed support the therapeutic effects of memantine by combating the persistent activation of N-methyl-D-aspartate (NMDA) receptors by the excitatory amino acid glutamate in the brain, a process thought to contribute to the symptomatology of AD. Approval of memantine was based on three randomized, double-blind, placebo-controlled clinical trials in moderate to severe AD (reviewed, Thomas and Grossberg, 2009). The clinical data indicate that memantine is a relatively safe drug with few major side effects and a negligible risk of drug-drug interactions, but only small clinically-relevant effects on cognition, global functioning and activities of daily living have been observed, mainly in patients with moderate AD (reviewed, van Marum, 2009). Memantine is recommended both as monotherapy and in combination with AChEIs, and the latter approach is very commonly used, despite a lack of clear evidence that one approach is superior to the other (reviewed, Thomas and Grossberg, 2009).

## 3. Novel Cholinergic-Based Strategies

Please see Table 1 for a summary of the variety of cholinergic-based strategies for cognitive enhancement.

### 3.1 New AChEIs

There is a variety of compounds that have been synthesized as analogs of the currently marketed AChEIs, or as analogs of older AChEIs with well-characterized pharmacological profiles such as physostigmine (originally derived as an alkaloid of the Calabar bean). One such compound, TAK-147, is a derivative of donepezil which has been shown to produce pro-cognitive effects in rats performing a scopolamine reversal task in the Morris water maze (Xu et al., 2002; Chen et al., 2002). The compound has also shown cognitive enhancing effects in a rat based chronic cerebral ischemia cognitive deficit model similar to donepezil (Xu et al., 2002). TAK-147 has also been shown to ameliorate deficits induced by diazepam on passive avoidance and restore performance in a scopolamine deficit model in

delayed-matching-to-position task in rats (Miyamoto, et al., 1996). BZYX is another donepezil derivative with a dialkylbenzyl amine moiety from rivastigmine that was designed to function as a dual binding site AChE inhibitor (Sheng et al., 2009). This compound is capable of reversing scopolamine, NaNO<sub>2</sub>, and ethanol induced deficits in rats in a radial arm maze (Zhang et al., 2009). Further, BZYX seems to have neuroprotective properties by scavenging reactive oxygen species produced by H<sub>2</sub>O<sub>2</sub> and preventing the loss of mitochondria membrane potential (Zhang et al., 2009).

Phenserine, a derivative of physostigmine is a long-acting AChEI and has shown considerable cognitive enhancing abilities. In the Morris water maze, phenserine has displayed pro-cognitive properties in rats by significantly decreasing travel distance and lowering latencies to reach the platform (Janas et al., 2005). In a dog delayed-non-matching-to-sample task, phenserine has demonstrated the ability to reverse scopolamine induced deficits on task accuracies (Araujo et al., 2005; Studzinski et al., 2005). The compound has also been shown to enhance cognition in rats using a 14 unit T-maze by reducing the number of errors it took to complete the maze (Ikari et al., 1995). Interestingly, phenserine is also able to overcome deficits induced by the NMDA receptor antagonist 3-(±) 2-carboxypiperzin-4-yl) propyl phosphonic acid (CPP) in the same 14-unit T-maze (Patel et al., 1998). Similarly, eptastigmine, a derivative of physostigmine, and ganstigmine (CHF2819), a derivative of geneserine (a closely related compound to physostigmine), have both shown cognitive enhancing properties by reversing scopolamine induced deficits (Dawson et al., 1991; Rupniak et al., 1992; Trabace et al., 2007).

Huperzine A, another alkaloid, isolated from the Chinese herb *Huperzia serrata*, is a potent inhibitor of AChE and has been shown to have a pro-cognitive effect in a wide number of preclinical animal models including the Morris water maze task (Wang et al., 2006; Wang et al., 2010) and the radial arm maze task (Wang and Tang, 1998). The compound has also shown cognitive enhancing properties in a delayed response task in monkeys (Ye et al., 1999; Ou et al., 2001). Further, Huperzine A has been used to reverse scopolamine induced deficits (Wang and Tang, 1998; Gao et al., 2000) and produce pro-cognitive effects in aged rodents and monkeys (Ye et al., 1999; Ye et al., 2000). Two other compounds of interest are loganin, an iridoid glycoside isolated from the *Cornus officinalis* fruit and pseudoberberine, isolated from *Corydalis turtschaninovii*. Both are AChE inhibitors and have displayed cognitive enhancing properties (Hung et al., 2008; Lee et al., 2009). Loganin displays cognitive enhancing properties when tested in a scopolamine induced deficit reversal task using the Y-maze and passive avoidance task (Kwon et al., 2009). Loganin has also been shown to improve learning and spatial memory in rats tested in the Morris water maze task (Lee et al., 2009). Pseudoberberine has also been shown to reverse scopolamine induced cognitive impairments in mice in both a passive avoidance task and in a water maze task (Hung et al., 2008).

Recently there has been renewed interest in AChE inhibitors as they have been shown to modulate the processing of APP away from production of the toxic amyloid beta peptide (De Ferrari et al., 2001; Racchi et al., 2005; Peng et al., 2007; Rizzo et al., 2010). This has led to the synthesis of novel AChE inhibitors many of which hit multiple targets relevant to AD (Fernández-Bachiller et al., 2010; Mohamed and Rao, 2010; Huang et al., 2010; Kikuchi et al., 2010; de Los Ríos et al., 2010; Samadi et al., 2010; Pisani et al., 2010). Many of these compounds are still in their infancies, requiring much more thorough examination before they prove their worth in the treatment of AD. Still, expect to see more interest in this area of research as these AChE inhibitors, as well as their derivatives, continue to be tested in the months and years ahead.

## 3.2 Muscarinic Acetylcholine Receptor (mAChRs) Ligands

mAChRs are metabotropic receptors that are coupled to G-proteins. Agonist binding to mAChRs results in decoupling of the G-protein prompting a signal transduction cascade that leads to either excitatory or inhibitory actions on the cell. There have been five types of mAChRs identified, labeled M1–M5 (Levey et al., 1991). The M1, M3, and M5 receptor subtypes are coupled to Gq proteins that activate the phospholipase C (PLC) pathway, which results in protein kinase C (PKC) activation leading to intracellular calcium release and excitatory actions on the cell. The M2 and M4 subtypes are coupled to Gi proteins that when activated inhibit adenylyl cyclase activity, resulting in a reduction of cAMP and inhibitory actions on the cell. The M1 receptor subtype has been the dominant target for cognition enhancement therapy, due to its wide expression in the cortex and hippocampus (Levey et al., 1991), and because the receptor subtype seems to be retained longer than other mAChRs in AD (Flynn et al., 1995).

**3.2.1 Muscarinic receptor agonists**—There have been numerous M1 receptor agonists evaluated in preclinical studies as possible treatments for cognitive impairment associated with AD and age-related memory decline (Clader and Wang, 2005). The first generation of M1 agonists were analogs of the mAChR agonist arecoline, a class of compounds that includes talsaclidine, sabomeline, xanomeline, and cevimeline, among others. Many of these compounds showed positive cognitive effects in preclinical tests. However, a lack of sufficient selectivity for the M1 receptor over the M3 and M5 receptors hindered further development efforts for many of these compounds for AD therapy, due to potential cholinergic side effects. Indeed, this problem has hindered much development of subtype selective mAChR agonists/antagonists, owing to the high amount of sequence homology shared by the orthosteric binding sites of mAChRs subtypes (Hulme et al., 1990). However, the M1 receptor continues to be an attractive target for cognition therapy.

Significant preclinical data supporting M1 receptors as valid targets for cognition enhancement were obtained through studies utilizing the “AF” compounds, a series of M1 partial agonists composed of derivatives of cevimeline. Early studies revealed that treatment with cevimeline, also known as AF102B, produced positive outcomes in a broad range of animal models of AD and cognitive impairment (Clader and Wang, 2005). The analogs developed from AF102B are more efficacious at M1 receptors with greater selectivity. Treatment with AF150(S) abolished cognitive impairments associated with ApoE knockout mice, a strain of mice that shows cognitive deficits resembling AD pathology (Fisher et al., 1998). Other studies with AF150(S) demonstrated memory and cognition improvements in young and old rodents in the delayed matching-to-position task, Morris water maze task, as well as the radial arm maze (Brandeis et al., 1995; Ruske and White, 1999). When administered to 3xTG-AD mice, a mouse model of AD characterized by A $\beta$  plaque buildup, neurofibrillary tangles, and deficits in cognition, the compound AF267B recovered deficits in spatial working memory seen in the Morris water maze test after 2 months of treatment (Caccamo et al., 2006). Treatment with AF267B was also effective in reducing cognitive deficits in animals with lesions in the nucleus basalis (Beach et al., 2003). Other analogs of AF102B include AF125 and AF151 (Fisher et al., 1993). Adding to the appeal of these more selective M1 compounds was an apparent lack of peripheral side effects in preclinical models as well as a longer half life and high bioavailability (Fisher et al., 2002).

Efforts continue to develop newer, more selective M1 agonists for cognition therapy in AD that also have a low potential for undesirable peripheral cholinergic effects. A newly synthesized class of arecoline analogs, N-alkyl/aryl substituted thiazolidinone arecoline derivatives, showed significant affinity for M1 receptors, and improved scopolamine-induced impairments in the passive avoidance step down task and elevated plus maze

transfer latency (Sadashiva et al., 2009). Other arecoline derivatives with M1 receptor selectivity are also currently being synthesized and tested in preclinical models of AD (Kumar et al., 2008; Malviya et al., 2008; Malviya et al., 2009). The novel derivative of xanomeline, EUK1001, improved novel object recognition and fear cognition in a passive avoidance task in aged mice (Cui et al., 2008a; Si et al., 2010). EUK1001 was also shown to have higher affinity for M1 receptors than xanomeline coupled with a significantly lower side effect profile (Cui et al., 2008b). It will remain to be seen whether these newer compounds succeed in any clinical setting. However, a great promise still holds for compounds in this pharmacological class, assuming a given new molecular entity carries sufficient selectivity for M1 receptors over other mAChR subtypes.

**3.2.2 Muscarinic receptor antagonists**—Both in vitro and in vivo (e.g., M2R<sup>-/-</sup> knockout mice) studies suggests that M2 muscarinic receptors in the mammalian hippocampus and cortex serve as presynaptic autoreceptors mediating the inhibition ACh release. This evidence supports the hypothesis that antagonists at central M2 autoreceptors might enhance cognition by increasing synaptic acetylcholine levels (see Wess et al., 2007 for review). Consistent with this notion are studies indicating that specific M2 receptor-preferring antagonists such as SCH 57790, SCH 72788, and BIBN-99 can improve learning and memory-related tasks in animal models (Carey et al., 2001; Lachowicz et al., 2001; Rowe et al., 2003). However, M2R<sup>-/-</sup> mice were found to exhibit deficits in some cognitive tasks suggesting that complete pharmacological blockade of central M2 receptors might (in fact) have negative effects on cognition (Tzavara et al., 2003). Several hypotheses have been proposed to explain these discrepant findings. One possibility is that the M2 receptor-preferring antagonists evaluated to date may only serve as partial antagonists of central M2 receptors. Another is that there are both presynaptic and postsynaptic M2 receptor-signalling pathways in the CNS that may have different sensitivities to the antagonists used (see Wess et al., 2007 for review). Thus, additional studies are needed to determine which strategy (M2 antagonism versus enhancement) will be likely to provide a therapeutic benefit in disorders of cognition.

**3.2.3 Muscarinic positive allosteric modulators**—The lack of sufficient selectivity of orthosteric ligands for targeted muscarinic receptor subtypes has driven research for compounds that act as positive allosteric modulators (PAMs) on the receptor, targeting binding sites away from the orthosteric site (see nicotinic positive allosteric modulator section below for a complete discussion). There have been several mAChR PAMs that have been identified in in vitro studies (Jones et al., 2008; Conn et al., 2009). Many of these compounds have been reported as possessing very high selectivity for the targeted receptor. However, to date only a small number of these compounds have been studied in animal models for cognition enhancement, though promising results have recently been published (Bradley et al., 2010). With the continuing need for new therapies addressing the cognitive decline associated with AD and advanced age, these compounds in this class will surely garner much attention for future preclinical cognition studies.

### 3.3 Nicotinic Acetylcholine Receptor (nAChR) Ligands

nAChRs are ionotropic receptors that open in response to agonist binding. The result of agonist binding to the receptor is the transfer of cations into, and out of, the cell, notably the influx of sodium and to a lesser degree calcium, and the efflux of potassium. The end result of nAChR activation is membrane depolarization. Nicotinic receptors are composed of five membrane-spanning subunits that form a pentameric arrangement around a water-filled pore (Jones et al., 1999; Karlin 2002; McGehee and Role, 1995). Neuronal nicotinic receptors are formed through a combination of  $\alpha$  and  $\beta$  subunits composing heteromeric receptors, or through homomeric  $\alpha$  configurations. Although there have been nine  $\alpha$  ( $\alpha 2$ – $\alpha 10$ ) and three  $\beta$

( $\beta 2$ – $\beta 4$ ) subunits currently identified that can form many different combinations, for the purposes of this review, the heteromeric  $\alpha 4\beta 2$  and homomeric  $\alpha 7$  nAChR subtypes will be primarily discussed. These two subtypes have been the targets for the majority of cognition enhancing drugs for AD and age-related memory decline, as both subtypes are highly expressed in the cerebral cortex and hippocampus. Targeting other nAChR subtypes has been associated with efficacy and side effect issues, owing to their distribution in other brain areas such as the nucleus accumbens (Buccafusco, 2004).

**3.3.1 Nicotinic receptor agonists**—The ability of nicotine to improve cognition in animal models as well as in humans has been known for over two decades, and has been extensively studied and reviewed (Elrod et al., 1988; Levin, 2002; Murray and Abeles, 2002). However, the use of nicotine clinically for any purpose other than smoking cessation was severely limited by concerns over serious side effects (Benowitz, 1986). The goal for cognition enhancement through nAChRs is to develop receptor subtype selective agonists that do not possess the undesirable side effects accompanied by the use of nicotine, particularly the abuse and addiction potential. Compounds selective for the  $\alpha 4\beta 2$  subtype were the first to be developed for cognition enhancement and studied in preclinical studies.

Two of the first  $\alpha 4\beta 2$  agonists tested for cognition enhancement were SIBIA Neuroscience's SIB-1508Y and Abbott's ABT-418. These compounds showed great promise in pre-clinical animal models of cognitive impairment (Schneider et al., 1999; Prendergast et al., 1998). However, a lack of selectivity for the  $\alpha 4\beta 2$  receptor hindered further development for these compounds. ABT-089, a follow-up compound to ABT-418, was shown to improve Morris water maze performance in aged rats, and when tested in young rats with septal lesions improved acquisition in a spatial discrimination water maze task (Decker et al., 1997). In the same study, ABT-089 improved cognitive performance in young and aged monkeys in the delayed matching-to-sample task. Although ABT-089 has less affinity for  $\alpha 4\beta 2$  nAChRs than either nicotine or ABT-418, the compound was shown to be as potent as nicotine in evoking hippocampal ACh release from rat synaptosomes (Sullivan et al., 1997). However, despite ABT-089's comparable effects to nicotine for ACh release, the compound does not share nicotine's potency at inducing dopamine release in the ventral tegmental area (VTA), perhaps owing to the compound's pharmacological profile as a partial agonist. ABT-594, an  $\alpha 4\beta 2$  agonist currently being studied for analgesic potential, was shown to improve delayed matching-to-sample task performance when studied in non-human primates (Buccafusco et al., 2007). The  $\alpha 4\beta 2$  partial agonist varenicline (Chantix<sup>®</sup>), already FDA approved for smoking cessation has also been considered a candidate for potential cognition enhancement. In pre-clinical tests, varenicline improved sustained attention task performance in rats, a model which tests top-down control of attention (Rollema et al., 2009). Deficits in top-down attention control are a prominent feature of age-related memory decline. Rats administered varenicline also demonstrated improvements in the novel object recognition task compared to vehicle treated animals in the same study. Ispronicline (AZD-3480), another  $\alpha 4\beta 2$  partial agonist, improved working memory in rodents, even after a single dose (Gatto et al., 2004). This improvement was observed for up to 18h despite a half life of only 2h, highlighting prolonged effects present even after elimination of the compound. Finally, a recently published study highlights the positive effects of the agonist S-38232 in a distractor version of the sustained attention task in rodents, though no effect was observed in a standard sustained attention task paradigm (Howe et al., 2010). These recent studies demonstrate the continuing appeal of targeting  $\alpha 4\beta 2$  nAChRs for cognition enhancement in preclinical research.

Compounds targeting the homomeric  $\alpha 7$  nicotinic receptor subtype have yielded some promising results for cognition enhancement and are a major focus of current preclinical research (Leiser et al., 2009). The distribution of  $\alpha 7$  nAChRs is limited primarily to the

hippocampus, certain cortical layers, and the hypothalamus, making these receptors intriguing targets for cognition enhancement (Freedman et al., 1995). Activation of  $\alpha 7$  nAChRs does not appear to be involved in the rewarding effects of nicotine, suggesting that targeting these receptors has limited abuse potential (Brioni et al., 1996). Also, while not the focus of this review, the extrasynaptic localization of  $\alpha 7$  nAChRs suggests a role for the receptor subtype in cell viability. The  $\alpha 7$  partial agonist and anabaseine synthetic derivative GTS-21 (3-[(2, 4 dimethoxy) benzylidene]-anabaseine, or DMXBA) is a well studied compound and was poised to perhaps become the first synthetic nicotinic compound to market for cognition enhancement. The compound has been shown in preclinical models to enhance cognitive processes, including studies in rodents and non-human primates (Buccafusco and Terry, 2009; Chen et al., 2010; Rodefer et al., 2007). This cognition enhancement by GTS-21 is accompanied by a low potential for side effects (Kem, 2000). What has made GTS-21 particularly interesting is the compound's relative specificity for cognition enhancement versus its inactivity in behavioral tasks in which nicotine is effective (Briggs et al, 1997). The most significant effects of GTS-21 reported to date have been observed in tasks of attention, working memory, and episodic secondary memory. The results of preclinical tests using compounds like GTS-21 have driven the development of newer, more selective  $\alpha 7$  receptor agonists for use in cognition enhancement. The Memory Pharmaceuticals (acquired by Roche) compound MEM 3454 (known as RG-3487) is an  $\alpha 7$  nAChR partial agonist with 5-HT<sub>3</sub> receptor antagonist properties that has been shown to enhance task performance in young and aged-cognitively impaired rodents as well as young and aged non-human primates (Callahan et al., 2006; Wallace-Boone et al., 2009; Wallace et al., 2011) across multiple cognition domains (episodic, attention, executive function, spatial reference and working memory). RG-3487 was also shown to improve sustained visual attention in rodents, a property shared by nicotine but not galanthamine (Rezvani et al., 2009). Moreover, the improvements in cognitive performance were shown to be reflective of the observed increases in acetylcholine and dopamine efflux in cortical and hippocampal brain regions (Huang et al., 2006). These positive preclinical observations translated into significant improvements in cognition, as measured by the cognitive drug research (CDR) test battery in healthy human volunteers and mild to moderate AD subjects during phase I and phase IIA clinical testing (Callahan et al., 2006). Another  $\alpha 7$  nAChR partial agonist from Memory Pharmaceuticals, MEM 64368, demonstrated robust cognitive enhancing properties in rodents and non-human primates (Rodefer et al., 2007; Wallace-Boone et al., 2007). This  $\alpha 7$  nAChR partial agonist was characterized as a low intrinsic activity agonist (~30%) with moderate affinity ( $K_i$  ~300 nM) at  $\alpha 7$  nAChR receptors compared to that of RG-3487 ( $K_i$ = 6nM; intrinsic activity ~68%) and displayed a 10-fold selectivity for  $\alpha 7$  nAChR over 5-HT<sub>3</sub> receptors (Wallace-Boone et al., 2007; Wallace et al., 2011). In addition, AstraZeneca's  $\alpha 7$  nAChR compound AZD-0328 improved spatial working memory in mice in a novel object recognition task, an improvement that was partly attributed to enhanced midbrain and cortical dopamine release (Sydserff et al., 2009). The novel  $\alpha 7$  agonist SEN12333 (WAY-317538) has also been shown to improve novel object recognition performance in rats (Roncarati et al., 2009). This improvement was observed even under scopolamine or MK-801-induced amnesia, representing relevant cholinergic and glutamatergic deficits commonly seen in AD. In the same study, SEN12333 recovered deficits in a passive avoidance task induced by scopolamine. Results similar to these have been observed in preclinical cognitive tests using a host of  $\alpha 7$  nAChR agonists (Table 1).

**3.3.2 Nicotinic Positive Allosteric Modulators**—Due to the historical difficulty in developing compounds with sufficient selectivity for the orthosteric ligand binding site of nAChR subtypes, there has been considerable interest directed toward allosteric modulators for targeting nAChRs. Positive allosteric modulators avoid direct receptor activation by utilizing binding sites away from the orthosteric site targeted by traditional agonists. Though



allosterically-acting compounds have been more difficult to discover than compounds acting at orthosteric sites, the existence of these compounds has been known for several years and their use has recently been suggested as a favorable strategy for targeting nicotinic receptors (Albuquerque et al., 2009). The emergence of more advanced techniques for investigating molecular site interactions, as well as *in silico* screening of large compound libraries for receptor targets, has generated interest in this approach in certain drug discovery arenas. However, only a few compounds in this class have been studied *in vivo*. The drug galantamine, an acetylcholinesterase inhibitor currently in use for AD treatment, has also been demonstrated to act as a positive allosteric modulator of nAChRs (Coyle et al., 2007).

The allosteric potentiating ligands (APLs) of nAChRs that have been tested in preclinical memory and cognition models have demonstrated great potential for this class of compounds. The compound NS-1738, classified as an  $\alpha 7$  selective positive allosteric modulator (PAM), recovered scopolamine-induced deficits in the Morris water maze task in rats as well as improving the animals performance in a social recognition model (Timmermann et al., 2007). The PAM known as compound 6 also targets  $\alpha 7$  nAChRs, and when administered to rats significantly improved their performance in the radial arm maze task compared to vehicle treated control animals (Ng et al., 2007). Compound 6 also showed positive results in sensory gating deficits, a property shared by another nAChR PAM, PNU-120596 (Hurst et al., 2005).

**3.3.3 Desensitization of nicotinic-acetylcholine Receptors (nAChRs)**—As is apparent in the preceding paragraphs, drug discovery efforts for pro-cognitive agents in the nicotinic field to date have focused on agonists, partial agonists and allosteric modulators (i.e., all efforts designed to increase activity of nAChRs). These approaches have (for the most part) been based on the idea that enhanced activity at nAChRs promotes the release of neurotransmitters in relevant brain regions which results in the desired behavioral effect (improved cognition). However, in a number of settings, nAChR agonists and antagonists can have very similar physiologic effects (see review, Buccafusco et al., 2009), and further, a large discrepancy often exists between the high concentrations of nicotinic drugs required to activate nicotinic responses *in vitro* and the lower doses used to induce various behavioral responses *in vivo*. Nicotine can both activate and desensitize its receptors over a relatively short time course leading to the question of whether (in fact) nAChR desensitization when compared to receptor activation, plays an equal if not more important role in the behavioral effects. Recent studies in our laboratories (Sood et al., 2007; Buccafusco et al., 2009) indicated that nicotine, the major nicotine metabolite, cotinine, and two novel analogs of choline, JWB1-84-1 [2-(4-(pyridin-3-ylmethyl)piperazin-1-yl)ethanol] and JAY2-22-33 [2-(methyl(pyridine-3-ylmethyl)amino)-ethanol], improved working memory in a delayed match to sample task (DMTS) in monkeys. JWB1-84-1 also reduced the number of errors involved in completing a radial arm water maze (RAWM) task in AD-Tg mice that over-express Abeta beginning around 7 months of age (Sood et al., 2007). The effectiveness of the four aforementioned compounds (nicotine, cotinine, JWB1-84-1 and JAY2-22-33) in the DMTS task was linearly related to their effectiveness in producing desensitization of a nAChR agonist response in rats. Only nicotine evoked an agonist-like action in these studies indicating that it is possible to develop new chemical entities (e.g., choline analogs, cotinine analogs) that have the ability to desensitize nAChRs without an antecedent agonist action. Since the side effects of nicotine (e.g., cardiovascular, gastrointestinal) are often associated with its agonist effects, such an approach could offer the advantage of better tolerability.

## 4. Phosphodiesterase Inhibitors

Please see Table 2 for a summary of the variety of phosphodiesterase-based strategies for cognitive enhancement.

Initially, phosphodiesterase (PDE) inhibitors (e.g., papaverine) were examined for their ability to increase cerebral blood flow in an attempt to improve cerebral metabolism and thereby, improve cognitive function in aged patients (for review see Yesavage et al., 1979). More recently, research has concentrated on the ability of PDE inhibitors to modulate the second messenger molecules cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP). Eleven subclasses of PDE inhibitors (PDE1–PDE11) have been identified with most having one or more isoforms (e.g., PDE4A, PDE4B, PDE4C and PDE4D) and each isoform may have multiple splice variants (e.g., PDE4D1–PDE4D9). Overall, more than 100 human PDEs exist (Bender and Beavo, 2006). All PDEs have a common catalytic domain, but PDE4, PDE7 and PDE8 hydrolyze cAMP whereas PDE5, PDE6 and PDE9 hydrolyze cGMP. PDE1, PDE2, PDE3, PDE10 and PDE11 are capable of hydrolyzing both cAMP and cGMP. PDE inhibitors (PDE-I) increase intracellular levels of cAMP and/or cGMP which can lead to gene transcription by activating the CREB signaling pathways (Impey et al., 1996; Lu and Hawkins, 2002; Tully et al., 2003). Activation of the cAMP/PKA/CREB and cGMP/PKG/CREB pathway has been implicated in synaptic plasticity such as long-term potentiation (LTP) and cognitive function (Frey et al., 1993; Son et al., 1998; Blokland et al., 2006). Despite 11 subclasses of PDEs, research has focused on PDE2, PDE4, PDE5, PDE9 and PDE10 and their potential involvement in learning and memory processes (see Rose et al., 2005 and Reneerkens et al., 2009 for a comprehensive review). To date, BAY 60–7550 is the only known selective PDE2-I that has been tested in animal models of cognition. BAY 60–7550 increased CA1 hippocampal LTP without affecting basal synaptic transmission (Boess et al., 2004). This PDE2-I improved object recognition acquisition and consolidation memory in rats and mice as well as social recognition memory performance in rats (Boess et al., 2004; Domek-Lopacinska and Strosznajder, 2008; Rutten et al., 2007). Cognitive enhancing effects of BAY 60–7550 was also observed in aged-impaired 24 month old rats in the object recognition task (Domek-Lopacinska and Strosznajder, 2008) and was shown to reverse the object recognition memory deficits of acute tryptophan depletion in rats (van Donkelaar et al., 2008). Moreover, BAY 60–7550 reversed the NMDA antagonist MK 801-induced working memory deficit in the spontaneous alternation T-maze task in mice (Boess et al., 2004). Of the PDE subtypes, PDE4 has been extensively characterized. PDE4-Is selectively inhibit the hydrolysis of cAMP and strong evidence exists for the cAMP/PKA/CREB pathway in mediating the cognitive effects of PDE4-I (Rose et al., 2005; Tully et al., 2003). Rolipram has been the most extensively studied PDE4-I. Rolipram has been shown to facilitate hippocampal LTP function in young and aged rodents (Barad et al., 1998; Navakkode et al., 2004), reverse  $\beta$ -amyloid (A $\beta$ 1-42) peptide impairment (Vitolo et al., 2002) and counteract the inhibitory effects of high levels of A $\beta$  on synaptic plasticity in the transgenic APP/PS1 mouse model of AD (Gong et al., 2004). Behaviorally, rolipram has shown improvements in object recognition, spatial reference, working and associative memory behavioral tasks in young and aged rodents (Bach et al., 1999; Barad et al., 1998; Monti et al., 2006; Ramos et al., 2003; Rutten et al., 2006. Rutten et al., 2007) as well as improvements in working and executive memory function in young (Ramos et al., 2003; Rutten et al., 2008) but not aged adult monkeys (Ramos et al., 2003). Rolipram also restores working and reference memory in the radial arm maze task and associative memory in the passive avoidance task following administration of scopolamine (Egawa et al., 1997; Ghelardini et al., 2002; Imanishi et al., 1997; Zhang and O'Donnell, 2000), MK-801 (Zhang et al., 2000, 2005), MAPK/ERK kinase (MEK) inhibition (Zhang et al., 2004) or anisomycin (Randt et al., 1982). Further, rolipram is capable of reversing cognitive impairments in genetic mouse models of human disorders such as Rubinstein-Taybi syndrome (Bourtchouladze et al., 2003) and AD (Gong et al., 2004; Comery et al., 2005). In addition to rolipram, several newly developed PDE4-Is have been evaluated for their potential involvement in learning and memory processes. HT-0712 has been shown to increase cAMP thereby enhancing the CREB signaling pathway leading to improved memory performance in young mice in the object recognition and fear

conditioning tasks as well as aged mice in a trace conditioning behavioral task (Bourtchouladze et al., 2003; Helicon Therapeutics.com). Moreover, HT-0712, like rolipram, was shown to reverse the object recognition deficit in a genetic mouse model of Rubinstein-Taybi syndrome (Bourtchouladze et al., 2003). L-454,560 improved cognitive performance in the delayed matching to position (DMTP) version of the Morris water maze task (Huang et al., 2007). MEM 1018 and MEM 1091 were efficacious in reversing the MK-801-induced working and reference memory deficits in the rat radial arm maze and each compound antagonized the amnesic effect of MK-801 on passive avoidance behavior (Zhang et al., 2005). Although PDE5-Is were originally developed for the treatment of erectile dysfunction (e.g., sildenafil (Viagra), vardenafil (Levitra) and tadalafil (Cialis); Setter et al., 2005), recent preclinical evidence has suggested a potential role in cognition. PDE5-Is selectively inhibit the enzymatic activity of PDE5 leading to increase intracellular levels of cGMP (Bender and Beavo, 2006). PDE5-Is have been shown to enhance memory function in the object recognition task in rats (Domek-Lopacinska and Strosznajder 2008; Prickaerts et al., 2002; 2005; Rutten et al., 2007) and mice (Rutten et al., 2005). The PDE5-I sildenafil also improved executive memory performance in monkeys using the prefrontal cognition object retrieval paradigm (Rutten et al., 2008). In pharmacological deficit models, PDE5-Is, sildenafil and vardenafil, have reversed recognition, spatial and associative memory performance impairments produced by acute tryptophan depletion (van Donkelaar et al., 2008), nitric oxide synthase (NOS) inhibition (Devan et al., 2006; Prickaerts et al., 1997), scopolamine (Devan et al., 2004; Shafiei et al., 2006) and diabetic neuropathy (Patil et al., 2004). Currently, only two PDE9-Is, SCH 81566 and BAY 73-6691, have been reported and only BAY 73-6691 has been characterized in animal models of cognition (van der Staay et al., 2008). PDE9-Is, like PDE5-Is, inhibit the degradation of the second messenger cGMP. BAY 73-6691 treatment increased hippocampal LTP function in young and aged rats and produced pro-cognitive effects in the object recognition and social recognition tasks in rats and mice. Further, BAY 73-6691 reversed the scopolamine-induced passive avoidance deficit as well as the MK-801-induced deficit in the working memory T-maze task (van der Staay et al., 2008). Lastly, PDE10A-Is have been suggested as a target for cognition particularly in the treatment of cognitive dysfunction associated with schizophrenia (Rodefer et al., 2005). PDE10A-Is regulate the activity of both cAMP and cGMP and while PDE10A-Is like papaverine, TP-10 (Schmidt et al., 2008) and MP-10 (Grauer et al., 2009) possess greater pharmacological similarity to that of the traditional antipsychotics (Grauer et al., 2009; Schmidt et al., 2008), there is evidence for a role in modulating cognitive function. Papaverine has been shown to reverse the executive memory deficits of subchronic phencyclidine following washout in the attentional set-shifting paradigm which is a rodent model analogous to the Wisconsin Card Sorting task in humans (Rodefer et al., 2005). Further, papaverine, but not MP-10 enhanced object recognition memory following a 48 hr delay interval in rats (Grauer et al., 2009) and both compounds attenuated the MK-801-induced social odor recognition deficits in mice (Grauer et al., 2009). Despite these encouraging results following acute administration of PDE10A-I, chronic administration of papaverine to mice disrupted spatial reference memory and reversal learning using the Morris water maze task (Hebb et al., 2008).

Overall, the preclinical evidence clearly demonstrates that PDE-Is for the PDE2, PDE4, PDE5, PDE9 and PDE10 subclasses are involved in learning and memory processes. It is now up to the pharmaceutical industry to develop safe and viable compounds to assess these preclinical effects on cognition in clinical trials in humans.

## 5. Serotonin Receptors

Please see Table 3 for a summary of the variety of serotonergic-based strategies for cognitive enhancement.

Serotonin (5-hydroxytryptamine; 5-HT) receptor systems are expressed throughout the peripheral and central nervous systems and are involved in a multitude of physiological functions such as cardiovascular and gastrointestinal regulation, thermoregulation, affective disorders as well as learning and memory (Barnes and Sharp, 1999; Cassel and Jeltsch, 1995). Within the 7 classes of 5-HT receptors (5-HT<sub>1-7</sub>) there are at least 13 distinct G protein-coupled receptors (GPCRs) and one ligand-gated ion channel (5-HT<sub>3</sub>; Barnes and Sharp, 1999; Hannon and Hoyer, 2008). Several of the 5-HT receptor subtypes (i.e., 5-HT<sub>1A</sub>, 5-HT<sub>2A</sub>, 5-HT<sub>3</sub>, 5-HT<sub>4</sub> and 5-HT<sub>6</sub>) have been implicated in learning and memory processes (for review see King et al., 2008; Terry et al., 2008) and are expressed in key brain regions (e.g., amygdala, cortex, hippocampus and striatum) where they modulate the neurotransmitter release of acetylcholine, dopamine, GABA and glutamate and noradrenaline (Barnes and Sharp, 1999).

### 5.1 5-HT<sub>1A</sub> receptors

The 5-HT<sub>1A</sub> receptor functions as both a somatodendritic inhibitory autoreceptor involved in raphe nuclei regulation as well as a postsynaptic heteroreceptor modulating neuronal release (Barnes and Sharp, 1999). For the most part, 5-HT<sub>1A</sub> receptor agonists (e.g., 8-OH-DPAT and buspirone) have been shown to disrupt learning and memory processes (Meneses and Perez-Garcia, 2007; Orgen et al., 2008), although there are reports of 5-HT<sub>1A</sub> receptor agonists (MDL 73005 and S15535) showing improvements in cognitive function (Bertrand et al., 2001; Millan et al., 2004). The real interest within 5-HT<sub>1A</sub> receptor pharmacology stems from the observations that selective “silent” 5-HT<sub>1A</sub> receptor antagonists (e.g., lecozotan (SRA-333), NAD-299, WAY 100635 and WAY 101405) reverse anatomical lesions (e.g., fornix and hippocampal) and pharmacological impairments (e.g., scopolamine and MK 801) as well as possess stand-alone improvements across a broad spectrum of behavioral models of cognition in rodents and non-human primates (Harder et al., 1996; Harder and Ridley, 2000; Hirst et al., 2008; Luttgen et al., 2005; Misane and Ogren, 2003; Pitsikas et al., 2003; Schechter et al., 2005). Although the exact mechanism underlying the observed pro-cognitive effects of 5-HT<sub>1A</sub> receptor antagonists is not completely known, it is thought that modulation of cholinergic and/or glutamatergic transmission is involved. Interestingly, the 5-HT<sub>1A</sub> receptor antagonist lecozotan (SRA-333) was shown to be safe and well tolerated in healthy young and elderly subjects (Patat et al., 2009) and a clinical phase II trial in mild to moderate AD patients was recently completed, though the outcome hasn't been reported (Sabbagh, 2009).

### 5.2 5-HT<sub>2A</sub> receptors

5-HT<sub>2A</sub> receptors are expressed in brain regions (e.g., cortex, striatum and hippocampus) known to be involved in cognition and perception and 5HT<sub>2A</sub> receptor antagonists appear to be involved in cognitive processes (Terry et al., 2005b; Williams et al., 2002; Winstanley et al., 2003). For example, medial prefrontal cortex infusions with the selective 5HT<sub>2A</sub> receptor antagonist MDL 100,907 improved visuospatial attention and decreased impulsivity in the rat 5-choice serial reaction time task (Winstanley et al., 2003) and oral administration of the 5HT<sub>2A</sub> receptor antagonist EMD 281014 improved medium and long retention interval performance in the delayed matching-to-sample task in young and aged rhesus monkeys (Terry et al., 2005b). However, the 5HT<sub>2A</sub> receptor antagonist ketanserin failed to show improvements in the radial arm maze or operant signal detection task in rats and co-administration of ketanserin with either acute or chronic administration of nicotine attenuated the enhanced attentional performance produced by nicotine (Levin and Rezvani, 2007). Interestingly, the 5HT<sub>2A</sub> receptor antagonists mianserin and MDL 100,907 have shown modest, but promising effects on cognition in schizophrenic patients (Roth et al., 2004). Additional research and interest from the pharmaceutical community is required to truly assess the potential of this drug target.

### 5.3 5-HT<sub>3</sub> receptors

The 5-HT<sub>3</sub> receptor and in particular, 5-HT<sub>3</sub> receptor antagonists have also generated interest as putative cognitive enhancers given their ability to increase acetylcholine release (Diez-Ariza et al., 2002; Ramirez et al., 1996) and improve certain aspects of cognitive function (Meneses, 1999; Terry et al., 2008). Overall, 5-HT<sub>3</sub> receptor antagonists (e.g., DAU 6215, granisetron, ondansetron, RS-56812, SEC 579 and WAY 100579) appear to be efficacious in reversing pharmacologically-induced impairments (e.g., atropine, scopolamine and ibotenate acid forebrain lesions) as well as produce stand-alone improvements in cognitive function across multiple cognition behavioral models in young and aged rats (Barnes et al., 1990; Carli et al., 1997; Fontana et al., 1995; Hodges et al., 1995; Pitsikas and Borsini 1996; 1997; Pitsikas et al., 1993) and non-human primates (Arnsten et al., 1997; Barnes et al., 1990; Carey et al., 1992; Domeney et al., 1991; Terry et al., 1996). Despite these encouraging results, several studies have failed to support the pro-cognitive effects of 5-HT<sub>3</sub> receptor antagonists (Bratt et al., 1994; Naghdi and Harooni, 2005; Pitsikas et al., 1993; Pitsikas and Borsini, 1997). Whereas ondansetron was found to be inactive in an AD clinical trial (Dysken et al., 2002), a renewed interest has been initiated for the cognitive dysfunction associated with schizophrenia (Akhondzadeh et al., 2009; Zhang et al., 2006).

### 5.4 5-HT<sub>4</sub> receptors

The concentration of 5-HT<sub>4</sub> receptors in limbic structures (e.g., cortex, amygdala, septum and hippocampus), along with modulation of cholinergic neuronal activity (Bockaert et al., 2004; Bonaventure et al., 2000; Vilaro et al., 2005) and neurochemical release of acetylcholine (Consolo et al., 2004; Matsumoto et al., 2001; Mohler et al., 2007) within these brain regions make it an attractive target for restoration of cognitive dysfunction. 5-HT<sub>4</sub> receptors are positively linked to adenylate cyclase and activation leads to increased intracellular levels of cAMP, a fundamental component in synaptic long-term potentiation (LTP; Frey et al., 1993). Indeed, 5-HT<sub>4</sub> receptor partial agonists (e.g., RS 67333, SC 53116 and SL 65.0155) enhance the activity of cAMP (Eglen et al., 1995) facilitate hippocampal (Kulla and Manahan-Vaughan, 2002; Marchetti et al., 2004; Matsumoto et al., 2001) and amygdala LTP function (Huang and Kandel, 2007). Moreover, 5-HT<sub>4</sub> receptor agonists were capable of reversing hippocampal deficient LTP synaptic transmission in transgenic mice overexpressing the amyloid  $\beta$  peptide (Spencer et al., 2004). Behaviorally, 5-HT<sub>4</sub> receptor agonists produce pro-cognitive effects in animal models of learning and memory (King et al., 2008; Meneses, 1999). The 5-HT<sub>4</sub> receptor partial agonists (e.g., BIMU 1, SC 53116, RS 67333, SL 65.0155 and VRX 03011) have been shown to reverse the amnesic effects of scopolamine and atropine on passive avoidance and spatial navigation performance in rats and mice (Fontana et al., 1997; Galeotti et al., 1998; Matsumoto et al., 2001; Moser et al., 2002), enhance place and object recognition memory in young and aged rats (Lamirault and Simon, 2001; Levallet et al., 2009; Moser et al., 2002), enhance delayed spontaneous alternation behavior (Mohler et al., 2007) and reverse the aged related deficit observed in the linear maze in rats (Moser et al., 2002). Additionally, SL 65.0155 improved attentional performance in the rat 5-choice serial reaction time task (Hille et al., 2008) and RS 17017 enhanced attentional and working memory performance in young and aged monkeys in the delayed matching-to-sample task (Terry et al., 1998). The ability of selective 5-HT<sub>4</sub> receptor antagonists (e.g., GR 125487, RS 67532 and SDZ 205,557) to block the pro-cognitive actions of 5-HT<sub>4</sub> receptor agonists across several behavioral models strengthens the role of 5-HT<sub>4</sub> receptors in cognitive processes (Fontana et al., 1997; Lamirault and Simon, 2001; Moser et al., 2002). Lastly, synergistic effects with cholinesterase inhibitors (e.g., donepezil, galanthamine and rivastigmine) and 5-HT<sub>4</sub> receptor agonists have been observed in animal models of cognition (Cachard-Chastel et al., 2008; Mohler et al., 2007; Moser et al., 2002) suggesting therapeutic utility for AD. In addition to providing potential symptomatic relief, activation of the 5-HT<sub>4</sub> receptor *in vitro* stimulates the secretion of the soluble form of

amyloid precursor protein (sAPP $\alpha$ ), a neuroprotective protein in the brain (Lezoualc'h 2007; Robert et al., 2001) and *in vivo*, 5-HT $_4$  receptor agonists have been shown to increase sAPP $\alpha$  levels in cortex and hippocampus in young adult and transgenic APP-overexpressing mice (Cachard-Chastel et al., 2007; 2008). Taken together, these data provide strong preclinical support for 5-HT $_4$  receptor involvement in the treatment of AD symptomatology and pathology.

### 5.5 5-HT $_6$ receptors

The 5-HT $_6$  G-protein coupled receptor was first cloned from rat striatum using RT-PCR techniques in the early 1990's and its activation stimulates cAMP production and protein kinase A (Kohen et al., 1996; Monsma et al., 1993; Ruat et al., 1993). 5-HT $_6$  receptor expression is almost exclusively within the central nervous system (CNS) thereby limiting any potential peripheral side effects. Receptor distribution resides within brain areas (e.g., striatum, cortex, hippocampus and hypothalamus) responsible for mediating many of its observed preclinical effects on anxiety and depression (Svenningsson et al., 2007; Wesolowska and Nikiforuk 2007), epilepsy (Routledge et al., 2000), obesity (Heal et al., 2008) and the current topic, learning and memory (Fone, 2008; King et al., 2008). Initial evidence supporting the involvement of 5-HT $_6$  receptors in cognitive processes was derived from the finding that receptor knockdown after intracerebroventricular treatment with 5-HT $_6$  receptor antisense oligonucleotides (AO) improved retention of the learned hidden platform position during probe trials in the water maze task in normal rats (Bentley et al., 1997; Woolley et al., 2001). The significance of this receptor blockade was later confirmed when administration of 5-HT $_6$  receptor antagonists Ro-04-6790, SB 271046 and SB 357134 led to improved probe trial, but not acquisition learning performance in normal adult rats (Marcos et al., 2008; Rogers and Hagan, 2001; Stean et al., 2002; Woolley et al., 2001). Conversely, in aged rats 5-HT $_6$  receptor antagonists are capable of enhancing both acquisition learning and retention probe trial performance (Foley et al., 2004; Hirst et al., 2006; Stean et al., 2002) suggesting that within the water maze task 5-HT $_6$  receptor antagonists may have a greater influence on declining cognitive function especially as it relates to cholinergic activity. Indeed, these aged-related findings were extended by classifying the aged rat population as either being aged cognitively-impaired or aged cognitively-unimpaired based upon their acquisition water maze performance to that of young adult rats (Rowe et al., 2007). In these cognitively-impaired aged rats, hippocampal acetylcholine levels are significantly decreased compared to their cognitively-unimpaired cohort and therefore, represent a model of "natural" aged-related cognitive decline, akin to that observed in humans suffering from mild cognitive impairment (MCI) or AD (Quirion et al., 1995). Administration of 5-HT $_6$  receptor antagonists (e.g., MEM 68626, MEM 34551, SB 271046 and SB 742457) completely reversed the acquisition learning and probe retention trial water maze performance deficits observed in this aged cognitively-impaired cohort (Callahan et al., 2004; Rowe et al., 2008, 2009). SB 742457 and MEM 68626 were also shown to reverse the spatial working memory deficits of aged cognitively-impaired rats in the 8-arm radial water maze task (Rowe et al., 2008). Moreover, synergistic effects were demonstrated with a sub-threshold dose of the acetylcholinesterase inhibitor donepezil in combination with SB 271046 on spatial memory performance (Callahan et al., 2004), whereby the drug combination enhanced acquisition and probe trial performance of the aged cognitively-impaired animal over that performance observed for either drug given alone. These results suggest that 5-HT $_6$  receptor antagonists administered along with cholinesterase inhibitors may provided added beneficial effects in the treatment of cognitive dysfunction in humans. In addition to spatial reference and working memory improvements, 5-HT $_6$  receptor antagonists are active across multiple behavioral models reflecting various cognitive domains (e.g., episodic, associative, executive function, and working memory) and brain regions (e.g., cortical, hippocampal, striatal and amygdalar). 5-HT $_6$  receptor antagonists

reverse pharmacologically-induced deficits (e.g., scopolamine, MK 801 and phencyclidine) as well as possess stand-alone effects in young and aged rats in the following tasks, object recognition (Callahan et al., 2004, 2009; Gannon et al., 2007; Hirst et al., 2006; King et al., 2004, 2005, 2009; Schreiber et al., 2007; Wolff et al., 2002), social recognition (Loiseau et al., 2008; Mitchell et al., 2006; Schreiber et al., 2007), social discrimination (Schreiber et al., 2007), operant autoshaping (Meneses 2001; Perez-Garcia and Meneses, 2005; Schreiber et al., 2007), passive avoidance (Bos et al., 2001; Callahan et al., 2004; Foley et al., 2004; Riemer et al., 2003; Schreiber et al., 2007), attentional set-shifting (Hatcher et al., 2005; Rodefer et al., 2008) and radial arm maze (Wolff et al., 2002). Moreover, the pro-cognitive effects observed in rodents have been extended to include non-human primates, though a limited number of studies have been published to date. SB 271046 was shown to improve accuracy in aged rhesus monkeys in the delayed matching-to-sample task and in marmosets, SB 271046 reversed the MK 801-induced deficits on both perceptual visual and visuospatial discrimination tasks (Upton et al., 2008). Many of the behavioral effects on cognition in young and aged animals can be attributed to the ability of 5-HT<sub>6</sub> receptor antagonists to increase the release of acetylcholine (Hirst et al., 2006; Riemer et al., 2003; Shirazi-Southall et al., 2002; Zhang et al., 2007), glutamate (Dawson et al., 2000; Dawson and Nguyen, 2001) and the monoamines (Lacroix et al., 2004; Li et al., 2007) in brain regions involved in learning and memory processes. Collectively, these preclinical findings along with the recent positive phase II clinical trial effects with SB 742457 in mild to moderate AD subjects (Maher-Edwards et al., 2010; Upton et al., 2008) suggest that 5-HT<sub>6</sub> receptor antagonists may be viable drug candidates for the symptomatic relief of cognitive dysfunction associated with age-related cognitive decline, AD and possibly, schizophrenia.

More recently, 5-HT<sub>6</sub> receptor agonists (e.g., EMD 386088, E-6801, R-13c, WAY 181187 and WAY 208466) have been identified (Cole et al., 2005; Mattsson et al., 2005; Romero et al., 2006; Schechter et al., 2008) and their potential involvement on cognitive processes assessed (Burnham et al., 2010; Callahan et al., 2009; Fone, 2008; Kendall et al., 2010; Loiseau et al., 2008; Meneses et al., 2008; Rowe et al., 2009). Initial studies have yielded controversy with reports showing pro-cognitive effects on attentional set-shifting (Burnham et al., 2010) and object recognition (Fone, 2008; Kendall et al., 2010) while others have demonstrated either cognitive impairment on social recognition (Loiseau et al., 2008) and autoshaping (Meneses et al., 2008) or no effects on cognition in object recognition, passive avoidance and water maze (Callahan et al., 2009; Rowe et al., 2009). In many cases, co-administration of a selective 5-HT<sub>6</sub> receptor antagonist has been used to either reverse the agonist-induced deficit (Loiseau et al., 2008) or agonist-induced improvement (Burnham et al., 2010). In the object recognition task, Kendall and colleagues (2010) showed that both 5-HT<sub>6</sub> receptor agonists and antagonists enhanced recognition memory at a 4 hr delay and that combining sub-optimal doses of each led to enhanced memory performance, whereas Callahan and colleagues (2009) showed that object recognition memory was neither affected at a short 1 hr delay or enhanced at a 48 hr delay by the 5-HT<sub>6</sub> receptor agonist WAY 181187. Moreover, the pro-cognitive effect observed at the 48 hr delay interval by the antagonist SB 742457 was dose-dependently blocked by co-administration of WAY 181187. Further, whereas SB 742457 dose-dependently reversed the age-related cognitive impairment in Fischer rats, WAY 181187 was ineffective in reversing the natural age-related cognitive deficit (Rowe et al., 2009). Given that these investigators have shown pro-cognitive effects with 5-HT<sub>6</sub> receptor antagonists (as well as other drug classes) in the same behavioral models used to characterize 5-HT<sub>6</sub> receptor agonists, simply stating methodological differences across laboratories appears to be an unlikely reason for the present disparity. Additionally, one might argue that the differences stem from the use of non-selective 5-HT<sub>6</sub> receptor agonists, however many of the investigators have used the same 5-HT<sub>6</sub> receptor agonist, WAY 181187 (Burnham et al., 2010; Callahan et al., 2009; Loiseau et al., 2008; Rowe et al., 2009), although a few have reported on less well

characterized compounds (e.g., EMD 386088, E-6801 and R-13c; Fone, 2008; Kendall et al., 2010; Meneses et al., 2008) especially with respects to neurochemical profiling. For example, the high affinity and receptor selective 5-HT<sub>6</sub> agonist WAY 181187 has been shown to increase GABA release in rat frontal cortex, hippocampus, striatum and amygdala. Effects that are blocked by SB 271046 administration (Schechter et al., 2008), thus supporting the proposed disinhibitory action by 5-HT<sub>6</sub> receptor antagonists on cholinergic and glutamatergic activity. WAY 181187 has also been shown to decrease dopamine and serotonin release in rat cortex and striatum and inhibit potassium-stimulated glutamate release from hippocampal slices (Schechter et al., 2008), effects opposite to 5-HT<sub>6</sub> receptor antagonists. Additionally, WAY 181187 has been shown to attenuate CA1 hippocampal LTP function, an effect blocked by SB 399885 (West et al., 2009). Generally, pro-cognitive drugs (e.g., 5-HT<sub>4</sub> receptor agonists, Matsumoto et al., 2001) enhance LTP activity which represents the synaptic basis for learning and memory (Bliss and Collingridge, 1993). Taken together, these data suggest that 5-HT<sub>6</sub> receptor agonists would either have no effect on cognition or impair cognitive performance. Conversely, it's been argued that the cognitive enhancing properties of 5-HT<sub>6</sub> receptor agonists occur by direct activation of 5-HT<sub>6</sub> receptors located on cholinergic and/or glutamatergic neurons, however autoradiography (Roberts et al., 2002), immunohistochemical (Woolley et al., 2004) and cholinergic lesion studies (Marcos et al., 2006) suggest little 5-HT<sub>6</sub> receptor expression on cholinergic neurons (as well as glutamate neurons) and that the observed increase in acetylcholine and glutamate release by 5-HT<sub>6</sub> receptor antagonists is due to an indirect disinhibitory effect on GABAergic neurons. If 5-HT<sub>6</sub> receptor agonists directly modulate cholinergic and/or glutamatergic release then microdialysis studies will provide the much needed proof of concept for the proposed mechanism of action. Alternatively, 5-HT<sub>6</sub> receptor agonists could be activating second messenger intraneuronal signaling cascades such as cAMP and/or cGMP which are widely known to be involved in learning and memory processes (Reneerkens et al., 2009; Rose et al., 2005). 5-HT<sub>6</sub> receptors are functionally coupled to cAMP and agonists would be expected to facilitate cAMP production triggering downstream protein kinase A (PKA) or mitogen-activated protein (MAP) kinase activity thereby leading to activation of transcription factor cAMP-response element binding protein (CREB) and subsequently gene expression and protein synthesis which is necessary for long-term memory formation (Tully et al., 2003). 5-HT<sub>4</sub> receptors, like 5-HT<sub>6</sub> receptors, trigger this cAMP/PKA signaling pathway which is regulated by phosphodiesterases (PDEs) and it was recently demonstrated that the cognitive improvement in object recognition memory by the 5-HT<sub>4</sub> receptor agonist RS 67333 was due to a direct stimulation of prefrontal cortex and hippocampal PDE activity (Levallet et al., 2009). It is therefore plausible to suggest that the observed pro-cognitive effects associated with 5-HT<sub>6</sub> agonists may involve modification of this second messenger system and in particular, PDE activity. Similar studies will need to be executed with 5-HT<sub>6</sub> agonists to verify this potential mechanism of action.

## 6. Histamine H<sub>3</sub> Receptor Antagonists

Histamine is an important neurotransmitter involved in many physiological functions (e.g., allergies, gastric acid secretion, immunomodulation and neurotransmitter release; Leurs et al., 2005). Histamine exerts its effects through four G-protein-coupled histamine receptor subtypes (H1, H2, H3 and H4). The histamine H3 receptor (H3R) and, in particular, H3R antagonists are being pursued as potential drug targets for the treatment of cognitive dysfunction associated with age-related cognitive decline, AD, attentional deficit hyperactivity disorder (ADHD) and schizophrenia (Esbenshade et al., 2008; Witkin and Nelson, 2004). H3Rs are expressed almost exclusively in mammalian brain with high densities in cortex, hippocampus, striatum and hypothalamus, areas involved in wakefulness, attention, vigilance and cognition (Martinez-Mir et al., 1990; Pollard et al.,



1993). H3Rs are presynaptic autoreceptors that regulate the synthesis and release of histamine from histaminergic neurons (Arrang et al., 1983; 1987). H3Rs also function as heteroreceptors and mediate the release of not only histamine but other neurotransmitters such as acetylcholine, dopamine, noradrenaline and serotonin (Leurs et al., 2005; Schlicker et al., 1994) and inhibition of H3Rs with selective antagonists can augment the release of neurotransmitters involved in cognition. (Fox et al., 2005; Galici et al., 2009; Medhurst et al., 2007a). Due to the high constitutive activity of native H3Rs and the ability of H3R antagonists to inhibit G-protein-coupled receptor signaling, some H3R antagonists have been termed “inverse agonist” (e.g., ciproxifan, thioperamide, ABT-239, BF 2.649, GSK 189254; Fox et al., 2005; Leurs et al., 2005; Ligneau et al., 2007; Medhurst et al., 2007a), however, not all H3R antagonists show this intrinsic activity and are considered to be “neutral” antagonists (e.g., JNJ- 5207852; JNJ-10181457; Barbier et al., 2004; Galici et al., 2009).

In preclinical behavioral models of cognition, a variety of H3R antagonists have proven to be efficacious in restoring and/or enhancing cognitive performance (for comprehensive reviews see Esbenshade et al., 2008; Leurs et al., 2005; Witkin and Nelson 2004). For example, studies with the first generation H3R antagonists such as thioperamide, ciproxifan or clobenpropit were shown to reverse passive avoidance (Giovannini et al., 1999; Miyazaki et al., 1997), object recognition (Giovannini et al., 1999; Pascoli et al., 2009), two-choice discrimination water maze, Barnes maze (Komater et al., 2005) and radial arm maze (Chen, 2000) impairments produced by scopolamine treatment. Additional support for cholinergic modulation came from the observation that newly developed H3R antagonists like BF 2.649, GSK 207040, GSK 334429, GSK 189254 and JNJ-10181457 were also efficacious in reversing scopolamine-induced passive avoidance (Medhurst et al., 2007a,b), object recognition (Ligneau et al., 2007) and delayed non-matching to position (DNMTP; Galici et al., 2009) performance deficits. Working and reference memory deficits produced by MK-801 administration have also been attenuated by clobenpropit in the rat radial arm maze task suggesting an interaction of histamine on NMDA function (Huang et al., 2004). H3R antagonists are also capable of enhancing cognitive performance of young and aged animals in the absence of pharmacological impairment. Administration of ciproxifan has been shown to improve attention and decrease impulsivity in the rat 5-choice serial reaction time test (5-CSRTT) which is analogous to the continuous performance test (CPT) widely used to assess attention in humans (Day et al., 2007; Ligneau et al., 1998). Thioperamide, BF 2.649 and GSK 189254 have been shown to enhance object recognition memory in young rats and mice when drug is given prior to training (Ligneau et al., 2007; Medhurst et al., 2007a; Rowe et al., 2006) but a recent report by Pascoli and colleagues (2009) using ciproxifan and thioperamide showed that recognition memory was only enhanced when drug was given immediately before the retention test session. The authors indicated that the H3R antagonists were unable to enhance recognition memory when the drugs were given prior to or immediately after the training trial. Additional work is required to clarify these observed differences. The H3R antagonists A-304121, A-317920, ABT-239 and JNJ-10181457 improved attention in a repeated acquisition inhibitory avoidance task in spontaneously hypertensive (SHR) rat pups (Bonaventure et al., 2007; Fox et al., 2003; 2005) and A-304121, A-317920 and ABT-239 also improved short-term memory recall in young adult (Fox et al., 2003; 2005) and aged rats (Fox et al., 2005) with a potency similar to that of ciproxifan. ABT-239 was also able to reverse the scopolamine-induced spatial memory deficit in a two-choice discrimination water maze task (Fox et al., 2005). Similarly, GSK 189254 improved water-maze acquisition latency and probe trial performance in aged rats (Medhurst, 2007a). Moreover, when aged rats are screened and classified as either aged cognitively-impaired or aged cognitively-unimpaired based upon their water-maze acquisition latency performance to that of young rats and testing occurs in the aged cognitively-impaired cohort, the H3R antagonist thioperamide is capable of completely

reversing the acquisition latency and probe trial performance deficits (Rowe et al., 2006). Rowe and colleagues (2006) extended this positive effect of thioperamide on cognitive restoration of spatial memory in aged cognitively-impaired mice and thioperamide also improved avoidance learning in senescence-accelerated mice (Meguro et al., 1995). In addition to the positive behavioral effects of H3R antagonists across multiple cognitive domains and the therapeutic possibilities to treat ADHD, age-related cognitive decline, AD and various sleep disorders, H3R antagonists may also play an important role in the treatment of schizophrenia. H3R antagonists have been demonstrated to reduce the hyperlocomotor effects of dopamine agonists (e.g., amphetamine, apomorphine and methamphetamine; Akhtar et al., 2006; Clapham and Kilpatrick, 1994; Fox et al., 2005), reduce apomorphine-induced climbing behavior (Akhtar et al., 2006) and in some cases (Akhtar et al., 2006; Pillot et al., 2002) but not all (Zhang et al., 2005) potentiate the cataleptic effects of antipsychotics. H3R antagonists have also been shown to reverse the inherent sensorimotor gating deficits of DBA/2J mice in prepulse inhibition of startle (PPI; Browman et al., 2004) and hippocampal N40 sensory gating (Fox et al. 2005). Additionally, the H3R antagonist GSK 207040 attenuated the isolation rearing-induced deficit in PPI in rats (Southam et al., 2009) and GSK 189254 enhanced cognitive flexibility and executive memory function in the attentional set-shifting paradigm (Medhurst et al., 2007a). Taken together, these data strongly support a role for H3R antagonists in modulating the neurophysiological attributes associated with schizophrenia.

## 7. Multiple Drug Targets and Multifunctional Compounds

In complex illnesses such as AD, it could be argued that a multiple drug target approach to therapy is necessary to address the complex pathophysiology of the disease and its diverse symptoms. However, even if the strategy of combining drugs with different therapeutic targets is feasible, the development of multi-functional compounds would circumvent the challenge of administering multiple drugs with potentially different degrees of bioavailability, pharmacokinetics, and metabolism (reviewed, Youdim and Buccafusco, 2005). An additional advantage to single drugs with multiple actions is the simplification of the therapeutic regimen and improved compliance (an important consideration, especially for individuals who suffer from memory-related disorders). Another advantage is the potential to treat multiple symptoms including both cognitive deficits and non-cognitive behavioral symptoms. This is especially important given that the currently available AD therapeutic agents (in addition to modest efficacy and adverse side effects), do not effectively treat the very problematic non-cognitive behavioral symptoms (e.g., agitation) of the illness. Such behavioral symptoms are thus; often managed clinically by potent antipsychotic drugs which are particularly problematic in older patients since they are associated with movement disorders and/or a wide variety of metabolic abnormalities (e.g., weight gain, hyperglycemia, hyperlipidemia, reviewed, Myamoto et al., 2005) as well as an increase mortality (Ballard et al., 2009). Thus, a compound exhibiting pro-cognitive and antipsychotic properties might be specialty valuable in AD as well as conditions like schizophrenia which is characterized by abnormal behavioral symptoms as well as cognitive deficits. There would also be a number of advantages to an agent that exhibits both pro-cognitive and neuroprotective activity or other disease modifying action. The obvious advantages would be enhanced cognition (and improved day to day quality of life) while at the same time delaying or preventing the progression of the illness (properties not observed in the currently available therapies). One potential advantage of a multifunctional compound over a disease-modifying agent that has no acute pro-cognitive actions pertain to the design and length of clinical trials (reviewed, Buccafusco, 2009). Clinical trials of disease modifying agents (by definition) require long evaluation periods which are often cost-prohibitive. A pro-cognitive compound (with disease-modifying actions) could be approved for cognition enhancement in shorter clinical trials, providing a return on the manufacturer's

investment while the more lengthy evaluations of disease-modifying actions could be carried out.

### 7.1 Ladostigil

Ladostigil is a multi-functional compound designed to incorporate the cognitive enhancing properties of rivastigmine (Ohara et al., 1997; Ballard and McAllister, 1999; Wesnes et al., 2002) with the neuroprotection properties of rasagiline (Maruyama et al., 2001; Weinreb et al., 2004). As such, the carbamate moiety of rivastigmine was introduced in the 6 position of rasagiline and a compound conveying inhibition at both acetylcholinesterase and monoamine oxidase (MAO) was born (Weinstock et al., 2000; Sterling et al., 2002; Buccafusco et al., 2003). MAO is a family of enzymes which catalyze the oxidation of biogenic amine neurotransmitters such as norepinephrine, dopamine and 5-HT. Ladostigil is a brain specific inhibitor of MAO-A and MAO-B and has thus been linked to an increase norepinephrine, dopamine and 5-HT levels in the brain after administration (Buccafusco et al., 2003; Sagi et al., 2003; Weinstock et al., 2003). Because of this MAO inhibition, ladostigil has been shown to be effective in animal models to assess antidepressant activity. (Weinstock et al., 2002) More importantly to the treatment of AD, ladostigil has been shown to be neuroprotective similar to other MAO inhibitors (Weinstock et al., 2001; Yogev-Falach et al., 2002; Sagi et al., 2003). Another interesting neuroprotective property of ladostigil centers on its ability to modulate APP processing through mitochondrial membrane potential stabilization and regulation of PKC and mitogen-activated protein kinase (MAPK) dependant signaling pathways (Yogev-Falach et al., 2002; Weinreb et al., 2008; Bar-Am et al., 2009) to reduce the production of neurotoxic amyloid beta (Yogev-Falach et al., 2006). Ladostigil also increases catalase activity and has thus been shown to protect against oxidative stress related apoptosis (Weinreb et al., 2008). Further, ladostigil can convey neuroprotection by increasing brain-derived nerve factor (BDNF) mRNA expression and production (Van der Schyf et al., 2007). Ladostigil has also shown pro-cognitive effects in a wide range of preclinical animal models (Buccafusco et al., 2003; Shoham et al., 2007; Luques et al., 2007; Weinstock et al., 2009) Administration of ladostigil was capable of preventing glial activation and cognitive deficits in object and place recognition tasks in a streptozotocin (STZ) model of AD (Shoham et al., 2007). More recently, ladostigil has been shown to enhance spatial memory in the Morris water maze by increasing proNGF, prevent age related increases in activation of astrocytes and microglia, and prevent age related reductions in cortical AChE activity in the hippocampus of aged rats. (Weinstock et al., 2009) Ladostigil has also shown pro-cognitive effects in a delayed matching-to-sample (DMTS) distracter task, as well as a DMTS titrating task in monkeys (Buccafusco et al., 2003). Similarly, ladostigil has shown positive cognitive effects in a cytochrome oxidase model of AD by preventing the decrease in choline acetyltransferase (ChAT) and restoring memory deficits in a object recognition task (Luques et al., 2007). Due to ladostigil's positive effects on such a wide range of Alzheimer's related targets (i.e. neuroprotective activity, regulatory effects on APP processing, stimulatory effects on PKC and MAPK, effects on BDNF production), it is easy to see why taking a more encompassing approach in future drug design could be beneficial in AD drug development.

### 7.2 Dimebolin

Dimebolin originally developed and used for its anti-histaminergic properties since the early 1980's in Russia, has recently enjoyed a new found interest as a potential treatment for AD. Dimebolin is a multifunctional drug enjoying a broad range activity profile on many targets relating to the treatment of AD. Dimebolin is a weak acetylcholinesterase ( $IC_{50} = 7.9 \mu M$ ) and butyrylcholinesterases ( $IC_{50} = 42 \mu M$ ) inhibitor (Bachurin et al., 2001). The compound also displays antagonistic properties at the glutamatergic NMDA (Bachurin et al., 2001; Wu and Li, 2008) receptor and has shown positive modulation of glutamate AMPA receptors

(Grigorev et al., 2003). Further, dimebolin administration is able to protect neurons from the damaging effects of excessive cell stimulation through antagonistic action ( $IC_{50} = 50 \mu M$ ) at L-type  $Ca^{2+}$  channels (Lermontova et al., 2001). Dimebolin is also an antagonist to the brain 5-HT<sub>6</sub> receptor and has been shown to improve social recognition memory similar to other 5-HT<sub>6</sub> receptor antagonists (Schaffhauser et al., 2009). Dimebolin has also shown to be neuroprotective through attenuating amyloid beta induced neurotoxicity in rat cortical neuron cultures (Lermontova et al., 2001) and through its regulation of mitochondrial pores after neurotoxic insult (Bachurin et al., 2003). Dimebolin has also shown pro-cognitive effects on behavior in a senescence-accelerated mouse prone 10 (SAMP 10) model of aging by enhancing short-term memory assessed by a passive avoidance task (Grigorev et al., 2003). Similarly, the compound has been shown to improve cognition in a rat novel object recognition task by improving retention punitively through 5-HT<sub>6</sub> receptor antagonism. However, the exact mechanism through which dimebolin produces its cognitive enhancing effects is not know. Still dimebolin is active at many targets (AChE inhibition, NMDA antagonism, H<sub>3</sub> receptor antagonism, and 5-HT<sub>6</sub> receptor antagonism) that related to cognitive enhancement and has shown positive results in several preclinical models of cognition as well as in AD patients (Doody et al., 2008.) Other reports have failed to see this improvement in AD patients treated with dimebolin (Miller, 2010). Weather or not dimebolin will prove itself as a viable treatment for AD is still uncertain. However, there remains an active interest in finding multi-functional drugs to treat AD.

### 7.3 JWS-USC-75-IX

Several years ago, our collaborators (Valli et al., 1992) synthesized a series of analogs of the ranitidine (an H<sub>2</sub> antagonist commonly used to treat duodenal ulcers) for the purpose of creating non-toxic AChEIs. The impetus for this work was the prior observation that ranitidine (a histamine H<sub>2</sub> antagonist commonly used to treat duodenal ulcers) had mild AChEI properties in vitro (Gwee and Cheah, 1986). Several of the compounds from this series of ranitidine analogs were subsequently found to possess potent AChEI properties as well as low toxicity profiles. Of this series, one compound, JWS-USC-75-IX (3-[[[2-[[[5-dimethylaminomethyl]-2-furanyl]methyl]thio]ethyl]amino]-4-nitropyridazine) was found to possess AChEI activity in vitro ( $IC_{50} \sim 0.47 \mu M$ ) as well as potent M<sub>2</sub> receptor antagonist activity in mouse cerebral cortex ( $IC_{50} \sim 60 \text{ nM}$ ). As had been suggested previously (Quirion, 1993; Mash et al., 1985), an M<sub>2</sub> (autoreceptor) antagonist might be very useful given in conjunction with an AChEI in AD to prevent the acetylcholine from decreasing its own release. JWS-USC-75-IX, thus, combined both features in a single molecule and offered a significant potential for improved reliability as a treatment of diseases where cholinergic function is impaired (e.g., AD). Later we (Terry et al., 1999) evaluated JWS-USC-75-IX for effects in memory-related tasks in rodents. The compound was found to enhance spatial learning, inhibitory avoidance, and working memory. More recent studies in our laboratories indicate that JWS-USC-75-IX has the potential to improve other domains of cognition in rodents (e.g., recognition memory) as well as attention and working memory in non-human primates (Terry et al., 2011).

### 7.4 Additional Multifunctional Compounds

The compound memoquin resulted from a multi-targeted approach, combining the radical scavenger moiety of coenzyme Q (CoQ) with the polyamine backbone of caproctamine conferring AChE inhibition properties. The rationale was to produce a single molecular entity that combined the neuroprotective qualities of CoQ with the cognitive enhancing properties of AChE inhibition. The result was a molecule that has been shown to simultaneously reduce the neurodegenerative pathology and rescue the behavioral impairment in an object recognition task in a mouse (AD11 mice) model of AD (Cavalli, 2007). Similarly, PMS777 is a multiple target AChE inhibitor that displays dual inhibition of

AChE and platelet-activating factor (PAF). PAF is thought to participate in the neurodegeneration related AD (Bazan, 2002; Bate, 2004). Thus, the rationale for PMS777 is to act through neuroprotection as well as cognitive enhancement. PMS777 has been shown to alleviate scopolamine induced impairment in mice (Li, 2006).

## 8. Other Targets

For several years chromatin modifications, especially histone-tail acetylation, have been implicated in modulating synaptic plasticity and long-lasting changes of neural circuits that contribute to memory formation. Moreover, increased histone-tail acetylation induced by inhibitors of histone deacetylases (HDACs) has been observed to facilitate learning and memory in wild-type mice as well as in mouse models of neurodegeneration (Guan et al., 2009). Accordingly, selective inhibitors HDAC (e.g., Bayer's compound EVP 0334) have been suggested as treatments for neurodegenerative diseases, including AD (Hannen et al., 2008).

Both angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blocker (ARB) drugs have been shown to improve cognitive function in animal models (reviewed, Gard, 2004) and as a result have been in various stages of development for AD over the last 15 years. While no such agent has progressed very far in clinical trials to date, recent observations that ARBs are associated with a significant reduction in the incidence and progression of AD and dementia (Li et al., 2010) may suggest that such a strategy remains viable.

While psychostimulant drugs (e.g., amphetamine, methylphenidate) have been employed for many years as effective treatments for ADHD and narcolepsy, their potential as pro-cognitive agents in other disorders has been equivocal. Interest in newer ADHD and narcolepsy-related treatments such as the norepinephrine transporter (NET) inhibitor atomoxetine and the psychostimulant modafinil, respectively, is accruing and based on recent published evidence. For example atomoxetine was observed to improve reversal learning in rats and monkeys (Seu et al., 2009) while modafinil has been observed to improve cognition in several domains including working memory and episodic memory in both rodent models and humans (reviewed, Minzenberg and Carter, 2008).

There is a host of additional therapeutic targets that have been investigated or are in various stages of investigation for the symptomatic treatment of AD, MCI and cognitive deficits associated with schizophrenia. Unfortunately, a full overview of these targets is beyond the scope of the present review and we refer the reader to additional articles (Gallagher et al., 1985; Minzenberg and Carter, 2008; Buccafusco, 2009; Lynch et al., 2011;) that discuss examples such as glycine transporter related drugs, mGluR2/3 agonists, somatostatin receptor ligands, estrogen receptor ligands, D2 agonists, calcium channel modulators (e.g., MEM 1003, nimodipine), adrenergic compounds (e.g., nicergoline), opioid antagonists (e.g., naloxone) and GABA-related compounds.

## 9. Animal Models and Their Translational Value

In the last several years there has been a growing concern over the discrepancies between preclinical results in AD-related animal models and clinical evaluations of novel AD-related compounds (Carlsson, 2008). A more general concern over the translation of preclinical results to clinical trial data is now evident in multiple health-related disciplines (see Lowenstein, 2009; Simon, 2008). It is clear that new and more rigorous efforts to develop animal models that meet the demands of face, construct, and predictive validity be undertaken so that preclinical endeavors become more translational. It is also our opinion that specific batteries of behavioral tasks that more closely map onto the domains of

cognition that are evaluated in clinical trials be developed and that some type of composite score approach to the animal tests be created that is analagous to those used in clinical studies (e.g., Mini-Mental Status Exam) so that an “overall” drug effect on cognition can be estimated. It is also very important that the full results of failed clinical trials be published in peer reviewed journals (a practice that is uncommon at present) so that the human and animal data can be carefully compared. Currently, such clinical trial results are often found only in abstract form or as summary data in reviews or lay publications, a practice that often makes it difficult for the science community to understand the reason for the so called “failure” of an investigational new drug (see review, McArthur et al., 2010).

## 10. Summary and Conclusions

Given the rapid growth of our elderly population and their concerns about memory loss and dementia it is incumbent on the scientific community to develop more effective pro-cognitive drugs. As diagnostic criteria for memory disorders evolve, the demand for pro-cognitive therapeutic agents is likely to move beyond AD and dementia to include MCI and potentially even less severe forms of memory decline. While there are a considerable number of compounds in various stages of preclinical development as pro-cognitive agents, due to the complexity of human cognition (and illness like AD that devastate cognition), the development of compounds that are both safe and effective (with sustained pro-cognitive effects over time) will likely require the diligent efforts of academia and private industry and the full armamentarium of a variety of scientific methods. Such methods include modern drug discovery techniques (e.g., molecular modeling, combinatorial chemistry, high throughput biological assays), the latest advances in molecular biology and transgenics, and more traditional research methods such as medicinal chemistry, behavioral neuroscience, and classical pharmacology.

## Acknowledgments

The authors would like to thank Ms. Ashley Davis for her administrative assistance in preparing this article. This corresponding author's laboratories are supported by the National Institute of Aging (AG032140 and AG029617), the National Institute of Environmental Health Sciences (ES012241), and the National Institute on Drug Abuse (DA029127). The corresponding author has also provided consultation or performed research (relevant to this article) in the last three years either contractually or in collaboration with a several pharmaceutical companies including Abbott Laboratories, Helicon, Merck, Neuralstem, Neurosearch, Roche Palo Alto, and Servier. This review is dedicated to our close personal friend, mentor, and colleague, the late Dr. Jerry J. Buccafusco whose contributions to the science of drug discovery and development for disorders of cognition spanned more than 25 years.

## References

- Akhondzadeh S, Mohammadi N, Noroozian M, Karamghadiri N, Ghoreishi A, Jamshidi AH, et al. Added ondansetron for stable schizophrenia: a double blind, placebo controlled trial. *Schizophr Res*. 2009; 107:206–212. [PubMed: 18789844]
- Akhtar M, Uma Devi P, Ali A, Pillai KK, Vohora D. Antipsychotic-like profile of thioperamide, a selective H3-receptor antagonist in mice. *Fundam Clin Pharmacol*. 2006; 20:373–378. [PubMed: 16867021]
- Albuquerque EX, Pereira EF, Alkondon M, Rogers SW. Mammalian nicotinic acetylcholine receptors: from structure to function. *Physiol Rev*. 2009; 89:73–120. [PubMed: 19126755]
- Araujo JA, Studzinski CM, Milgram NW. Further evidence for the cholinergic hypothesis of aging and dementia from the canine model of aging. *Prog Neuropsychopharmacol Biol Psychiatry*. 2005; 29(3):411–422. [PubMed: 15795050]
- Arnsten AF, Lin CH, Van Dyck CH, Stanhope KJ. The effects of 5-HT3 receptor antagonists on cognitive performance in aged monkeys. *Neurobiol Aging*. 1997; 18:21–28. [PubMed: 8983029]

- Arrang JM, Garbarg M, Schwartz JC. Auto-inhibition of brain histamine release mediated by a novel class (H3) of histamine receptor. *Nature*. 1983; 302:832–837. [PubMed: 6188956]
- Arrang JM, Garbarg M, Schwartz JC. Autoinhibition of histamine synthesis mediated by presynaptic H3-receptors. *Neuroscience*. 1987; 23:149–157. [PubMed: 2446202]
- Auld DS, Kornecook TJ, Bastianetto S, Quirion R. Alzheimer's disease and the basal forebrain cholinergic system: relations to beta-amyloid peptides, cognition, and treatment strategies. *Prog Neurobiol*. 2002; 68:209–245. [PubMed: 12450488]
- Bach ME, Barad M, Son H, Zhuo M, Lu YF, Shih R, et al. Age-related deficits in spatial memory are correlated with deficits in the late phase of hippocampal long-term potentiation in vitro and are attenuated by drugs that enhance the cAMP signaling pathway. *Proc Natl Acad Sci USA*. 1999; 96:5280–5285. [PubMed: 10220457]
- Bachurin S, Bukatina E, Lermontova N, Tkachenko S, Afanasiev A, Grigoriev V, et al. Antihistamine agent Dimebon as a novel neuroprotector and a cognition enhancer. *Ann N Y Acad Sci*. 2001; 939:425–435. [PubMed: 11462798]
- Ballard C, Hanney ML, Theodoulou M, Douglas S, McShane R, Kossakowski K, et al. The dementia antipsychotic withdrawal trial (DART-AD): long-term follow-up of a randomised placebo-controlled trial. *Lancet Neurol*. 2009 Feb; 8(2):151–157. [PubMed: 19138567]
- Ballard J. Forgetfulness and older adults: concept analysis. *J Adv Nurs*. 2010 Jun; 66(6):1409–1419. [PubMed: 20384638]
- Ballard TM, McAllister KH. The acetylcholinesterase inhibitor, ENA 713 (Exelon), attenuates the working memory impairment induced by scopolamine in an operant DNMT task in rats. *Psychopharmacology (Berl)*. 1999; 146:10–18. [PubMed: 10485959]
- Barad M, Bourtchouladze R, Winder DG, Golan H, Kandel E. Rolipram, a type IV-specific phosphodiesterase inhibitor, facilitates the establishment of long-lasting long-term potentiation and improves memory. *Proc Natl Acad Sci USA*. 1998; 95:15020–15025. [PubMed: 9844008]
- Bar-Am O, Weinreb O, Amit T, Youdim MB. The novel cholinesterase-monoamine oxidase inhibitor and antioxidant, ladostigil, confers neuroprotection in neuroblastoma cells and aged rats. *J Mol Neurosci*. 2009; 37:135–145. [PubMed: 18751929]
- Barbier AJ, Berridge C, Dugovic C, Laposky AD, Wison SJ, Boggs J, et al. Acute wake-promoting actions of JNJ-5207852, a novel, diamine-based H3 antagonist. *Brit J Pharmacol*. 2004; 143:649–661. [PubMed: 15466448]
- Barnes JM, Costall B, Coughlan J, Domeney AM, Gerrard PA, Kelly ME, et al. The effects of ondansetron, a 5-HT3 receptor antagonist, on cognition in rodents and primates. *Pharmacol Biochem Behav*. 1990; 35:955–962. [PubMed: 2140610]
- Barnes NM, Sharp T. A review of central 5-HT receptors and their function. *Neuropharmacology*. 1999; 38:1083–1152. [PubMed: 10462127]
- Bartolomeo AC, Morris H, Buccafusco JJ, Kille N, Rosenzweig-Lipson S, Husbands MG, et al. The preclinical pharmacological profile of WAY-132983, a potent M1 preferring agonist. *J Pharmacol Exp Ther*. 2000; 292(2):584–596. [PubMed: 10640295]
- Bartus RT, Dean RL 3rd, Beer B, Lippa AS. The cholinergic hypothesis of geriatric memory dysfunction. *Science*. 1983; 217(4558):408–414. [PubMed: 7046051]
- Bartus RT. On neurodegenerative diseases, models, and treatment strategies: lessons learned and lessons forgotten a generation following the cholinergic hypothesis. *Exp Neurol*. 2000 Jun; 163(2):495–529. [PubMed: 10833325]
- Bate C, Salmona M, Williams A. The role of platelet activating factor in prion and amyloid-b neurotoxicity. *Neuroreport*. 2004; 15:509–513. [PubMed: 15094513]
- Bazan NG, Colangelo V, Lukiw WJ. Prostaglandins and other lipid mediators in Alzheimer's disease. *Prostaglandins & Other Lipid Mediators*. 2002:68–69. 197–210.
- Beach T, Walker D, Potter L, Sue L, Scott S, Layne K, et al. Immunotoxin lesion of the cholinergic nucleus basalis causes Aβ deposition: towards a physiologic animal model of Alzheimer's disease. *Curr Med Chem*. 2003; 3:233–243.
- Bender AT, Beavo JA. Cyclic nucleotide phosphodiesterases: molecular regulation to clinical use. *Pharmacol Rev*. 2006; 58:488–520. [PubMed: 16968949]
- Benowitz NL. Clinical pharmacology of nicotine. *Ann Rev Med*. 1986; 37:21–32. [PubMed: 3518606]

- Bentley J, Sleight AJ, Marsden CA, Fone KCF. 5-HT<sub>6</sub> antisense oligonucleotide i.c.v. affects rat performance in the water maze and feeding. *J Psychopharmacol.* 1997; 11:A64.
- Benzi G, Moretti A. Is there a rationale for the use of acetylcholinesterase inhibitors in the therapy of Alzheimer's disease? *Eur J Pharmacol.* 1998; 346(1):1–13. [PubMed: 9617746]
- Bertrand F, Lehmann O, Galani R, Lazarus C, Jeltsch H, Cassel JC. Effects of MDL 73005 on water-maze performances and locomotor activity in scopolamine-treated rats. *Pharmacol Biochem Behav.* 2001; 68:647–660. [PubMed: 11526961]
- Bliss TV, Collingridge GL. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature.* 1993; 361:31–39. [PubMed: 8421494]
- Boada-Rovira M, Brodaty H, Cras P, Baloyannis S, Emre M, Zhang R, et al. Efficacy and safety of donepezil in patients with Alzheimer's disease: results of a global, multinational, clinical experience study. *Drugs Aging.* 2004; 21(1):43–53. [PubMed: 14715043]
- Bockaert J, Claeysen S, Compan V, Dumuis A. 5-HT<sub>4</sub> receptors. *Curr Drug Targets CNS Neurol Diord.* 2004; 3:39–51.
- Boess FG, Hendrix M, van der Staay FJ, Erb C, Schreiber R, van Staveren W, et al. Inhibition of Phosphodiesterase 2 increases neuronal cGMP, synaptic plasticity and memory performance. *Neuropharmacology.* 2004; 47:1081–1092. [PubMed: 15555642]
- Bonaventure P, Hall H, Gommeren W, Cras P, Langlois X, Jurzak M, et al. Mapping of serotonin 5-HT<sub>4</sub> receptor mRNA and ligand binding sites in the post-mortem human brain. *Synapse.* 2000; 36:35–46. [PubMed: 10700024]
- Bonaventure P, Letavic M, Dugovic C, Wilson S, Aluisio L, Pudiak C, et al. Histamine H<sub>3</sub> receptor antagonist: from target identification to drug leads. *Biochem Pharmacol.* 2007; 73:1084–1096. [PubMed: 17129577]
- Bos M, Sleight AJ, Godel T, Martin JR, Reimer C, Stadler H, 5-HT<sub>6</sub> receptor antagonists: lead optimization and biological evaluation of N-aryl and N-heteroaryl 4-amino-benzene sulfonamides. *Eur J Med Chem.* 2001; 36:165–178. [PubMed: 11311747]
- Bourtchouladze R, Lidge R, Catapano R, Stanley J, Gossweiler S, Romashko D, et al. A mouse model of Rubinstein-Taybi syndrome: defective long-term memory is ameliorated by inhibitors of Phosphodiesterase 4. *Proc Natl Acad Sci USA.* 2003; 100:10518–10522. [PubMed: 12930888]
- Bradley SR, Lameh J, Ohrmund L, Thomas S, Bajpai A, Nguyen D, et al. AC-260584, an orally bioavailable M<sub>1</sub> muscarinic receptor allosteric agonist, improves cognitive performance in an animal model. *Neuropharm.* 2010; 58:365–373.
- Brandeis R, Sapir M, Hafif N, Abraham S, Oz N, Stein E, et al. AF150(S): a new functionally selective M<sub>1</sub> agonist improves cognitive performance in rats. *Pharmacol Biochem Behav.* 1999; 51(4):667–674. [PubMed: 7675841]
- Bratt AM, Kelly ME, Domeney AM, Naylor RJ, Costall B. Ondansetron fails to attenuate a scopolamine-induced deficit in a stone maze task. *Neuroreport.* 1994; 5:1921–1924. [PubMed: 7841377]
- Briggs CA, Anderson DJ, Brioni JD, Buccafusco JJ, Buckley MJ, Campell JE, et al. Functional characterization of the novel neuronal nicotinic receptor ligand GTS-21 in vitro and in vivo. *Pharmacol Biochem Behav.* 1997; 57:231–241. [PubMed: 9164577]
- Brioni JD, Kim DJ, O'Neill AB. Nicotine cue: lack of effect of the alpha 7 nicotinic receptor antagonist methyllycaconitine. *Eur J Pharmacol.* 1996; 301:1–5. [PubMed: 8773440]
- Browman KE, Komater VA, Curzon P, Rueter LE, Hancock AA, Decker MW, et al. Enhancement of prepulse inhibition of startle in mice by the H<sub>3</sub> receptor antagonists thioperamide and ciproxifan. *Behav Brain Res.* 2004; 153:69–76. [PubMed: 15219708]
- Buccafusco JJ. Neuronal nicotinic receptor subtypes: defining therapeutic targets. *Mol Interv.* 2004; 4(5):285–295. [PubMed: 15471911]
- Buccafusco JJ, Terry AV Jr, Decker MW, Gopalakrishnan M. Profile of nicotinic acetylcholine receptor agonists ABT-594 and A-582941, with differential subtype selectivity, on delayed matching accuracy by young monkeys. *Biochem Pharmacol.* 2007; 74:1202–1211. [PubMed: 17706609]



- Buccafusco JJ, Terry AV Jr. A reversible model of the cognitive impairment associated with schizophrenia in monkeys: potential therapeutic effects of two nicotinic acetylcholine receptor agonists. *Biochem Pharmacol.* 2009; 78(7):852–862. [PubMed: 19577545]
- Buccafusco JJ, Terry AV Jr, Goren T, Blaugrun E. Potential cognitive actions of (n-propargyl-(3r)-aminoindan-5-yl)-ethyl, methyl carbamate (tv3326), a novel neuroprotective agent, as assessed in old rhesus monkeys in their performance of versions of a delayed matching task. *Neuroscience.* 2003; 119:669–678. [PubMed: 12809688]
- Buccafusco JJ, Beach JW, Terry AV Jr. Desensitization of nicotinic acetylcholine receptors as a strategy for drug development. *J Pharmacol Exp Ther.* 2009 Feb; 328(2):364–370. [PubMed: 19023041]
- Buccafusco JJ. Emerging cognitive enhancing drugs. *Expert Opin Emerg Drugs.* 2009 Dec; 14(4):577–589. [PubMed: 19772371]
- Burnham KE, Baxter MG, Bainton JR, Southam E, Dawson LA, Bannerman DM, et al. Activation of 5-HT<sub>6</sub> receptors facilitates attentional set shifting. *Psychopharmacology.* 2010; 208:13–21. [PubMed: 19902184]
- Caccamo A, Oddo S, Billings LM, Green KN, Martinez-Coria H, Fisher A, et al. M1 receptors play a central role in modulating AD-like pathology in transgenic mice. *Neuron.* 2006; 49:671–682. [PubMed: 16504943]
- Cachard-Chastel M, Lezoualc'h F, Dewachter I, Delomenie C, Croes S, Devijver H, et al. 5-HT<sub>4</sub> receptor agonists increase sAPP $\alpha$  levels in the cortex and hippocampus of male C57BL/6j mice. *Brit J Pharmacol.* 2007; 150:883–892. [PubMed: 17325649]
- Cachard-Chastel M, Devers S, Sicsic S, Langlois M, Lezoualc'h F, Gardier AM, et al. Prucalopride and donepezil act synergistically to reverse scopolamine-induced memory deficit in C57BL/6j mice. *Behav Brain Res.* 2008; 187:455–461. [PubMed: 18061284]
- Callahan PM, Ilch CP, Rowe WB, Tehim A. Characterization of the selective 5-HT<sub>6</sub> receptor antagonist SB-271046 in behavioral models of cognition. *Soc Neurosci Abstr.* 2004 776.19.
- Callahan PM, Wang S, Xie W, Dragan S, Sun S, Michael T, Herbert B, Rowe WB, Tehim A, Lowe DA, Barrett JE. Pharmacological characterization of MEM 3454, a novel nicotinic  $\alpha$ 7 receptor partial agonist: Therapeutic potential for the cognitive deficits associated with Alzheimer's disease and schizophrenia. *Neuropsychopharmacology.* 2006; 31 Supl 1:S198.
- Callahan PM, Rowe WB, Hsu CC, Tehim A. Characterization of the 5-HT<sub>6</sub> receptor agonist, WAY 181187 and antagonist, SB 742457: effects on recognition, associative and spatial reference memory function in young animals. *Soc Neurosci Abstr.* 2009 883.3.
- Carey GJ, Costall B, Domeney AM, Gerrard PA, Jones DN, Naylor RJ, et al. Ondansetron and arecoline prevent scopolamine-induced cognitive deficits in the marmoset. *Pharmacol Biochem Behav.* 1992; 42:75–83. [PubMed: 1388279]
- Carey GJ, Billard W, Binch H 3rd, Cohen-Williams M, Crosby G, Grzelak M, Guzik H, Kozlowski JA, Lowe DB, Pond AJ, Tedesco RP, Watkins RW, Coffin VL. SCH 57790, a selective muscarinic M2 receptor antagonist, releases acetylcholine and produces cognitive enhancement in laboratory animals. *Eur. J. Pharmacol.* 2001; 16:189–200. [PubMed: 11728425]
- Carli M, Lushi R, Sammanin R. Dose related impairments of spatial learning by intrahippocampal scopolamine and antagonism by ondansetron, a 5-HT<sub>3</sub> receptor antagonist. *Behav Brain Res.* 1997; 82:185–194. [PubMed: 9030400]
- Carlsson CM. Lessons learned from failed and discontinued clinical trials for the treatment of Alzheimer' disease: future directions. *J Alzheimers Dis.* 2008; 15:327–338. [PubMed: 18953117]
- Cassel JC, Jeltsch H. Serotonergic modulation of cholinergic function in the central nervous system: cognitive implications. *Neuroscience.* 1995; 69:1–41. [PubMed: 8637608]
- Cavalli A, Bolognesi ML, Capsoni S, Andrisano V, Bartolini M, Margotti E, et al. A small molecule targeting the multifactorial nature of Alzheimer's disease. *Angew Chem Int Ed Engl.* 2007; 46(20): 3689–3692. [PubMed: 17397121]
- Centers for Disease Control and Prevention and The Merck Company Foundation. *The State of Aging and Health in America, 2007.* Whitehouse Station, NJ: The Merck Company Foundation; 2007.

- Chen L, Wang H, Zhang Z, Li Z, He D, Sokabe M, et al. DMXB (GTS-21) ameliorates the cognitive deficits in beta amyloid(25–35(–)) injected mice through preventing the dysfunction of alpha7 nicotinic receptor. *J Neurosci Res*. 2010; 88(8):1784–1794. [PubMed: 20127813]
- Chen Z. Effect of histamine H3-receptor antagonist clobenpropit on spatial memory of radial maze performance in rats. *Acta Pharmacol Sin*. 2000; 21:905–910. [PubMed: 11501041]
- Chen Z, Xu AJ, Li R, Wei EQ. Reversal of scopolamine-induced spatial memory deficits in rats by TAK-147. *Acta Pharmacol Sin*. 2002 Apr; 23(4):355–360. [PubMed: 11931694]
- Clader JW, Wang Y. Muscarinic receptor agonists and antagonists in the treatment of Alzheimer's disease. *Curr Pharmaceutical Des*. 2005; 11:3353–3361.
- Clapham J, Kilpatrick GJ. Thioperamide, the selective histamine H3 receptor antagonist, attenuates stimulant-induced locomotor activity in the mouse. *Eur J Pharmacol*. 1994; 259:107–114. [PubMed: 7957603]
- Cole DC, Lennox WJ, Lombardi S, Ellingboe JW, Bernotas RC, Tawa GJ, et al. Discovery of 5-arylsulfonamido-3(pyrrolidin-2-ylmethyl)-1H-indole derivatives as potent, selective 5-HT6 receptor agonists and antagonists. *J Med Chem*. 2005; 48:353–356. [PubMed: 15658848]
- Comery TA, Martone RL, Aschmies S, Atchison KP, Diamantidis G, Gong X, et al. Acute  $\gamma$ -secretase inhibition improves contextual fear conditioning in the Tg2576 mouse model of Alzheimer's disease. *J Neurosci*. 2005; 25:8898–8902. [PubMed: 16192379]
- Conn PJ, Jones CK, Lindsley CW. Subtype-selective allosteric modulators of muscarinic receptors for the treatment of CNS disorders. *Trends Pharmacol Sci*. 2009; 30:148–155. [PubMed: 19201489]
- Consolo S, Amaboldi S, Giorgi S, Russi G, Ladinsky H. 5-HT4 receptor stimulation facilitates acetylcholine release in rat frontal cortex. *Neuroreport*. 2004; 5:1230–1232. [PubMed: 7919171]
- Coyle JT, Price DL, DeLong MR. Alzheimer's disease: a disorder of cortical cholinergic innervation. *Science*. 1983; 219(4589):1184–1190. [PubMed: 6338589]
- Coyle JT, Geerts H, Sorra K, Amatniek J. Beyond in vitro data: a review of in vivo evidence regarding the allosteric potentiating effect of galantamine on nicotinic acetylcholine receptors in Alzheimer's neuropathology. *J Alzheimers Dis*. 2007; 11:491–507. [PubMed: 17656829]
- Cui YH, Si W, Yin L, An SM, Jin J, Deng SN, et al. A novel derivative of xanomeline improved memory function in aged mice. *Neurosci Bull*. 2008a; 24(4):251–257. [PubMed: 18668154]
- Cui Y, Wang D, Si W, Lv W, Niu Y, Lei X, et al. Enhancement of memory function in aged mice by a novel derivative of xanomeline. *Cell Res*. 2008b; 18:1151–1153. [PubMed: 18936783]
- Davies P, Maloney AJ. Selective loss of central cholinergic neurons in Alzheimer's disease. *Lancet*. 1976; 2(8000):1403. [PubMed: 63862]
- Dawson GR, Bentley G, Draper F, Rycroft W, Iversen SD, Pagella PG. The behavioral effects of heptyl physostigmine, a new cholinesterase inhibitor, in tests of long-term and working memory in rodents. *Pharmacol Biochem Behav*. 1991; 39(4):865–871. [PubMed: 1763105]
- Dawson LA, Nguyen HQ, Li P. In vivo effects of the 5-HT(6) antagonist SB-271046 on striatal and frontal cortex extracellular concentrations of noradrenaline, dopamine, 5-HT, glutamate and aspartate. *Br J Pharmacol*. 2000; 130:23–26.
- Dawson LA, Nguyen HQ, Li P. The 5-HT(6) receptor antagonist SB-271046 selectively enhances excitatory neurotransmission in the rat frontal cortex and hippocampus. *Neuropsychopharmacology*. 2001; 25:662–668. [PubMed: 11682249]
- Day M, Pan JB, Buckley MJ, Cronin E, Hollingsworth PR, Hirst WD, et al. Differential effects of ciproxifan and nicotine on impulsivity and attention measures in the 5-choice serial reaction time test. *Biochem Pharmacol*. 2007; 73:1123–1134. [PubMed: 17214974]
- Decker MW, Bannon AW, Curzon P, Gunther KL, Brioni JD, Holladay MW, et al. ABT-089 [2-methyl-3-(2-(S)-pyrrolidinylmethoxy)pyridine dihydrochloride]: II. A novel cholinergic channel modulator with effects on cognitive performance in rats and monkeys. *J Pharmacol Exp Ther*. 1997; 283:247–258. [PubMed: 9336330]
- De Ferrari GV, Canales MA, Shin I, Weiner LM, Silman I, Inestrosa NC. A structural motif of acetylcholinesterase that promotes amyloid beta-peptide fibril formation. *Biochemistry*. 2001; 40:10447–10457. [PubMed: 11523986]

- de Filippi G, Mogg AJ, Phillips KG, Zwart R, Sher E, Chen Y. The subtype-selective nicotinic acetylcholine receptor positive allosteric modulator 2087101 facilitates neurotransmission in the brain. *Eur J Pharmacol.* 2010; 643(2–3):218–224. [PubMed: 20624387]
- de Los Ríos C, Egea J, Marco-Contelles J, León R, Samadi A, Iriepa I, et al. Synthesis, inhibitory activity of cholinesterases, and neuroprotective profile of novel 1,8-naphthyridine derivatives. *J Med Chem.* 2010; 53(14):5129–5143. [PubMed: 20575555]
- Devan BD, Sierra-Mercado D Jr, Jimenez M, Bowker JL, Duffy KB, Spangler EL, et al. Phosphodiesterase inhibition by sildenafil citrate attenuates the learning impairment induced by blockade of cholinergic muscarinic receptors in rats. *Pharmacol Biochem Behav.* 2004; 79:691–699.
- Devan BD, Bowker JL, Duffy KB, Bharati IS, Jimenez M, Sierra-Mercado D Jr, et al. Phosphodiesterase inhibition by sildenafil citrate attenuates a maze learning impairment in rats induced by nitric oxide synthase inhibition. *Psychopharmacology.* 2006; 183:439–445. [PubMed: 16320087]
- Diez-Ariza M, Garcia-Alloza M, Lasheras B, Del Rio J, Ramirez MJ. GABA(A) receptor antagonists enhance cortical acetylcholine release induced by 5-HT(3) receptor blockade in freely moving rats. *Brain Res.* 2002; 956:81–85. [PubMed: 12426049]
- Doody RS, GavriloVA SI, Sano M, Thomas RG, Aisen PS, Bachurin SO, et al. Effect of dimebon on cognition, activities of daily living, behaviour, and global function in patients with mild-to-moderate Alzheimer's disease: a randomised, double-blind, placebo-controlled study. *Lancet.* 2006; 372(9634):207–215. [PubMed: 18640457]
- Domek-Lopacinska K, Strosznajder JB. The effect of selective inhibitor of cyclic GMP hydrolyzing phosphodiesterase 2 and 5 on learning and memory processes and nitric oxide synthase activity in brain during aging. *Brain Res.* 2008; 1216:68–77. [PubMed: 18499090]
- Domeney AM, Costall B, Gerrard PA, Jones DN, Naylor RJ, Tyers MB. The effect of ondansetron on cognitive performance in the marmoset. *Pharmacol Biochem Behav.* 1991; 38:169–175. [PubMed: 1826787]
- Dysken M, Kuskowski M, Love S. Ondansetron in the treatment of cognitive decline in Alzheimer's dementia. *Am J Geriatr Psychiatry.* 2002; 10:212–215. [PubMed: 11925283]
- Egawa T, Mishima K, Matsumoto Y, Iwasaki K, Fujiwara M. Rolipram and its optical isomers, Phosphodiesterase 4 inhibitors, attenuated the scopolamine-induced impairments of learning and memory in rats. *Jpn J Pharmacol.* 1997; 75:275–281. [PubMed: 9434259]
- Elrod K, Buccafusco JJ, Jackson WJ. Nicotine enhances delayed matching-to-sample performance by primates. *Life Sci.* 1988; 43:277–287. [PubMed: 2899825]
- Eglen RM, Wong EHF, Dumuis A, Bockaert J. Central 5-HT4 receptors. *Trends Pharmacol Sci.* 1995; 16:391–398. [PubMed: 8578609]
- Esbenshade TA, Browman KE, Bitner RS, Strakhova M, Cowart MD, Brioni JD. The histamine H3 receptor: an attractive target for the treatment of cognitive disorders. *Brit J Pharmacol.* 2008; 154:1166–1181. [PubMed: 18469850]
- Fernández-Bachiller MI, Pérez C, González-Muñoz GC, Conde S, López MG, Villarroya M, et al. Novel tacrine-8-hydroxyquinoline hybrids as multifunctional agents for the treatment of Alzheimer's disease, with neuroprotective, cholinergic, antioxidant, and copper-complexing properties. *J Med Chem.* 2010; 53(13):4927–4937. [PubMed: 20545360]
- Fisher A, Karton Y, Heldman E, Gurwitz D, Haring R, Meshulam H, et al. Progress in medicinal chemistry of novel selective muscarinic agonists. *Drug Des Discov.* 1993; 9(3–4):221–235. [PubMed: 8400004]
- Fisher A, Brandeis R, Chapman S, Pittel Z, Michaelson DM. M1 muscarinic agonist treatment reverses cognitive and cholinergic impairments of apolipoprotein E-deficient mice. *J Neurochem.* 1998; 70:1991–1997. [PubMed: 9572284]
- Fisher A, Brandeis R, Haring Nira Bar-Ner R, Kliger-Spatz M, Natan N, Sonogo H, et al. AF150(S) and AF267B: M1 muscarinic agonists as innovative therapies for Alzheimer's disease. *J Mol Neurosci.* 2002; 19(1–2):145–153. [PubMed: 12212772]

- Flynn DD, Ferrari-DiLeo G, Levey AI, Mash DC. Differential alterations in muscarinic receptor subtypes in Alzheimer's disease: implications for cholinergic-based therapies. *Life Sci.* 1995; 56:869–876. [PubMed: 10188787]
- Foley AG, Murphy KJ, Hirst WD, Gallagher HC, Hagan JJ, Upton N, et al. The 5-HT<sub>6</sub> receptor antagonist SB-271046 reverses scopolamine-disrupted consolidation of a passive avoidance task and ameliorates spatial task deficits in aged rats. *Neuropsychopharmacology.* 2004; 29:58–67.
- Fone KCF. An update on the role of the 5-hydroxytryptamine<sub>6</sub> receptor in cognitive function. *Neuropharmacology.* 2008; 55:1015–1022. [PubMed: 18655798]
- Fontana DJ, Daniels SE, Henderson C, Eglen RM, Wong EH. Ondansetron improves cognitive performance in the Morris water maze spatial navigation task. *Psychopharmacology.* 1995; 120:408–417.
- Fontana DJ, Daniels SE, Wong EH, Clark RD, Eglen RM. The effects of novel, selective 5-hydroxytryptamine (5-HT)<sub>4</sub> receptor ligands in rat spatial navigation. *Neuropharmacology.* 1997; 36:689–696. [PubMed: 9225295]
- Fox GB, Pan JB, Radek RJ, Lewis AM, Bitner RS, Esbenshade TA, et al. Two novel and selective nonimidazole H<sub>3</sub> receptor antagonists A-304121 and A-317920: II. In vivo behavioral and neurophysiological characterization. *J Pharmacol Exp Ther.* 2003; 305:897–908. [PubMed: 12606600]
- Fox GB, Esbenshade TA, Pan JB, Radek RJ, Krueger KM, Yao BB, et al. Pharmacological properties of ABT-239 [4-(2-{2-[(2R)-2-methylpyrrolidinyl]ethyl}-benzofuran-5-yl)benzonitril e]: II. Neurophysiological characterization and broad preclinical efficacy in cognition and schizophrenia of a potent and selective histamine H<sub>3</sub> receptor antagonist. *J Pharmacol Exp Ther.* 2005; 313:176–190. [PubMed: 15608077]
- Freedman R, Hall M, Adler LE, Leonard S. Evidence in post-mortem brain tissue for decreased numbers of hippocampal nicotinic receptors in schizophrenia. *Biologic Psychiat.* 1995; 38:22–33.
- Frey U, Huang YY, Kandel ER. Effects of cAMP simulate a late stage pf LTP in hippocampal CA1 neurons. *Science.* 1993; 260:1661–1664. [PubMed: 8389057]
- Frolich L, Eckerwall G, Jonas N. Sirocco-Investigators. A multicenter, double-blind, placebo-controlled phase IIB proof-of-concept dose-ranging study of AZD3480 and donepezil over 12 weeks in patients with mild to moderate Alzheimer's disease. *Alzheimers Dement.* 2009; 5 suppl 1(4):P85.
- Galeotti N, Ghelardini C, Bartolini A. Role of 5HT<sub>4</sub> receptors in mouse passive avoidance test. *J Pharmacol Exp Ther.* 1998; 286:1115–1121. [PubMed: 9732367]
- Galici R, Boggs JD, Aluisio L, Frazer IC, Bonaventure P, Lord B, et al. JNJ-10181457, a selective non-imidazole histamine H<sub>3</sub> receptor antagonist, normalizes acetylcholine neurotransmission and has efficacy in translational rat models of cognition. *Neuropharmacology.* 2009; 56:1131–1137. [PubMed: 19345233]
- Gallagher M, Rapp PR, Fanelli RJ. Opiate antagonist facilitation of time-dependent memory processes: dependence upon intact norepinephrine function. *Brain Res.* 1985; 347:284–290. [PubMed: 3933763]
- Gannon KS, King M, Fone KCF, Shacham S, Fichman M, Melendez R, Cheruku S, Lobera M, Orbach P. PRX-07034, a potent and selective 5-HT<sub>6</sub> antagonist, enhances novel object discrimination (NOD) memory and augments the effect of Aricept. *Soc Neurosci Abstr.* 2006:64.9.
- Gao Y, Tang XC, Guan LC, Kuang PZ. Huperzine A reverses scopolamine- and uscimol-induced memory deficits in chick. *Acta Pharmacol Sin.* 2000; 21:1169–1173.
- Gard PR, Rusted JM. Angiotensin and Alzheimer's disease: therapeutic prospects. *Expert Rev Neurotherapeut.* 2004; 4:87–96.
- Gatto GJ, Bohme GA, Caldwell WS, Letchworth SR, Traina VM, Obinu MC, et al. TC-1734: an orally active neuronal nicotinic acetylcholine receptor modulator with antidepressant, neuroprotective, and long-lasting cognitive effects. *CNS Drug Rev.* 2004; 10(2):147–166. [PubMed: 15179444]
- Ghelardini C, Galeotti N, Gualtieri F, Romanelli MN, Bucherelli C, Baldi E, et al. DM235 (sunifiram): a novel nootropic with potential as a cognitive enhancer. *Naunyn Schmiedebergs Arch Pharmacol.* 2002; 365:419–426. [PubMed: 12070754]

- Giovannini MG, Bartolini L, Bacciottini L, Greco L, Blandina P. Effects of histamine H3 receptor agonists and antagonists on cognitive performance and scopolamine-induced amnesia. *Behav Brain Res.* 1999; 104:147–155. [PubMed: 11125734]
- Gong B, Vitolo OV, Trinchese F, Liu S, Shelanski M, Arancio O. Persistent improvement in synaptic and cognitive functions in an Alzheimer mouse model after rolipram treatment. *J Clin Invest.* 2004; 114:1624–1634. [PubMed: 15578094]
- Grauer SM, Pulito VL, Navarra RL, Kelly MP, Kelley C, Graf R, et al. Phosphodiesterase 10A inhibitor activity in preclinical models of positive, cognitive, and negative symptoms of schizophrenia. *J Pharmacol Exper Ther.* 2009; 331:574–590. [PubMed: 19661377]
- Grigorev VV, Dranyi OA, Bachurin SO. Comparative study of action mechanisms of dimebon and memantine on AMPA- and NMDA-subtypes glutamate receptors in rat cerebral neurons. *Bull Exp Biol Med.* 2003; 136:474–477. [PubMed: 14968164]
- Grigorev VV, Garibova TL, Voronina TA, Litvinova SA, Bachurin SO. Effects of chronic administration of dimebon on behavior and memory of SAMP 10 mice. *Eksp Klin Farmakol.* 2009; 72(1):52–56.
- Grön G, Brandenburg I, Wunderlich AP, Riepe MW. Inhibition of hippocampal function in mild cognitive impairment: targeting the cholinergic hypothesis. *Neurobiol Aging.* 2006; (1):78–87.
- Guan JS, Haggarty SJ, Giacometti E, Dannenberg JH, Joseph N, Gao J, et al. HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature.* 2009 May 7; 459(7243):55–60. [PubMed: 19424149]
- Gwee MC, Cheah LS. Actions of cimetidine and ranitidine at some cholinergic sites: implications in toxicology and anesthesia. *Life Sci.* 1986; 39:383–388. [PubMed: 2874464]
- Hahnen E, Hauke J, Tränkle C, et al. Histone deacetylase inhibitors: possible implications for neurodegenerative disorders. *Expert Opin Investig Drugs.* 2008; 17:169–184.
- Hannon J, Hoyer D. Molecular biology of 5-HT receptors. *Behav Brain Res.* 2008; 195:198–213. [PubMed: 18571247]
- Harder JA, Maclean CJ, Alder JT, Francis PT, Ridley RM. The 5-HT1A antagonist, WAY-100635, ameliorates the cognitive impairment induced by fornix transection in the marmoset. *Psychopharmacology.* 1996; 127:245–254. [PubMed: 8912403]
- Harder JA, Ridley RM. The 5-HT1A antagonist, WAY 100635, alleviates cognitive impairments induced by dizocilpine (MK-801) in monkeys. *Neuropharmacology.* 2000; 39:547–552. [PubMed: 10728875]
- Hasselmo ME. The role of acetylcholine in learning and memory. *Curr Opin Neurobiol.* 2006; 16(6): 710–715. Epub 2006 Sep 29. [PubMed: 17011181]
- Hatcher PD, Brown VJ, Tait DS, Bate S, Overend P, Hagan JJ, Jones DNC. 5-HT6 receptor antagonists improve performance in an attentional set shifting task in rats. *Psychopharmacology.* 2005; 181:253–259. [PubMed: 15846482]
- Heal DJ, Smith SL, Fias A, Codony X, Buschmann H. Selective 5-HT6 receptor ligands: progress in the development of a novel pharmacological approach to the treatment of obesity and related metabolic disorders. *Pharmacol Ther.* 2008; 117:207–231. [PubMed: 18068807]
- Hebb AL, Robertson HA, Denovan-Wright EM. Phosphodiesterase 10A inhibition is associated with locomotor and cognitive deficits and increased anxiety in mice. *Eur Neuropsychopharmacol.* 2008; 18:339–363. [PubMed: 17913473]
- Hebert LE, Scherr PA, Bienias JL, Bennett DA, Evans DA. “Alzheimer’s disease in the U.S. population: Prevalence estimates using the 2000 census.”. *Archives of Neurology.* 2003; 60:1119–1122. [PubMed: 12925369]
- Hille C, Bate S, Davis J, Gonzalez MI. 5-HT4 receptor agonism in the five-choice serial reaction time task. *Behav Brain Res.* 2008; 195:180–186. [PubMed: 18765258]
- Hilt, D.; Gawryl, M.; Koenig, G. EVP-6124\_StudyGroup. EVP-6124: safety, tolerability, and cognitive effects of a novel A7 nicotinic receptor agonist in Alzheimer’s disease patients on stable donepezil or rivastigmine therapy; International Conference on Alzheimer’s disease; Vienna, Austria. July 11–16, 2009;

- Hirst WD, Stean TO, Rogers DC, Sunter D, Pugh P, Moss SF, et al. SB-399885 is a potent, selective 5-HT<sub>6</sub> receptor antagonist with cognitive enhancing properties in aged rat water maze and novel object recognition models. *Eur J Pharmacol.* 2006; 553:109–119. [PubMed: 17069795]
- Hirst WD, Andree TH, Aschmies S, Childers WE, Comery TA, Dawson LA, et al. Correlating efficacy in rodent cognition models with in vivo 5-hydroxytryptamine<sub>1A</sub> receptor occupancy by a novel antagonist, (R)-N-(2-methyl-(4-indolyl-1-piperazinyl)ethyl)-N-(2-pyridinyl)-cyclohexane carboxamide (WAY-101405). *J Pharmacol Exp Ther.* 2008; 325:134–145. [PubMed: 18182558]
- Hodges H, Sowinski P, Sinden JD, Netto CA, Fletcher A. The selective 5HT<sub>3</sub> antagonist, WAY-100289, enhances spatial memory in rats with ibotenate lesions of the forebrain cholinergic projection system. *Psychopharmacology.* 1995; 117:318–332. [PubMed: 7770608]
- Howe WM, Ji J, Parikh V, Williams S, Mocaer E, Trocme-Thibierge C, et al. Enhancement of attentional performance by selective stimulation of  $\alpha 4\beta 2^*$  nAChRs: Underlying cholinergic mechanisms. *Neuropsychopharm.* 2010; 35:1391–1401.
- Huang L, Luo Z, He F, Lu J, Li X. Synthesis and biological evaluation of a new series of berberine derivatives as dual inhibitors of acetylcholinesterase and butyrylcholinesterase. *Bioorg Med Chem.* 2010; 18(12):4475–4484. [PubMed: 20471843]
- Huang M, Li Z, Prus AJ, Dai J, Callahan PM, Meltzer HY. The nicotinic  $\alpha 7$  receptor agonist MEM 3454 increases dopamine and acetylcholine release in rat medial prefrontal cortex and hippocampus alone and in combination with risperidone. *Neuropsychopharmacology.* 2006; 31 Suppl. 1:S185.
- Huang YW, Hu WW, Chen Z, Zhang LS, Shen HQ, Timmerman H, et al. Effect of the histamine H<sub>3</sub>-antagonist clobenpropit on spatial memory deficits induced by MK-801 as evaluated by radial arm maze in Sprague-Dawley rats. *Behav Brain Res.* 2004; 151:287–293. [PubMed: 15084444]
- Huang YY, Kandel ER. Low-frequency stimulation induces a pathway-specific late phase of LTP in the amygdala that is mediated by PKA and dependent on protein synthesis. *Learn Mem.* 2007; 14:497–503. [PubMed: 17626908]
- Huang Z, Dias R, Jones T, Liu S, Styhler A, Claveau D, et al. L-454,560, a potent and selective PDE4 inhibitor with in vivo efficacy in animal models of asthma and cognition. *Biochem Pharmacol.* 2007; 73:1971–1981. [PubMed: 17428447]
- Hung TM, Na M, Dat NT, Ngoc TM, Youn U, Kim HJ, et al. Cholinesterase inhibitory and anti-amnesic activity of alkaloids from *Corydalis turtschaninovii*. *J Ethnopharmacol.* 2008 Sep 2; 119(1):74–80. Epub 2008 Jun 13. [PubMed: 18601993]
- Hulme EC, Lu ZL, Bee MS. Scanning mutagenesis studies of the M1 muscarinic acetylcholine receptor. *Receptors Channels.* 1990; 9:215–228. [PubMed: 12893535]
- Hurst RS, Hajos M, Raggenbass M, Wall TM, Higdon NR, Lawson JA, et al. A novel positive allosteric modulator of the  $\alpha 7$  neuronal nicotinic acetylcholine receptor: in vitro and in vivo characterization. *J Neurosci.* 2005; 25(17):4396–4405. [PubMed: 15858066]
- Ikari H, Spangler EL, Greig NH, Pei XF, Brossi A, Speer D, et al. Maze learning in aged rats is enhanced by phenserine, a novel anticholinesterase. *Neuroreport.* 1995; 6(3):481–484. [PubMed: 7766848]
- Imanishi T, Sawa A, Ichimaru Y, Miyashiro M, Kato S, Yamamoto T, et al. Ameliorating effects of rolipram on experimentally induced impairments of learning and memory in rodents. *Eur J Pharmacol.* 1997; 321:272–278.
- Imhof L, Wallhagen MI, Mahrer-Imhof R, Monsch AU. Becoming forgetful: how elderly people deal with forgetfulness in everyday life. *Am J Alzheimers Dis Other Dement.* 2006 Oct–Nov; 21(5): 347–353. [PubMed: 17062554]
- Impey S, Mark M, Villacres EC, Poser S, Chavkin C, Storm DR. Induction of CRE-mediated gene expression by stimuli that generate long-lasting LTP in area CA1 of the hippocampus. *Neuron.* 1996; 16:973–982. [PubMed: 8630255]
- Janas AM, Cunningham SC, Duffy KB, Devan BD, Greig NH, Holloway HW, et al. The cholinesterase inhibitor, phenserine, improves Morris water maze performance of scopolamine-treated rats. *Life Sci.* 2005; 76(10):1073–1081. [PubMed: 15620572]

- Jones CK, Brady AE, Davis AA, Xiang Z, Bubser M, Tantawy MN, et al. Novel selective allosteric activator of the M1 muscarinic acetylcholine receptor regulates amyloid processing and produces antipsychotic-like activity in rats. *J Neurosci*. 2008; 28:10422–10433. [PubMed: 18842902]
- Jones S, Sudweeks S, Yakel JL. Nicotinic receptors in the brain: correlating physiology with function. *Trends Neurosci*. 1999; 22:555–561. [PubMed: 10542436]
- Karlin A. Emerging structure of the nicotinic acetylcholine receptors. *Nat Rev Neurosci*. 2002; 3:102–114. [PubMed: 11836518]
- Kem WR. The brain alpha7 nicotinic receptor may be an important therapeutic target for the treatment of Alzheimer's disease: studies with DMXBA (GTS-21). *Behav Brain Res*. 2000; 113:169–181. [PubMed: 10942043]
- Kendall I, Slotten HA, Codony X, Burgueno J, Pauwels PJ, Vela JM, et al. E-6801, a 5-HT6 receptor agonist, improves recognition memory by combined modulation of cholinergic and glutamatergic neurotransmission in the rat. *Psychopharmacology*. 2010 (Epub ahead of print).
- Kennedy D. Just treat, or enhance? *Science*. 2004; 304:317.
- Kikuchi T, Okamura T, Fukushi K, Irie T. Piperidine-4-methanthal ester derivatives for a selective acetylcholinesterase assay. *Biol Pharm Bull*. 2010; 33(4):702–706. [PubMed: 20410609]
- King MV, Sleight AJ, Woolley ML, Topham IA, Marsden CA, Fone KCF. 5-HT6 receptor antagonists reverse delay-dependent deficits in novel object discrimination by enhancing consolidation—an effect sensitive to NMDA receptor antagonism. *Neuropharmacology*. 2004; 47:195–204. [PubMed: 15223298]
- King MV, Sleight AJ, Fone KCF, Marsden CA. The selective 5-HT6 receptor antagonist Ro4-6790 reverses age-related and chronic intermittent PCP-induced cognitive dysfunction in rats. *J Psychopharmacol*. 2005; 19:A12.
- King MV, Marsden CA, Fone KCF. A role for the 5-HT1A, 5-HT4 and 5-HT6 receptors in learning and memory. *Trends Pharmacol Sci*. 2008; 29:482–492. [PubMed: 19086256]
- King MV, Spicer CH, Sleight AJ, Marsden CA, Fone KCF. Impact of regional 5-HT depletion on the cognitive enhancing effects of a typical 5-HT6 receptor antagonist, Ro 04-6790, in the novel object discrimination task. *Psychopharmacology*. 2009; 202:111–123. [PubMed: 18839151]
- Kinsella, K.; He, W. Washington, D.C.: U.S. Government Printing Office; 2009. *An Aging World*. U.S. Census Bureau, International Population Reports, P95/09–1.
- Knapp MJ, Knopman DS, Solomon PR, Pendlebury WW, Davis CS, Gracon SI. A 30-week randomized controlled trial of high-dose tacrine in patients with Alzheimer's disease. The Tacrine Study Group. *JAMA*. 1994; 271(13):985–991. [PubMed: 8139083]
- Kohen R, Metcalf MA, Khan N, Druck T, Huebner K, Lachowicz JE, et al. Cloning, characterization, and chromosomal localization of a human 5-HT6 serotonin receptor. *J Neurochem*. 1996; 66:47–56. [PubMed: 8522988]
- Komater VA, Buckley MJ, Browman KE, Pan JB, Hancock AA, Decker MW, et al. Effects of histamine H3 receptor antagonists in two models of spatial learning. *Behav Brain Res*. 2005; 159:295–300. [PubMed: 15817192]
- Kulla A, Manahan-Vaughan D. Modulation by serotonin 5-HT(4) receptors of long-term potentiation and depotentiation in the dentate gyrus of freely moving rats. *Cereb Cortex*. 2002; 12:150–162. [PubMed: 11739263]
- Kumar YC, Malviya M, Chandra JN, Sadashiva CT, Kumar CS, Prasad SB, et al. Effect of novel N-aryl sulfonamide substituted 3-morpholino arecoline derivatives as muscarinic receptor 1 agonists in Alzheimer's dementia models. *Bioorg Med Chem*. 2008; 16(9):5157–5163. [PubMed: 18359231]
- Kwon SH, Kim HC, Lee SY, Jang CG. Loganin improves learning and memory impairments induced by scopolamine in mice. *Eur J Pharmacol*. 2009; 619(1–3):44–49. [PubMed: 19666019]
- Lachowicz JE, Duffy RA, Ruperto V, Kozlowski J, Zhou G, Clader J, Billard W, Binch H 3rd, Crosby G, Cohen-Williams M, Strader CD, Coffin V. Facilitation of acetylcholine release and improvement in cognition by a selective M2 muscarinic antagonist, SCH 72788. *Life Sci*. 2001; 68:2585–2592. [PubMed: 11392630]

- Lacroix LP, Dawson LA, Hagan JJ, Heidebreder CA. 5-HT<sub>6</sub> receptor antagonist SB-271046 enhances extracellular levels of monoamines in the rat medial prefrontal cortex. *Synapse*. 2004; 51:158–164. [PubMed: 14618683]
- Lamirault L, Simon H. Enhancement of place and object recognition memory in young adult and old rats by RS 67333, a partial agonist of 5-HT<sub>4</sub> receptors. *Neuropharmacology*. 2001; 41:844–853. [PubMed: 11684148]
- Lanni C, Lenzken SC, Pascale A, Del Vecchio I, Racchi M, Pistoia F, et al. Cognition enhancers between treating and doping the mind. *Pharmacol Res*. 2008 Mar; 57(3):196–213. [PubMed: 18353672]
- Lee KY, Sung SH, Kim SH, Jang YP, Oh TH, Kim YC. Cognitive-enhancing activity of loganin isolated from *Cornus officinalis* in scopolamine-induced amnesic mice. *Arch Pharm Res*. 2009; 32(5):677–683. [PubMed: 19471881]
- Lee I, Choe YS, Ryu EK, Choi BW, Choi JY, Choi Y, et al. Synthesis and evaluation of radioiodine-labelled CP-118,954 for the in-vivo imaging of acetylcholinesterase. *Nucl Med Commun*. 2007; 28(7):561–566. [PubMed: 17538398]
- Leiser SC, Bowlby MR, Comery TA, Dunlop J. A cog in cognition: how the  $\alpha 7$  nicotinic acetylcholine receptor is geared towards improving cognitive deficits. *Pharmacol & Ther*. 2009; 122:302–311. [PubMed: 19351547]
- Lermontova NN, Redkozubov AE, Shevtsova EF, Serkova TP, Kireeva EG, Bachurin SO. Dimebon and tacrine inhibit neurotoxic action of beta-amyloid in culture and block L-type Ca(2+) channels. *Bull Exp Biol Med*. 2001; 132(5):1079–1083. [PubMed: 11865327]
- Leurs R, Bakker RA, Timmerman H, de Esch IJP. The histamine H<sub>3</sub> receptor: from gene cloning to H<sub>3</sub> receptor drugs. *Nature Rev Drug Discov*. 2005; 4:107–120. [PubMed: 15665857]
- Levallet G, Hotte M, Boulouard M, Dauphin F. Increased particulate Phosphodiesterase 4 in the prefrontal cortex supports 5-HT<sub>4</sub> receptor-induced improvement of object recognition memory in the rat. *Psychopharmacology*. 2009; 202:125–139. [PubMed: 18712363]
- Levey AI, Kitt CA, Simonds WF, Price DL, Brann MR. Identification and localization of muscarinic acetylcholine receptor proteins in brain with subtype-specific antibodies. *J Neurosci*. 1991; 11:3218–3226. [PubMed: 1941081]
- Levin ED. Nicotinic receptor subtypes and cognitive function. *J Neurobiol*. 2002; 53:633–640. [PubMed: 12436426]
- Levin ED, Rezvani AH. Nicotinic interactions with antipsychotic drugs, models of schizophrenia and impacts on cognitive function. *Biochem Pharmacol*. 2007; 74:1182–1191. [PubMed: 17714691]
- Lezoualc'h F. 5-HT<sub>4</sub> receptor and Alzheimer's disease: the amyloid connection. *Exper Neurol*. 2007; 205:325–329. [PubMed: 17346704]
- Li J, Huang H, Miezian Ezoulin JM, Gao XL, Massicot F, et al. Pharmacological profile of PMS777, a new AChE inhibitor with PAF antagonistic activity. *Int J Neuropsychopharmacol*. 2006; (1):21–29. Epub 2006 Jan 23. [PubMed: 16426477]
- Li NC, Lee A, Whitmer RA, Kivipelto M, Lawler E, Kazis LE, et al. Use of angiotensin receptor blockers and risk of dementia in a predominantly male population: prospective cohort analysis. *BMJ*. 2010 Jan 12.340:b5465. [PubMed: 20068258]
- Li Z, Huang M, Prus AJ, Dai J, Meltzer HY. 5-HT<sub>6</sub> receptor antagonist SB-399885 potentiates haloperidol and risperidone-induced dopamine efflux in the medial prefrontal cortex or hippocampus. *Brain Res*. 2007; 1134:70–78. [PubMed: 17207474]
- Ligneau X, Lin J-S, Vanni-Mercier G, Jouvet M, Muir JL, Ganellin CR, et al. Neurochemical and behavioral effects of ciproxifan, a potent histamine H<sub>3</sub> receptor antagonist. *J Pharmacol Exp Ther*. 1998; 287:658–666. [PubMed: 9808693]
- Ligneau X, Perrin D, Landais L, Camelin J-C, Calmels TPG, Berrebi-Bertrand I, et al. BF2.649 [1-{3-[3-(4-chlorophenyl)propoxy]propyl}piperidine, hydrochloride], a nonimidazole inverse agonist/antagonist at the human histamine H<sub>3</sub> receptor: preclinical pharmacology. *J Pharmacol Exp Ther*. 2007; 320:365–375. [PubMed: 17005916]
- Loiseau F, Dekeyne A, Millan MJ. Pro-cognitive effects of 5-HT<sub>6</sub> receptor antagonists in the social recognition procedure in rats: implication of the frontal cortex. *Psychopharmacology*. 2008; 196:93–104. [PubMed: 17922111]



- Lopez-Hernandez G, Placzek AN, Thinschmidt JS, Lestage P, Trocme-Thibierge C, Morain P, et al. Partial agonist and neuromodulatory activity of S 24795 for alpha7 nAChR responses of hippocampal interneurons. *Neuropharmacology*. 2007; 53(1):134–144. [PubMed: 17544457]
- Lowenstein PR, Castro MG. Uncertainty in the translation of preclinical experiments to clinical trials. Why do most phase III clinical trials fail? *Curr Gene Ther*. 2008; 9:368–374. [PubMed: 19860651]
- Lu YF, Hawkins RD. Ryanodine receptors contribute to cGMP-induced late-phase LTP and CREB phosphorylation in the hippocampus. *J Neurophysiol*. 2002; 88:1270–1278. [PubMed: 12205148]
- Luck T, Luppia M, Briel S, Riedel-Heller SG. Incidence of mild cognitive impairment: a systematic review. *Dement Geriatr Cogn Disord*. 2010; 29(2):164–175. [PubMed: 20150735]
- Luques L, Shoham S, Weinstock M. Chronic brain cytochrome oxidase inhibition selectively alters hippocampal cholinergic innervation and impairs memory: prevention by ladostigil. *Exp Neurol*. 2007; 206(2):209–219. Epub 2007 Apr 27. [PubMed: 17580085]
- Luttgen M, Elvander E, Madjid N, Ogren SO. Analysis of the role of 5-HT1A receptors in spatial and aversive learning in the rat. *Neuropharmacology*. 2005; 48:830–852. [PubMed: 15829255]
- Lynch G, Palmer LC, Gall CM. The likelihood of cognitive enhancement. *Pharmacol Biochem Behavior*. 2011; Jan(6) [Epub ahead of print].
- Maher-Edwards G, Zvartau-Hind M, Hunter AJ, Gold M, Hopton G, Jacobs G, et al. Double-blind, controlled phase II study of a 5-HT6 receptor antagonist, SB-742457, in Alzheimer's disease. *Curr Alzheimer Res*. 2010; 7:374–385. [PubMed: 20043816]
- Malviya M, Kumar YC, Asha D, Chandra JN, Subhash MN, Rangappa KS. Muscarinic receptor 1 agonist activity of novel N-arylthioureas substituted 3-morpholino arecoline derivatives in Alzheimer's presenile dementia models. *Bioorg Med Chem*. 2008; 16(15):7095–7101. [PubMed: 18640043]
- Malviya M, Sunil Kumar YC, Mythri RB, Venkateshappa C, Subhash MN, Rangappa KS. Muscarinic receptor 1 agonist activity of novel N-aryl carboxamide substituted 3-morpholino arecoline derivatives in Alzheimer's presenile dementia models. *Bioorg Med Chem*. 2009; 17:5526–5534. [PubMed: 19595599]
- Marchetti E, Chaillan FA, Dumuis A, Bockaert J, Soumireu-Mourat B, Roman FS. Modulation of memory processes and cellular excitability in the dentate gyrus of freely moving rats by a 5-HT4 receptor partial agonist, and an antagonist. *Neuropharmacology*. 2004; 47:1021–1035. [PubMed: 15555636]
- Marcos B, Gil-Bea FJ, Hirst WD, Garcia-Alloza M, Ramirez MJ. Lack of localization of 5-HT6 receptors on cholinergic neurons: implication of multiple neurotransmitter systems in 5-HT6 receptor-mediated acetylcholine release. *Eur J Neurosci*. 2006; 24:1299–1306. [PubMed: 16987217]
- Marcos B, Chuang TT, Gil-Bea FJ, Ramirez MJ. Effects of 5-HT6 receptor antagonism and cholinesterase inhibition in models of cognitive impairment in the rat. *Brit J Pharmacol*. 2008; 155:434–440. [PubMed: 18622410]
- Martinez-Mir MI, Pollard H, Moreay J, Arrang JM, Ruat M, Traiffort E, et al. Three histamine receptors (H1, H2 and H3) visualized in the brain of human and non-human primates. *Brain Res*. 1990; 526:322–327. [PubMed: 1979518]
- Maruyama W, Akao Y, Youdim MB, Davis BA, Naoi M. Transfection-enforced Bcl-2 overexpression and an anti-Parkinson drug, rasagiline, prevent nuclear accumulation of glyceraldehyde-3-phosphate dehydrogenase induced by an endogenous dopaminergic neurotoxin, N-methyl(R)salsolinol. *Journal of Neurochemistry*. 2001; 78:727–735. [PubMed: 11520893]
- Mash DC, Flynn DD, Potter LT. Loss of M2 muscarine receptors in the cerebral cortex in Alzheimer's disease and experimental cholinergic denervation. *Science*. 1985; 228:1115–1117. [PubMed: 3992249]
- Matsumoto M, Togashi H, Mori K, Ueno KI, Ohashi S, Kojima T, et al. Evidence for involvement of central 5-HT4 receptors in cholinergic function associated with cognitive processes: behavioral electrophysiological, and neurochemical studies. *J Pharmacol Exp Ther*. 2001; 296:676–682. [PubMed: 11181892]

- Mattsson C, Sonesson C, Sandahl A, Greiner HE, Gassen M, Plaschke J, et al. 2-Alkyl-3-(1,2,3,6-tetrahydropyridin-4-yl)-1H-indoles as novel 5-HT<sub>6</sub> receptor agonists. *Bioorg Med Chem Lett*. 2005; 15:4230–4234. [PubMed: 16055331]
- McArthur RA, Gray J, Schreiber R. Cognitive effects of muscarinic M1 functional agonists in non-human primates and clinical trials. *Curr Opin Investig Drugs*. 2010; 11:740–760.
- McGehee DS, Role LW. Physiological diversity of nicotinic acetylcholine receptors expressed by vertebrate neurons. *Annu Rev Physiol*. 1995; 57:521–546. [PubMed: 7778876]
- Medhurst AD, Atkins AR, Beresford IJ, Brackenborough K, Briggs MA, Calver AR, et al. GSK189254, a novel H<sub>3</sub> receptor antagonist that binds to histamine H<sub>3</sub> receptors in Alzheimer's disease brain and improves cognitive performance in preclinical models. *J Pharmacol Exp Ther*. 2007a; 321:1032–1045. [PubMed: 17327487]
- Medhurst AD, Briggs MA, Bruton G, Calver AR, Chessell I, Crook B, et al. Structurally novel histamine H<sub>3</sub> receptor antagonists GSK207040 and GSK334429 improve scopolamine-induced memory impairment and capsaicin-induced secondary allodynia in rats. *Biochem Pharmacol*. 2007b; 73:1182–1194. [PubMed: 17276409]
- Meguro K, Yanai K, Sakai N, Sakurai E, Maeyama K, Sasaki H, et al. Effects of thioperamide, a histamine H<sub>3</sub> antagonist, on the step-through passive avoidance response and histidine decarboxylase activity in senescence-accelerated mice. *Pharmacol Biochem Behav*. 1995; 50:321–325. [PubMed: 7617668]
- Meneses A. 5-HT system and cognition. *Neurosci Biobehav Rev*. 1999; 23:1111–1125. [PubMed: 10643820]
- Meneses A. Role of 5-HT<sub>6</sub> receptors in memory formation. *Drug News Perspect*. 2001; 4:396–400. [PubMed: 12813582]
- Meneses A, Perez-Garcia G. 5-HT<sub>1A</sub> receptors and memory. *Neurosci Biobehav Rev*. 2007; 31:705–727. [PubMed: 17418894]
- Meneses A, Perez-Garcia G, Lly-Salmeron G, Flores-Galvez D, Castillo C, Castillo E. The effects of the 5-HT<sub>6</sub> receptor agonist EMD and the 5-HT<sub>7</sub> receptor agonist AS19 on memory formation. *Behav Brain Res*. 2008; 195:112–119. [PubMed: 18191236]
- Millan MJ, Gobert A, Roux S, Porsolt R, Meneses A, Carli M, et al. The serotonin<sub>1A</sub> receptor partial agonist S15535 [4-(benzodioxan-5-yl)-1-(indan-2-yl)piperazine] enhances cholinergic transmission and cognitive function in rodents: a combined neurochemical and behavioral analysis. *J Pharmacol Exp Ther*. 2004; 311:190–203. [PubMed: 15146031]
- Miller G. Pharmacology. The puzzling rise and fall of a dark-horse Alzheimer's drug. *Science*. 2010; 327(5971):1309. [PubMed: 20223954]
- Mintzer JE, Kershaw P. The efficacy of galantamine in the treatment of Alzheimer's disease: comparison of patients previously treated with acetylcholinesterase inhibitors to patients with no prior exposure. *Int J Geriatr Psychiatry*. 2003; 18:292–297. [PubMed: 12673604]
- Minzenberg MJ, Carter CS. Modafinil: a review of neurochemical actions and effects on cognition. *Neuropsychopharmacology*. 2008; 33:1477–1502. [PubMed: 17712350]
- Misane I, Ogren SO. Selective 5-HT<sub>1A</sub> antagonists WAY 100635 and NAD-299 attenuate the impairment of passive avoidance caused by scopolamine in the rat. *Neuropsychopharmacology*. 2003; 28:253–264. [PubMed: 12589378]
- Mitchell ES, Hoplight BJ, Lear SP, Neumaier JF. BGC20-761, a novel tryptamine analog, enhances memory consolidation and reverses scopolamine-induced memory deficit in social and visuospatial memory tasks through a 5-HT<sub>6</sub> receptor-mediated mechanism. *Neuropharmacology*. 2006; 50:412–420. [PubMed: 16298400]
- Miyamoto M, Takahashi H, Kato K, Hirai K, Ishihara Y, Goto G. Effects of 3-[1-(phenylmethyl)-4-piperidinyl]-1-(2,3,4,5-tetrahydro-1-H-1-benzazepin-8-yl)-1-propanone fumarate (TAK-147), a novel acetylcholinesterase inhibitor, on impaired learning and memory in animal models. *J Pharmacol Exp Ther*. 1996; 277(3):1292–1304. [PubMed: 8667190]
- Miyazaki S, Onodera K, Imaizumi M, Timmerman H. Effects of clobenpropit (VUF-9153), a histamine H<sub>3</sub>-receptor antagonist, on learning and memory, and on cholinergic and monoaminergic systems in mice. *Life Sci*. 1997; 61:355–361. [PubMed: 9244361]

- Mohamed T, Rao PP. Design, synthesis and evaluation of 2,4-disubstituted pyrimidines as cholinesterase inhibitors. *Bioorg Med Chem Lett*. 2010; 20(12):3606–3609. [PubMed: 20472431]
- Mohler EG, Shacham S, Noiman S, Lezoualc'h F, Robert S, Gastineau M, et al. VRX-03011, a novel 5-HT<sub>4</sub> agonist, enhances memory and hippocampal acetylcholine efflux. *Neuropharmacology*. 2007; 53:563–573. [PubMed: 17692343]
- Monsma FJ, Shen Y, Ward RP, Hamblin MW, Sibley DR. Cloning and expression of a novel serotonin receptor with high-affinity for tricyclic psychotropic-drugs. *Mol Pharmacol*. 1993; 43:320–327. [PubMed: 7680751]
- Monti B, Berteotti C, Contestabile A. Subchronic rolipram delivery activates hippocampal CREB and Arc, enhances retention and slows down extinction of conditioned fear. *Neuropsychopharmacology*. 2006; 31:278–286.
- Moser PC, Bergis OE, Jegham S, Lochead A, Duconseille E, Terranova JP, et al. SL65.0155, a novel 5-hydroxytryptamine<sub>4</sub> receptor partial agonist with potent cognition-enhancing properties. *J Pharmacol Exp Ther*. 2002; 302:731–741. [PubMed: 12130738]
- Murray KN, Abeles N. Nicotine's effect on neural and cognitive functioning in an aging population. *Aging & Mental Health*. 2002; 6:129–138. [PubMed: 12028881]
- Naghdi N, Harooni HE. The effect of intrahippocampal injections of ritanserin (5-HT<sub>2A/2C</sub> antagonist) and granisetron (5-HT<sub>3</sub> antagonist) on learning as assessed in the spatial version of the water maze. *Behav Brain Res*. 2005; 157:205–210. [PubMed: 15639171]
- Navakkode S, Sajikumar S, Frey JU. The type IV-specific phosphodiesterase inhibitor rolipram and its effect on hippocampal long-term potentiation and synaptic tagging. *J Neurosci*. 2004; 24:7740–7744. [PubMed: 15342741]
- Ng HJ, Whitemore ER, Tran MB, Hogenkamp DJ, Broide RS, Johnstone TB, et al. Nootropic alpha7 nicotinic receptor allosteric modulator derived from GABAA receptor modulators. *Proc Natl Acad Sci*. 2007; 104(19):8059–8064. [PubMed: 17470817]
- Ohara T, Fukaya H, Itanaka K, Seno N. Ameliorating effects of SDZ ENA 713 on age-associated decreases in learning performance and brain choline acetyltransferase activity in rats. *Brain Res Bull*. 1997; 43:39–42. [PubMed: 9205792]
- Ohman A, Josephsson S, Nygård L. Awareness through interaction in everyday occupations: experiences of people with Alzheimer's disease. *Scand J Occup Ther*. 2008 Mar; 15(1):43–51. [PubMed: 17852957]
- Ogren SO, Eriksson TM, Elvander-Tottie E, D'Addario C, Ekstrom JC, Svenningsson P, et al. The role of 5-HT<sub>1A</sub> receptors in learning and memory. *Behav Brain Res*. 2008; 195:54–77. [PubMed: 18394726]
- Ou LY, Tang XC, Cai JX. Effect of huperzine A on working memory in reserpine- or yohimbine-treated monkeys. *Eur J Pharmacol*. 2001; 433:151–156. [PubMed: 11755146]
- Parkinson study group. Randomized placebo-controlled study of the nicotinic agonist SIB-1508Y in Parkinson disease. *Neurology*. 2006; 66(3):408–410. [PubMed: 16476941]
- Pascoli V, Boer-Saccomani C, Hermant J-F. H<sub>3</sub> receptor antagonists reverse delay-dependent deficits in novel object discrimination by enhancing retrieval. *Psychopharmacology*. 2009; 202:141–152. [PubMed: 18493749]
- Patat A, Parks V, Raje S, Plotka A, Chassard D, Coz FL. Safety, tolerability, pharmacokinetics and pharmacodynamics of ascending single and multiple doses of lecozotan in healthy young and elderly subjects. *Brit J Clin Pharmacol*. 2008; 67:299–308. [PubMed: 19523013]
- Patel N, Spangler EL, Greig NH, Yu QS, Ingram DK, Meyer RC. Phenserine, a novel acetylcholinesterase inhibitor, attenuates impaired learning of rats in a 14-unit T-maze induced by blockade of the N-methyl-D-aspartate receptor. *Neuroreport*. 1998; 9(1):171–176. [PubMed: 9592071]
- Patil CS, Singh VP, Singh S, Kulkarni SK. Modulatory effect of the PDE-5 inhibitor sildenafil in diabetic neuropathy. *Pharmacology*. 2004; 72:192–195.
- Prendergast MA, Jackson WJ, Terry AV Jr, Decker MW, Arneric SP, Buccafusco JJ. Central nicotinic receptor agonists ABT-418, ABT-089, and (–)-nicotine reduce distractibility in adult monkeys. *Psychopharmacol (Berl)*. 1998; 136(1):50–58.

- Peng Y, Lee DY, Jiang L, Ma Z, Schachter SC, Lemere CA. Huperzine A regulates amyloid precursor protein processing via protein kinase C and mitogenactivated protein kinase pathways in neuroblastoma SK-N-SH cells over-expressing wild type human amyloid precursor protein 695. *Neuroscience*. 2007; 150:386–395. [PubMed: 17945434]
- Perez-Garcia G, Meneses A. Oral administration of the 5-HT<sub>6</sub> receptor antagonists SB-357134 and SB-399885 improves memory formation in an autoshaping learning task. *Pharmacol Biochem Behav*. 2005; 81:673–682. [PubMed: 15964617]
- Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med*. 2004 Sep; 256(3):183–194. [PubMed: 15324362]
- Pillot C, Ortiz J, Heron A, Ridray S, Schwartz JC, Arrang JM. Ciproxifan, a histamine H<sub>3</sub>-receptor antagonist/inverse agonist, potentiates neurochemical and behavioral effects of haloperidol in the rat. *J Neurosci*. 2002; 22:7272–7280. [PubMed: 12177222]
- Pisani L, Catto M, Giangreco I, Leonetti F, Nicolotti O, Stefanachi A, et al. Design, Synthesis, and Biological Evaluation of Coumarin Derivatives Tethered to an Edrophonium-like Fragment as Highly Potent and Selective Dual Binding Site Acetylcholinesterase Inhibitors. *ChemMedChem*. 2010 Aug 2. [Epub ahead of print].
- Pitsikas N, Bramibilla A, Borsini F. DAU 6215, a novel 5-HT<sub>3</sub> receptor antagonist, improves performance in the aged rat in the Morris water maze task. *Neurobiol Aging*. 1993; 14:561–564. [PubMed: 8295658]
- Pitsikas N, Borsini F. Itasetron (DAU 6215) prevents age-related memory deficits in the rat in a multiple choice avoidance task. *Eur J Pharmacol*. 1996; 311:115–119. [PubMed: 8891590]
- Pitsikas N, Borsini F. Different effects of tropisetron and ondansetron in learning and memory paradigms. *Pharmacol Biochem Behav*. 1997; 56:571–576. [PubMed: 9130279]
- Pitsikas N, Rigamonti AE, Cella SG, Muller EE. The 5-HT<sub>1A</sub> receptor antagonist WAY 100635 improves rats performance in different models of amnesia evaluated by the object recognition task. *Brain Res*. 2003; 983:215–222. [PubMed: 12914983]
- Pollard H, Moreau J, Arrang JM, Schwartz JC. A detailed autoradiographic mapping of histamine H<sub>3</sub> receptors in rat brain areas. *Neuroscience*. 1993; 52:169–189. [PubMed: 8381924]
- Prendergast MA, Jackson WJ, Terry AV Jr, Decker MW, Arneric SP, Buccafusco JJ. Central nicotinic receptor agonists ABT-418, ABT-089, and (-)-nicotine reduce distractibility in adult monkeys. *Psychopharmacol (Berl)*. 1998; 136(1):50–58.
- Prickaerts J, Steinbusch HW, Smits JF, de Vente J. Possible role of nitric oxide-cyclic GMP pathway in object recognition memory: effects of 7-nitroindazole and zaprinast. *Eur J Pharmacol*. 1997; 337:125–136. [PubMed: 9430406]
- Prickaerts J, van Staveren WC, Sik A, Markerink-van Ittersum M, Niewohner U, van der Staay FJ, et al. Effects of two selective Phosphodiesterase type 5 inhibitors, sildenafil and vardenafil, on object recognition memory and hippocampal cyclic GMP levels in the rat. *Neuroscience*. 2002; 113:351–361. [PubMed: 12127092]
- Prickaerts J, Sik A, van der Staay FJ, de Vente J, Blokland A. Dissociable effects of acetylcholinesterase inhibitors and phosphodiesterase type 5 inhibitors on object recognition memory: acquisition versus consolidation. *Psychopharmacology*. 2005; 177:381–390. [PubMed: 15630588]
- Qiu C, Kivipelto M, von Strauss E. Epidemiology of Alzheimer's disease: occurrence, determinants, and strategies toward intervention. *Dialogues Clin Neurosci*. 2009; 11(2):111–128. [PubMed: 19585947]
- Quirion R. Cholinergic markers in Alzheimer disease and the autoregulation of acetylcholine release. *J Psychiatr Neurosci*. 1993; 18:226–234.
- Quirion R, Wilson A, Rowe W, Aubert I, Richard J, Doods H, et al. Facilitation of acetylcholine release and cognitive performance by an M<sub>2</sub>-muscarinic receptor antagonist in aged memory-impaired rats. *J Neurosci*. 1995; 15:1455–1462. [PubMed: 7869110]
- Racchi M, Mazzucchelli M, Lenzken SC, Porrello E, Lanni C, Govoni S. Role of acetylcholinesterase inhibitors in the regulation of amyloid beta precursor protein (AβetaPP) metabolism. *Chem. Biol. Interact*. 2005; 157–158. 335–338.

- Ramirez MJ, Cenarruzabeitia E, Lasheras B, Del Rio J. Involvement of GABA systems in acetylcholine release induced by 5-HT<sub>3</sub> receptor blockade in slices from rat entorhinal cortex. *Brain Res.* 1996; 712:274–280. [PubMed: 8814902]
- Ramos BP, Birnbaum SG, Lindenmeyer I, Newton SS, Duman RS, Arnsten AF. Dysregulation of protein kinase A signaling in the aged prefrontal cortex: new strategy for treating age-related cognitive decline. *Neuron.* 2003; 40:835–845. [PubMed: 14622586]
- Randt CT, Judge ME, Bonnet KA, Quartermain D. Brain cyclic AMP and memory in mice. *Pharmacol Biochem Behav.* 1982; 17:677–680. [PubMed: 6294676]
- Raskind MA, Peskind ER, Wessel T, Yuan W. Galantamine in AD: A 6-month randomized, placebo-controlled trial with a 6-month extension. The Galantamine USA-1 Study Group. *Neurology.* 2000; 54(12):2261–2268. [PubMed: 10881250]
- Reisine TD, Yamamura HI, Bird ED, Spokes E, Enna SJ. Pre- and postsynaptic neurochemical alterations in Alzheimer's disease. *Brain Res.* 1978 Dec 29; 159(2):477–481. [PubMed: 215272]
- Reneerkens OAH, Rutten K, Steinbusch HWM, Blokland A, Prickaerts J. Selective Phosphodiesterase inhibitors: a promising target for cognition enhancement. *Psychopharmacology.* 2009; 202:419–443. [PubMed: 18709359]
- Rezvani AH, Kholdebarin E, Brucato FH, Callahan PM, Lowe DA, Levin ED. Effect of R3487/MEM3454, a novel nicotinic  $\alpha 7$  receptor partial agonist and 5-HT<sub>3</sub> antagonist on sustained attention in rats. *Prog Neuropsychopharmacol Biol Psychiatry.* 2009; 33(2):269–275. [PubMed: 19110025]
- Riemer C, Borroni E, Levet-Trafit B, Martin JR, Poli S, Porter RHP, et al. Influence of the 5-HT<sub>6</sub> receptor on acetylcholine release in the cortex: pharmacological characterization of 4-(2-bromo-6-pyrrolidin-1-ylpyridine-4-sulfonyl)phenylamine, a potent, selective 5-HT<sub>6</sub> receptor antagonist. *J Med Chem.* 2003; 46:1273–1276. [PubMed: 12646038]
- Rizzo S, Bartolini M, Ceccarini L, Piazzini L, Gobbi S, Cavalli A, et al. Targeting Alzheimer's disease: Novel indanone hybrids bearing a pharmacophoric fragment of AP2238. *Bioorg Med Chem.* 2010 Mar 1; 18(5):1749–1760. Epub 2010 Feb 4. [PubMed: 20171894]
- Robert SJ, Zugaza JL, Fischmeister R, Gardier AM, Lezoualc'h F. The human serotonin 5-HT<sub>4</sub> receptor regulates secretion of non-amyloidogenic precursor protein. *J Biol Chem.* 2001; 276:44881–44888. [PubMed: 11584021]
- Roberts JC, Reavill C, East SZ, Harrison PJ, Patel S, Routledge C, et al. The distribution of 5-HT<sub>6</sub> receptors in rat brain: an autoradiographic binding study using the radiolabelled 5-HT<sub>6</sub> receptor antagonist I-125 SB-258585. *Brain Res.* 2002; 934:49–57. [PubMed: 11937069]
- Rodefer JS, Murphy ER, Baxter MG. PDE10A inhibition reverses subchronic PCP-induced deficits in attentional set-shifting in rats. *Eur J Neurosci.* 2005; 21:1070–1076. [PubMed: 15787711]
- Rodefer JS, Baseile JL, Callahan PM, Van Natta K, Yeo H, Lowe DA, Santarelli L, Vivian JA. Attentional set-shifting in rats and monkeys: Effects of cholinergic challenges. *Soc Neurosci Abstr.* 2007 746.4.
- Rodefer JS, Nguyen TN, Karlsson JJ, Arnt J. Reversal of subchronic PCP-induced deficits in attentional set shifting in rats by sertindole and a 5-HT<sub>6</sub> receptor antagonist: comparison among antipsychotics. *Neuropsychopharmacology.* 2008; 33:2657–2666. [PubMed: 18094666]
- Rogers DC, Hagan JJ. 5-HT<sub>6</sub> receptor antagonists enhance retention of a water-maze task in the rat. *Psychopharmacology.* 2001; 158:114–119. [PubMed: 11702084]
- Rollema H, Hajos M, Seymour PA, Kozak R, Majchrzak MJ, Guanowsky V, et al. Preclinical pharmacology of the  $\alpha 4\beta 2$  nAChR partial agonist varenicline related to effects on reward, mood and cognition. *Biochem Pharmacol.* 2009; 78:813–824. [PubMed: 19501054]
- Romero G, Sanchez E, Pujol M, Perez P, Codony X, Holenz J, et al. Efficacy of selective 5-HT<sub>6</sub> receptor ligands determined by monitoring 5-HT<sub>6</sub> receptor-mediated cAMP signaling pathways. *Brit J Pharmacol.* 2006; 148:1133–1143. [PubMed: 16865095]
- Roncarati R, Scali C, Comery TA, Grauer SM, Aschmi S, Bothmann H, et al. Pro-cognitive and neuroprotective activity of a novel  $\alpha 7$  nicotinic acetylcholine receptor agonist for treatment of neurodegenerative and cognitive disorders. *J Pharmacol Exp Ther.* 2009; 329(2):459–468. [PubMed: 19223665]

- Rose GM, Hopper AH, De Vivo M, Tehim A. Phosphodiesterase inhibitors for cognitive enhancement. *Curr Pharm Design*. 2005; 11:3329–3334.
- Roth BL, Hanizavareh SM, Blum AE. Serotonin receptors represent highly favorable molecular targets for cognitive enhancement in schizophrenia and other disorders. *Psychopharmacology*. 2004; 174:17–24.
- Routledge C, Bromidge SM, Moss SF, Price GW, Hirst W, Newman H, et al. Characterization of SB-271046: a potent, selective and orally active 5-HT<sub>6</sub> receptor antagonist. *Brit J Pharmacol*. 2000; 130:1606–1612. [PubMed: 10928964]
- Rowe WB, O'Donnell JP, Pearson D, Rose GM, Meaney MJ, Quirion R. Long-term effects of BIBN-99, a selective muscarinic M<sub>2</sub> receptor antagonist, on improving spatial memory performance in aged cognitively impaired rats. *Behav Brain Res*. 2003; 145:171–178. [PubMed: 14529815]
- Rowe WB, DiSomma RM, Hsu CC, Solomon K, Callahan PM, Wang D, et al. Histamine H<sub>3</sub> receptor antagonists elicit pro-cognitive effects on episodic and spatial reference memories in young and aged animals. *Alzheimer's and Dementia*. 2006; 2:S647.
- Rowe WB, Blalock EM, Chen KC, Kadish I, Wang D, Barrett JE, et al. Hippocampal expression analyses reveal selective association of immediate-early, neuroenergetic, and myelinogenic pathways with cognitive impairment in aged rats. *J Neurosci*. 2007; 27:3098–3110. [PubMed: 17376971]
- Rowe WB, Callahan PM, Xie W, Hsu CC, Dunn R, Murray M, et al. Serotonin 5-HT<sub>6</sub> receptor antagonists modulate episodic, spatial and working memory highlighting a potential therapeutic role in alleviating deficits associated with mild cognitive impairment and Alzheimer's disease. *Soc Neurosci Abstr*. 2008; 685:9.
- Rowe WB, Callahan PM, Hsu CC, Tehim A. Characterization of the 5-HT<sub>6</sub> receptor agonist, WAY 181187 and antagonist, SB 742457: effects on spatial reference and spatial working memory function in aged F344 rats. *Soc Neurosci Abstr*. 2009; 883:2.
- Ruat M, Traffort E, Arrang JM, Tardivellacombe J, Diaz J, Leurs R, et al. A novel rat serotonin (5-HT<sub>6</sub>) receptor-molecular cloning, localization and stimulation of cAMP accumulation. *Biochem Biophys Res Comm*. 1993; 193:268–276. [PubMed: 8389146]
- Rupniak NM, Tye SJ, Brazell C, Heald A, Iversen SD, Pagella PG. Reversal of cognitive impairment by heptyl physostigmine, a long-lasting cholinesterase inhibitor, in primates. *J Neurol Sci*. 1992; 107(2):246–249. [PubMed: 1564524]
- Ruske AC, White KG. Facilitation of memory performance by a novel muscarinic agonist in young and old rats. *Pharmacol Biochem Behav*. 1999; 63(4):663–667. [PubMed: 10462196]
- Rutten K, Basile JL, Prickaerts J, Blokland A, Vivian JA. Selective PDE inhibitors rolipram and sildenafil improve object retrieval performance in adult cynomolgus macaques. *Psychopharmacology*. 2008; 196:643–648. [PubMed: 18034336]
- Rutten K, Prickaerts J, Hendrix M, van der Staay FJ, Sik A, Blokland A. Time-dependent involvement of cAMP and cGMP in consolidation of object memory: studies using selective phosphodiesterase type 2, 4 and 5 inhibitors. *Eur J Pharmacol*. 2007; 558:107–112. [PubMed: 17207788]
- Rutten K, Prickaerts J, Blokland A. Rolipram reverses scopolamine-induced and time-dependent memory deficits in object recognition by different mechanisms of action. *Neurobiol Learn Mem*. 2006; 85:132–138. [PubMed: 16242977]
- Rutten K, Vente JD, Sik A, Ittersum MM, Prickaerts J, Blokland A. The selective PDE5 inhibitor, sildenafil, improves object memory in Swiss mice and increases cGMP levels in hippocampus. *Behav Brain Res*. 2005; 164:11–16. [PubMed: 16076505]
- Sabbagh MN. Drug development for Alzheimer's disease: where are we now and where are we headed? *Am J Geriatr Pharm*. 2009; 7:167–185.
- Sadashiva CT, Narendra Sharath Chandra JN, Kavitha CV, Thimmegowda A, Subhash MN, Rangappa KS. Synthesis and pharmacological evaluation of novel N-alkyl/aryl substituted thiazolidinone arecoline analogues as muscarinic M<sub>1</sub> receptor agonists in Alzheimer's dementia models. *Eur J Med Chem*. 2009; 44:4848–4854. [PubMed: 19717214]

- Sagi Y, Weinstock M, Youdim MB. Attenuation of MPTP-induced dopaminergic neurotoxicity by TV3326, a cholinesterase-monoamine oxidase inhibitor. *J. Neurochem.* 2003; 86:290–297. [PubMed: 12871570]
- Samadi A, Marco-Contelles J, Soriano E, Alvarez-Pérez M, Chioua M, Romero A, et al. Multipotent drugs with cholinergic and neuroprotective properties for the treatment of Alzheimer and neuronal vascular diseases. I. Synthesis, biological assessment, and molecular modeling of simple and readily available 2-aminopyridine-, and 2-chloropyridine-3,5-dicarbonitriles. *Bioorg Med Chem.* 2010; 18(16):5861–5872. [PubMed: 20656495]
- Schaffhauser H, Mathiasen JR, Dicamillo A, Huffman MJ, Lu LD, McKenna BA, et al. Dimebolin is a 5-HT<sub>6</sub> antagonist with acute cognition enhancing activities. *Biochem Pharmacol.* 2009; 78:1035–1042. [PubMed: 19549510]
- Schechter LE, Lin Q, Smith DL, Zhang G, Shan Q, Platt B, et al. Neuropharmacological profile of novel and selective 5-HT<sub>6</sub> receptor agonists: WAY-181187 and WAY-208466. *Neuropsychopharmacology.* 2008; 33:1323–1335. [PubMed: 17625499]
- Schechter LE, Smith DL, Rosenzweig-Lipson S, Sukoff SJ, Dawson LA, Marquis K, et al. Lecozotan (SRA-333): a selective serotonin 1A receptor antagonist that enhances the stimulated release of glutamate and acetylcholine in the hippocampus and possesses cognitive-enhancing properties. *J Pharmacol Exp Ther.* 2005; 314:1274–1289. [PubMed: 15951399]
- Schlicker E, Malinowska B, Kathmann M, Gothert M. Modulation of neurotransmitter release via histamine H<sub>3</sub> heteroreceptors. *Fund Clin Pharmacol.* 1994; 8:128–137.
- Schmidt CJ, Chapin DS, Cianfrogna J, Corman ML, Hajos M, Harms JF, et al. Preclinical characterization of selective phosphodiesterase 10A inhibitors: a new therapeutic approach to the treatment of schizophrenia. *J Pharmacol Exper Ther.* 2008; 325:681–690. [PubMed: 18287214]
- Schneider JS, Tinker JP, Van Velson M, Menzaghi F, Lloyd GK. Nicotinic acetylcholine receptor agonist SIB-1508Y improves cognitive functioning in chronic low-dose MPTP-treated monkeys. *J Pharmacol Exp Ther.* 1999; 290(2):731–739. [PubMed: 10411585]
- Scheider JS, Tinker JP, Menzaghi F, Lloyd GK. The subtype selective nicotinic acetylcholine receptor agonist SIB-1553A improves both attention and working memory components of a spatial working memory task in chronic low dose 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine-treated monkeys. *J Pharmacol Exp Ther.* 2003; 306(1):401–406. [PubMed: 12721323]
- Schreiber R, Vivian J, Hedley L, Szczepanski K, Secchi RL, Zuzow M, et al. Effects of the novel 5-HT<sub>6</sub> receptor antagonist RO4368554 in rat models for cognition and sensorimotor gating. *Eur Neuropsychopharm.* 2007; 17:277–288.
- Setter SM, Iltz JL, Fincham JE, Campbell RK, Baker DE. Phosphodiesterase 5 inhibitors for erectile dysfunction. *Ann Pharmacother.* 2005; 39:1286–1295. [PubMed: 15941818]
- Seu E, Lang A, Rivera RJ, Jentsch JD. Inhibition of the norepinephrine transporter improves behavioral flexibility in rats and monkeys. *Psychopharmacology (Berl).* 2009; 202:505–519. [PubMed: 18604598]
- Sheng R, Xu Y, Hu C, Zhang J, Lin X, Li J, et al. Design, synthesis and AChE inhibitory activity of indanone and aurone derivatives. *Eur. J. Med. Chem.* 2009; 44:7–17. [PubMed: 18436348]
- Shafiei M, Mahmoudian M, Rostami P, Nemati F. Effect of sildenafil (Viagra) on memory retention of a passive avoidance response in rats. *Acta Physiol Hung.* 2006; 93:53–59. [PubMed: 16830693]
- Shirazi-Southall S, Rodriguez DE, Nomikos GG. Effects of typical and atypical antipsychotics and receptor selective compounds on acetylcholine efflux in the hippocampus of the rat. *Neuropsychopharmacology.* 2002; 26:583–594. [PubMed: 11927183]
- Shoham S, Bejar C, Kovalev E, Schorer-Apelbaum D, Weinstock M. Ladostigil prevents gliosis, oxidative-nitrative stress and memory deficits induced by intracerebroventricular injection of streptozotocin in rats. *Neuropharmacology.* 2007; 52(3):836–843. Epub 2006 Nov 22. [PubMed: 17123555]
- Si W, Zhang X, Niu Y, Yu H, Lei X, Chen H, et al. A novel derivative of xanomeline improves fear cognition in aged mice. *Neurosci Lett.* 2010; 473:115–119. [PubMed: 20178835]
- Simon R. Lost in translation: problems and pitfalls in translating laboratory observations to clinical utility. *Eur J Cancer.* 2008; 44:2707–2713. [PubMed: 18977655]

- Son H, Lu YF, Zhuo M, Arancio O, Kandel ER, Hawkins RD. The specific role of cGMP in hippocampal LTP. *Learn Mem.* 1998; 5:231–245. [PubMed: 10454367]
- Sood A, Beach WJ, Webster SJ, Terry AV, Buccafusco JJ. The effects of JWB1-84-1 on memory related task performance by amyloid Abeta transgenic mice and by young and aged monkeys. *Neuropharm.* 2007; 53(5):588–600.
- Southam E, Cilia J, Gartlon JE, Woolley ML, Lacroix LP, Jennings CA, et al. Preclinical investigations into the antipsychotic potential of the novel histamine H3 receptor antagonist GSK207040. *Psychopharmacology.* 2009; 201:483–494. [PubMed: 18762914]
- Spalding TAMAJN, Ott TR, Friberg M, Bajpai A, Bradley SR, et al. Structural requirements of transmembrane domain 3 for activation by the M1 muscarinic receptor agonists AC-42, AC-260584, clozapine, and N-desmethylozapine: evidence for three distinct modes of receptor activation. *Mol Pharmacol.* 2006; 70(6):1974–1983. [PubMed: 16959945]
- Spencer JP, Brown JT, Richardson JC, Medhurst AD, Sehmi SS, Calver AR, et al. Modulation of hippocampal excitability by 5-HT4 receptor agonists persists in a transgenic model of Alzheimer's disease. *Neuroscience.* 2004; 129:49–54. [PubMed: 15489027]
- Standridge JB. Pharmacotherapeutic approaches to the treatment of Alzheimer's disease. *Clin Ther.* 2004 May; 26(5):615–630. [PubMed: 15220008]
- Stean TO, Hirst WD, Thomas DR, Price GW, Rogers D, Riley G, et al. Pharmacological profile of SB-357134: a potent, selective brain penetrant, and orally active 5-HT(6) receptor antagonist. *Pharmacol Biochem Behav.* 2002; 71:645–654. [PubMed: 11888556]
- Sterling J, Herzig Y, Goren T, Finkelstein N, Lerner D, Goldenberg W, et al. Novel dual inhibitors of AChE and MAO derived from hydroxy aminoindan and phenethylamine as potential treatment for Alzheimer's disease. *J. Med. Chem.* 2002; 45:5260–5279. [PubMed: 12431053]
- Studzinski CM, Araujo JA, Milgram NW. The canine model of human cognitive aging and dementia: pharmacological validity of the model for assessment of human cognitive-enhancing drugs. *Prog Neuropsychopharmacol Biol Psychiatry.* 2005; 29(3):489–498. [PubMed: 15795058]
- Sullivan JP, Donnelly-Roberts D, Briggs CA, Anderson DJ, Gopalakrishnan M, Xue IC, et al. ABT-089 [2-methyl-3-(2-(S)-pyrrolidinylmethoxy)pyridine]: I. A potent and selective cholinergic channel modulator with neuroprotective properties. *J Pharmacol Exp Ther.* 1997; 283(1):235–246. [PubMed: 9336329]
- Svenningsson P, Tzavara ET, Qi HS, Carruthers R, Witkin JM, Nomikos GG, et al. Biochemical and behavioral evidence for antidepressant-like effects of 5-HT6 receptor stimulation. *J Neurosci.* 2007; 27:4201–4209. [PubMed: 17428998]
- Sydeserff S, Sutton EJ, Song D, Quirk MC, Maciag C, Li C, et al. Selective alpha7 nicotinic receptor activation by AZD0328 enhances cortical dopamine release and improves learning and attentional processes. *Biochem Pharmacol.* 2009; 78(7):880–888. [PubMed: 19615981]
- Taly A, Corringer PJ, Guedin D, Lestage P, Changeux JP. Nicotinic receptors: allosteric transitions and therapeutic targets in the nervous system. *Nat Rev Drug Dis.* 2009; 8(9):733–750.
- Teclé H, Schwarz RD, Barrett SD, Callahan MJ, Caprathe BW, Davis RE, et al. CI-1017, a functionally M1-selective muscarinic agonist: design, synthesis, and preclinical pharmacology. *Pharm Acta Helv.* 2000; 74(2–3):141–148. [PubMed: 10812951]
- Terry AV Jr, Buccafusco JJ, Prendergast MA, Jackson WJ, Fontana DL, Wong EH, et al. The 5-HT3 receptor antagonist, RS-56812, enhances delayed matching performance in monkeys. *Neuroreport.* 1996; 8:49–54. [PubMed: 9051751]
- Terry AV Jr, Buccafusco JJ, Jackson WJ, Prendergast MA, Fontana DJ, Wong EHF, et al. Enhanced delayed matching performance in younger and older macaques administered the 5HT4 receptor agonist, RS 17017. *Psychopharmacology.* 1998; 135:407–415. [PubMed: 9539266]
- Terry AV Jr, Gattu M, Buccafusco JJ, Sowell JW, Kosh JW. Ranitidine Analog, JWS USC 75IX, Enhances Memory Related Task Performance in Rats. *Drug Development Research.* 1999; 47:97–106.
- Terry AV Jr, Buccafusco JJ. The Cholinergic Hypothesis of Age and Alzheimer's Disease-Related Cognitive Deficits: Recent Challenges and Their Implications for Novel Drug Development. *J Pharmacol Exper Ther.* 2003; 306(3):821–827. [PubMed: 12805474]



- Terry AV Jr, Hernandez CM, Hohnadel EJ, Bouchard KP, Buccafusco JJ. Cotinine. A Neuroactive Metabolite of Nicotine: Potential for Treating Disorders of Impaired Cognition. *CNS Drug Reviews*. 2005a; 11:229–252. [PubMed: 16389292]
- Terry AV Jr, Buccafusco JJ, Bartoszyk GD. Selective serotonin 5-HT<sub>2A</sub> receptor antagonist EMD 281014 improves delayed matching performance in young and aged rhesus monkeys. *Psychopharmacology*. 2005b; 179:725–732. [PubMed: 15619109]
- Terry AV Jr, Buccafusco JJ, Wilson C. Cognitive dysfunction in neuropsychiatric disorders: selected serotonin receptor subtypes as therapeutic targets. *Behav Brain Res*. 2008; 195:30–38. [PubMed: 18241938]
- Terry AV, Buccafusco JJ, Herman EJ, Callahan PM, Beck WD, Warner S, Vandenhuerk L, Bouchard K, Schwarz GM, Gao J, Chapman JM. Prototypical ranitidine analog, JWS-USC-75-IX, improves information processing and cognitive function in animal models. *J Pharmacol Exper Ther*. 2011 (in press).
- Tzavara ET, Bymaster FP, Felder CC, Wade M, Gomeza J, Wess J, McKinzie DL, Nomikos GG. Dysregulated hippocampal acetylcholine neurotransmission and impaired cognition in M2, M4 and M2/M4 muscarinic receptor knockout mice. *Mol. Psychiatry*. 2003; 8:673–679. [PubMed: 12874603]
- The Centers for Disease Control and Prevention. Public health and aging: trends in aging – United States and worldwide. *JAMA*. 2003; 289:1371–1373. [PubMed: 12636453]
- Thomas SJ, Grossberg GT. Memantine: a review of studies into its safety and efficacy in treating Alzheimer's disease and other dementias. *Clin Interv Aging*. 2009; 4:367–377. [PubMed: 19851512]
- Timmermann DB, Gronlien JH, Kohlhaas KL, Neilsen EO, Dam E, Jorgensen TD, et al. An allosteric modulator of the alpha7 nicotinic acetylcholine receptor possessing cognition-enhancing properties in vivo. *J Pharmacol Exp Ther*. 2007; 323:294–307. [PubMed: 17625074]
- Trabace L, Cassano T, Colaianna M, Castrignanò S, Giustino A, Amoroso S, et al. Neurochemical and neurobehavioral effects of ganstigmine (CHF2819), a novel acetylcholinesterase inhibitor, in rat prefrontal cortex: an in vivo study. *Pharmacol Res*. 2007; 56(4):288–294. Epub 2007 Jul 27. [PubMed: 17822918]
- Tully T, Bourtchouladze R, Scott R, Tallman J. Targeting the CREB pathway for memory enhancers. *Nat Rev Drug Discov*. 2003; 2:267–277. [PubMed: 12669026]
- United Nations. New York: United Nations; 2007. *World Population Ageing 2007*.
- Upton N, Chuang TT, Hunter AJ, Virley DJ. 5-HT<sub>6</sub> receptor antagonists as novel cognitive enhancing agents for Alzheimer's disease. *Neurotherapeutics*. 2008; 5:458–469. [PubMed: 18625457]
- Valli MJ, Tang Y, Kosh JW, Chapman JM Jr, Sowell JW Sr. Synthesis and cholinergic properties of N-aryl-2-[[[5-[(dimethylamino)methyl]-2-furanyl]methyl]thio]ethylamino analogs of ranitidine. *J Med Chem*. 1992; 35:3141–3147. [PubMed: 1507203]
- Van der Schyf CJ, Mandel S, Geldenhuys WJ, Amit T, Avramovich Y, Zheng H, et al. Novel multifunctional anti-Alzheimer drugs with various CNS neurotransmitter targets and neuroprotective moieties. *Curr Alzheimer Res*. 2007; 4:522–536. [PubMed: 18220515]
- van der Staay FJ, Rutten K, Barfacker L, DeVry J, Erb C, Heckroth H, et al. The novel selective PDE9 inhibitor BAY 73–6691 improves learning and memory in rodents. *Neuropharmacology*. 2008; 55:908–918. [PubMed: 18674549]
- van Donkelaar EL, Rutten K, Blokland A, Akkerman S, Steinbusch HWM, Prickaerts J. Phosphodiesterase 2 and 5 inhibition attenuates the object memory deficit induced by acute tryptophan depletion. *Eur J Pharmacol*. 2008; 600:98–104. [PubMed: 18957291]
- van Kampen M, Selbach K, Schneider R, Schiegel E, Boess F, Schreiber R. AR-R 17779 improves social recognition in rats by activation of nicotinic alpha7 receptors. 2004; 172(4):375–383.
- van Marum RJ. Update on the use of memantine in Alzheimer's disease. *Neuropsychiatr Dis Treat*. 2009; 5:237–247. [PubMed: 19557118]
- Vilario MT, Cortes R, Mengod G. Serotonin 5-HT<sub>4</sub> receptors and their mRNAs in rat and guinea pig brain: distribution and effects of neurotoxic lesions. *J Comp Neurol*. 2005; 484:418–439. [PubMed: 15770652]

- Vitolo OV, Sant'Angelo A, Costanzo V, Battaglia F, Arancio O, Shelanski M. Amyloid beta-peptide inhibition of the PKA/CREB pathway and long-term potentiation: reversibility by drugs that enhance cAMP signaling. *Pro Natl Acad Sci USA*. 2002; 99:13217–13221.
- Wallace TL, Callahan PM, Tehim A, Bertrand D, Tombaugh G, Wang S, Xie W, Rowe WB, Ong V, Graham E, Terry AV Jr, Rodefer JS, Herbert B, Murray M, Porter R, Santarelli L, Lowe DA. RG3487, a novel nicotinic  $\alpha 7$  receptor partial agonist, improves cognition and sensorimotor gating in rodents. *J Pharmacol Exp Ther*. 2011; 336:242–253. [PubMed: 20959364]
- Wallace-Boone TL, Chiu G, Dao H, Lowe D, Porter R, Santarelli L. R3487/MEM3454, a novel nicotinic alpha 7 receptor partial agonist, improves attention and working memory performance in cynomolgus macaques. *Soc Neurosci Abstr*. 2009:887.5.
- Wallace-Boone TL, Yeo H, Callahan PM, Tombaugh G, Misner DL, Grovey B, Thompson J, Van Natta K, Sahdeo S, Tehim A, Maag H, Santarelli L. Pro-cognitive properties of MEM 64368, a novel nicotinic alpha 7 receptor partial agonist, following acute and chronic administration. *Biochem Pharmacol*. 2007; 74 SMA-45.
- Wang T, Tang XC. Reversal of scopolamine-induced deficits in radial maze performance by (-)-huperzine A: comparison with E2020 and tacrine. *Eur J Pharmacol*. 1998; 349:137–142. [PubMed: 9671090]
- Wang ZF, Tang LL, Yan H, Wang YJ, Tang XC. Effects of huperzine A on memory deficits and neurotrophic factors production after transient cerebral ischemia and reperfusion in mice. *Pharmacol Biochem Behav*. 2006; 83(4):603–611. Epub 2006 May 9. [PubMed: 16687166]
- Wang J, Zhang HY, Tang XC. Huperzine a improves chronic inflammation and cognitive decline in rats with cerebral hypoperfusion. *J Neurosci Res*. 2010; 88(4):807–815. [PubMed: 19795377]
- Watkins PB, Zimmerman HJ, Knapp MJ, Gracon SI, Lewis KW. Hepatotoxic effects of tacrine administration in patients with Alzheimer's disease. *JAMA*. 1994; 271(13):992–998. [PubMed: 8139084]
- Weinreb O, Bar-Am O, Amit T, Chillag-Talmor O, Youdim MBH. Neuroprotection via pro-survival protein kinase C isoforms associated with Bcl-2 family members. *FASEB Journal*. 2004; 18:1471–1473. [PubMed: 15247150]
- Weinreb O, Amit T, Bar-Am O, Yogev-Falach M, Youdim MB. The neuroprotective mechanism of action of the multimodal drug ladostigil. *Front Biosci*. 2008; 13:5131–5137. [PubMed: 18508575]
- Weinstock M, Behar C, Wang RH, Poltyrev T, Gross A, Finberg JP, et al. TV3326, a novel neuroprotective drug with cholinesterase and monoamine oxidase inhibitory activities for the treatment of Alzheimer's disease. *J. Neural Transm*. 2000; 60:157–169.
- Weinstock M, Kirschbaum-Slager N, Lazarovici P, Bejar C, Youdim MBH, Shoham S. Neuroprotective effects of novel cholinesterase inhibitors derived from rasagiline as potential anti- Alzheimer drugs. *Ann NY Acad Sci*. 2001; 939:148–162. [PubMed: 11462767]
- Weinstock M, Poltyrev T, Bejar C, Youdim MB. Effect of TV3326, a novel monoamineoxidase cholinesterase inhibitor, in rat models of anxiety and depression. *Psychopharmacology (Berl.)*. 2002; 160:318–324. [PubMed: 11889501]
- Weinstock M, Gorodetsky E, Poltyrev T, Gross A, Sagi Y, Youdim M. A novel cholinesterase and brain-selective monoamine oxidase inhibitor for the treatment of dementia comorbid with depression and Parkinson's disease. *Prog. Neuropsychopharmacol. Biol. Psychiatry*. 2003; 27:555–561. [PubMed: 12787840]
- Weinstock M, Luques L, Poltyrev T, Bejar C, Shoham S. Ladostigil prevents age-related glial activation and spatial memory deficits in rats. *Neurobiol Aging*. 2009 Jul 20. [Epub ahead of print].
- Wesnes KA, McKeith IG, Ferrara R, Emre M, del Ser T, Spano PF, et al. Effects of rivastigmine on cognitive function in dementia with lewy bodies: a randomised placebo-controlled international study using the cognitive drug research computerised assessment system. *Dement Geriatr Cogn Disord*. 2002; 13:183–192. [PubMed: 11893841]
- Wesolowska A, Nikiforuk A. Effects of the brain-penetrant and selective 5-HT<sub>6</sub> receptor antagonist SB-399885 in animal models of anxiety and depression. *Neuropharmacology*. 2007; 52:1274–1283. [PubMed: 17320917]

- Wess J, Eglen RM, Gautam D. Muscarinic acetylcholine receptors: mutant mice provide new insights for drug development. *Nature Reviews Drug Discovery*. 2007; 6:721–733.
- West PJ, Marcy VR, Marino MJ, Schaffhauser H. Activation of the 5-HT<sub>6</sub> receptor attenuates long-term potentiation and facilitates GABAergic neurotransmission in rat hippocampus. *Neuroscience*. 2009; 164:692–701. [PubMed: 19660530]
- Wilens TE, Decker MW. Neuronal nicotinic receptor agonists for the treatment of attention-deficit/hyperactivity disorder: focus on cognition. *Biochem Pharmacol*. 2007; 74:1212–1223. [PubMed: 17689498]
- Wilkinson DG, Francis PT, Schwam E, Payne-Parrish J. Cholinesterase inhibitors used in the treatment of Alzheimer's disease: the relationship between pharmacological effects and clinical efficacy. *Drugs Aging*. 2004; 21(7):453–478. [PubMed: 15132713]
- Williams GV, Rao SG, Goldman-Rakic PS. The physiological role of 5-HT<sub>2A</sub> receptors in working memory. *J Neurosci*. 2002; 22:2843–2854. [PubMed: 11923449]
- Winblad B, Engedal K, Soininen H, Verhey F, Waldemar G, Wimo A, et al. A 1-year, randomized, placebo-controlled study of donepezil in patients with mild to moderate AD. *Neurology*. 2001; 57(3):489–495. [PubMed: 11502918]
- Winstanley CA, Chudasama Y, Dalley JW, Theobald DEH, Glennon JC, Robbins TW. Intra-prefrontal 8-OH-DPAT and M100907 improve visuospatial attention and decrease impulsivity on the five-choice serial reaction time task in rats. *Psychopharmacology*. 2003; 167:304–314. [PubMed: 12677356]
- Witkin JM, Nelson DL. Selective histamine H<sub>3</sub> receptor antagonists for treatment of cognitive deficiencies and other disorders of the central nervous system. *Pharmacol Ther*. 2004; 103:1–20. [PubMed: 15251226]
- Wolff MC, Overshiner C, Leander JD. 5-Ht<sub>6</sub> receptor antagonists enhance memory in radial maze and object recognition tasks. *Soc Neurosci Abstr*. 2002:586.1.
- Woolley ML, Bentley JC, Sleight AJ, Mardsen CA, Fone KCF. A role for 5-Ht(6) receptors in retention of spatial learning in the Morris water maze. *Neuropharmacology*. 2001; 41:210–219. [PubMed: 11489457]
- Woolley ML, Marsden CA, Fone KCF. 5-ht<sub>6</sub> receptors. *Curr Drug Targets CNS Neurol Disord*. 2004; 3:59–79. [PubMed: 14965245]
- World Health Organization. The World is Fast Aging – Have We Noticed?. 2008. <http://www.who.int/ageing/en/index.html>
- Wu J, Li Q, Bezprozvanny I. Evaluation of Dimebon in cellular model of Huntington's disease. *Mol Neurodegener*. 2008 Oct 21.3:15. [PubMed: 18939977]
- Xu AJ, Chen Z, Li R, Zhu CY, Wei EQ. Effects of TAK-147, a novel acetylcholinesterase inhibitor, on spatial memory deficit as evaluated by Morris water maze of rats. *Zhejiang Da Xue Xue Bao Yi Xue Ban*. 2002a Apr; 31(2):98–102. [PubMed: 12539269]
- Xu AJ, Chen Z, Yanai K, Huang YW, Wei EQ. Effect of 3-[1-(phenylmethyl)-4-piperidiny]-1-(2,3,4,5-tetrahydro-1H-1-benzazepin-8-yl)-1-propanone fumarate, a novel acetylcholinesterase inhibitor, on spatial cognitive impairment induced by chronic cerebral hypoperfusion in rats. *Neurosci Lett*. 2002b; 331(1):33–36. [PubMed: 12359317]
- Ye JW, Cai JX, Wang LM, Tang XC. Improving effects of huperzine A on spatial working memory in aged monkeys and young adult monkeys with experimental cognitive impairment. *J Pharmacol Exp Ther*. 1999; 288:814–819. [PubMed: 9918593]
- Ye JW, Shang YZ, Wang ZM, Tang XC. Huperzine A ameliorates the impaired memory of aged rat in the Morris water maze performance. *Acta Pharmacol Sin*. 2000; 21:65–69. [PubMed: 11263250]
- Yesavage JA, Tinklenberg JR, Hollister LE, Berger PA. Vasodilators in senile dementias: a review of the literature. *Arch Gen Psychiatry*. 1979; 36:220–223. [PubMed: 420543]
- Yogev-Falach M, Amit T, Bar-Am O, Sagi Y, Weinstock M, Youdim MBH. The involvement of mitogen-activated protein (MAP) kinase in the regulation of amyloid precursor protein processing by novel cholinesterase inhibitors derived from rasagiline. *FASEB J*. 2002; 16(12):1674–1676. [PubMed: 12206996]

- Yogev-Falach M, Bar-Am O, Amit T, Weinreb O, Youdim MB. A multifunctional, neuroprotective drug, ladostigil (TV3326), regulates holo-APP translation and processing. *FASEB J.* 2006; 20:2177–2179. [PubMed: 16935943]
- Youdim MB, Buccafusco JJ. Multi-functional drugs for various CNS targets in the treatment of neurodegenerative disorders. *Trends Pharmacol Sci.* 2005 Jan; 26(1):27–35. [PubMed: 15629202]
- Zhang HT, O'Donnell JM. Effects of rolipram on scopolamine-induced impairment of working and reference memory in the radial-arm maze test in rats. *Psychopharmacology.* 2000a; 150:311–316. [PubMed: 10923759]
- Zhang HT, Crissman AM, Dorairaj NR, Chandler LJ, O'Donnell JM. Inhibition of cyclic AMP phosphodiesterase (PDE4) reverses memory deficits associated with NMDA receptor antagonism. *Neuropsychopharmacology.* 2000b; 23:198–204. [PubMed: 10882846]
- Zhang HT, Zhao Y, Huang Y, Dorairaj NR, Chandler LJ, O'Donnell JM. Inhibition of the phosphodiesterase (PDE4) enzyme reverses memory deficits produced by infusion of the MEK inhibitor U0126 into the CA1 subregion of the rat hippocampus. *Neuropsychopharmacology.* 2004; 29:1432–1439. [PubMed: 15114341]
- Zhang HT, Huang Y, Suvarna NU, Deng C, Crissman AM, Hopper AT, et al. Effects of the novel PDE4 inhibitors MEM1018 and MEM 1091 on memory in the radial-arm maze and inhibitory avoidance tests in rats. *Psychopharmacology.* 2005; 179:613–619. [PubMed: 15672274]
- Zhang J, Zhu D, Sheng R, Wu H, Hu Y, Wang F, et al. BZYX, a novel acetylcholinesterase inhibitor, significantly improved chemicals-induced learning and memory impairments on rodents and protected PC12 cells from apoptosis induced by hydrogen peroxide. *Eur J Pharmacol.* 2009; 613(1–3):1–9. [PubMed: 19345205]
- Zhang M, Ballard ME, Pan L, Roberts S, Faghih R, Cowart M, et al. Lack of cataleptogenic potentiation with non-imidazole H3 receptor antagonists reveals potential drug-drug interactions between imidazole-based H3 receptor antagonists and antipsychotic drugs. *Brain Res.* 2005; 1045:142–149. [PubMed: 15910772]
- Zhang M-Y, Hughes ZA, Kerns EH, Lin Q, Beyer CE. Development of a liquid chromatography/tandem mass spectrometry method for the quantitation of acetylcholine and related neurotransmitters in brain microdialysis samples. *J Pharm Biomed Anal.* 2007; 44:586–593. [PubMed: 17383138]
- Zhang ZJ, Kang WH, Li Q, Wang XY, Yao SM, Ma AQ. Beneficial effects of ondansetron as an adjunct to haloperidol for chronic, treatment-resistant schizophrenia: a double-blind, randomized, placebo-controlled study. *Schizophr Res.* 2006; 88:102–110. [PubMed: 16959472]

Table 1

## Cholinergic-Based Compounds

Compound Name	Receptor	Mechanism	References
<b>AChE Inhibitors</b>			
TAK-147	AChE	Inhibitor	Xu, 2002
BZYX	AChE	Inhibitor	Shen, 2009
Phenserine	AChE	Inhibitor	Janas, 2005
CHF2819	AChE	Inhibitor	Trabace, 2007
Eptastigmine	AChE	Inhibitor	Imbimbo, 2001
Huperzine A	AChE	Inhibitor	Wang, 2010
Loganin	AChE	Inhibitor	Lee, 2009
NP-0361	AChE	Inhibitor	Buccafusco, 2009
Pseudoberberine	AChE	Inhibitor	Hung, 2008
Memoquin	AChE	Inhibitor	Cavalli, 2007
PMS777	AChE	Inhibitor	Bate, 2004
<b>Nicotinic</b>			
SIB-1508Y	$\alpha 4\beta 2$	agonist	Schneider et al., 1999
SIB-1553A	$\alpha 2\beta 4$	agonist	Schneider et al., 2003
ABT-418	$\alpha 4\beta 2$	agonist	Prendergast et al., 1998
ABT-594	$\alpha 4\beta 2$	agonist	Buccafusco et al., 2007
S-38232	$\alpha 4\beta 2$	agonist	Howe et al., 2010
ABT-089	$\alpha 4\beta 2$	partial agonist	Decker et al., 1997
Ispronicline/AZD-3480	$\alpha 4\beta 2$	partial agonist	Gatto et al., 2004
Varenicline/Chantix <sup>®</sup>	$\alpha 4\beta 2/\alpha 7$	partial agonist/full agonist	Rollema et al., 2009
GTS-21	$\alpha 7$	partial agonist	Buccafusco and Terry, 2009b
MEM-3454/RG-3487	$\alpha 7$	partial agonist	Callahan et al., 2006; Rezvani et al., 2009
MEM-63908/RG4996	$\alpha 7$	partial agonist	Taly et al., 2009
AZD-0328	$\alpha 7$	partial agonist	Sydserff et al., 2009
AR-R17779	$\alpha 7$	agonist	Van Kampen et al., 2004
S-24795	$\alpha 7$	partial agonist	Lopez-Hernandez et al., 2007
SSR-180711	$\alpha 7$	partial agonist	Taly et al., 2009
SEN12333 (WAY-317538)	$\alpha 7$	agonist	Roncarati et al., 2009
NS-1738	$\alpha 7$	positive allosteric modulator	Timmermann et al., 2007
Compound 6	$\alpha 7$	positive allosteric modulator	Ng et al., 2007
PNU-120596	$\alpha 7$	positive allosteric modulator	Hurst et al., 2005
2087101	$\alpha 4\beta 2/\alpha 7$	positive allosteric modulator	de Filippi et al., 2010
Cotinine	$\alpha 4\beta 2/\alpha 7$	nAChR desensitization	Buccafusco et al., 2009; Terry et al., 2005a
JWB1-84-1	$\alpha 4\beta 2/\alpha 7$	nAChR desensitization	Buccafusco et al., 2009; Sood et al., 2007
JAY2-22-33	$\alpha 4\beta 2/\alpha 7$	nAChR desensitization	Buccafusco et al., 2009; Sood et al., 2007
<b>Muscarinic</b>			
Talsaclidine	M1	agonist	Clader and Wang, 2005
Xanomeline	M1	agonist	Clader and Wang, 2005

<b>Compound Name</b>	<b>Receptor</b>	<b>Mechanism</b>	<b>References</b>
Cevimeline/AF102B	M1	agonist	Fisher et al, 2002
AF150(S)	M1	agonist	Fisher et al, 1998;
AF267B	M1	agonist	Cacomo et al, 2006
CI-1017	M1	agonist	Tecle et al, 2000
WAY-132983	M1	agonist	Bartolomeo et al, 2000
EUK-1001	M1	agonist	Cui et al, 2008a
MCD-386	M1	agonist	Buccafusco, 2009a
AC-42	M1	positive allosteric modulator	Spalding et al, 2006
77-LH-28-1	M1	positive allosteric modulator	Fisher, 2008
TBPB	M1	positive allosteric modulator	Jones et al, 2008
SCH 57790	M2	antagonist	Carey et al., 2001
SCH 72788	M2	antagonist	Lachowicz et al., 2001
BIBN-99	M2	antagonist	Rowe et al., 2003

**Table 2**

## Phosphodiesterase Inhibitors

Compound Name/Identifier	Reference
<b>PDE2 Inhibitors</b>	
BAY 60-7550	Boess et al., 2004; Domek-Lopacinska and Strosznajder 2008; Rutten et al., 2007; van Donkelaar et al., 2008
<b>PDE4 Inhibitors</b>	
HT-0712	Bourtchouladze et al., 2003
L-454,560	Huang et al., 2007
MEM1018	Zhang et al., 2005
MEM 1091	Zhang et al., 2005
Rolipram	Barad et al., 1998; Comery et al., 2005; Egawa et al., 1997; Gong et al., 2004; Rutten et al., 2007, 2008; Zhang and O'Donnell, 2000; Zhang et al., 2000
<b>PDE5 Inhibitors</b>	
Sildenafil	Devan et al., 2004, 2006; Domek-Lopacinska and Strosznajder, 2008; Prickaerts et al., 2002, 2005; Rutten et al., 2008; Shafiei et al., 2006
Vardenafil	Prickaerts et al., 2002; Rutten et al., 2007; van Donkelaar et al., 2008
Zaprinast	Prickaerts et al., 1997; Domek-Lopacinska and Strosznajder, 2008
<b>PDE9 Inhibitors</b>	
BAY 73-6691	Van der Staay et al., 2008
<b>PDE10 Inhibitors</b>	
MP-10	Grauer et al., 2009
Papaverine	Hebb et al., 2008; Rodefer et al., 2005
TP-10	Schmidt et al., 2008

Table 3

## Serotonergic-Based Compounds

Compound Name	Receptor	Reference
<b>5-HT<sub>1A</sub></b>		
8-OH-DPAT	agonist	Meneses and Perez-Garcia, 2007; Orgen et al., 2008
Buspirone	partial agonist	Meneses, 1999; Orgen et al., 2008
MDL 73005	agonist	Bertrand et al., 2001
S15535	agonist	Millan et al., 2004
lecozotan/SRA-333	antagonist	Schechter et al., 2005
NAD-299	antagonist	Luttgen et al., 2005; Misane and Ogren, 2003
WAY 100635	antagonist	Harder and Ridley, 2000; Misane and Ogren, 2003; Pitsikas et al., 2003
WAY 101405	antagonist	Hirst et al., 2008
<b>5-HT<sub>2A</sub></b>		
MDL 100,907	antagonist	Meneses, 1999; Roth et al., 2004; Williams et al., 2002; Winstanley et al., 2003
EMD 281014	antagonist	Terry et al., 2005b
Ketanserin	antagonist	Levin and Rezvani, 2007
Mianserin	antagonist	Roth et al., 2004
<b>5-HT<sub>3</sub></b>		
DAU 6215	antagonist	Pitsikas et al., 1996
Granisetron	antagonist	Naghdi and Harooni, 2005; Pitsikas and Borsini, 1997;
Ondansetron	antagonist	Arnsten et al., 1997; Barnes et al., 1999; Carli et al., 1997; Domeneq et al., 1991; Pitsikas and Borsini, 1997
RS-56812	antagonist	Terry et al., 1996
SEC 579	antagonist	Meneses, 1999
WAY 100289	antagonist	Hodges et al., 1995
<b>5-HT<sub>4</sub></b>		
RS 67333	partial agonist	Chaillan et al., 2004; Fontana et al., 1997; Kulla and Manahan-Vaughan, 2002; Levallet et al., 2009
SC 53116	partial agonist	Matsumoto et al., 2001
SL 65.0155	partial agonist	Hille et al., 2008; Moser et al., 2002; Spencer et al., 2004
BIMU1	partial agonist	Galeotti et al., 1998
VRX 03011	partial agonist	Mohler et al., 2007
RS 17017	agonist	Terry et al., 1998
GR 125487	antagonist	Galeotti et al., 1998; Lamirault and Simon, 2001; Spencer et al., 2004
RS 67532	antagonist	Chaillan et al., 2004; Fontana et al., 1997
SDZ 205,557	antagonist	Galeotti et al., 1998
<b>5-HT<sub>6</sub></b>		
Ro-04-6790	antagonist	King et al., 2004, 2005, 2009; Meneses, 2001; Woolley et al., 2004
Ro-4368554	antagonist	Lschreiber et al., 2007
SB 271046	antagonist	Callahan et al., 2004; Foley et al., 2004; King et al., 2004; Loiseau et al., 2008; Marcos et al., 2008; Rodefer et al., 2008; Routledge et al., 2000
SB 357134	antagonist	Rogers and Hagan, 2001; Stean et al., 2002
SB 399885	antagonist	Hatcher et al., 2005; Hirst et al., 2006; Rowe et al., 2008
SB 742457	antagonist	Callahan et al., 2009; Maher-Edwards et al., 2010; Rowe et al., 2009; Upton et al., 2008



---

<b>Compound Name</b>	<b>Receptor</b>	<b>Reference</b>
MEM 68626	antagonist	Rowe et al., 2008
MEM 34551	antagonist	Rowe et al., 2008
PRX-07034	antagonist	Gannon et al., 2006
SB 742457	antagonist	Callahan et al., 2009; Maher-Edwards et al., 2010; Rowe et al., 2009; Upton et al., 2008
EMD 386088	agonist	Kendall et al., 2010; Meneses et al., 2008
E-6801	agonist	Kendall et al., 2010; Romero et al., 2006
R-13c	agonist	Fone, 2008
WAY 181187	agonist	Burnham et al., 2010; Callahan et al., 2009; Loiseau et al., 2008; Rowe et al., 2009; Schechter et al., 2008; West et al., 2009
WAY 208466	agonist	Schechter et al., 2008
SB 399885	antagonist	Hatcher et al., 2005; Hirst et al., 2006; Rowe et al., 2008

---

**Table 4****Histamine (H<sub>3</sub>) Receptor Ligands and Miscellaneous Categories**

<b>Compound Name</b>	<b>Receptor/Mechanism</b>	<b>Reference</b>
<b>H<sub>3</sub> Receptor Ligands</b>		
ciproxifan	inverse agonist/antagonist	Day et al., 2007; Ligneau et al., 1998; Pascoli et al., 2009
thioperamide	inverse agonist/antagonist	Giovannini et al., 1999; Meguro et al., 1995; Rowe et al., 2006
ABT-239	inverse agonist/antagonist	Esbenshade et al., 2005, 2008; Fox et al., 2005
BF 2.649	inverse agonist/antagonist	Ligneau et al., 2007
GSK 189254	inverse agonist/antagonist	Medhurst et al., 2007a
GSK 207040	Antagonist/antagonist	Medhurst et al., 2007b; Southam et al., 2009
GSK 334429	Antagonist/antagonist	Medhurst et al., 2007b
JNJ- 5207852	neutral antagonist	Barbier et al., 2004; Bonaventure et al., 2007
JNJ-10181457	neutral antagonist	Bonaventure et al., 2007
clobenpropit	inverse agonist/antagonist	Giovannini et al., 1999; Huang et al., 2004
A-304121	inverse agonist/antagonist	Esbenshade et al., 2008; Fox et al., 2003
A-317920	inverse agonist/antagonist	Esbenshade et al., 2008; Fox et al., 2003
ABT-239	inverse agonist/antagonist	Esbenshade et al., 2005, 2008; Fox et al., 2005
<b>Multi-Functional Compounds</b>		
Ladostigil	AChEI/MAOI	Reviewed, Buccafusco, 2003; Reviewed, Youdim, 2006
Dimebolin	AChEI/ NMDA, H <sub>3</sub> , and 5HT <sub>6</sub> antagonist	Bachurin, 2001; Grigorev, 2003; Schaffhauser, 2009
JWS-USC-75IX	AChEI/M <sub>2</sub> antagonist	Terry et al., 1999; Terry et al., 2011
memoquin	AChEI/Coenzyme Q activity	Cavalli, 2007; Bolognesi, 2009
PM5777	AChEI/Platelet activating factor inhibitors	Li, 2007
<b>Other Agents and Targets</b>		
EVP 0334	HDAC Inhibitor	Hannen et al., 2008
	Angiotensin converting enzyme inhibitors and receptor blockers	Reviewed, Gard, 2004
	glycine transporter, mGluR2/3, somatostatin receptor, estrogen receptor, D2 receptor, calcium channels receptors, GABA	Reviewed, Buccafusco, 2009