

Further evidence from the effect of fungi on breaking *Opuntia* seed dormancy

Pablo Delgado-Sánchez,¹ María Azucena Ortega-Amaro,² Aída Araceli Rodríguez-Hernández,² Juan Francisco Jiménez-Bremont² and Joel Flores^{1,*}

¹División de Ciencias Ambientales; ²División de Biología Molecular; Instituto Potosino de Investigación Científica y Tecnológica; San Luis Potosí, México

Recently, we found that fungi are involved in breaking seed dormancy of *Opuntia streptacantha*, and that the effect of fungi on seeds is species-specific. However, the effect of fungi on seed germination from other *Opuntia* spp. has not been evaluated. Thus, we evaluated the effect of four fungal species (*Penicillium chrysogenum*, *Phoma* sp., *Trichoderma harzianum*, *Trichoderma koningii*) on the germination of *Opuntia leucotricha*, an abundant species in the Chihuahuan Desert, Mexico. We found that seeds inoculated with the four fungal species had higher germination than control seeds. *Trichoderma* spp. were the most effective. Our results strongly indicate that fungi are involved in breaking seed dormancy of *O. leucotricha*. Thus, we suggest that these fungi could promote seed germination from other *Opuntia* species.

Seeds in the soil interact with microorganisms that could help them break seed dormancy. Fungi attack the testa, eroding or cracking the hard/stony endocarp, and could reduce the mechanical resistance to germination in seeds with physiological dormancy.¹ In arid environments, the effects of fungi on breaking seed dormancy in cacti have received very little attention. Recently, our work group found that *Phoma* sp. and *Trichoderma koningii*, and in less proportion *Penicillium chrysogenum*, help break seed dormancy of *Opuntia streptacantha*, maybe by the action of enzymes that degrade the testa.² However, the effect of fungi on seed

germination from other *Opuntia* species has not been evaluated.

In this study, we test the effects of four fungal species (two isolated from *O. streptacantha* testa) in breaking seed dormancy of *Opuntia leucotricha*; a perennial arborescent cactus of economic interest distributed on the semiarid lands of central Mexico.

Since seeds of *Opuntia* spp. have physiological dormancy, they need a period of after-ripening to break dormancy, and the embryos have low growth potential; we used two-year-old seeds, assuming that old seeds have broken physiological seed dormancy and that fungi can reduce mechanical resistance to germination.² *O. leucotricha* seeds were collected from mature fruits in 2008 and stored in paper bags at room temperature during two years.

Penicillium chrysogenum, *Phoma* sp., *Trichoderma harzianum* and *T. koningii* were grown on PDA plates at 28°C for three days. The spores (*P. chrysogenum*, *T. harzianum* and *T. koningii*) and mycelia (*Phoma* sp.) were collected in sterile distilled water and counted in a Neubauer chamber for later inoculation of *O. leucotricha* seeds. Sterilized seeds were grown on water-agar plates and inoculated with 2 µl of spore solution or mycelium (6×10^7 ml⁻¹) from each fungus. Seeds were incubated in water-agar plates for 35 days in an automatic germination chamber with a 16 h light and 8 h dark photoperiod at 25°C ± 2°C. There were five replicates per treatment and 20 seeds per replicate.

Key words: cactaceae, *Opuntia leucotricha*, *Penicillium chrysogenum*, *Phoma* sp., physiological dormancy, prickly pear, seed germination, *Trichoderma* spp.

Submitted: 06/27/10

Accepted: 06/28/10

Previously published online:
www.landesbioscience.com/journals/psb/
article/12835

DOI: 10.4161/psb.5.10.12835

*Correspondence to: Joel Flores;
Email: joel@ipicyt.edu.mx

Addendum to: Delgado-Sánchez P, Ortega-Amaro MA, Jiménez-Bremont JF, Flores J. Are fungi important for breaking seed dormancy in desert species? Experimental evidence in *Opuntia streptacantha* (Cactaceae). Plant Biol 2010; 13:154-9; PMID: 21143736; DOI: 10.1111/j.1438-8677.2010.00333.

Table 1. Effect of fungal species on germination of *Opuntia leucotricha* seeds

Treatment	Germination percentage (\pm S.E.)
Control	0% ^c
<i>Penicillium chrysogenum</i>	15% (\pm 3.35) ^b
Phoma sp.	10% (\pm 2.23) ^b
<i>Trichoderma harzianum</i>	40% (\pm 8.94) ^a
<i>Trichoderma koningii</i>	37% (\pm 8.27) ^a

Significant differences ($p < 0.0001$) between fungal species are indicated by different lower-case letters.

After one-way ANOVA, we found a significant effect of fungal species ($F = 52.198$, $p < 0.0001$) on *O. leucotricha* seed germination. Seeds inoculated with the four fungal species had higher germination than control, although *Trichoderma* spp. promoted higher seed germination than the other fungi examined (Table 1).

Opuntia species have hard to break dormancy in their seeds.²⁻⁸ Seed dormancy is a common plant strategy in arid and semiarid environments, which has been found in several plant families,⁹⁻¹¹ including Cactaceae.¹²⁻¹⁶

In our experiment, we found that four fungal species erode the endocarp and break seed dormancy of *O. leucotricha*. Since seeds of *Opuntia* are known to have physiological dormancy, i.e., the embryo has low growth potential,⁶ it is possible that fungal attack of the tests would reduce the mechanical resistance of the testa, thus promoting embryo growth.

Arredondo et al.¹⁶ found that *Rhizopus* sp. moderately breaks seed dormancy of *Thelocactus hexahedrophorus*, another cactus species from the Chihuahuan Desert. Olvera-Carrillo et al.⁸ found that seven-month-old exhumed seeds from *O. tomentosa* showed fungal hyphae penetrating the funicular envelope through the openings, favoring germination but with a

weak embryo (an embryo with low growth potential).

Conclusions

We found that *O. leucotricha* seeds inoculated with the four fungal species had higher germination than control seeds, similar to findings for *O. streptacantha*.² These results show that fungi play an important role in breaking seed dormancy of *Opuntia* species; they contribute to understanding germination biology of cactus species, opening new insights regarding the effect of fungi on breaking seed dormancy of arid and semiarid plants.

Acknowledgements

Enrique Jurado kindly gave excellent suggestions that greatly benefited the manuscript. Financial support was obtained from SEMARNAT (2006-S-3406 to J.F.) and SAGARPA (2004-COI-216 to J.F.J-B.).

References

- Morpeth DR, Hall AM. Microbial enhancement of seed germination in *Rosa corymbifera* 'Laxa'. *Seed Sci Res* 2000; 10:489-94.
- Delgado-Sánchez P, Ortega-Amaro M, Jimenez-Bremont JF, Flores J. Are fungi important for breaking seed dormancy in desert species? Experimental evidence in *Opuntia streptacantha* (Cactaceae). *Plant Biol* 2010; 13:154-9.

- Wang X, Felker P, Paterson A, Mizrahi Y, Nerd A, Mondragón-Jacobo C. Cross-hybridization and seed germination in *Opuntia* species. *J Prof Assoc Cactus Develop* 1996; 1:49-60.
- Mandujano MC, Montaña C, Rojas-Aréchiga M. Breaking seed dormancy in *Opuntia rastrojera* from the Chihuahuan desert. *J Arid Environ* 2005; 62:15-21.
- Mandujano MC, Golubov J, Rojas-Aréchiga M. Efecto del ácido giberélico en la germinación de tres especies del género *Opuntia* (Cactaceae) del Desierto Chihuahuense. *Cact Suc Mex* 2007; 52:46-52.
- Orozco-Segovia A, Márquez-Guzmán J, Sánchez-Coronado ME, Gamboa De Buen A, Baskin JM, Baskin CC. Seed anatomy and water uptake in relation to seed dormancy in *Opuntia tomentosa* (Cactaceae, Opuntioideae). *Ann Bot* 2007; 99:581-92.
- Ochoa-Alfaro AE, Silva-Ortega CO, Becerra-Flora A, Flores J, Jiménez-Bremont JF. Effect of salt stress, proline and polyamines on seed germination of *Opuntia streptacantha*. *J Prof Assoc Cactus Develop* 2008; 10:56-70.
- Olvera-Carrillo Y, Márquez-Guzmán J, Sánchez-Coronado ME, Barradas VL, Rincón E, Orozco-Segovia A. Effect of burial on the germination of *Opuntia tomentosa*'s (Cactaceae, Opuntioideae) seeds. *J Arid Environ* 2009; 73:421-7.
- Baskin CC, Baskin JM. *Seeds: Ecology, Biogeography and Evolution of Dormancy and Germination*. San Diego, USA Academic Press 2001.
- Jurado E, Moles AT. Germination deferment strategies. In: Nicolás G, Bradford KJ, Côme D, Curie M, Pritchard HW, Eds. *The Biology of Seeds: Recent Research Advances*. Wallingford UK. CABI Publishing 2003; 381-8.
- Jurado E, Flores J. Is seed dormancy under environmental control or bound to plant traits? *J Veg Sci* 2005; 16:559-64.
- Rojas-Aréchiga M, Vázquez-Yanes C. Cactus seed germination: A review. *J Arid Environ* 2000; 44:85-104.
- Flores J, Arredondo A, Jurado E. Comparative seed germination in species of *Turbinicarpus*: an endangered cacti genus. *Nat Areas J* 2005; 25:183-7.
- Flores J, Jurado E, Arredondo A. Effect of light on germination of seeds of Cactaceae from the Chihuahuan Desert, México. *Seed Sci Res* 2006; 16:149-55.
- Flores J, Jurado E, Jiménez-Bremont JF. Breaking seed dormancy in specially protected *Turbinicarpus lophophoroides* and *Turbinicarpus pseudopectinatus* (Cactaceae). *Plant Species Biol* 2008; 23:44-7.
- Arredondo A, Rocha-Ruiz A, Flores J. Rompimiento de Latencia en Semillas de Cinco Especies de Cactáceas del Desierto Chihuahuense. Folleto Técnico No. 32. Campo Experimental San Luis. San Luis Potosí, México. CIRNE-INIFAP 2007.