

Published in final edited form as:

Int J Eat Disord. 2011 May ; 44(4): . doi:10.1002/eat.20865.

Diet and Physical Activity in Women Recovered from Anorexia Nervosa: A Pilot Study

Jocilyn E. Dellava, PhD^{1,2,*}, Robert M. Hamer, PhD^{1,3}, Akansha Kanodia¹, Mae Lynn Reyes-Rodríguez, PhD¹, and Cynthia M. Bulik, PhD^{1,4}

¹Department of Psychiatry, University of North Carolina at Chapel Hill

²Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University

³Department of Biostatistics, University of North Carolina at Chapel Hill

⁴ Department of Nutrition, University of North Carolina at Chapel Hill

Abstract

Objective—After recovery, women with anorexia nervosa (AN) tend to maintain lower body mass indices (BMI) than women in the general population. Reasons for this are unknown as little is known about diet, food choices, physical activity levels (PAL), and reasons for exercise in women recovered from AN.

Method—Diet, reasons for food choice, PAL, and reasons for exercise were measured in an exploratory study of 15 women recovered from AN and 22 women with no eating disorder history.

Results—In these hypotheses generating analyses, mean BMI in recovered women was numerically lower than control women [21.4kg/m² (2.0) and 23.6 kg/m² (4.4); respectively ($p < 0.06$)]. Recovered women were more likely to base food choice on health benefits ($p < 0.04$) compared with control women.

Discussion—Pathological behaviors that are pathognomonic of AN may resolve into healthy food and activity choices that help maintain BMIs lower in the healthy range in recovered individuals.

Keywords

Anorexia nervosa; Recovery; Body mass index; Diet; Physical activity; Reasons for food choice

Introduction

Women recovered from anorexia nervosa (AN) appear to maintain a lower body mass index (BMI) compared with comparably aged women with no history of an eating disorder.¹⁻⁴ Diet and physical activity level (PAL) are two factors that are important in the attainment and maintenance of BMIs; however, little is known about the long-term diet and PAL of individuals who have recovered from AN.

Dietary intake

Several studies suggest that dietary fat intake comprises a lower percent of total energy intake⁵⁻¹¹ and dietary protein a higher percent of total energy intake^{5, 10, 12} in women with

*Correspondence to: Dr. Dellava, Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, PO Box 980126 MCV; Richmond, VA 23298; Voice: (804) 828-3782 Fax: (804) 828-1471; jdellava@vcu.edu.

None of the authors have any financial or conflicts of interest to disclose.

AN compared with women without an eating disorder. Both lower^{5, 13} and higher^{6, 8} percent of total calories from carbohydrates were reported in individuals with AN relative to those without an eating disorder.

Few studies have addressed diet in women after recovery from AN. In a study of 14 women in treatment for AN, macronutrient composition of the diet did not change from intake to one year follow-up, at which time the participants were still below 85% of ideal body weight, and the women's diets were not compared to diets of individuals without a history of an eating disorder.¹⁴ In a separate study, following hospitalization for AN, 29 women deemed to have "successful" outcomes consumed a higher percent of energy from fat than 12 women considered "treatment failures."¹⁵ This study was conducted less than one year post-hospitalization and diet was not compared with diets of control women nor to the patient's diet while acutely ill with AN.¹⁵ Therefore it remains unknown if dietary intake differs in women who are long-term recovered from AN compared with women without a history of an eating disorder. A related heretofore unaddressed question, concerns factors that influence food choice in women recovered from AN. Although food choice might be directly related to macronutrient content of food items, it could also reflect alternative motivations (i.e., eating less trans fat, less saturated fat, or eating fewer processed food items), which might have health benefits.

Physical activity

PAL is another possible mechanism that could contribute to low BMIs in women recovered from AN. In several studies, PAL was higher in individuals who were acutely ill with AN compared with control participants.^{16, 19} PAL might remain relatively stable during initial recovery from AN²⁰ and persist after recovery.¹⁶

Women with AN report different reasons for exercise than women without eating disorders. In a study of 59 women with AN and 53 women without an eating disorder, using the Reasons for Exercise Inventory (REI), women with AN were less likely to cite health/fitness and more likely to cite negative affect regulation as reasons for exercise.²¹ However, when applying a different set of subscales to the REI, 102 women with an eating disorder were more likely to cite weight control, attractiveness, body tone, and mood and less likely to cite enjoyment as reasons for exercise compared with 184 women without an eating disorder.²²

The reasons for increased PAL during acute AN are not entirely clear. It is often assumed that increased PAL is consciously motivated to increase energy expenditure with a resolute goal of facilitating weight loss or maintenance of low weight, but alternative explanations have also been posited including thermoregulation,^{23, 24} anxiety,^{25, 26} and mood regulation.^{21, 22} If recovery is characterized by improved thermoregulation, and reduced anxiety and negative affect these motivators could cease to be operative and reasons for physical activity after recovery could differ from reasons for physical activity during the acute phase of the illness. Therefore it is important to explore reasons for exercise in women who are long-term recovered from AN.

Diet, reasons for food choice, PAL, and reasons for exercise are all items which might independently be important to BMI maintenance along with the long-term health of women recovered from AN, but to the best of the authors' knowledge, no study has examined these items in recovered women. The aim of the current study is to further understand characteristics of women recovered from AN for three or more years by determining whether they differ from control women on energy intake, dietary macronutrient intake, reasons for food choice, PAL, and reasons for exercise.

Methods

Participants

All recovered and control participants were recruited from university list-serves and advertisements. Participants were women between the ages of 18 and 65 years with no current medical conditions which would influence weight, diet, or PAL. All women in the recovered group met lifetime diagnostic criteria for AN, including amenorrhea, and had not met diagnostic criteria for any eating disorder during the three years prior to the study. Women also reported not having BMIs below 17.5 kg/m² nor engaging in any binge eating or purging in the last three years. Control women reported never having met diagnostic criteria for any eating disorder and not having a BMI below 17.5 kg/m² in the three years prior to the study. Participants were not excluded based on maximum BMI. All but one participant were non-smokers. Following initial screening, three potential recovered women and 10 potential control participants declined participation in this study.

Complete data for questionnaires were available for 15 women recovered from AN and 22 control women. Of these, two control women did not complete food journals and two control women did not complete a weekend day food journal and were removed from dietary analyses yielding a final sample of 15 women recovered from AN and 18 control women for dietary analyses. Sufficient hours of wear time for the physical activity monitors were not logged on three week days and one weekend day for two women recovered from AN and five control women; these women were removed from physical activity analyses yielding complete physical activity data for 13 women recovered from AN and 15 control women. One control woman was a BMI outlier, but left in analyses given the small sample size and removal of the participant did not impact results.

Assessments

Eating disorder diagnosis—Module H of Structured Clinical Interview for DSM-IV Axis I Disorders - Patient Edition (SCID-I/P)²⁷ was used to establish presence or absence of current and lifetime eating disorders. The SCID-I/P is a semi-structured interview which was conducted over the telephone by a trained clinical interviewer.

Anthropometric data—Current height was measured in meters and weight was measured in kilograms, on a calibrated Scale-Tronix 50220 scale, while the participant was wearing light clothing and socks. Current, measured, height and weight were used to calculate BMI.

Dietary intake—Women were instructed to keep detailed food journals for three weekdays and one weekend day. Women were asked to indicate items consumed, brand names, and amounts of each item consumed. For combination food items, such as salads and sandwiches, women were asked to list each component separately and to give precise amount of each item consumed. Women were instructed to use common household items as a guide to how much was consumed (i.e., a fist is approximately one cup) and asked to measure items when possible. All data were entered into the Nutritionist Pro SoftwareTM Version 4.3.0.²⁸ This program generated the total kilocalorie (kcal) and macronutrient data used for analyses.

Food Choice Questionnaire—The food choice questionnaire (FCQ) was used to determine what factors impact the participants' food choices. The FCQ is a 36-item self report questionnaire that has the following subscales: convenience, ethical concern, familiarity, health, mood, natural content, price, sensory, and weight control and is used to assess different factors that influence food preferences.²⁹ The FCQ has been shown to have test-retest reliability and internal consistency; Cronbach's α scores for each subscale are as

follows: convenience=0.84; ethical concern=0.74; familiarity=0.72; health =0.81; mood=0.83; natural content=0.86; price=0.83; sensory appeal=0.72; and weight control=0.85.²⁹

Physical activity—Participants were asked to wear an Actigraph™ activity monitor for three weekdays and one weekend day. The epoch length of the monitors was set to one minute and 60 consecutive zero counts was considered indicative of the monitor not being worn. In order to be considered worn for a day the participant needed to log ten hours of wear time. Participants were instructed to not wear the monitors while swimming or showering. Data reduction programs were used to determine wear time and number of counts per day; these were based on programs used by the National Health and Nutrition Examination Survey.³⁰

Reasons for Exercise Inventory—The REI was used to assess reasons for exercise. This questionnaire contains 24 questions and subscales to assess weight control, fitness, mood, health, attractiveness, enjoyment, and tone.³¹ Cronbach's scores for each subscale are as follows: weight control=0.81; fitness=0.71; mood=0.79; health=0.73; attractiveness=0.78; enjoyment=0.67; and tone=0.74.³¹

Statistical Analyses

All data were cleaned and analyzed in SAS/STAT® 9.2.³² Means and standard deviations are presented for all continuous measures. Student's t-tests were used to determine differences between group for total energy intake, percent total intake from fat, percent total intake from carbohydrate, percent total intake from protein, subscales from the FCQ, total physical activity counts, and subscales from the REI. A general linear model was used to determine if group differences in BMI existed after controlling for age. Age was included as a covariate given the broad age range of participants in the current study and the established relation between age and BMI.³³

Proc TTEST was used for student's t-tests and Proc GLM was used for the general linear model. All significance tests were two-tailed and p -value of <0.05 was considered significant. Given that this was a pilot hypothesis-generating study and our statistical power was low we did not correct for multiple testing.

Results

Age did not differ significantly between recovered and control women [32.5 (14.3) years vs. 28.9 (10.7) years, respectively]. Recovered women were recovered for a mean of 12.9 (9.0) years; (range: 3.0 years to 32.0 years) (data not shown).

BMI

BMI was over two units lower in recovered women compared with control women [21.4 kg/m² (2.0) vs. 23.6 kg/m² (4.4), respectively]. However, when controlling for age the small group comparison was not significantly different ($p<0.06$). Of the women in this study, one recovered woman (6.7%) and four control women (18.2%) had a BMI in the overweight range (25.0 BMI <30.0) and no recovered women (0.0%) and two control women (9.1%) had a BMI in the obese range (BMI ≥ 30.0).

Energy Intake and Macronutrient Content of Diet

Table 1 presents energy intake and macronutrient breakdown of reported diet. Reported macronutrient intake for both groups is within suggested ranges and groups did not differ on total energy intake or macronutrient composition of the diet.

Reasons for Food Choice

Reasons for food choice are presented in Table 2. Women recovered from AN were significantly more likely than control women to report selecting food items based on perceived health benefits ($p < 0.04$). Between group differences were not found for the other scales of this questionnaire.

Physical Activity and Reasons for Exercise

Total daily activity and reasons for exercise did not differ between groups as shown in Table 3.

Discussion

Overall, we found that energy intake, macronutrient composition of the diet, and PAL did not significantly differ in women recovered from AN compared with women without an eating disorder history. This was, however, a pilot investigation, and we were underpowered to find between group differences for the variables observed in this study.

This pilot study provides a preliminary glimpse into eating and activity behaviors of individuals who have recovered from AN. Our data suggest that after recovery, women with histories of AN focus on health benefits of foods more than non eating disordered peers, although overall energy intake did not differ between the groups. Women recovered from AN had BMIs over two units lower than women without a history of an eating disorder and a numerically lower percent of recovered women were considered overweight or obese. If our observations regarding lower BMIs in the presence of non-significantly different energy intake and PAL were to be replicated in a larger sample, there are several alternative biological, genetic and behavioral explanations for lower BMIs that would be worthy of further exploration. If biologic or genetic mechanisms are the operative factors in the maintenance of lower BMIs post-recovery, it would also be important to determine whether the same biological or genetic mechanisms might also have contributed to the ability to attain low BMIs during acute AN in the first place.

Considering potentially relevant biological mechanisms, previous research indicated women recovered from AN have higher fat oxidation than women without a history of an eating disorder.³⁴ In Pima Indians, low fat oxidation is associated with weight gain,³⁵ and obese individuals who lost weight have lower fat oxidation than individuals of the same BMI who were never obese.³⁶ Therefore it follows that high fat oxidation, present in women recovered from AN, might be associated with an ability to attain and maintain a lower BMI.

A second biologically plausible explanation for the ability to maintain a lower healthy BMI might result from differences in uncoupling proteins (UCPs). UCPs are positively associated with resting metabolic rate and negatively associated with fat mass;³⁷⁻⁴⁰ higher than expected resting metabolic rate could account for lower BMIs. Lower percent fat mass would contribute to higher resting metabolic rate and therefore also contribute to lower BMIs. To add further support to this notion, obese women who are “diet responsive” have higher mRNA expression of UCP3 compared with obese women who are “diet resistant”.³⁸ In addition, individuals with AN might be more likely to have a mutation in UCPs 2 and 3⁴¹ and this could result in an ability to attain and maintain a lower BMI compared with their peers in the general population. It is also possible that AN is associated with other genetic variations that influence the development and maintenance of a low BMI prior to, during, and following acute AN.

Third, non-exercise activity thermogenesis (NEAT) may be higher in recovered individuals than in individuals without a history of an eating disorder. NEAT shows inter-person

variation by up to 2000 kcal/day⁴² and would include such factors as fidgeting and sitting as opposed to standing, items unable to be measured by the current study. Therefore differences in NEAT could contribute to the differences in BMI.

An alternative, behavioral explanation for the ability of women recovered from AN to maintain lower BMIs than women without a history of an eating disorder could result from subtle and sustained non-statistically significant differences in energy balance (energy intake relative to energy expenditure). For example, albeit in this small study, both energy intake and PAL were slightly higher in women recovered from AN compared with control women. Despite the higher values for both energy intake and PAL, even a sustained 50 kcal/day difference in energy balance would result in drastic differences in BMI over time. Therefore, even with apparent similarities in energy intake and PAL, an energy deficit might be present in recovered women and contribute to lower BMIs.

In addition, it has been hypothesized that the pathology of AN may resolve into healthy behaviors which may be protective against obesity.⁴³ In the current study, recovered women did indicate a higher preference for food choice selection based on health benefits compared with control women. Selecting food items based on perceived health benefits could result in lower consumption of unhealthy, unnecessary components of food items (i.e., trans fat and added sugars). Higher importance of food selection based on health benefits in recovered women might serve as an indication that individuals who are able to recover from AN develop healthy eating and lifestyle habits that may be protective against obesity and obesity related chronic diseases.

Limitations

Due to small sample size, both the significant and non-significant results must be interpreted with caution. First, the overall sample size for this study was small and was further reduced by the ability to obtain useable diet and physical activity data from those data collected in this study. Therefore this study was underpowered to find significant differences between groups. For example, this study had approximately 39% power to find a 2.2 kg/m² difference in BMI. Although mean values for energy intake and percent macronutrient composition of the diets did not differ significantly between groups, it should be noted that more control than recovered participants were removed from analyses due to incomplete food journals or physical activity data. We are unable to determine how this may have impacted our results. Second, it is probable that participants did not accurately estimate dietary consumption. It is also possible that women altered diet and physical activity habits on the days that records were kept. It is unknown whether accuracy and alterations would differ between groups. Third, there are inherent short-comings in using activity monitors for measurement of PAL including failure of monitors to detect motion in all directions, failure of monitors to detect certain types of activity (i.e., lifting weights), failure to detect NEAT, and failure to detect other factors associated with activity (i.e., running with weights or while pushing a stroller). In addition, swimming was unable to be included in total counts; few participants reported swimming during the data collection period. Fourth, it is possible that women who were satisfied with diet and physical activity habits were more likely to respond to advertisements for this study. Therefore, results of this study might not generalize to all women who are recovering from AN; especially those who are less comfortable with their current diet and physical activity patterns. Fifth, results of this study might not generalize to other populations. Participants in this study were highly educated, largely Caucasian, and largely non-smokers.

Conclusion

The current study provides additional evidence to support the hypothesis that pathologic behaviors of AN resolve into healthy and adaptive behaviors after recovery. The preliminary results of this exploratory study encourage future, larger, investigations examining differences in diet, PAL, and overall energy balance using more precise methodology (i.e., multiple methods to assess usual diet, multi-axial and water-proof activity monitors accompanied by activity logs to assess PAL, devices that can estimate NEAT, and doubly labeled water to assess total energy expenditure), in order to help better understand factors associated with recovered women's tendency to maintain lower BMIs. Further exploration of biological mechanisms and genetic variants which may contribute to lower BMIs in recovered women would also be useful and provide a better understanding of long-term health in individuals recovered from AN.

Acknowledgments

We express our gratitude to Millie Maxwell, Chrysecolla Taico, Stephanie Zerwas, and the participants who made this research possible. No conflicts of interest have been declared.

Funding Source: Dr. Dellava was supported by T32MH076694-03 (PI: Bulik) and T32MH20030 (PI: Michael C. Neale). This project was partially supported by a Student Research award from the Academy for Eating Disorders (AED). We also thank the Epidemiology Core at the University of North Carolina at Chapel Hill (NIH DK56350).

References

1. Bulik CM, Sullivan PF, Fear JL, Pickering A. Outcome of anorexia nervosa: eating attitudes, personality, and parental bonding. *Int J Eat Disord.* 2000; 28:139–147. [PubMed: 10897075]
2. Misra M, Prabhakaran R, Miller KK, Goldstein MA, Mickley D, Clauss L, et al. Weight gain and restoration of menses as predictors of bone mineral density change in adolescent girls with anorexia nervosa-1. *J Clin Endocrinol Metab.* 2008; 93:1231–1237. [PubMed: 18089702]
3. Wagner A, Aizenstein H, Venkatraman VK, Fudge J, May JC, Mazurkewicz L, et al. Altered reward processing in women recovered from anorexia nervosa. *Am J Psychiatry.* 2007; 164:1842–1849. [PubMed: 18056239]
4. Ehrlich S, Salbach-Andrae H, Weiss D, Burghardt R, Goldhahn K, Craciun EM, et al. S100B in underweight and weight-recovered patients with anorexia nervosa. *Psychoneuroendocrinology.* 2008; 33:782–788. [PubMed: 18423888]
5. Hetherington MM, Rolls BJ. Eating behavior in eating disorders: response to preloads. *Physiol Behav.* 1991; 50:101–108. [PubMed: 1946701]
6. Misra M, Tsai P, Anderson EJ, Hubbard JL, Gallagher K, Soyka LA, et al. Nutrient intake in community-dwelling adolescent girls with anorexia nervosa and in healthy adolescents. *Am J Clin Nutr.* 2006; 84:698–706. [PubMed: 17023694]
7. Fernstrom MH, Weltzin TE, Neuberger S, Srinivasagam N, Kaye WH. Twenty-four-hour food intake in patients with anorexia nervosa and in healthy control subjects. *Biol Psychiatry.* 1994; 36:696–702. [PubMed: 7880939]
8. Hadigan CM, Anderson EJ, Miller KK, Hubbard JL, Herzog DB, Klibanski A, et al. Assessment of macronutrient and micronutrient intake in women with anorexia nervosa. *Int J Eat Disord.* 2000; 28:284–292. [PubMed: 10942914]
9. Rolls BJ, Andersen AE, Moran TH, McNelis AL, Baier HC, Fedoroff IC. Food intake, hunger, and satiety after preloads in women with eating disorders. *Am J Clin Nutr.* 1992; 55:1093–1103. [PubMed: 1595580]
10. Beumont PJ, Chambers TL, Rouse L, Abraham SF. The diet composition and nutritional knowledge of patients with anorexia nervosa. *J Hum Nutr.* 1981; 35:265–273. [PubMed: 7276554]
11. Affenito SG, Dohm FA, Crawford PB, Daniels SR, Striegel-Moore RH. Macronutrient intake in anorexia nervosa: The National Heart, Lung, and Blood Institute Growth and Health Study. *J Pediatr.* 2002; 141:701–705. [PubMed: 12410201]

12. Thibault L, Roberge AG. The nutritional status of subjects with anorexia nervosa. *Int J Vitam Nutr Res.* 1987; 57:447–452. [PubMed: 3440720]
13. Soh NL, Touyz S, Dobbins T, Surgenor L, Clarke S, Kohn M, et al. Cross-cultural differences in the macronutrient intakes of women with anorexia nervosa in Australia and Singapore. *Eur Eat Disord Rev.* 2008; 16:427–435. [PubMed: 18567064]
14. Nova E, Varela P, Lopez-Vidriero I, Toro O, Cenal MJ, Casas J, et al. A one-year follow-up study in anorexia nervosa. Dietary pattern and anthropometrical evolution. *Eur J Clin Nutr.* 2001; 55:547–554. [PubMed: 11464228]
15. Schebendach JE, Mayer LE, Devlin MJ, Attia E, Contento IR, Wolf RL, et al. Dietary energy density and diet variety as predictors of outcome in anorexia nervosa. *Am J Clin Nutr.* 2008; 87:810–816. [PubMed: 18400701]
16. Kron L, Katz JL, Gorzynski G, Weiner H. Hyperactivity in anorexia nervosa: A fundamental clinical feature. *Compr Psychiatry.* 1978; 19:433–440. [PubMed: 679677]
17. Davis C, Kennedy SH, Ravelski E, Dionne M. The role of physical activity in the development and maintenance of eating disorders. *Psychol Med.* 1994; 24:957–967. [PubMed: 7892363]
18. Casper RC, Schoeller DA, Kushner R, Hnilicka J, Gold ST. Total daily energy expenditure and activity level in anorexia nervosa. *Am J Clin Nutr.* 1991; 53:1143–1150. [PubMed: 1850575]
19. Davis C, Katzman DK, Kaptein S, Kirsh C, Brewer H, Kalmbach K, et al. The prevalence of high-level exercise in the eating disorders: etiological implications. *Compr Psychiatry.* 1997; 38:321–326. [PubMed: 9406737]
20. Falk JR, Halmi KA, Tryon WW. Activity measures in anorexia nervosa. *Arch Gen Psychiatry.* 1985; 42:811–814. [PubMed: 4015326]
21. Bratland-Sanda S, Sundgot-Borgen J, Ro O, Rosenvinge JH, Hoffart A, Martinsen EW. “I’m not physically active - I only go for walks”: physical activity in patients with longstanding eating disorders. *Int J Eat Disord.* 2010; 43:88–92. [PubMed: 19728373]
22. Mond JM, Calogero RM. Excessive exercise in eating disorder patients and in healthy women. *Aust N Z J Psychiatry.* 2009; 43:227–234. [PubMed: 19221911]
23. Gutierrez E, Vazquez R, Boakes RA. Activity-based anorexia: ambient temperature has been a neglected factor. *Psychon Bull Rev.* 2002; 9:239–249. [PubMed: 12120785]
24. Gutierrez E, Cerrato M, Carrera O, Vazquez R. Heat reversal of activity-based anorexia: implications for the treatment of anorexia nervosa. *Int J Eat Disord.* 2008; 41:594–601. [PubMed: 18446833]
25. Norris R, Carroll D, Cochrane R. The effects of physical activity and exercise training on psychological stress and well-being in an adolescent population. *J Psychosom Res.* 1992; 36:55–65. [PubMed: 1538350]
26. Sexton H, Maere A, Dahl NH. Exercise intensity and reduction in neurotic symptoms. A controlled follow-up study. *Acta Psychiatr Scand.* 1989; 80:231–235. [PubMed: 2816466]
27. First, MB.; Spitzer, R.; Gibbon, M.; Williams, JB. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition (SCID-I/P). New York: Biometrics Research, New York State Psychiatric Institute; 2002.
28. Axxya Systems. Nutritionist Pro®: Version 4.3.0. Stafford, TX: Axxya Systems; 2010.
29. Steptoe A, Pollard TM, Wardle J. Development of a measure of the motives underlying the selection of food: the food choice questionnaire. *Appetite.* 1995; 25:267–284. [PubMed: 8746966]
30. Center for Disease Control. National Health and Nutrition Examination Survey. 2010 cited; Available from: <http://www.cdc.gov/nchs/nhanes.htm>
31. Silberstein LR, Striegel-Moore RH, Timko C, Rodin J. Behavioral and psychological implications of body dissatisfaction: Do men and women differ? *Sex Roles.* 1988; 19:219–232.
32. SAS Institute Inc. SAS/STAT® Software: Version 9. Cary, NC: SAS Institute, Inc; 2004.
33. Chambers JA, Swanson V. A health assessment tool for multiple risk factors for obesity: age and sex differences in the prediction of body mass index. *Br J Nutr.* 2010:1–10.
34. Dellava JE, Policastro P, Hoffman DJ. Energy metabolism and body composition in long-term recovery from anorexia nervosa. *Int J Eat Disord.* 2009; 42:415–421. [PubMed: 19107831]

35. Zurlo F, Lillioja S, Esposito-Del Puente A, Nyomba BL, Raz I, Saad MF, et al. Low ratio of fat to carbohydrate oxidation as predictor of weight gain: study of 24-h RQ. *Am J Physiol.* 1990; 259:E650–657. [PubMed: 2240203]
36. Filozof CM, Murua C, Sanchez MP, Brailovsky C, Perman M, Gonzalez CD, et al. Low plasma leptin concentration and low rates of fat oxidation in weight-stable post-obese subjects. *Obes Res.* 2000; 8:205–210. [PubMed: 10832762]
37. Bouchard C, Perusse L, Chagnon YC, Warden C, Ricquier D. Linkage between markers in the vicinity of the uncoupling protein 2 gene and resting metabolic rate in humans. *Hum Mol Genet.* 1997; 6:1887–1889. [PubMed: 9302267]
38. Harper ME, Dent R, Monemdjou S, Bezaire V, Van Wyck L, Wells G, et al. Decreased mitochondrial proton leak and reduced expression of uncoupling protein 3 in skeletal muscle of obese diet-resistant women. *Diabetes.* 2002; 51:2459–2466. [PubMed: 12145158]
39. Rolfe DF, Newman JM, Buckingham JA, Clark MG, Brand MD. Contribution of mitochondrial proton leak to respiration rate in working skeletal muscle and liver and to SMR. *Am J Physiol.* 1999; 276:C692–699. [PubMed: 10069997]
40. Mori S, Satou M, Kanazawa S, Yoshizuka N, Hase T, Tokimitsu I, et al. Body fat mass reduction and up-regulation of uncoupling protein by novel lipolysis-promoting plant extract. *Int J Biol Sci.* 2009; 5:311–318. [PubMed: 19421341]
41. Campbell DA, Sundaramurthy D, Gordon D, Markham AF, Pieri LF. Association between a marker in the UCP-2/UCP-3 gene cluster and genetic susceptibility to anorexia nervosa. *Mol Psychiatry.* 1999; 4:68–70. [PubMed: 10089012]
42. Levine JA. Nonexercise activity thermogenesis--liberating the life-force. *J Intern Med.* 2007; 262:273–287. [PubMed: 17697152]
43. Gendall K, Bulik C. The long term biological consequences of anorexia nervosa. *Curr Nutr and Food Sci.* 2005; 1:87–96.

Table 1

Average daily dietary intake of women recovered from anorexia nervosa (AN) and control women.

Macronutrient	Recovered AN	Control	<i>p</i>-value
	Mean (SD)	Mean (SD)	
Total kilocalories	1942 (543)	1915 (460)	0.89
Percent Fat	29.5 (7.1)	30.1 (6.5)	0.80
Percent Protein	14.3 (2.9)	14.0 (1.9)	0.69
Percent Carbohydrate	54.1 (9.0)	54.2 (8.1)	0.97

Note: Percentages do not add to 100 due to a small amount of alcohol consumption not included.

SD, standard deviation

Table 2

Reasons for food choice in women recovered from anorexia nervosa (AN) and control women.

Reason food choice	Recovered AN	Control	<i>p</i> -value
	Mean (SD)	Mean (SD)	
Convenience	14.5 (3.1)	14.5 (3.2)	1.00
Ethical Concern	5.2 (1.6)	5.1 (2.2)	0.82
Familiarity	5.9 (2.2)	6.1 (2.2)	0.79
Health	19.3 (3.2)	16.9 (3.4)	0.04
Mood	13.1 (3.0)	11.4 (3.3)	0.11
Natural Content	5.9 (2.6)	6.8 (2.7)	0.35
Price	8.1 (2.3)	8.4 (2.2)	0.70
Sensory	10.7 (2.6)	10.8 (2.2)	0.85
Weight Control	7.9 (2.6)	7.4 (2.2)	0.58

SD, standard deviation

Table 3

Average daily total activity and reasons for exercise in women recovered from anorexia nervosa (AN) and control women. Means and standard deviations (SD) are presented.

	Recovered AN	Control	<i>p</i> -value
Activity	Mean (SD)	Mean (SD)	
Counts	91762 (34575)	85397 (25070)	0.58
Reason for exercise	Mean (SD)	Mean (SD)	
Weight	14.5 (4.0)	14.1 (5.6)	0.82
Fitness	20.1 (5.2)	21.3 (3.5)	0.42
Mood	19.3 (5.2)	18.1 (5.3)	0.50
Health	22.0 (4.4)	23.5 (3.3)	0.26
Attractiveness	12.9 (4.9)	13.9 (5.3)	0.58
Enjoyment	10.9 (5.4)	11.0 (4.9)	0.96
Tone	11.3 (4.1)	13.1 (4.7)	0.24