



Published in final edited form as:

*Psychol Med.* 2011 August ; 41(8): 1775–1783. doi:10.1017/S0033291710002345.

## Trajectories of Marijuana Use and Psychological Adjustment Among Urban African American and Puerto Rican Women

Kerstin Pahl, Ph.D.<sup>\*</sup>, Judith S. Brook, Ed.D., and Jonathan Koppel, M.A.

Department of Psychiatry, New York University School of Medicine

### Abstract

**Background**—The current longitudinal study examined the developmental patterns of marijuana use, as well as their relationship with subsequent psychological adjustment in a community-based sample of urban African American and Puerto Rican women.

**Method**—Participants were interviewed five times over a period ranging from adolescence (mean age 14.0 years) to adulthood (mean age 32.5). Outcome measures included depressive symptoms, anger/hostility, as well as the presence of a substance use disorder (abuse/dependence).

**Results**—Three distinct trajectories of marijuana use were identified: nonusers, increasers, and quitters. Increasers reported higher levels of depressive symptoms and anger/hostility than did nonusers, and were more likely to meet criteria for a substance use disorder at age 32.5 years.

**Conclusions**—Findings indicated that early-starting long-term use of marijuana is associated with psychological maladjustment among women. Prevention efforts should emphasize the long-term cost associated with marijuana use, and that the best psychological health is reported by those who abstain from the drug.

### Keywords

Women's Health; Psychological adjustment; Marijuana Use; Longitudinal

### Introduction

A relationship between marijuana use and reduced psychological adjustment, in the form of affective disorders and symptomatology, has been established in both cross-sectional (Degenhardt *et al.* 2001, Chen *et al.* 2002) and longitudinal (Patton *et al.* 2002 Friedman *et al.*, 2004) research. The majority of the research suggests that marijuana use is negatively associated with measures of psychological adjustment (for exceptions see Shedler & Block, 1990; Galaif *et al.* 1998).

Recent longitudinal research, which has focused on developmental patterns of marijuana use from adolescence to young adulthood, has found that differential trajectories of marijuana use are related to different levels of adjustment in young adulthood (Brown *et al.* 2004; Ellickson *et al.* 2004; Flory *et al.* 2004; Windle & Wiesner, 2004; Schulenberg *et al.* 2005). Generally, usage patterns characterized by earlier onset, longer duration, and higher frequency of marijuana use are associated with less desirable outcomes.

---

Address for correspondence: Dr. Kerstin Pahl, Department of Psychiatry, New York University School of Medicine, 215 Lexington Avenue, 15<sup>th</sup> Floor, New York, NY 10016. kerstin.pahl@nyumc.org.

#### Declaration of Interest

None.

Although marijuana use is more prevalent in men than in women (Stinson *et al.* 2006), the areas of psychological adjustment most often affected by marijuana use are those on which women generally do worse than men. For example, women exhibit a higher prevalence of mood disorders (Hasin *et al.* 2005; Kessler *et al.* 2005) and higher sub-clinical levels of depression and distress than their male counterparts (McDonough & Walters, 2001; Denton *et al.* 2004). Marijuana use may, therefore, have especially strong implications for psychological adjustment in women. This consideration takes on further weight from research, which has documented a stronger relationship between marijuana use and psychological adjustment in females as compared with males (Thomas, 1996; Patton *et al.* 2002; Friedman *et al.* 2004).

There are several mechanisms, which may be responsible for the link between psychological functioning and marijuana use. One explanation, postulated by Problem Behavior Theory (e.g., Jessor *et al.* 1991), may be an underlying third factor, which predisposes individuals for both problem behaviors (including substance use) and psychological maladjustment. Thus, a common cause, possibly in the form of a personality dimension, such as unconventionality, shared by “high-risk” individuals would be responsible for a broad range of adjustment problems, including drug use, deviant behaviors, and psychological maladjustment (Jessor *et al.* 1980; Donovan & Jessor 1985). Similarly, it is possible, that other confounding factors, such as family dysfunction, may underlie both marijuana use and later psychological adjustment (Fergusson & Horwood, 1997; Degenhardt *et al.* 2003).

Another possibility is that marijuana use is precipitated by lower levels of psychological adjustment (Macleod *et al.* 2004). Specifically, those suffering from psychological maladjustment may be more inclined to use marijuana, possibly in an attempt to ameliorate their psychological symptoms. However, most longitudinal research has not supported this theory of self-medication (e.g., Bovasso, 2001).

Finally, marijuana use, particularly at high levels and over a long period of time, may have a negative impact on psychological adjustment (e.g., Kandel, 1989), either directly or via its relationship with other psychosocial constructs. Research has established that marijuana use in adolescence and young adulthood is related not only to later psychological maladjustment, but also to a host of other undesirable outcomes, including lower levels of education (Fergusson *et al.* 2003; Lynskey, Coffey *et al.* 2003); reduced occupational attainment (Schuster *et al.* 2001); impaired cognitive functioning (Solowij *et al.* 2002; Pope *et al.* 2003); and failure to make a transition to adult roles and behaviors (Brook *et al.* 2002). Failing at important developmental tasks of adolescence and young adulthood (e.g., graduating from high school, obtaining employment) may give rise to psychological maladjustment in young adults (Degenhardt, *et al.* 2003).

In addition, some longitudinal research has established a link between earlier marijuana use and later substance use disorders, including alcohol and other illicit drug use disorders (Fergusson & Horwood 1997; Windle & Wiesner, 2004). As part of psychological adjustment, this research therefore assessed the relationship between trajectories of marijuana use and later substance use disorder.

To date, the majority of the research on marijuana use and related outcome variables has been conducted on samples that were primarily or exclusively of European American descent (but see Brown *et al.* 2004). However, given that marijuana use patterns, as well as the outcomes associated with disparate patterns, may differ across race and/or ethnicity (Brown *et al.* 2004), it is important to evaluate developmental patterns of marijuana use and associated outcome measures among ethnic and racial minority groups.

The current study thus had three goals: (1) to identify distinct homogenous subgroups of marijuana use from adolescence (age 14.0 years) to young adulthood (age 26.1 years) among African American and Puerto Rican women, (2) to examine potential racial/ethnic differences in marijuana usage patterns, and (3) to identify differences among trajectory groups in self-reported levels of psychological adjustment, including symptoms of depression, anger/hostility, and the presence of a substance use disorder, at mean age 32.5 years. We hypothesized that individuals in trajectories characterized by an earlier onset, longer duration, and/or higher frequency of marijuana use would report significantly higher levels of symptoms of depression and/or anger/hostility, and be at greater risk for substance use disorder than individuals in trajectory groups characterized by later onset, abstinence, shorter durations, and/or lower frequencies of marijuana use. To rule out the possibility that earlier psychological maladjustment may be underlying both marijuana usage patterns, as well as later levels of psychological adjustment, we controlled for earlier levels of psychological maladjustment. In addition, we controlled for a measure of earlier unconventionality to see whether this construct may be underlying the relationship between marijuana use patterns and psychological functioning, as suggested by Problem Behavior Theory. We also included measures representing conflict in the participants' relationships with their mothers at the beginning of the study to control for earlier levels of family dysfunction.

## Methods

### Participants

Participants in the current research were 474 females who had participated in at least three waves of a five-wave longitudinal study of African American and Puerto Rican adolescents/young adults. Data for the Time 1 (T1) sample (N = 716) were collected in 1990, Time 2 (T2, N = 654) in 1995, Time 3 (T3, N = 335) in 2000–2001, Time 4 (T4, N = 241) in 2001–2003, and Time 5 (T5, N = 382) in 2008–2009 for a study investigating tobacco use among African American and Puerto Rican young adults<sup>1</sup>. The Institutional Review Board at Mount Sinai School of Medicine approved the study's procedures for all data collections and the Institutional Review Board at New York University School of Medicine has approved the study annually since 2004.

Participants came from grades 7–10 in school districts serving the East Harlem area of New York City. Because of budget limitations, the T3 data collection did not target the entire T2 sample, but over-sampled those respondents who reported using marijuana or other illicit drugs at T2. The main reason for this strategy was to sample a sufficient number of drug users and stay within budgetary limitations. At T4, a stratified random sample of T3 participants were invited to participate in the study. Sampling was proportionate to sex and ethnicity. At T5, we sought the participation of all those who had participated at least at two previous time points. Data were collected by trained interviewers who were matched on sex and ethnicity, whenever possible.

Of the 474 female participants eligible for inclusion in this research, 39% had participated in three waves of data collection, 20% had participated in four waves, and 41% had participated in all five waves. Of these 474 females, 52.3% (n=248) were African American, 47.7% (n=226) were Puerto Rican. Their mean ages were 14.0 years (SD = 1.3) at T1, 19.1 years (SD = 1.5) at T2, 24.5 years (SD = 1.4) at T3, 26.1 years (SD = 1.4) at T4, and 32.5 years (SD = 1.4) at T5. Attrition analyses revealed that those who participated at both T1 and T5 compared to those who participated at T1 and not at T5 (1) used marijuana more

---

<sup>1</sup>Numbers reflect the female participants in the study.

frequently at T2 ( $\bar{x}_{T1, T5} = 0.58$ ;  $\bar{x}_{T1} = 0.35$ ;  $t = -2.88$ ,  $p < .05$ ), (2) reported greater unconventionality at T1 ( $\bar{x}_{T1, T5} = 10.06$ ;  $\bar{x}_{T1} = 9.56$ ;  $t = -2.05$ ,  $p < .05$ ), and (3) were younger at T1 ( $\bar{x}_{T1, T5} = 13.9$ ;  $\bar{x}_{T1} = 14.5$ ;  $t = 5.59$ ,  $p < .0001$ ). No statistically significant differences were found with regard to marijuana use at T1, T3, or T4, levels of conflict in the participants' relationships with their mothers, or T1 levels of psychological maladjustment (depression, anger/hostility) ( $p > 0.05$ ).

For determining the trajectories of marijuana use, all 474 participants were included in the analyses. In the next step, linking trajectory group membership to T5 outcomes, we only included those women who participated at T5 (N=382).<sup>2</sup>

## Measures

Marijuana use at each wave (T1 – T4) was measured by an item asking how often in the past year the respondent had used marijuana. Answering options were (0) “never,” (1) “a few times a year or less,” (2) “about once a month,” (3) “several times a month,” and (4) “once a week or more.” Parallel items were used to assess other illegal drug use (cocaine, Ecstasy) and alcohol use in the past 12 months at T5. Answering options for how often and how much alcohol was consumed in the past year ranged from (0) “None at all” to (5) “Five drinks or more every day.” Variables representing psychological adjustment in young adulthood (T5) included depression, anger/hostility, and the presence of a substance use disorder (including marijuana, other illicit drugs, and alcohol) in the past 12 months. Scales measuring depression and anger/hostility were adapted from the Hopkins Symptom Check List (HSCL) (Derogatis *et al.* 1974). The stem for all items from the HSCL read, “Over the past few years, how much were you bothered by the following?” Answering options ranged from (0) “not at all” to (4) “extremely.” Sample items for the depression and anger/hostility scales were “feeling hopeless about the future” and “temper outbursts you cannot control,” respectively. Mean scores were computed across the six items measuring depression and the five items measuring anger/hostility. Cronbach's alpha was acceptable for both scales; 0.78 for depression, and 0.68 for anger/hostility.

To control for the presence of depressive symptoms and anger/hostility at T1, we employed partial/similar measures of these constructs, which were available at T1. Depressive symptoms were measured by two items assessing 1) feelings of hopelessness and 2) depressive affect (“sad, unhappy, depressed,”  $r = 0.61$ ). Two items reflecting anger (“losing temper,” “feeling like swearing”) were combined to form a measure of anger/hostility at T1 ( $r = 0.39$ ). Earlier unconventionality was assessed with a six-item measure of tolerance of deviance/risk taking. A sample item read: “You like to live dangerously” ( $\alpha = 0.67$ ). Two measures were used to assess the degree of conflict in the women's relationships with their mothers at T1. One measure assessed the degree to which the adolescent daughter resisted the mother's control (3 items,  $\alpha = 0.77$ ), while the other measure reflected the mother's punitive behavior toward her daughter (two items,  $r = 0.42$ ). Sample items for the two measures read “You try to see what you can get away with” and “She acts cold and unfriendly when you do something she does not like,” respectively.

Presence of a substance use disorder in the past 12 months was assessed by the Mini-International Neuropsychiatric Interview (M.I.N.I.; Sheehan, *et al.* 1998). Questions assessed all seven criteria specified by DSM-IV-TR (American Psychiatric Association, 2000). A diagnosis of dependence (on marijuana, other illicit drugs, and/or alcohol) was assigned if three or more of the criteria were met. Abuse of these substances was assessed by

<sup>2</sup>Trajectory analyses were conducted with the entire sample (n=474) as well as with the reduced sample (n=382). Results did not differ appreciably.

four criteria specified by DSM-IV-TR (American Psychiatric Association, 2000). A diagnosis of abuse was assigned if one or more of the criteria were met. Marijuana, other illicit drug, and alcohol use disorders were assessed separately according to the following logic: If a participant did not meet criteria for dependence, the presence of abuse was assessed. Scoring positively for marijuana dependence, marijuana abuse, other illicit drug dependence, other illicit drug abuse, alcohol dependence, or alcohol abuse resulted in being classified as having a substance use disorder. The M.I.N.I. has demonstrated good reliability and validity (Sheehan *et al.* 1997; Otsubo *et al.* 2005).

## Data Analysis

We employed growth mixture modeling (GMM) using the MPlus software (Muthén & Muthén, 2007) to identify longitudinal trajectories of marijuana use from adolescence (mean age 14.0 years) to young adulthood (mean age 26.1 years). We established the number of trajectories using several criteria: 1) the Bayes information criterion (BIC) (Schwarz 1978; Raftery 1985) (the most parsimonious model has the smallest absolute value of the BIC, with a BIC difference of 2.0 or greater representing positive evidence in favor of the lower-scoring model; Raftery, 1995), 2) the entropy (values closer to 1 indicate better fit; McLachlan & Peel, 2000), 3) the Lo-Mendell-Rubin (2001) likelihood ratio test (LMR-LRT; Lo *et al.* 2001), 4) the parametric bootstrap likelihood ratio test (BLRT; Nylund *et al.* 2007), 5) a minimum 5% proportion in each latent class, and 6) theoretical considerations and interpretability. Using the LMR-LRT and the BLRT, a low *p* value indicates the model with one less class is rejected in favor of the estimated model.

In addition, we considered the accuracy of group classification indicated by the average modal Bayesian posterior probability (BPP) for each latent class. Average modal posterior probabilities of .70 or higher are considered sufficient to avoid classification error (Nagin, 2005).

Next, we conducted multiple and logistic regression analyses with depressive symptoms and symptoms of anger/hostility, and the presence of a substance use disorder, respectively, at T5 as the dependent variables. To address the issue of uncertainty of assignment to a latent class, we used the BPPs, rather than the class assignments, when predicting adjustment outcomes in the regression analyses. Race/ethnicity (dummy coded), age, total household income (measured at T5), education (T5), earlier measures of the respective outcomes, a measure of tolerance of deviance/risk taking, and the two measures of conflict with one's mother were included as covariates when predicting symptoms of depression and anger/hostility. When predicting the presence of a substance use disorder, we controlled for race/ethnicity, age, household income, education, past-year alcohol and other illicit drug use at age 32.5 years, T1 tolerance of deviance/risk taking, and the two measures of conflict with one's mother (T1).

## Results

### Preliminary Analyses

*T* tests comparing African American and Puerto Rican women indicated no statistically significant differences in marijuana at any time point (see Table 1) or in the age of onset of marijuana use. In addition, there were no ethnic/racial differences in levels of anger/hostility and depressive symptoms (Table 1), or in the prevalence of substance use disorders at T5 ( $\chi^2 = 0.83, p > 0.05$ ). A table of the frequencies of other illicit drug use at T5 revealed that there was no use of illegal drugs other than marijuana, ecstasy, and cocaine. In addition, only one person reported use of cocaine at T5. Therefore, the logistic regressions predicting

the presence of a substance use disorder at T5 only controlled for concurrent alcohol and ecstasy use.

### Trajectories of Marijuana Use

Given the similarities in age of onset and levels of marijuana use between African American and Puerto Rican women, we performed the trajectory analyses on the combined sample of women. We fitted the data using the categorical data option in MPlus to accommodate the ordered categorical (ordinal) nature of the data. We specified a quadratic model, because the developmental progression of marijuana use typically entails an increase of use between adolescence and young adulthood, followed by a leveling off (O'Malley *et al.* 2004). A likelihood ratio test indicated the superior fit of the quadratic vis-a-vis a linear model ( $\chi^2(3) = 52.46, p > 0.001$ ).

We fitted one-group, two-group, three-group, four-group, and five-group models, respectively. A three-group model was chosen based on the following results: it had the lowest BIC (2276.2 vs. 2279.5 for the two-group model, and 2287.2 for the four-group model); the LMR-LRT and the BLRT for the three-group model were statistically significant, compared to the two-group model; the LMR-LRT was not statistically significant when comparing the 4-group model to the 3-group model. Finally, the entropy (0.68 for the three-group model) while lower than for the two-group model (0.74), was higher than for the four-group model (0.56). All BPPs for the three-group model were above 0.75, indicating adequate classification (Nagin, 2005).

We named the three trajectory groups “nonusers” (participants who reported no or negligible marijuana use at all four times), “increasers” (participants who reported some use at age 14.0 years which continued to increase until age 24.5 years and then leveled off slightly), and “quitters” (participants who reported some marijuana use at age 14.0, rapidly increased their use until age 19.1 years, and had essentially ceased use by age 24.5 years) (see Figure 1). The occurrence probabilities were 73.6% nonusers, 17.9% increasers, and 8.5% quitters.

Including race/ethnicity as a predictor of trajectory group membership did not result in improved model fit, nor did the ethnicity dummy code reach statistical significance. Thus, no racial/ethnic differences in group membership were present, confirming our preliminary finding that there were no racial/ethnic group differences in marijuana use over time.

### Group differences in psychological adjustment

Results from the multiple regression analyses showed that increasers, compared to nonusers, displayed higher levels of depression ( $b = 1.31, SE = 0.60, t = 2.20, p < 0.05$ ) and anger/hostility ( $b = 1.40, SE = 0.38, t = 3.71, p < 0.001$ ) at age 32.5 years, controlling for age, race/ethnicity, T5 household income and education, T1 tolerance of deviance/risk taking, T1 level of conflict in the mother-daughter relationship, and T1 levels of the respective outcome. In addition, increasers also displayed higher levels of anger/hostility than quitters ( $b = 1.48, SE = 0.63, t = 2.37, p < 0.05$ ). Adjusted means for the three groups are displayed in Table 2.

A multiple logistic regression, controlling for the participant's age, race/ethnicity, T5 household income and education, T1 intolerance of deviance/risk-taking, T1 level of conflict in the mother-daughter relationship, and alcohol and ecstasy use at age 32.5 (T5) revealed that increasers were more likely than nonusers to meet criteria for a substance use disorder at age 32.5 years (AOR = 4.86, CI = 1.52–15.57). There were no statistically significant differences between quitters and nonusers, or between quitters and increasers.

## Discussion

### Trajectories of Marijuana Use

Our longitudinal study of African American and Puerto Rican females identified three trajectories of marijuana use over a period extending from adolescence (mean age = 14.0 years) to young adulthood (mean age = 26.1 years): nonusers, increasers, and quitters. Race/ethnicity was not related to marijuana trajectory group membership in the analyses, suggesting that urban African American and Puerto Rican women follow similar developmental patterns of marijuana use. Possibly, their substance use patterns were similar because these urban Puerto Rican and African American adolescents/young adults shared similar socioeconomic backgrounds, attended the same schools, and lived in the same neighborhoods. These contextual similarities may have given rise to similar patterns of substance use, including marijuana use (Brook *et al.* 2006).

Similar to other researchers, we identified a trajectory of increasers, a group of women who started using marijuana early and continued to increase their use until their mid-twenties. Though women tend to use marijuana less frequently than men (Stinson *et al.* 2006), almost 20% of these urban African American and Puerto Rican women engaged in relatively frequent use (more than once a month) into their mid-twenties, a time during which marijuana use typically declines. Future research should try to identify the risk factors for this high-risk group of women.

Women in the quitters group sharply increased their use between early and late adolescence in a manner similar to the increasers, but, by age 24.5 years, levels of use in this group had dropped to the same level as that of the nonusers. While other authors have also identified groups of decreasers (e.g., Ellickson *et al.* 2004; Windle & Wiesner, 2004; Schulenberg *et al.* 2005), this pattern of rapid increase followed by rapid decrease is somewhat different and may well be gender-specific. Possibly, women in this group stopped using marijuana because they had children. Indeed, a greater percentage of women who quit (75.0%) had a child by age 24.5 (T3), compared to women in the other two trajectory groups (42.1% among nonusers; 44.7% among increasers). Becoming a parent is a powerful incentive for ceasing substance use among women (Bachman *et al.* 2002).

### Trajectories of Marijuana Use and Later Psychological Adjustment

As predicted, among the three trajectory groups, increasers reported more depressive symptoms and anger/hostility, and were at greater risk of having a substance use disorder than nonusers at age 32.5 years. While, perhaps unexpectedly, the mean depression values reported by the quitters were also higher than those of the nonusers and increasers, this difference was not statistically significant. The lack of statistical significance may have been due to the limited statistical power, as the group of quitters for whom T5 data were available was rather small.

Increasers also reported higher levels of anger/hostility than did quitters at age 32.5. This finding is consistent with previous research which has found that the psychosocial sequelae of marijuana use do not seem to persist over time (Schulenberg *et al.*, 2005). However, in light of the modest size of the group of quitters, and the finding that the mean value of depression for the quitters was higher than those of the increasers at age 32.5, this result should be replicated before final conclusions can be drawn.

Overall, our results are in agreement with those of Ellickson *et al.* (2004) who found that trajectory classes of increased marijuana use, compared to abstainers, reported reduced psychological adjustment. However, some authors who have investigated trajectories of marijuana use and their association with psychological adjustment have not identified a link

between membership in a high-frequency marijuana use trajectory group and lower levels of psychological adjustment (e.g., Brown *et al.* 2004; Flory *et al.* 2004). Given the clear association found in this study, it is possible that the membership in a high-frequency marijuana use trajectory is associated with less psychological adjustment mainly among women. This notion is supported by research, which finds a stronger relationship between marijuana use and psychological adjustment in females than in males (Thomas, 1996; Patton *et al.* 2002; Friedman *et al.* 2004).

Our finding that increasers reported higher levels of depression and anger/hostility substance use disorders at age 32.5 than nonusers, and, in the case of anger/hostility, than quitters, clearly shows a link between early-starting continued use of marijuana and psychological adjustment in adulthood among women. This finding is supported by the fact that our analyses included variables representing earlier (T1) levels of psychological adjustment. Marijuana use in adolescence and young adulthood is associated with a number of other negative correlates and outcomes, most notably reduced functioning in age-appropriate roles (for a review, see Macleod *et al.* 2004). Failure to fulfill age-appropriate developmental tasks (e.g., graduating from high school, obtaining employment), in turn, may be linked to reduced psychological adjustment in women (Degenhardt *et al.*, 2003).

The relationship between membership in a marijuana trajectory group and depressive symptoms and anger/hostility, respectively, was also maintained despite statistical control of a variable representing an early (T1) propensity for unconventional and risk taking behavior. Thus, it is less likely that such a personality dimension explains fully the relationship between membership in a marijuana use group and later psychological adjustment in women.

Similarly, controlling for earlier levels of conflict in the relationship between the participant and her mother, a construct reflecting family dysfunction, did not render the association between marijuana use group membership and later psychological adjustment statistically insignificant. However, it should be noted that there are many other psychosocial influences, not measured in the current study, which may account for the relationship between marijuana use and women's psychological maladjustment. In addition, it is possible that a common genetic liability predisposes individuals to both marijuana use and psychological maladjustment (e.g., Sullivan *et al.* 2000).

Women who followed trajectories of high and increasing marijuana use into their mid-twenties were more likely than non-users to have a substance use disorder (including marijuana, other illicit drug, and alcohol use disorders) at age 32.5. This association was maintained despite control on concurrent (T5) use of alcohol and ecstasy. Long-term marijuana use, especially when started early, results in addiction for some users (Iversen, 2003; Haney *et al.* 2004). High levels of marijuana use have also been shown to be related to increased use of other substances, possibly due to a biochemical effect (Lynskey, Heath, *et al.* 2003; Fergusson *et al.* 2006). It is also possible that those following trajectories of high and increased use are more likely to suffer from an underlying diathesis to addiction (e.g., Agrawal *et al.* 2004; Kreek *et al.* 2005).

## Limitations

Limitations of the current study included the oversampling of participants who displayed more deviant behaviors at T2. Possibly, our results would have differed had we been able to recruit participants representing a broader spectrum of risk-taking behaviors. Another limitation was our sole reliance on self-report measures. However, a recent study by Harrison and colleagues (2007) showed that most marijuana users reported their use accurately. Furthermore, despite our attempts to control for other underlying factors which



may explain the relationship between membership in a latent class of elevated marijuana use and psychological maladjustment, our results cannot be considered causal. Finally, our study is limited by the small size of the group of quitters at T5. Nevertheless, our results regarding the differences in psychological adjustment between non-users and increasers were consistent across all three outcomes.

## Conclusions

The findings of the present longitudinal study of women by and large supported our hypotheses. Among women, longer-duration and more frequent use of marijuana was associated with poorer outcomes on measures of psychological adjustment. While the relationships between marijuana use trajectories and psychological adjustment cannot be considered causal, the fact that we chose to assess adjustment in young adulthood (measurement wave 5) and the inclusion of variables representing earlier levels of the outcomes as well as measures of unconventionality and family conflict, support the idea that marijuana use may contribute to women's psychological maladjustment over time. This relationship may well be mediated by other psychosocial factors not included in this research.

Practitioners who observe symptoms frequently reported by women, such as depressive mood and feelings of anger/hostility, are well-advised to take into account the possibility of marijuana use. Clinicians may consider screening women who present with symptoms of depression and/or anger for marijuana and other substance use.

Our findings also suggest that high-frequency marijuana use over time among women should be considered as a serious risk factor for the development of substance abuse and dependence. In treating marijuana users reporting such a pattern of long-term high-frequency use, practitioners should be aware of the associated diminished adjustment. Similarly, prevention programs should focus on the long-term psychological cost associated with marijuana use, pointing out that the best psychological health is reported by those who abstain from the drug.

## Acknowledgments

This research was supported by NIH Grant DA005702, and Research Scientist Award DA00244, both from the National Institute on Drug Abuse; and Grant 7R01CA84063 from the National Cancer Institute. The grants were awarded to Judith S. Brook.


## References

- Agrawal A, Neale MC, Prescott CA, Kendler KS. A twin study of early cannabis use and subsequent use and abuse/dependence of other illicit drugs. *Psychological Medicine*. 2004; 34:1227–1237. [PubMed: 15697049]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR*. American Psychiatric Association; Washington, DC: 2000.
- Bachman, JG.; O'Malley, PM.; Schulenberg, JE.; Johnston, LD.; Bryant, AL.; Merline, AC. *The Decline of Substance Use in Young Adulthood: Changes in Social Activities*. Lawrence Erlbaum; Mahwah, NJ: 2002.
- Bovasso GB. Cannabis abuse as risk factor for depressive symptoms. *American Journal of Psychiatry*. 2001; 158:2033–2037. [PubMed: 11729021]
- Brook JS, Adams RE, Balka EB, Johnson E. Early adolescent marijuana use: Risks for the transition to young adulthood. *Psychological Medicine*. 2002; 32:79–91. [PubMed: 11883732]
- Brook JS, Pahl K, Ning Y. Peer and parental influences on longitudinal trajectories of smoking among African Americans and Puerto Ricans. *Nicotine & Tobacco Research*. 2006; 8:639–651. [PubMed: 17008191]

- Brown TL, Flory K, Lyman DR, Leukefeld C, Clayton RR. Comparing the developmental trajectories of marijuana use of African American and Caucasian adolescents: Patterns, antecedents, and consequences. *Experimental and Clinical Psychopharmacology*. 2004; 12:47–56. [PubMed: 14769099]
- Chen CY, Wagner FA, Anthony JC. Marijuana use and the risk of major depressive episode: Epidemiological evidence from the United States National Comorbidity Survey. *Social Psychiatry and Psychiatric Epidemiology*. 2002; 37:199–206. [PubMed: 12107710]
- Degenhardt L, Hall W, Lynskey M. Alcohol, cannabis, and tobacco use among Australians: A comparison of their associations with other drug use and use disorders, affective and anxiety disorders, and psychosis. *Addiction*. 2001; 96:1603–1614. [PubMed: 11784457]
- Degenhardt L, Hall W, Lynskey M. Exploring the association between cannabis use and depression. *Addiction*. 2003; 98:1493–1504. [PubMed: 14616175]
- Denton M, Prus S, Walters V. Gender differences in health: A Canadian study of the psychosocial, structural, and behavioral determinants of health. *Social Science & Medicine*. 2004; 58:2585–2600. [PubMed: 15081207]
- Derogatis LR, Lipman RS, Rickels K, Uhlenhuth EH, Covi L. The Hopkins Symptom Checklist (HCSL): A self-report symptom inventory. *Behavioral Science*. 1974; 19:1–15. [PubMed: 4808738]
- Donovan JE, Jessor R. Structure of problem behavior in adolescence and young adulthood. *Journal of Consulting and Clinical Psychology*. 1985; 53:890–904. [PubMed: 4086689]
- Ellickson PL, Martino SC, Collins RL. Marijuana use from adolescence to young adulthood: Multiple developmental trajectories and their associated outcomes. *Health Psychology*. 2004; 23:299–307. [PubMed: 15099171]
- Fergusson DM, Boden JM, Horwood LJ. Cannabis use and other illicit drug use: Testing the cannabis gateway hypothesis. *Addiction*. 2006; 101:556–569. [PubMed: 16548935]
- Fergusson DM, Horwood LJ. Early onset cannabis use and psychosocial adjustment in young adults. *Addiction*. 1997; 98:279–296. [PubMed: 9219390]
- Fergusson DM, Horwood LJ, Beautrais AL. Cannabis and educational achievement. *Addiction*. 2003; 98:1681–1692. [PubMed: 14651500]
- Flory KD, Lynam R, Milich C, Leukefeld C, Clayton RR. Early adolescent through young adult alcohol and marijuana use trajectories: Early predictors, young adult outcomes, and predictive utility. *Developmental Psychopathology*. 2004; 16:193–213.
- Friedman AS, Terras A, Zhu W, McCallum J. Depression, negative self-image, and suicide attempts as effects of substance use and substance dependence. *Journal of Addictive Diseases*. 2004; 23:55–71. [PubMed: 15339714]
- Galaif E, Chou CP, Sussman S, Dent CW. Depression, suicidal ideation, and substance use among continuation high school students. *Journal of Youth and Adolescence*. 1998; 27:275–299.
- Haney M, Hart CL, Vosburg SK, Nasser J, Bennett A, Zubarán, Nasser J, Bennett A, Zubarán C, Foltin RW. Marijuana withdrawal in humans: Effects of oral THC or dival proex. *Neuropsychopharmacology*. 2004; 29:158–170. [PubMed: 14560320]
- Harrison, LD.; Martin, SS.; Enev, T.; Harrington, D. DHHS Population No. SMA 07-4249, Methodology Series M-7. Substance Abuse and Mental Health Services Administration, Office of Applied Studies; Rockville, MD: 2007. Comparing Drug Testing and Self-Report of Drug Use Among Youths and Young Adults in the General Population.
- Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: Results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Archives of General Psychiatry*. 2005; 62:1097–1106. [PubMed: 16203955]
- Iversen L. Cannabis and the brain. *Brain*. 2003; 126:1252–1270. [PubMed: 12764049]
- Jessor R, Chase JA, Donovan JE. Psychosocial correlates of marijuana use and problem drinking in a national sample of adolescents. *American Journal of Public Health*. 1980; 70:604–610. [PubMed: 7377436]
- Jessor, R.; Donovan, JE.; Costa, FM. *Beyond Adolescence: Problem Behavior and Young Adult Development*. Cambridge University Press; Cambridge, UK: 1991.

- Kandel DB. Issues of sequencing of adolescent drug use and other problem behaviors. *Drugs & Society*. 1989; 3:55–76.
- Kessler RC, Chiu WT, Demler O, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*. 2005; 6:617–627. [PubMed: 15939839]
- Kreek MJ, Nielsen DA, Butelman ER, LaForge KS. Genetic influences on impulsivity, risk taking, stress responsivity and vulnerability to drug abuse and addiction. *Nature Neuroscience*. 2005; 8:1450–1457.
- Lo Y, Mendell NR, Rubin DB. Testing the number of components in a normal mixture. *Biometrika*. 2001; 88:767–778.
- Lynskey MT, Coffey C, Degenhardt L, Carlin JB, Patton GA. A longitudinal study of the effects of adolescent cannabis use on high school completion. *Addiction*. 2003; 98:685–692. [PubMed: 12751986]
- Lynskey MT, Heath AC, Bucholz KK, Slutske WS, Madden PA, Nelson EC, Statham DJ, Martin NG. Escalation of drug use in early onset cannabis users vs. co-twin controls. *JAMA*. 2003; 289:427–433. [PubMed: 12533121]
- MacLeod J, Oakes R, Copello A, Crome I, Egger M, Hickman M, Oppenkowski T, Stokes-Lampard H, Smith G. Psychological and social sequelae of cannabis and other illicit drug use by young people: a Systematic review of longitudinal general population studies. *Lancet*. 2004; 363:1579–1588. [PubMed: 15145631]
- McDonough P, Walters V. Gender and health: Reassessing patterns and explanations. *Social Science & Medicine*. 2001; 52:547–559. [PubMed: 11206652]
- McLachlan, GJ.; Peel, S. *Finite Mixture Modeling*. John Wiley & Sons; New York: 2000.
- Muthén, LK.; Muthén, BO. *Mplus: Statistical analysis with latent variables: User's guide*. Muthén & Muthén; Los Angeles: 2007.
- Nagin, DS. *Group-Based Modeling of Development*. Harvard University Press; Cambridge, MA: 2005.
- Nylund KL, Asparouhov T, Muthén BO. Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study. *Structural Equation Modeling*. 2007; 14:535–569.
- O'Malley, PM.; Bachman, JG.; Johnston, LD.; Schulenberg, J. Studying the transition from youth to adulthood: Impacts on substance use and abuse. In: Juster, TF.; Kahn, RL., editors. *A Telescope on Society: Survey Research and Social Science at the University of Michigan and Beyond*. The University of Michigan Press; Ann Arbor: 2004. p. 305-329.
- Otsubo T, Tanaka K, Koda R, Shinoda J, Sano N, Tanaka S, Aoyama H, Mimura M, Kamijima K. Reliability and validity of Japanese version of the Mini-International Neuropsychiatric Interview. *Psychiatry and Clinical Neurosciences*. 2005; 59:517–526. [PubMed: 16194252]
- Patton GC, Coffey C, Carlin JB, Degenhardt L, Lynskey M, Hall W. Cannabis use and mental health in young people: A cohort study. *British Medical Journal*. 2002; 23:1199–1212.
- Pope HG, Gruber AJ, Hudson JI, Cohane G, Huestis MA, Yurgelun-Todd D. Early-onset cannabis use and cognitive deficits: What is the nature of the association? *Drug and Alcohol Dependence*. 2003; 69:303–310. [PubMed: 12633916]
- Raftery AE. A model for high-order Markov chains. *Journal of the Royal Statistical Society Series B (Methodological)*. 1985; 47:528–530.
- Raftery AE. Model selection in social research. *Sociological Methodology*. 1995; 25:111–163.
- Schulenberg JE, Merline AC, Johnston LD, O'Malley PM, Bachman JG, Laetz VB. Trajectories of marijuana use during the transition to adulthood: The big picture based on national panel data. *Journal of Drug Issues*. 2005; 35:255–279. [PubMed: 16534532]
- Schuster C, O'Malley PM, Bachman JG, Johnston LD, Schulenberg J. Adolescent marijuana use and adult occupational attainment: A longitudinal study from age 18 to 28. *Substance Use & Misuse*. 2001; 36:997–1014. [PubMed: 11504156]
- Schwarz G. Estimating the dimension of a model. *The Annals of Statistics*. 1978; 6:461–464.
- Shedler J, Block J. Adolescent drug use and psychological health: A longitudinal inquiry. *American Psychologist*. 1990; 45:612–630. [PubMed: 2350080]

- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R, Dunbar GC. The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*. 1998; 59:22–33. [PubMed: 9881538]
- Sheehan DV, Lecrubier Y, Sheehan KH, Janavs J, Weiller E, Keskiner A, Schinka J, Knapp E, Sheehan MF, Dunbar GC. The validity of the the Mini-International Neuropsychiatric Interview (MINI) according to the SCID-P and its reliability. *European Psychiatry*. 1997; 12:232–241.
- Solowij N, Stephens RS, Roffman RA, Babor T, Kadden R, Miller M, Christiansen K, McRee B, Vendetti J. for the Marijuana Treatment Project Research Group. Cognitive functioning of long-term heavy cannabis users s seeking treatment. *JAMA*. 2002; 287:1123–1131. [PubMed: 11879109]
- Stinson FS, Ruan WJ, Pickering R, Grant BF. Cannabis use disorders in the USA: Prevalence, correlates, and co-morbidity. *Psychological Medicine*. 2006; 36:1447–1460. [PubMed: 16854249]
- Sullivan PF, Neale MC, Kendler KS. Genetic epidemiology of major depression: Review and meta-analysis. *American Journal of Psychiatry*. 2000; 157:1552–1662. [PubMed: 11007705]
- Thomas H. A community survey of adverse effects of cannabis use. *Drug and Alcohol Dependence*. 1996; 42:201–207. [PubMed: 8912803]
- Windle M, Wiesner M. Trajectories of marijuana use from adolescence to young adulthood: Predictors and outcomes. *Developmental Psychopathology*. 2004; 16:1007–1027.



**Figure 1.** Trajectories of Marijuana Use for African American and Puerto Rican Women (N=474).

**Table 1**

Means and standard deviations for African American and Puerto Rican Women

	African American (Mean, SD)	Puerto Rican (Mean, SD)	T
Marijuana Use T1	0.10 ( <b>0.55</b> )	0.19 ( <b>0.68</b> )	1.52
Marijuana Use T2	0.48 ( <b>1.10</b> )	0.68 ( <b>1.25</b> )	1.61
Marijuana Use T3	0.78 ( <b>1.34</b> )	0.65 ( <b>1.31</b> )	-0.78
Marijuana Use T4	0.77 ( <b>1.34</b> )	0.63 ( <b>1.27</b> )	-0.73
Age at Initiation of Marijuana Use (T2)	15.6 ( <b>2.45</b> )	15.3 ( <b>2.30</b> )	-0.83
Depressive Symptoms (T5)	3.89 ( <b>3.37</b> )	4.11 ( <b>3.69</b> )	0.62
Anger/Hostility (T5)	3.82 ( <b>2.07</b> )	3.89 ( <b>2.38</b> )	0.31
Mean Age (T5)	32.7 ( <b>0.55</b> )	32.3 ( <b>0.55</b> )	-2.74**
Household Income (T5)	\$56,920.5 ( <b>40,176.6</b> )	\$50,208.6 ( <b>34,547.2</b> )	-1.74
Education (T5)	3.76 ( <b>2.26</b> )	2.42 ( <b>2.17</b> )	-5.92***

**Table 2**

Means and Standard Deviations of Psychological Adjustment for Women's Marijuana Trajectory Latent Classes (N=382)

<b>Psychological Adjustment Outcomes at Age 32.5</b>	<b>Nonusers (n=294)</b>	<b>Quitters (n=25)</b>	<b>Increases (n=63)</b>
Depressive Symptoms	3.72 (3.31)	5.49 (4.71)	4.74 (3.55)
Anger/Hostility	3.65 (2.05)	4.02 (2.68)	4.77 (2.57)