

Selective inactivation of adenosine A_{2A} receptors in striatal neurons enhances working memory and reversal learning

Catherine J. Wei,^{1,2} Philipp Singer,³ Joana Coelho,¹ Detlev Boison,⁴ Joram Feldon,³ Benjamin K. Yee,^{3,5} and Jiang-Fan Chen^{1,2,5}

¹Molecular Neuropharmacology Laboratory, Department of Neurology, Boston University School of Medicine, Boston, Massachusetts 02118, USA; ²Department of Pharmacology and Experimental Therapeutics, Boston University School of Medicine, Boston, Massachusetts 02118, USA; ³Laboratory of Behavioural Neurobiology, Swiss Federal Institute of Technology (ETH) Zurich, Schorenstrasse 16, 8603 Schwerzenbach, Switzerland; ⁴R.S. Dow Neurobiology Laboratories, Legacy Research, Portland, Oregon 97232, USA

The adenosine A_{2A} receptor (A_{2A}R) is highly enriched in the striatum where it is uniquely positioned to integrate dopaminergic, glutamatergic, and other signals to modulate cognition. Although previous studies support the hypothesis that A_{2A}R inactivation can be pro-cognitive, analyses of A_{2A}R's effects on cognitive functions have been restricted to a small subset of cognitive domains. Furthermore, the relative contribution of A_{2A}R in distinct brain regions remains largely unknown. Here, we studied the regulation of multiple memory processes by brain region-specific populations of A_{2A}R. Specifically, we evaluated the cognitive impacts of conditional A_{2A}R deletion restricted to either the entire forebrain (i.e., cerebral cortex, hippocampus, and striatum, fb-A_{2A}R KO) or to striatum alone (st-A_{2A}R KO) in recognition memory, working memory, reference memory, and reversal learning. This comprehensive, comparative analysis showed for the first time that depletion of A_{2A}R-dependent signaling in either the entire forebrain or striatum alone is associated with two specific phenotypes indicative of cognitive flexibility—enhanced working memory and enhanced reversal learning. These selective pro-cognitive phenotypes seemed largely attributed to inactivation of striatal A_{2A}R as they were captured by A_{2A}R deletion restricted to striatal neurons. Neither spatial reference memory acquisition nor spatial recognition memory were grossly affected, and no evidence for compensatory changes in striatal or cortical D₁, D₂, or A₁ receptor expression was found. This study provides the first direct demonstration that targeting striatal A_{2A}R may be an effective, novel strategy to facilitate cognitive flexibility under normal and pathologic conditions.

[Supplemental material is available for this article.]

Adenosine modulates neurotransmission and synaptic function in the central nervous system (e.g., de Mendonca and Ribeiro 1994, 1997; Sebastiao and Ribeiro 1996) and is increasingly thought to play an important role in learning and memory (e.g., Fredholm et al. 2005; Boison 2007; Yee et al. 2007). Adenosine A_{2A} receptors (A_{2A}R) are uniquely positioned in the brain to modulate cognitive functions that depend on the complex integration of dopaminergic, glutamatergic, and other neuronal signals. They are abundantly expressed in the striatum and are largely found on post-synaptic striatopallidal medium spiny neurons (MSNs) of the “indirect” pathway (e.g., Schiffmann et al. 1991a,b; Rosin et al. 1998; Rebola et al. 2005) where they colocalize and interact with receptors of other neurotransmitter and neuromodulatory systems to influence synaptic plasticity and behavior (Ferre et al. 2007; Wei et al. 2011). Specifically, striatal A_{2A}R interact antagonistically with dopamine D₂ receptors (D₂R) (Hillion et al. 2002; Canals et al. 2003) and oppose N-methyl-D-aspartate receptor (NMDAR) function (Norenberg et al. 1998; Wirkner et al. 2000; Gerevich et al. 2002). In contrast,

striatal A_{2A}R may also interact synergistically with metabotropic glutamate 5 receptors (mGlu₅R) (Ferre et al. 2002; Kachroo et al. 2005) and cannabinoid CB₁ receptors (CB₁R) (Tebano et al. 2009; Lerner et al. 2010). Moreover, A_{2A}R activity on MSNs has been shown to critically modulate long-term potentiation (LTP) at cortico-accumbal synapses (d'Alcantara et al. 2001) and spike-timing-dependent LTP at glutamatergic synapses onto striatopallidal MSNs (Shen et al. 2008b), a process thought to subserve learning and memory (Bliss and Collingridge 1993). Outside the striatum, A_{2A}R are also weakly expressed in neurons in the hippocampus and the cortex (Cunha et al. 1994; Dixon et al. 1996; Rosin et al. 1998). Such extra-striatal A_{2A}R similarly interact with other receptors such as mGlu₅R (Rodrigues et al. 2005; Tebano et al. 2005). Within the hippocampus, A_{2A}R appear to be essential for LTP at mossy fiber-CA3 (Rebola et al. 2008) and CA3 → CA1 (Fontinha et al. 2009) synapses. A_{2A}R are also found at pre-synaptic terminals in cortical afferent neurons projecting onto striatal MSNs (Rosin et al. 2003; Rebola et al. 2005), and these strategically located cortical A_{2A}R can modulate glutamate release in the striatum (Rebola et al. 2005; Ciruela et al. 2006). Hence, A_{2A}R are uniquely positioned to fine-tune, at the circuitry level, the complex integration of dopaminergic, glutamatergic, and other neuronal signals that underlie cognitive processes including learning and memory. In transgenic models, A_{2A}R overexpression

⁵Corresponding authors.

E-mail chenjf@bu.edu; fax (617) 638-5354.

E-mail byee@ethz.ch; fax 41-44-6557203.

Article is online at <http://www.learnmem.org/cgi/doi/10.1101/lm.2136011>.

impaired short-term object recognition memory and working memory (Gimenez-Llort et al. 2007), whereas A_{2A}R inactivation enhanced spatial recognition memory (Wang et al. 2006) and working memory (Zhou et al. 2009). The latter might suggest that suppression of A_{2A}R activity could be pro-cognitive as A_{2A}R inactivation also counteracted age-related and pathologic memory loss in rodents (Prediger et al. 2005a,b,c; Dall'Igna et al. 2007).

However, existing studies have been limited in scope, with analyses restricted to a small subset of cognitive domains. A comprehensive characterization of A_{2A}R's effects on particular cognitive functions is lacking. Furthermore, the relative contribution of A_{2A}Rs within distinct brain regions to these A_{2A}R-dependent cognitive outcomes has only begun to be recognized. Yu et al. (2009) recently identified a critical role for striatal A_{2A}Rs in habit formation during instrumental learning, a finding that is consistent with the well-known striatal locus for habit formation (Packard and Knowlton 2002). The relevance of these abundant striatal A_{2A}Rs to higher cognitive processes such as spatial learning, which is typically attributable to hippocampus and association cortices (for review, see Aggleton et al. 2000; Kesner 2009; Save and Poucet 2009), however, has not yet been explored. Despite their relatively low expression levels, hippocampal and cortical A_{2A}Rs have been shown to modulate important synaptic functions (like LTP, as mentioned previously) thought to critically underlie learning and memory (Bliss and Collingridge 1993; Lynch 2004). These extra-striatal A_{2A}Rs might therefore also regulate such higher cognitive functions. A functional dissociation between striatal and extra-striatal (i.e., cortical and hippocampal) A_{2A}Rs in the regulation of behavior is highlighted by our recent study suggesting that they exert opposite control over dopamine- and glutamate/NMDAR-mediated psychomotor activity (Shen et al. 2008a).

The present study builds upon this initial finding and aims to dissect the complex, complementary contributions of striatal and cortical A_{2A}Rs to higher cognitive functions. We assessed multiple learning and memory domains with brain region specification of A_{2A}R's effects by determining the cognitive impacts of conditional deletion of *Adora2a* (A_{2A}R) restricted to forebrain (i.e., cortex, hippocampus, and striatum, fb-A_{2A}R KO) (Bastia et al. 2005) or to striatum (st-A_{2A}R KO) (Shen et al. 2008a) in recognition memory, working memory, reference memory, and reversal learning. This comparison allowed us to dissociate the effects of striatal A_{2A}Rs from extra-striatal (but within forebrain) A_{2A}Rs on specific cognitive processes. This comprehensive assessment revealed two previously underappreciated phenotypes of improved cognitive flexibility following targeted brain regional A_{2A}R deletion. Specifically, deleting A_{2A}Rs in forebrain neurons led to a selective enhancement of working memory and reversal learning. Restricting the deletion to A_{2A}Rs only in striatal neurons still captured these same pro-cognitive phenotypes, thus suggesting that striatal A_{2A}R inactivation alone is sufficient to enhance working memory and reversal learning. Targeting striatal A_{2A}Rs may therefore represent a novel approach for facilitating cognitive flexibility underlying effective goal-directed behavior when environmental demands or conditions change.

Results

Validation of brain region-specific A_{2A}R deletion restricted to forebrain or striatum

Two separate conditional A_{2A}R knockout mouse lines with targeted A_{2A}R deletion in either the entire forebrain (i.e., fb-A_{2A}R KO, *Camk2a-cre(+)-Adora2a^{flox/flox}*) or striatum only (i.e., st-A_{2A}R KO, *Dlx5/6-cre(+)-Adora2a^{flox/flox}*) were generated using the Cre-*loxP* strategy. This standard strategy uses promoters with brain region-specific activity to drive the localized expression of the Cre

protein, which subsequently recombines/deletes the particular gene sequence flanked by *loxP* sequences. In this study, the *Camk2a* promoter or the *Dlx5/6* enhancer elements was used to produce Cre-mediated A_{2A}R deletion restricted to the forebrain or striatum, respectively.

Fb-A_{2A}R KO and st-A_{2A}R KO mice have been characterized for their selective A_{2A}R deletion in the forebrain (i.e., the cortex, the hippocampus, and the striatum) (Bastia et al. 2005; Yu et al. 2008) or exclusively in striatal (Shen et al. 2008a) neurons, as shown in our previous studies. To further characterize the Cre-mediated effects on A_{2A}R expression in different brain regions in fb-A_{2A}R KO and st-A_{2A}R KO mice, we determined Cre expression by X-gal staining of LacZ in a Rosa26 reporter transgenic line, PCR analysis of Cre-mediated A_{2A}R deletion in brain, and ³H-ZM241385 (a selective A_{2A}R antagonist) radioligand binding of A_{2A}R density in the brain.

X-gal staining revealed high levels of Cre recombination in the cortex and the hippocampus and moderate to high levels in the striatum of *Camk2a-cre(+)-Rosa26^{flox/flox}* mice (Fig. 1A, upper panel). This expression pattern is consistent with our previous detection of Cre-mediated A_{2A}R deletion in fb-A_{2A}R KO mice by PCR analysis and by ³H-SCH58261 radioligand binding assay (Yu et al. 2008). In contrast, in *Dlx5/6-cre(+)-Rosa26^{flox/flox}* mice, Cre recombination was robust and localized mainly to dorsal and ventral striatum, although sparse recombination was detected in extra-striatal brain regions such as the hippocampus and the cortical mantle (Fig. 1A, lower panel). This pattern of scattered and weak staining in extra-striatal brain regions did not appear to include the principal excitatory neurons in which A_{2A}Rs are found, but might represent instead GABA-ergic interneurons (Batista-Brito et al. 2008).

To evaluate the time course of Cre-mediated A_{2A}R deletion, we performed PCR analysis of genomic DNA extracted from the striatum, the cortex, the hippocampus, and/or the cerebellum of fb-A_{2A}R KO or st-A_{2A}R KO mice. As shown in Figure 1B (upper panel) Cre-mediated recombination was first detected at postnatal day 23 in fb-A_{2A}R KO mice in the striatum and the cortex, but not in the cerebellum, thus demonstrating the forebrain specificity of the A_{2A}R deletion at the DNA level. In contrast, in st-A_{2A}R KO mice, Cre-mediated recombination was evident as early as postnatal day 5 (the earliest postnatal day examined) in the striatum, but was largely absent in the cortex or the hippocampus (Fig. 1B, lower panel), thereby confirming striatum-specific A_{2A}R deletion. In both fb-A_{2A}R KO and st-A_{2A}R KO mice, it appeared that the recombination might be incomplete as shown by the persistence of the flox band.

As Cre activity may not directly reflect the extent of A_{2A}R protein loss, ³H-ZM241385 radioligand binding assays were also conducted to quantify A_{2A}R protein expression in the striatum, the cortex, the hippocampus, and the olfactory bulb of fb-A_{2A}R KO and st-A_{2A}R KO mice (Fig. 1C). One-way ANOVA ($n = 5-7$ mice per group) of ³H-ZM241385 binding in fb-A_{2A}R KO and fb-WT mice revealed significant loss of A_{2A}R expression in the striatum ($F_{(1,10)} = 64.48$, $P < 0.0001$) and the olfactory bulb ($F_{(1,9)} = 12.87$, $P < 0.01$), as well as a trend of reduction in the hippocampus and the cortex (Fig. 1C, upper panel). These results are entirely consistent with the pattern previously reported using ³H-SCH58261 (another selective A_{2A}R antagonist) as the radioligand (Yu et al. 2008). In contrast, analysis of st-A_{2A}R KO and st-WT mice ($n = 6-7$ per group) demonstrated that A_{2A}R expression was completely lost in the striatum ($F_{(1,11)} = 177.47$, $P < 0.0001$), but preserved in the cortex and the hippocampus (F 's < 1), as well as olfactory bulb ($P = 0.16$) of st-A_{2A}R KO mice (Fig. 1C, lower panel). Thus, forebrain-specific or striatum-specific A_{2A}R deletion was also achieved at the protein level in fb-A_{2A}R KO and st-A_{2A}R KO mice, respectively.




Figure 1. Forebrain- or striatum-specific A_{2A}R KO is selective. (A) X-gal staining for brain regions of Cre expression demonstrates strong staining throughout the entire forebrain (i.e., cortex, hippocampus, and striatum) in *Camk2a-cre(+)/Rosa26^{flox/flox}* mice (upper panel). X-gal staining also shows strong staining throughout the entire striatum (that is largely absent in the cortex or the hippocampus) in *Dlx5/6-cre(+)/Rosa26^{flox/flox}* mice (lower panel). (B) Postnatal developmental time course of Cre recombination and deletion of the “floxed” A_{2A}R allele. Representative PCR analysis of genomic DNA isolated from (1) the striatum, (2) the cortex, and (3) the cerebellum of fb-A_{2A}R KO mice at the various developmental stages (P15, P23, and P35) demonstrates a forebrain-specific pattern of A_{2A}R deletion in the striatum and the cortex, but not in the cerebellum beginning around P23 (upper panel). Similar PCR analysis of (1) the striatum, (2) the cortex, and (3) the hippocampus from st-A_{2A}R KO mice at different developmental stages indicates a striatum-specific pattern of A_{2A}R deletion in the striatum, but not in the cortex or the hippocampus as early as P5 (lower panel). (C) ³H-ZM241385 (selective A_{2A}R antagonist) radioligand binding to quantify Cre-mediated loss of A_{2A}R expression in the striatum, the cortex, the hippocampus, and the olfactory bulb in fb-A_{2A}R KO mice and st-A_{2A}R KO mice. ³H-ZM241385 binding is reduced in all forebrain regions examined in fb-A_{2A}R KO mice (upper panel), but only in the striatum of st-A_{2A}R KO mice (lower panel). *n* = 5–7 per group. Mean ± SEM are plotted. **P* = 0.06, ***P* < 0.01, ****P* < 0.0001, KO vs. WT.

In summary, analyses of the spatial and temporal patterns of Cre-recombination and Cre-mediated A_{2A}R deletion at both the genomic and receptor binding levels in both knockout mouse lines collectively demonstrated that fb-A_{2A}R KO had achieved selective loss of A_{2A}R in the entire forebrain (i.e., cortex, hippocampus, and striatum) beginning at postnatal day 23, whereas st-A_{2A}R KO had achieved a highly selective loss of A_{2A}R in the striatum only (but not in the cortex or the hippocampus) as early as postnatal day 5. Thus, both fb-A_{2A}R KO and st-A_{2A}R KO mice shared a common feature of deficient A_{2A}R expression in the striatum, although they differed in the timing of deletion. In fb-A_{2A}R KO mice, as expected, this regional deletion extended from striatum to also include the cortex and the hippocampus and it occurred later.

Lack of compensatory changes in adenosine A₁ or dopamine D₁ and D₂ receptors

A_{2A}Rs are closely linked to the adenosinergic (i.e., A₁Rs) and dopaminergic (i.e., D₁Rs and D₂Rs) receptor systems (Fredholm et al. 2007). Compensatory changes in the expression of these receptors might affect behavior independently of A_{2A}R deletion (Fredholm et al. 2005; El-Ghundi et al. 2007). We therefore performed quantitative assays of binding densities in striatal and/or cortical total membranes from fb-A_{2A}R KO and st-A_{2A}R KO mice for A₁Rs (³H-DPCPX, a selective A₁R antagonist), D₁Rs (³H-SCH23390, a selective D₁R antagonist), and D₂Rs (³H-raclopride, a selective D₂R antagonist). No significant changes emerged in the binding assays for A₁Rs or D₁Rs in both striatal and cortical membranes, or for D₂Rs in striatal membranes from fb-A_{2A}R KO and st-A_{2A}R KO mice in comparison with their respective WT littermates (Fig. 2A,B). These data suggest that compensatory changes in A₁R, D₁R, or D₂R expression were not a concern in our A_{2A}R KO

mouse models. However, we cannot exclude the possibility of compensatory shifts in cell surface expression of these receptors, G-protein-coupling, or other downstream signaling pathways.

Spatial recognition memory in the Y-maze is unaffected in both fb-A_{2A}R KO and st-A_{2A}R KO mice

Recognition memory for distant spatial cues across a broad range of retention delays was evaluated in the nonaversive Y-maze test. Preferential exploration of the novel arm during the test phase performed 2 min, 30 min, 3.5 h, or 1 d after the sample phase was evaluated as an index of spatial recognition memory. A clear and comparable preference for the novel arm was apparent between genotypes in both forebrain and striatum cohorts, with all groups showing the expected decline in preference with increasing delay between sample and test phases (Fig. 3A,B).

These interpretations were supported by separate 2 × 2 × 4 (Genotype × Sex × Delays) ANOVAs of the percent time spent in the novel arm, each of which only yielded a significant effect of Delays (fb-A_{2A}R KO, *F*_(3,63) = 3.92, *P* < 0.05; st-A_{2A}R KO, *F*_(3,81) = 7.99, *P* < 0.001). An additional analysis of the time spent in each arm using separate 2 × 2 × 4 × 3 (Genotype × Sex × Delays × Arms) ANOVAs further confirmed the overall preference for the novel arm (fb-A_{2A}R KO, *F*_(2,42) = 71.19, *P* < 0.001; st-A_{2A}R KO, *F*_(2,54) = 93.73, *P* < 0.001) and its dependency on the delay (Delays × Arms interaction: fb-A_{2A}R KO, *F*_(6,126) = 2.80, *P* < 0.05; st-A_{2A}R KO, *F*_(6,162) = 5.91, *P* < 0.001). It was therefore concluded that spatial recognition across a range of retention delays was not significantly altered by loss of forebrain or striatal A_{2A}Rs. This conclusion was strengthened by supplementary analyses of the distance traveled in each arm, which yielded a parallel pattern of results, thus arguing against any potential confound of




Figure 2. Forebrain- and striatum-specific A_{2A}R KO is without compensatory changes in A₁, D₁, or D₂ receptor levels. (A,B) Quantitative analysis of ³H-DPCPX (selective A₁R antagonist), ³H-SCH23390 (selective D₁R antagonist), and ³H-raclopride (selective D₂R antagonist) in total membranes from the striatum and/or cortex of fb-A_{2A}R KO and fb-WT mice ($n = 3-6$ per group) (A), and of st-A_{2A}R KO and st-WT mice ($n = 5-8$ per group) (B). No differences in radioligand binding densities were demonstrated for these receptors (one-way ANOVA, P 's > 0.13). Mean \pm SEM are plotted.

locomotor activity on the time measure. Similarly, novel arm preference during the test phase was not confounded by any difference in arm exposure during the sample phase: familiar arm exploration was highly comparable between genotypes across each delay condition following either fb-A_{2A}R KO or st-A_{2A}R KO.

Spatial working memory performance is facilitated in both fb-A_{2A}R KO and st-A_{2A}R KO mice

Visually guided escape behavior

Initial training using the visible cued platform yielded no evidence of any genotype difference (Fig. 4A). All mice acquired the swimming and escape response, with performance improving over the four trials performed across two consecutive days, thus demonstrating their ability to utilize the local visual cues to locate the escape platform. A $2 \times 2 \times 4$ (Genotype \times Sex \times Trials) ANOVA of escape latency confirmed the presence of a significant trials effect (fb-A_{2A}R KO, $F_{(3,75)} = 20.78$, $P < 0.001$; st-A_{2A}R KO, $F_{(3,87)} = 22.57$, $P < 0.001$). An identical pattern of results was obtained in the separate analysis of path length (Trials effect: fb-A_{2A}R KO, $F_{(3,75)} = 18.25$, $P < 0.001$; st-A_{2A}R KO, $F_{(3,87)} = 22.01$, $P < 0.001$), which is consistent with the impression that swim speed remained relatively stable and comparable between KO and WT mice [mean swim speed: fb-A_{2A}R KO: 14.47 ± 0.49 cm/sec, fb-WT: 15.57 ± 0.58 cm/sec, $F_{(1,25)} = 2.09$, $P = 0.16$; st-A_{2A}R KO: 16.13 ± 0.67 cm/sec, st-WT: 17.57 ± 0.71 cm/sec, $F_{(1,29)} = 2.17$, $P = 0.151$]. An identical analysis of these parameters during the visible cue task also yielded a similar pattern of results reflecting similar performance and behaviors

among st-A_{2A}R KO and st-WT mice of the second striatal cohort (Experiment set II, data not shown). Thus, neither the motor nor motivational component of the water-maze escape task was significantly affected in fb-A_{2A}R KO or st-A_{2A}R KO mice.

Working memory performance in fb-A_{2A}R KO mice

In this test, improved escape performance from trial 1 to 2 provides a measure of working memory (Hodges et al. 1995). The working memory task taxes the application of a daily matching-to-sample rule where the animal's experience on trial 1 (i.e., when the mouse has located the novel unknown position of a hidden platform) is used to guide the subsequent search for an identically located escape platform on trial 2 on a given day. Hence, only the same day's trial 1 experience is effective in guiding search performance during trial 2 in the working memory task. Initial analysis of both escape latency and path length by separate $2 \times 2 \times 3 \times 4 \times 2$ (Genotype \times Sex \times Delays \times Days \times Trials) ANOVAs yielded a significant four-way interaction (Genotype \times Sex \times Delays \times Trials interaction: escape latency: $F_{(2,46)} = 3.57$, $P < 0.05$; path length: $F_{(2,46)} = 3.19$, $P = 0.05$), which was suggestive of a sex-dependent genotype effect (see Fig. 4B). This was investigated by additional analyses restricted to either sex alone, which clearly demarcated the presence of a genotype effect on working memory performance in the female but not in the male sex. It is worth noting that the phenotypic difference in working memory observed between male and female fb-A_{2A}R KO mice is unlikely attributable to a differential loss of forebrain A_{2A}Rs in these mice as X-gal staining showed a similar Cre-expression pattern between male and female fb-A_{2A}R KO mice (data not shown).




Figure 3. Fb-A_{2A}R KO and st-A_{2A}R KO mice show normal spatial recognition memory in the Y-maze. To evaluate memory for distant spatial cues, mice were first allowed 5 min to explore the start and familiar arms of the maze (sample phase) and then returned to the maze after a delay period (2 min, 30 min, 3.5 h, and then 1 d) and given 3 min to explore these same two arms plus an additional, novel arm (test phase). Preference for the novel arm, expressed as the percentage of time spent in the novel arm [(time in novel arm/time in all arms) × 100%], during each test phase was used to index spatial recognition memory. (A) Fb-A_{2A}R KO and fb-WT mice showed a similar preference for the novel arm, which progressively weakened at a comparable rate with increasing retention demands. (B) St-A_{2A}R KO and st-WT mice also showed a strong and comparable preference for the novel arm that gradually declined at a similar rate toward chance performance upon increasing the delay interval. Values depicted are mean ± SEM. Chance level = 33.33%, dashed line.

Significant Genotype × Trials (escape latency: $F_{(1,12)} = 7.61$, $P < 0.05$; path length: $F_{(1,12)} = 4.58$, $P = 0.05$) and Genotype × Delays × Trials (escape latency: $F_{(2,24)} = 0.057$; path length: $F_{(2,24)} = 3.76$, $P < 0.05$) interactions were evident in the females, but clearly absent in the males (F 's < 1). As illustrated in Figure 4B (left panel), these interaction terms, which were unique to the females, arose because fb-WT mice showed the expected delay-dependent deterioration of performance, whereas fb-A_{2A}R KO mice continued to show a clear improvement in escape during trial 2 relative to trial 1, regardless of the delay. Forebrain A_{2A}R deletion thus led to a delay-dependent improvement of working memory performance: fb-A_{2A}R KO and fb-WT mice were highly comparable at the minimal delay, but diverged only when the increasing retention demand of the delay effectively reduced performance in fb-WT mice to chance level. Post-hoc pair-wise comparisons confirmed that female fb-A_{2A}R KO and fb-WT mice never significantly differed from each other during trial 1 (P 's > 0.30). A clear difference, however, was detected during trial 2 under the delay conditions of 10 min ($P < 0.05$) and 15 min ($P < 0.005$), and only fb-A_{2A}R KO mice demonstrated significant performance improvement from trial 1 to 2 at the longest delay ($P < 0.02$). This genetic knockout effect on working memory performance was completely absent in the males (Fig. 4B, right panel).

Working memory performance in st-A_{2A}R KO mice

Experiment set I. Striatal A_{2A}R KO similarly led to a facilitation of working memory performance (Fig. 4C), but this effect was (1) already evident at the minimal delay condition when st-WT mice failed to consistently improve performance from trial 1 to 2 and (2) comparable between sexes, i.e., not sex-dependent. Separate $2 \times 2 \times 4 \times 2$ (Genotype × Sex × Days × Trials) ANOVAs of the two performance measures yielded a significant Genotype × Trials interaction (escape latency: $F_{(1,29)} = 6.80$, $P < 0.05$; path length: $F_{(1,29)} = 6.41$, $P < 0.05$) that was independent of sex (Genotype × Sex × Trials interaction: escape latency:

$F_{(1,29)} = 1.55$, $P = 0.22$; path length: $F_{(1,29)} = 1.15$, $P = 0.29$). Post-hoc comparisons indicated the presence of a significant improvement from trial 1 to 2 in st-A_{2A}R KO mice ($P < 0.005$), but not in st-WT mice ($P = 0.64$), and performance on trial 2 differed significantly between genotypes ($P < 0.05$) despite comparable performance on trial 1 ($P = 0.15$).

Experiment set II. The lack of evidence of effective working memory function among st-WT mice in Experiment set I raised the possibility that st-WT mice might have failed to master the matching rule of the task, thus leaving open the interpretation that the performance enhancement in st-A_{2A}R KO mice might not represent an enhancement in working memory ability. To exclude this alternative interpretation, we conducted a separate experiment in a separate cohort of st-A_{2A}R KO and st-WT mice using a slightly modified training protocol to ensure satisfactory learning performance in the st-WT controls (see Materials and Methods).

To capture the key element of working memory, in particular, we focused our analyses to the first two trials of each day. As depicted in Figure 4D (first panel), both st-A_{2A}R KO and st-WT mice similarly demonstrated a clear improvement in trial 2 compared to trial 1 when matching-to-sample learning was promoted by using the four-trial-per-day training protocol in conjunction with a minimal retention load of 20 sec. This impression was confirmed by separate $2 \times 2 \times 4 \times 2$ (Genotype × Sex × Days × Trials) ANOVAs of the two performance measures. As expected, all mice demonstrated significant improvement from trial 1 to 2 (main effect of Trials: escape latency: $F_{(1,13)} = 20.55$, $P < 0.001$; path length: $F_{(1,13)} = 7.71$, $P < 0.05$), which did not differ between st-A_{2A}R KO and st-WT mice (Genotype × Trials interaction: escape latency: $F_{(1,13)} = 1.28$, $P = 0.28$; path length: $F_{(1,13)} = 2.60$, $P = 0.13$). These results confirmed that both st-A_{2A}R KO and st-WT mice were able to learn the procedures and acquire the day-dependent matching rule of the task.

Following initial training with the four-trial-per-day training protocol in which performance was comparable between genotypes, mice were then returned to a two-trial-per-day testing protocol as in Experiment set I. As shown in Figure 4D (last three panels), performance began to diverge again with st-A_{2A}R KO mice (but not st-WT mice) showing significant improvement from trial 1 to 2. This genotype effect was evident with the short ITI (20-sec delay) and persisted as the retention load was increased with the extended ITI (10-min delay) in both the females and males. Thus, this outcome was entirely consistent with that in Experiment set I in which st-A_{2A}R KO mice consistently outperformed st-WT mice in both sexes. A $2 \times 2 \times 2 \times 4 \times 2$ (Genotype × Sex × Delay × Days × Trials) ANOVA of escape latency confirmed this observation yielding only a main effect of Trials ($F_{(1,13)} = 8.12$, $P < 0.05$) and a significant Genotype × Trials interaction ($F_{(1,13)} = 9.18$, $P < 0.01$), which were both independent of the delay intervals (Delay × Trials interaction: $F_{(1,13)} = 3.26$, $P = 0.09$; Genotype × Delay × Trials interaction: $F < 1$) and of sex (Sex × Trials and Genotype × Sex × Trials interactions: F 's < 1). Post-hoc comparisons indicated a significant improvement in escape latency from trial 1 to 2 in st-A_{2A}R KO mice ($P < 0.005$), but not in st-WT mice ($P = 0.92$) despite their having demonstrated similar baseline performance on trial 1 ($P = 0.11$).

Collectively, these data from both sets of experiments indicate that st-A_{2A}R KO mice consistently showed improved spatial working memory compared to st-WT mice, and this was not due to st-WT mice failing to acquire the matching rule. The fact that Experiment set II successfully replicated the phenotype previously observed in Experiment set I also strongly argues against any critical impact of prior trainings in the first striatal cohort on the original observation of working memory enhancement




Figure 4. Fb-A_{2A}R KO (female only) and st-A_{2A}R KO mice demonstrate normal visually guided escape behavior but improved spatial working memory. Mice were first pretrained in the water maze for up to two consecutive days (two trials per day) to swim directly to a visible platform for escape. Working memory was subsequently evaluated by examining over blocks of 4 d the improvement in escape latency (s) from trial 1 to 2 to reach a hidden platform whose location changed every day. (A) Both fb-A_{2A}R KO (left panel) and st-A_{2A}R KO (right panel) mice showed a comparable reduction in escape latency with each trial compared to fb-WT and st-WT mice, respectively. This indicates that A_{2A}R deletion did not alter motivation or sensory and motor capabilities required to successfully learn and execute an escape onto a visible platform. (B,C) Experiment set I: The forebrain cohort was tested over three blocks corresponding to a 20-sec, 10-min, or 15-min delay between trials. The striatal cohort was tested in a single block at the minimal delay of 20 sec. Among the female mice (B, left panel), clear evidence for a delay-dependent enhancement of working memory was observed in fb-A_{2A}R KO mice: these mice demonstrated a consistent and marked reduction in escape latency from trial 1 to 2 at all delays, whereas fb-WT mice only readily showed such improvement at the shorter delay, suggesting improved working memory capabilities in these female fb-A_{2A}R KO mice. In contrast, among the male mice (B, right panel), fb-A_{2A}R KO and fb-WT mice performed remarkably similarly, showing comparable improvements in escape latency from trial 1 to 2 at all delays. St-A_{2A}R KO mice, like female fb-A_{2A}R KO mice, readily showed improvement across the two trials, whereas st-WT mice did not (C). (D) Experiment set II: A separate striatal cohort was first trained for 4 d using a four-trial-per-day training protocol to facilitate learning and ensure mastery of the matching rule. Mice were then returned in the testing phase to a two-trial-per-day protocol as in Experiment set I and tested at 20-sec and 10-min delays. St-A_{2A}R KO and st-WT mice demonstrated comparable improvement from trial 1 to 2 during training, but as the task demands increased during testing, st-A_{2A}R KO mice again continued to out-perform st-WT mice, an effect that was not sex-dependent. Despite this lack of sex dependency in st-A_{2A}R KO mice, the data are further plotted split by sex in order to provide a parallel comparison with the enhanced working memory phenotype in fb-A_{2A}R KO mice, which did show a sex effect. Values depicted are mean ± SEM. **P* < 0.05, ****P* < 0.005, KO vs. WT. ^*P* < 0.02, ^^*P* < 0.005, trial 1 vs. 2 in A_{2A}R KO mice; ##*P* < 0.01, Genotype × Trials interaction.

because the two striatal cohorts differed substantially in their respective experimental history. Moreover, unlike the phenotype that had emerged in fb-A_{2A}R KO mice, the phenotype of enhanced working memory in st-A_{2A}R KO mice was demonstrably independent of sex in both Experiment set I and Experiment set II (see Fig. 4D, last two panels).

Reference memory acquisition performance is unaffected in both fb-A_{2A}R KO and st-A_{2A}R KO mice

In contrast to working memory, when the platform is fixed in a constant position across days as in the reference memory procedure, the gradual accumulation of a relevant memory trace connecting the fixed location with the opportunity to escape from the water contributes to (1) performance improvement over days and (2) development of a search preference for the relevant spatial location.

Across the 10 d of acquisition training, all groups showed a gradual improvement in their escape performance. This was evidenced by a reduction in escape latency (Fig. 5A,D) and path length, with KO and WT mice demonstrating near-identical performance levels at the end of training. Separate 2 × 2 × 10 × 2

(Genotype × Sex × Days × Trials) ANOVAs of both performance measures in the two cohorts yielded a highly significant Days effect (fb-A_{2A}R KO: escape latency: *F*_(9,225) = 6.33, *P* < 0.001; path length: *F*_(9,225) = 10.27, *P* < 0.001; st-A_{2A}R KO: escape latency: *F*_(9,243) = 2.58, *P* < 0.01; path length: *F*_(9,243) = 3.35, *P* < 0.001). Neither the main effect of Genotype nor its interactions approached statistical significance. Thus, A_{2A} receptor knockout in the forebrain or the striatum did not significantly affect acquisition performance.

Post-acquisition probe test performance is reduced in st-A_{2A}R KO but not fb-A_{2A}R KO mice

Spatial search pattern in Probe tests 1 and 2

Two probe tests, in which the platform was removed from the maze, were conducted 24 and 72 h following the conclusion of acquisition training to assess the strength and retention of the spatial search preference that had developed as a result of acquisition training. The two tests yielded parallel outcomes, suggesting stable retention of the spatial preference (Fig. 5B,E). Comparison between fb-A_{2A}R KO and fb-WT mice indicated that both groups




Figure 5. Reference memory performance is spared in fb-A_{2A}R KO and st-A_{2A}R KO mice. Mice were trained for 10 consecutive days (two trials per day) to acquire the fixed spatial location of a hidden escape platform. Following acquisition, two probe tests were conducted on days 11 (Probe test 1) and 13 (Probe test 2) during which the escape platform was removed from the pool and the spatial search pattern of mice was observed for 60 sec. (A) Fb-A_{2A}R KO mice, like fb-WT mice, demonstrated a steady decline in escape latency with each training day, showing highly comparable escape performance by the end of acquisition training (day 10). (B) Both fb-A_{2A}R KO and fb-WT mice showed a strong and similar search preference for the spatial location of the escape platform (i.e., target quadrant) in both probe tests. (C) Search accuracy defined by the number of annular crossings was comparable between fb-A_{2A}R KO and fb-WT mice. (D) Similarly, st-A_{2A}R KO and st-WT mice showed gradual performance improvement during acquisition, achieving near-identical performance at the end of training (day 10). (E) Both st-A_{2A}R KO and st-WT mice demonstrated an above-chance level search preference for the target quadrant in both probe tests; however, this preference was weaker in st-A_{2A}R KO mice. (F) St-A_{2A}R KO and st-WT mice did not significantly differ in the number of annular crossings, indicating comparable search accuracy in both probe tests. Values depicted are mean \pm SEM. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. chance level (=25%, dashed line). ^ $P < 0.05$, st-A_{2A}R KO vs. st-WT.

showed a strong preference for the quadrant area (i.e., target quadrant) centering on the platform's location and comparable search accuracy of the precise platform location during acquisition training (Fig. 5B,C). In contrast, this preference, albeit still above chance level in st-A_{2A}R KO mice, was significantly weaker in comparison with that in st-WT mice (Fig. 5E). These interpretations were confirmed by separate $2 \times 2 \times 4$ (Genotype \times Sex \times Quadrants) ANOVAs of the percent time spent per quadrant in each probe test. The analyses comparing fb-A_{2A}R KO and fb-WT mice only yielded a main effect of Quadrants (Probe test 1: $F_{(3,75)} = 28.69$, $P < 0.001$; Probe test 2: $F_{(3,75)} = 18.42$, $P < 0.001$). In contrast, the analyses comparing st-A_{2A}R KO and st-WT mice revealed in addition, a significant Genotype \times Quadrants interaction in both probe tests (Probe test 1: $F_{(3,81)} = 3.18$, $P < 0.05$; Probe test 2: $F_{(3,81)} = 3.76$, $P < 0.05$). This was attributable to different target quadrant preference between genotypes as indicated by post-hoc pair-wise comparisons ($P < 0.05$). One-sample *t*-tests further confirmed that all groups displayed a target quadrant preference significantly exceeding that expected by chance, i.e., $>25\%$ (P 's < 0.05).

Normal memory retention and temporal search pattern during probe testing in st-A_{2A}R KO mice

Differences in the amount of time spent in the target quadrant during probe testing can reflect differences in memory retention as well as in temporal search patterns. Therefore, we performed a set of additional analyses to test whether the observed relative performance impairment on Probe tests 1 and 2 in st-A_{2A}R KO mice might be attributable to impaired memory retention or an altered temporal search pattern during probe testing in st-A_{2A}R

KO mice. First, to address whether st-A_{2A}R KO mice differed in their ability to retain the memory for the spatial location of the platform, we examined trial 1 performance during acquisition training, which can indicate memory retention from the previous day's training. Separate $2 \times 2 \times 10$ (Genotype \times Sex \times Days) ANOVAs of trial 1 escape latency and path length did not yield any significant Genotype effect (F 's < 1) or its interactions (Genotype \times Days interaction: escape latency: $F_{(9,243)} = 1.34$, $P = 0.24$; path length: $F_{(9,243)} = 1.38$, $P = 0.22$), a result that is consistent with the impression based on the overall analysis in the acquisition phase (see Fig. 5D). To further clarify this issue, we also analyzed the number of annular crossings (i.e., the frequency at which the swim path crossed the former spatial location of the platform) and the latency of the first annular crossing in each probe test to provide both, respectively, an index of search accuracy and an estimate of what the escape latency would have been if the platform were present. Separate two-way (Genotype \times Sex) ANOVAs of the number of annular crossings (Fig. 5F) or the latency of the first annular crossing (data not shown) in each of the two probe tests did not reveal any significant differences between st-A_{2A}R KO and st-WT mice. Together with the above-mentioned findings that trial 1 performance across acquisition days showed gradual improvement and was indistinguishable between genotypes, these data suggest that st-A_{2A}R KO mice were not consistently impaired in their ability to retain the memory of the precise spatial location of the escape platform compared to st-WT mice.

Second, to address whether a different temporal search pattern (e.g., possibly reflective of enhanced within-session extinction) in st-A_{2A}R KO mice could account for the apparent probe test deficit reported previously, we examined the temporal profile

(across successive 15-sec bins) of the preference for the target quadrant in each probe test. Separate $2 \times 2 \times 4$ (Genotype \times Sex \times Bins) ANOVAs of the percent time spent in the target quadrant suggested that the preference was relatively stable over time and without evidence for extinction in both st-A_{2A}R KO and st-WT mice (Bins effect and Genotype \times Bins interaction: F 's < 1), but consistent with the initial probe test analyses comparing all four quadrants (see Fig. 5E), this preference was reduced in st-A_{2A}R KO mice (main effect of Genotype: Probe test 1: $F_{(1,27)} = 5.42$, $P < 0.05$; Probe test 2: $F_{(1,27)} = 4.71$, $P < 0.05$). Therefore, the probe test performance deficit observed in st-A_{2A}R KO mice was not associated with a difference in their temporal search pattern in the target quadrant during probe testing, when actual escape onto the platform was not available. This suggests that enhanced extinction of the target quadrant preference is an unlikely explanation for their weaker probe test performance. The exact reason for the weaker performance in Probe tests 1 and 2 in st-A_{2A}R KO mice remains unclear but may stem from a weakening of habit formation (see Discussion) rather than from an impairment in reference memory retention or acquisition as such.

Reversal phase learning occurs more rapidly in both fb-A_{2A}R KO and st-A_{2A}R KO mice

Reversal learning began 4 d after the end of acquisition on experimental day 14 (see Fig. 6). This lasted for 4 d in the forebrain cohort when fb-WT mice showed a clear preference for the

new target quadrant in Probe test 3. This preference, however, was not observed in st-WT mice, and thus, reversal training continued for another 4 d in the striatal cohort. By the end of this additional training, st-WT mice had exhibited a clear preference for the new target quadrant. This pattern of results aligned with the impression obtained from the working memory test, which suggested that the fb-WT mice tended to learn more quickly in general relative to the st-WT mice.

Fb-A_{2A}R KO facilitated reversal learning performance without affecting probe test performance

The change of platform location (from day 10 to day 14) disrupted the efficiency of locating the escape platform in fb-A_{2A}R KO and fb-WT mice. To gauge this reversal effect, analyses contrasting the last acquisition day and the first reversal day were performed, which yielded only a main effect of Days (three-way ANOVA: escape latency: $F_{(1,25)} = 34.99$, $P < 0.001$; path length: $F_{(1,25)} = 34.91$, $P < 0.001$). Although the initial impact of reversal appeared comparable between fb-A_{2A}R KO and fb-WT mice, their performance appeared to diverge as reversal training continued (Fig. 6A).

Indeed, fb-A_{2A}R KO mice rapidly improved over the 4 d of reversal training, achieving a performance level (i.e., on day 17) that was comparable with that at the end of acquisition (i.e., on day 10). In contrast, fb-WT mice performed relatively poorly during the reversal phase. Separate $2 \times 2 \times 4 \times 2$ (Genotype \times




Figure 6. Faster reversal learning in fb-A_{2A}R KO and st-A_{2A}R KO mice. Reversal learning began once the mice had learned the fixed position of the hidden escape platform during acquisition. It lasted for 4 d (days 14–17) in the forebrain cohort and for 8 d (days 14–21) in the striatal cohort. During reversal, the location of the platform was shifted 180° to the opposite quadrant, and mice were required to learn the new escape location. Probe tests were conducted 24 h after a training session to evaluate the progress of reversal learning on day 18 (Probe test 3) in both cohorts, and additionally on days 20 and 22 (Probe tests 4 and 5) in the striatal cohort only. (A) Fb-A_{2A}R KO mice were less affected by the shift in platform location as indicated by faster escape latencies during the reversal phase. This effect appeared more pronounced with additional reversal training as demonstrated by a divergence in escape latencies. (B) Both fb-A_{2A}R KO and fb-WT mice exhibited a comparable search preference for the new target quadrant after 4 d of reversal training. (C) St-A_{2A}R KO mice also escaped more quickly throughout reversal learning compared to st-WT mice, indicating that they were less disrupted by the sudden change in platform location. (D) St-A_{2A}R KO mice also demonstrated a strong preference for the new target quadrant in all three probe tests. This preference was significantly greater in st-A_{2A}R KO mice during Probe test 3 compared to that in st-WT mice. A significant target preference was not observed in st-WT mice until the end of reversal training (i.e., Probe test 5). Values depicted are mean \pm SEM. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.005$ vs. chance level (=25%, dashed line). ^ $P < 0.05$, st-A_{2A}R KO vs. st-WT.

Sex × Days × Trials) ANOVAs of escape latency and path length yielded similar patterns of results, although the latency measure appeared to be more powerful in detecting a genotypic difference. Significant effects of Genotype (escape latency: $F_{(1,25)} = 4.39$, $P < 0.05$; path length: $F_{(1,25)} = 3.92$, $P = 0.06$) and of Days (escape latency: $F_{(3,75)} = 10.02$, $P < 0.001$; path length: $F_{(3,75)} = 9.49$, $P < 0.001$) were found.

A probe test conducted to evaluate the pattern of spatial search on day 18 yielded highly comparable outcomes between fb-A_{2A}R KO and fb-WT mice (Fig. 6B). After 4 d of reversal training, both groups had developed an overall preference for the new target quadrant that significantly exceeded chance level (P 's < 0.01) with limited evidence of any residual preference for the former (now opposite) target quadrant. A $2 \times 2 \times 4$ (Genotype × Sex × Quadrants) ANOVA of the percent time spent in each quadrant only yielded a highly significant Quadrants effect ($F_{(3,75)} = 10.64$, $P < 0.001$).

St-A_{2A}R KO facilitated reversal learning and showed a less persistent preference for the previously acquired platform location during probe testing

The impact of relocating the escape platform impaired performance as expected (Fig. 6C). Analyses contrasting the last acquisition day and the first reversal day confirmed the impact of reversal on both performance measures, yielding a significant Days effect (escape latency: $F_{(1,27)} = 25.13$, $P < 0.001$; path length: $F_{(1,27)} = 19.36$, $P < 0.001$). There was no statistical support that the initial impact of reversal differed between st-A_{2A}R KO and st-WT mice (Genotype × Days interaction: escape latency: $F_{(1,27)} = 2.02$, $P = 0.17$; path length: $F < 1$).

However, st-A_{2A}R KO mice out-performed st-WT mice over the course of reversal learning. Separate $2 \times 2 \times 8 \times 2$ (Genotype × Sex × Days × Trials) ANOVAs of the two performance measures yielded similar outcomes, although escape latency was again more powerful in detecting a genotypic difference. Significant effects of Genotype (escape latency: $F_{(1,27)} = 7.49$, $P = 0.01$; path length: $F_{(1,27)} = 3.78$, $P = 0.06$) and of Days (escape latency: $F_{(7,189)} = 4.64$, $P < 0.001$; path length: $F_{(7,189)} = 3.54$, $P < 0.01$) emerged, but their interaction was not significant (F 's < 1). Thus, the statistical outcomes resembled those obtained in the forebrain cohort although the two experiments differed in the number of training days.

To allow an effective assessment of the competition between any residual preference for the previous platform location (i.e., during acquisition) and the newly acquired preference for the platform location used during reversal training, three probe tests (Probe tests 3, 4, and 5) were performed over the course of reversal training. The first of these probe tests was performed after 4 d of reversal training (just before reversal training on day 5). St-A_{2A}R KO mice already showed a preference for the new target quadrant during this probe test, whereas st-WT mice still exhibited an overall preference for the previous target quadrant used during acquisition (see Probe test 3) (Fig. 6D). This impression was confirmed by the emergence of a near-significant Genotype × Quadrants interaction ($F_{(3,81)} = 2.70$, $P = 0.05$) in a $2 \times 2 \times 4$ (Genotype × Sex × Quadrants) ANOVA of percent time spent per quadrant. The same interaction term achieved clear statistical significance in a separate $2 \times 2 \times 2$ ANOVA contrasting solely the percent time spent between the (diagonally opposite) new and previous target quadrants ($F_{(1,27)} = 5.76$, $P = 0.025$). This analysis provides a meaningful and specific assessment of the expression of the reversal effect by focusing on the search behavior in the previously reinforced quadrant and the presently (newly) reinforced quadrant in exclusion of the never-reinforced quadrants. Post-hoc pairwise comparisons further confirmed that st-A_{2A}R KO mice already

preferred the new target quadrant more than did st-WT mice on the first reversal probe test (Probe test 3, $P < 0.05$).

By the second and third probe tests in the reversal phase, however, performance between st-A_{2A}R KO and st-WT mice had become increasingly comparable (see Probe tests 4 and 5) (Fig. 6D). In st-WT mice, preference for the new target quadrant continued to increase as their preference for the previous target quadrant waned. On the other hand, performance across these three probe tests was relatively stable in st-A_{2A}R KO mice. Separate $2 \times 2 \times 4$ (Genotype × Sex × Quadrants) ANOVAs of the percent time spent per quadrant revealed only a main effect of Quadrant in these two probe tests (Probe test 4: $F_{(3,81)} = 4.02$, $P = 0.01$; Probe test 5: $F_{(3,81)} = 12.81$, $P < 0.001$) without any evidence for genotypic differences. The latter result was in agreement with the analysis of annular crossings variables (data not shown). Additional one-sample t -tests confirmed that target quadrant preference exceeded chance level in all probe tests in st-A_{2A}R KO mice (P 's < 0.006), but st-WT mice only achieved that in the final probe test ($P < 0.02$). Increasing preference for the new target quadrant across the three probe tests in st-WT mice ($F_{(2,24)} = 5.88$, $P < 0.01$), but not in st-A_{2A}R KO mice ($F_{(2,30)} = 1.77$, $P = 0.20$), was confirmed by a repeated measures ANOVA that directly compared preference for the new target quadrant across the three probe tests.

Discussion

This study provides the first direct demonstration that targeted inactivation of A_{2A}Rs on intrinsic striatal neurons is sufficient to facilitate two forms of cognitive-behavioral flexibility—working memory and reversal learning. These phenotypes arose largely in the absence of any gross or persistent effects on other behavioral parameters, such as spatial reference memory acquisition or spatial recognition memory. They were also free from compensatory changes in D₁, D₂, or A₁ receptor expression in the striatum or the cortex. Moreover, comparative analysis of fb-A_{2A}R KO and st-A_{2A}R KO revealed largely similar phenotypes of enhanced working memory and reversal learning, leading to the conclusion that A_{2A}R inactivation in the striatum (rather than in the cortex or the hippocampus) plays a predominant role in the expression of cognitive flexibility by A_{2A}Rs. Thus, targeting striatal A_{2A}Rs alone may be sufficient to facilitate effective behavioral adaptations in response to changing environmental contingencies. Striatal A_{2A}Rs might therefore represent an attractive, novel strategy to restore cognitive flexibility in neuropsychiatric conditions, such as Parkinson's disease, schizophrenia, or related neuropsychiatric disorders in which striatal dysfunction is implicated.

Selective inactivation of striatal neuronal A_{2A}Rs is sufficient to enhance working memory

Working memory captures important elements of cognitive flexibility, notably the capacity to maintain or update information held online and select appropriate behavioral responses in accordance to shifting positive and negative stimulus-response contingencies (Goldman-Rakic 1995; Marie and Defer 2003; Dalley et al. 2004). A_{2A}R deletion on striatal neurons facilitated spatial working memory performance as evidenced by consistent trial 1-to-2 improvement in st-A_{2A}R KO but not st-WT mice in two independent experiments (i.e., Experiment sets I and II). We observed similar results in female fb-A_{2A}R KO mice, but this difference was absent in the males. This sex-dependent phenotypic difference is unlikely attributable to a differential loss of forebrain A_{2A}Rs between sexes given that Cre expression did not reveal any such sex difference (data not shown). Although the precise origin or mechanism underlying this sex-dependent working

memory phenotype in fb-A_{2A}R KO mice is presently unclear, it is worth noting that global A_{2A}R knockout has previously been reported to influence other brain-relevant processes in a sex-dependent manner, including body temperature regulation (Yang et al. 2009) and ethanol sensitivity and consumption (Naassila et al. 2002). Interestingly, estrogen has been shown to modify the expression of A_{2A}R transcript in several different brain regions (Ribeiro et al. 2009); thus, one might venture to speculate a degree of sex hormonal influence over A_{2A}R-dependent phenotypes. Here, because the expression of the working memory phenotype was only dependent on sex in the forebrain but not striatal A_{2A}R knockout mouse line, any such modulation by sex hormones might be more critical and/or more likely occurring in areas outside the striatum (e.g., in the cortex and/or the hippocampus where A_{2A}Rs were deleted in fb-A_{2A}R KO but spared in st-A_{2A}R KO mice). Regardless of the precise mechanism underlying the sex-dependent effect unique to fb-A_{2A}R KO mice, it does not undermine the major finding that working memory enhancement can be induced by A_{2A}R inactivation (which led to a similar overall impact between the two conditional A_{2A}R knockout lines) and that striatal A_{2A}R inactivation alone is sufficient to produce this selective enhancement, which incidentally was not significantly modified by sex.

Improved working memory in st-A_{2A}R KO and female fb-A_{2A}R KO mice unlikely reflects ineffective mastery of the matching rule by their respective controls: both control groups clearly demonstrated successful task performance when the cognitive load was minimal (i.e., Experiment set II training in the striatal cohort under the initial four-trial-per-day protocol and 20-sec delay in the forebrain cohort). Thus, performance between knockout and control only diverged when task difficulty increased (see Fig. 4B,D). In contrast, spatial recognition memory and reference memory acquisition were largely unaffected by A_{2A}R inactivation in either knockout line. This pattern of outcomes closely resembles that seen following global A_{2A}R inactivation in mice (Zhou et al. 2009), which also selectively enhanced working memory without affecting reference memory learning. Conversely, overexpression of A_{2A}Rs in the brain impaired working memory performance, but also did not affect reference memory function in transgenic rats (Gimenez-Llort et al. 2007). The selectivity of these findings is consistent with the view that striatal function is not critical to forming spatial representations as such (Packard and McGaugh 1992; McDonald et al. 2008; Berke et al. 2009).

The novel finding of our study is that selective inactivation of striatal A_{2A}Rs alone was sufficient to reproduce the pro-cognitive phenotypes resulting from A_{2A}R deletion extending to the entire forebrain. An earlier transgenic study suggested that A_{2A}Rs in the cortex and the hippocampus modulate working memory (Gimenez-Llort et al. 2007), which is consistent with A_{2A}R's well-documented functional effects on neuronal plasticity at the cortico-striatal (Schiffmann et al. 2007) and the hippocampal mossy fiber-CA3 (Rebola et al. 2008) pathways and with the identification of cortical regions (e.g., prefrontal and parietal cortices), in particular as the key structures subserving effective working memory based on electrophysiology and imaging (Goldman-Rakic 1995; Rowe et al. 2000).

Our comparative analysis, however, revealed a largely similar pro-cognitive phenotype in both fb-A_{2A}R KO and st-A_{2A}R KO mice. This profile was similar both in the type and selectivity of the cognitive functions that were enhanced, as well as in the magnitude of the observed cognitive enhancements, particularly in the case of working memory. This strongly suggests that striatal A_{2A}Rs (not cortical or hippocampal A_{2A}Rs) play a critical role in A_{2A}R-dependent modulation of cognition. Extending the deletion of A_{2A}Rs beyond the striatum to include the hippocampus

and the cortex (i.e., extra-striatal) as in fb-A_{2A}R KO mice did not produce any phenotypes that were distinguishable from those already present in st-A_{2A}R KO mice. Thus, striatal A_{2A}R activity appears to assume a more prominent influence on these cognitive functions compared with extra-striatal cortical/hippocampal A_{2A}R activity. The exact contribution of cortical/hippocampal A_{2A}Rs to cognition, however, remains to be defined, e.g., through the use of cortex- and/or hippocampal-specific A_{2A}R KO mouse models. Last, it would be prudent to point out that there are extra-striatal regions where A_{2A}Rs were apparently deleted (e.g., hypothalamus) in both A_{2A}R KO mouse lines examined. The possible loss of A_{2A}Rs in hypothalamus, where expression is normally low, might be predicted to affect performance via its control of arousal rather than of learning as such. Given that striatal A_{2A}Rs represent the majority of the A_{2A}Rs commonly lost between both knockout lines (due to its sheer number), it is both reasonable and parsimonious to infer that striatal A_{2A}Rs play a key role in yielding the observed working memory enhancement.

Striatal D₂R activity is important for effective working memory as shown in animals and humans (e.g., Kellendonk et al. 2006; Mehta et al. 2008). Inactivation of striatal A_{2A}Rs might potentiate striatal dopaminergic signaling via D₂Rs to produce the working memory enhancement, given the well-documented antagonistic A_{2A}R–D₂R interaction in striatopallidal MSNs of the “indirect” pathway (Ferre et al. 1997; Fredholm et al. 2007). In keeping with this notion, we have recently shown that st-A_{2A}R KO potentiated the motor-stimulant effect induced by dopaminergic stimulation (Shen et al. 2008a). Interestingly, a form of striatal LTD (i.e., eCB-LTD) that is dependent on endocannabinoid release and D₂R activation (Kretzner and Malenka 2005, 2007) was found to be restricted to this very same population of striatopallidal MSNs (Kretzner and Malenka 2007). Moreover, this critical release of endocannabinoids from striatal MSNs was previously reported to require dopamine release in conjunction with up-state-dependent activation of mGlu₅Rs and L-type calcium channels (Kretzner and Malenka 2005). Consistent with this observation, D₂R, mGlu₅R, and Cav1.3 stimulation was shown to induce LTD using a protocol capable of eliciting spike-timing-dependent plasticity (STDP) (Shen et al. 2008b). Although the impact of post-synaptic striatal A_{2A}R activity on striatal LTD has not yet been directly examined, concomitant stimulation of A_{2A}Rs and D₂Rs shifted the striatopallidal MSN plasticity response from that of a D₂R-induced LTD response to a LTP response instead (Shen et al. 2008b). It is noteworthy that activation of A_{2A}Rs, in the setting of FGFR co-activation, has also been demonstrated to promote LTP at these cortico-striatopallidal synapses (Flajolet et al. 2008), and suppression of A_{2A}Rs by pharmacologic blockade or by global A_{2A}R deletion has been shown to impair LTP at these synapses (d'Alcantara et al. 2001; Shen et al. 2008b). Therefore, it is possible that A_{2A}R deficiency in the striatal “indirect” pathway MSNs modifies working memory by modulating dopamine-dependent signaling and plasticity, LTD, and/or LTP in the striatum.

A_{2A}R modulation of NMDAR current at post-synaptic striatal neurons (Norenberg et al. 1998), as well as A_{2A}R heterodimerization and functional interaction with mGlu₅Rs (Ferre et al. 2002), further suggest that striatal A_{2A}R activity may also modify working memory by modulating striatal glutamatergic signaling. The modification of the striatal function potentially through dopaminergic and/or glutamatergic mechanisms by striatal A_{2A}Rs might be expected to alter the processing of cortical information entering the striatum, as well as striatal projections to efferent targets such as the prefrontal cortex (Simpson et al. 2010). The latter is supported by the recent demonstration that manipulation of striatal D₂R activity is sufficient to modify the prefrontal cortical function and working memory (Kellendonk et al. 2006).

Selective inactivation of striatal neuronal A_{2A}Rs enhances reversal learning

Reversal learning is an accepted measure of cognitive flexibility because efficient task performance requires the ability to rapidly switch from one response pattern to a conflicting one in response to the complete reversal of environmental contingencies between the relevant action and goal. This study provides the first demonstration that striatal A_{2A}R inactivation facilitates spatial reversal learning largely without affecting spatial reference memory acquisition or recognition memory. Both fb-A_{2A}R KO and st-A_{2A}R KO reversed faster, as evidenced by their efficient escape performance during reversal training (fb-A_{2A}R KO and st-A_{2A}R KO) and the earlier emergence of a spatial bias for the novel target quadrant across successive probe tests (st-A_{2A}R KO). Thus, compared to their respective control groups, these knockout mice more effectively and more rapidly redirected their search behavior in response to the relocation of the escape platform from its initial constant position. This outcome might reflect that fb-A_{2A}R KO and st-A_{2A}R KO mice were more ready to inhibit their previously acquired (but no longer effective) response and extend their search to alternative areas, thereby allowing them to establish the newly adaptive response more quickly than their controls. This phenotype cannot, however, be solely attributed to poorer initial learning of the original platform location because st-A_{2A}R KO and st-WT mice performed comparably during acquisition, achieving near-identical performance on the last training day, and demonstrated equivalent search accuracy and above-chance performance during probe testing (see Fig. 5D,F). This interpretation is reinforced by the observation of a similar enhanced reversal learning phenotype in fb-A_{2A}R KO mice that had clearly demonstrated normal probe test performance.

Although the water-maze reversal procedure differs from reversal learning based on the two-alternative forced choice paradigm by the fact that there were areas of the maze that were consistently not associated with the escape platform across both acquisition and reversal, it is still highly effective in taxing the ability to suppress or inhibit the previously reinforced response. As mentioned previously, the reversal learning phenotype may reflect a facilitation of the ability to suppress a learned response in general. This possibility is strengthened by our recent demonstration showing that st-A_{2A}R KO mice remained sensitive to the devaluation of a reinforcer's incentive value when overtraining had instilled a resistance in the controls whose responding had become habitual (Yu et al. 2009). Similarly, st-A_{2A}R KO mice were also demonstrably more responsive to an "omission" procedure, whereby lever pressing was discouraged by lowering the frequency and increasing the delay of food delivery (Yu et al. 2009). Consistent with this finding is the evidence suggesting that the striatum plays an important role in active response suppression (e.g., Zandbelt and Vink 2010).

All together, it is apparent that facilitated reversal learning and cognitive flexibility (observed in this study) may come at the expense of weakened habit formation (as shown in Yu et al. 2009). The balance and trade-off between the two strategies of behavioral control might therefore be effectively modulated by striatal A_{2A}Rs. For example, under conditions in which persistence is maladaptive and unproductive, the promotion of alternative behavioral responses might lead to a more rapid adaptation to changing environmental contingencies, leading to "enhanced" performance outcomes. Such conditions were prominent in both the reversal test in the present study as well as in the instrumental tests reported by Yu et al. (2009). One interpretation of these findings is that st-A_{2A}R KO mice remained goal-directed with a corresponding weakening of habit formation. The latter may imply that habit formation depends on the induction of

A_{2A}R-dependent striatal LTP (Lovinger 2010) as LTP in the striatum is reduced in global A_{2A}R KO mice (d'Alcantara et al. 2001; Shen et al. 2008b).

Neural circuits in several different brain regions including the prefrontal cortex, the hippocampus, the amygdala, and the striatum are collectively involved in reversal learning. Genetic knockout studies have implicated glutamatergic and dopaminergic signaling in reversal learning, but have not determined the relative contributions of cortical vs. subcortical receptors. The similar enhanced reversal phenotype in both knockout mouse lines argues that modulating A_{2A}R signaling in the striatum alone is sufficient for reversal learning enhancement and suggests that striatal A_{2A}Rs play a critical role in modulating this form of cognitive flexibility. This interpretation agrees with converging evidence that the striatum is pivotal in reversal learning (Bellebaum et al. 2008; Clarke et al. 2008).

Our finding of enhanced reversal learning, without disrupting acquisition during water-maze testing, is reminiscent of the impaired reversal learning phenotype in mutant mice with glutamatergic signaling deficiency including mGlu₅R (Xu et al. 2009), NMDA receptor subunit NR2A (Bannerman et al. 2008), and AMPA receptor subunit GluR-A (Bannerman et al. 2003) knockout mice and of the enhanced phenotype in forebrain neuron-specific glycine transporter (GlyT1) knockout mice (Singer et al. 2009). These gene knockout studies suggest that the potentiation of glutamatergic and/or NMDAR-mediated signaling may be linked to the enhancement of reversal learning. In light of A_{2A}R's functional antagonism with NMDARs (Wirkner et al. 2004) in the striatum, the enhanced reversal phenotype might also be mediated by heightened striatal NMDAR/glutamatergic signaling resulting from deficient striatal A_{2A}R activity.

Notwithstanding, D₂Rs are also important for reversal learning. In humans, reversal performance correlated positively with D₂R binding in the caudate nucleus (Clatworthy et al. 2009), whereas pharmacologic D₂R blockade in nonhuman primates (Lee et al. 2007), genetic D₂R deletion in mice (Kruzich et al. 2006; De Steno and Schmauss 2009), or striatal D₂R overexpression in mice (Kellendonk et al. 2006) impaired reversal learning. Thus, A_{2A}R's interaction with D₂Rs offers another mechanism whereby striatal A_{2A}Rs may influence reversal learning.

Last, recent evidence has shown that A_{2A}R antagonism can increase levels of the retrograde endocannabinoid messenger, 2-arachidonoylglycerol, in the striatum and facilitate endocannabinoid-dependent LTD at excitatory synapses onto post-synaptic striatopallidal MSNs (Lerner et al. 2010). Moreover, the motor stimulating effect of A_{2A}R antagonism, which is known to occur through a blockade of post-synaptic striatal A_{2A}Rs (Shen et al. 2008a), was markedly attenuated by global CB1-receptor antagonism or knockout (Lerner et al. 2010), thus highlighting the cross-talk between post-synaptic striatal A_{2A}Rs and endocannabinoid-CB1 receptor pathways. Along with evidence that pharmacologic manipulation (Hill et al. 2006) or genetic deletion (Varvel and Lichtman 2002) of CB1 receptors can impair reversal performance in rodents, one might speculate that suppression of striatal A_{2A}R activity may also facilitate reversal learning by potentiating endocannabinoid-CB1 receptor signaling in excitatory (e.g., cortical glutamatergic) synapses in the striatum.

Conclusion

In summary, we provide the first direct demonstration that A_{2A}R inactivation on intrinsic striatal neurons where A_{2A}R expression is most abundant is sufficient to selectively enhance working memory performance and facilitate reversal learning. This suggests that striatal A_{2A}Rs may be an effective, novel target to enhance cognition under physiological conditions. As A_{2A}R

antagonists are in clinical phase II–III trials for Parkinson's disease and cognitive inflexibility is a core cognitive disturbance in neuropsychiatric disorders including Parkinson's disease and schizophrenia, A_{2A}R antagonists might ameliorate these associated cognitive deficits. Our results support an important role for the striatum in cognition (Simpson et al. 2010) and lend a partial explanation for the prominent cognitive changes associated with neuropathologic disorders, like Huntington's disease and Parkinson's disease, where neuronal degeneration and loss is largely restricted to the striatum, without significant pathologic changes in the cortex.

Materials and Methods

Approval from the Institutional Animal Care and Use Committee at Boston University School of Medicine and the Zurich Cantonal Veterinarian Office had been previously granted for all experiments conducted in Boston and Zurich, respectively. They adhered to the NIH *Guide for the Care and Use of Laboratory Animals* (1982), the Swiss Federal Law and Ordinance on Animal Protection, and European Council Directive 86/609/EEC (1986).

Generation of forebrain-specific and striatum-specific

A_{2A}R knockout mice

Two conditional A_{2A}R knockout mouse lines with brain region-specific deletion of a critical region (i.e., exon 2) of *Adora2a*, the gene encoding the adenosine A_{2A} receptor (A_{2A}R), were generated using the Cre/*loxP* strategy. This particular strategy uses a region-specific promoter to drive the expression of Cre, which subsequently mediates the deletion of a gene flanked by *loxP* sites. Although we refer to our engineered mouse lines as gene “knock-out” to maintain consistency with our previous reports using these engineered mouse lines, our model might more precisely be considered a gene “knockdown” as Cre expression and/or Cre-mediated recombination of *Adora2a* may be incomplete (see Results).

Forebrain-specific A_{2A}R knockout mice (fb-A_{2A}R KO, *Camk2a-cre(+)-Adora2a^{lox/lox}*, congenic C57BL/6 genetic background) were generated and genotyped as previously detailed (Bastia et al. 2005). Briefly, *Camk2a-cre(+)* mice (*L7ag#13* line, C57BL/6 genetic background) (Dragatsis and Zeitlin 2000) and *Adora2a^{lox/lox}* mice (mixed Sv129 × C57BL/6 genetic background) were independently backcrossed to C57BL/6 mice for 10 generations at the Laboratory of Dr. Michael Schwarzschild (Massachusetts General Hospital) and then interbred to generate congenic fb-A_{2A}R KO mice and their WT littermates.

Striatum-specific A_{2A}R knockout mice (st-A_{2A}R KO, *Dlx5/6-cre(+)-Adora2a^{lox/lox}*, of a mixed FVB × C57BL/6 genetic background) were generated and genotyped as previously described (Shen et al. 2008a). Briefly, *Dlx5/6-cre(+)* transgenic mice in a FVB genetic background were provided (Ohtsuka et al. 2008) and cross-bred to *Adora2a^{lox/lox}* mice in a congenic C57BL/6 genetic background. A *Dlx5/6* intron regulatory element drove the embryonic, striatal neuron-specific Cre-mediated deletion of the “floxed” allele (Zerucha et al. 2000; Ghanem et al. 2003; Ohtsuka et al. 2008).

In earlier pilot studies, both *Adora2a^{lox/lox}* mice (i.e., without the *cre* transgene) and *Adora2a^{-/-}* mice (i.e., with *Camk2a-cre* or *Dlx5/6-cre* transgene, but without floxed allele) responded similarly to amphetamine. Thus, for this study, we used only the *Adora2a^{lox/lox}* mice for our “wild-type” littermate control groups (fb-WT or st-WT for the fb-A_{2A}R KO or st-A_{2A}R KO lines, respectively).

Generation of *Camk2a-cre(+)**Rosa26^{lox/lox}* and *Dlx5/6-cre(+)**Rosa26^{lox/lox}* reporter mice and X-gal staining of Cre expression in brain

Rosa26 (R26R) reporter mice (B6.129S4.*Gt(ROSA)26Sor^{tm1Sor}/J*, Jackson Laboratories, Bar Harbor, ME) were crossed with

Camk2a-cre(+) (fb-A_{2A}R KO line) or *Dlx5/6-cre(+)* (st-A_{2A}R KO line) transgenic mice to generate *Camk2a-cre(+)**Rosa26^{lox/lox}* or *Dlx5/6-cre(+)**Rosa26^{lox/lox}* mice for visualization of Cre expression in the brain by X-gal staining.

Briefly, naive mice were anesthetized with tribromoethanol (Avertin), transcardially perfused with ice-cold phosphate buffered saline (PBS) followed by 2% paraformaldehyde. The brains were removed in toto, post-fixed overnight, and then cryoprotected in a sucrose solution (10%–20%–30%) until processing. Parasagittal sections of 50 μm were incubated in X-gal solution (5 mM K₄Fe(CN)₆ · 3H₂O, 5 mM K₃Fe(CN)₆, 2 mM MgCl₂, 1 mg/mL X-gal (5-bromo-4-chloro-3-indoyl-β-D-galactoside, Invitrogen) in PBS and 2.5% dimethyl formamide) for 30 min at 37°C.

PCR analysis of Cre-mediated recombination and A_{2A}R deletion in various brain regions during postnatal development

PCR was conducted on striatal, cortical, and cerebellar tissue isolated from fb-A_{2A}R KO and fb-WT mice at postnatal days 15, 23, and 35 and on striatal, cortical, and hippocampal tissue isolated from st-A_{2A}R KO and st-WT mice at postnatal days 5 and 33. Whole tissue samples were isolated from each hemisphere, and PCR on extracted genomic DNA was conducted according to the procedure described in full previously (Bastia et al. 2005).

Total membrane binding assessment of adenosine and dopamine receptors in various brain regions

Total membranes were prepared from the striatum, the cortex, the hippocampus, and/or the olfactory bulb, and single-point saturation binding assays were performed in duplicate to quantify total A_{2A}R, A₁R, D₁R, and D₂R levels as described earlier (Dewar et al. 1989; Lidow et al. 1989; Cunha et al. 1996; Lopes et al. 2004; Houchi et al. 2005; Rebola et al. 2005) with slight modifications. Each 300-μL binding assay, consisting of assay buffer (50 μL), radioligand (50 μL), and membranes (200 μL, 100–200 μg), was incubated at room temperature (or 30°C for D₁R) for 1 h (or 2 h for A₁R). The radioligands and final concentrations were: (1) A_{2A}R: ³H-ZM241385 (3 nM, s.a. 27.4 Ci/mmol, American Radiolabeled Chemicals, Inc.); (2) A₁R: ³H-DPCPX (2 nM, s.a. 111.6 Ci/mmol, GE Healthcare); (3) D₁R: ³H-SCH23390 (3 nM, s.a. 73.1 Ci/mmol, PerkinElmer); and (4) D₂R: ³H-raclopride (5 nM, s.a. 82.8 Ci/mmol, PerkinElmer). Nonspecific binding was determined using 2 μM xanthine amine congener (A_{2A}R and A₁R), 5 μM fluphenazine dihydrochloride (D₁R), or 300 μM (S)-(–)-sulpiride (D₂R) (Sigma).

Behavioral evaluation

Experiment set 1 (fb-A_{2A}R KO and st-A_{2A}R KO mice)

Two cohorts of mice, referred to hereon as the “forebrain” and “striatal cohorts,” were used exclusively for behavioral evaluations. The forebrain cohort comprised 17 fb-A_{2A}R KO mice (nine male and eight female) and 13 fb-WT mice (six male and seven female), and the striatal cohort comprised 17 st-A_{2A}R KO mice (nine male and eight female) and 17 st-WT mice (seven male and 10 female). They were bred at Boston University School of Medicine (Boston, MA, USA) and transported to ETH-Zurich (Schwerzenbach, Switzerland) 1 mo before experimentation began, when they were 3.5–4.5 mo old. All mice were individually housed with ad libitum food and water in a temperature- and humidity-controlled vivarium maintained under a 12-h reversed light–dark cycle (lights on at 7 pm). Behavioral testing was always conducted in the dark phase.

A total of 11 tests were conducted and their chronologic sequence is listed in Supplemental Table 1, Experiment set I. This within-subjects approach was essential to assess potential confounding neurologic changes and other nonspecific effects. It also facilitated comparisons across tests and avoided excessive

animal use. To minimize transfer effects that might complicate data interpretation, the tests were ordered in terms of the severity of experimental stress that animals might incur. The present report includes results derived from the (1) Y-maze test of spatial recognition and (2) water-maze tests of visual discrimination, working memory, reference memory, and reversal learning. Results of the remaining tests (i.e., elevated plus maze test of anxiety, open field and home cage activity, prepulse inhibition test of sensorimotor gating, and Pavlovian fear conditioning) are to be reported elsewhere (CJ Wei, P Singer, D Boison, J Feldon, BK Yee, and JF Chen, in prep.).

Experiment set II (st-A_{2A} R KO mice)

A separate, independent striatal cohort comprising nine st-A_{2A}R KO mice (five male and four female) and eight st-WT mice (five male and three female) was subsequently tested in a follow-up experiment to further examine working memory performance in the water maze. These mice had previously undergone active avoidance testing followed by conditioned taste aversion testing (P Singer, CJ Wei, JF Chen, and BK Yee, unpubl.). The testing sequence of the five total behavioral tests along with the subject number are detailed in Supplemental Table 1, Experiment set II.

Spatial recognition

This was assessed by measuring the spontaneous preference for novel over familiar places using a Y-maze as fully described before (Pietropaolo et al. 2009). Briefly, mice were placed in the maze in a randomly selected start arm and allowed to explore one other arm (familiar arm). After 5 min, mice were removed from the maze for a variable duration (2 min, 30 min, 3.5 h, or 1 d) and then re-introduced to the maze for an additional 3 min during which they could explore all three maze arms. Preferential exploration of the never-visited novel arm during this test phase was indexed by the following ratio: (time spent in novel arm/time spent in all three arms) × 100%. A novel testing room with unique distal spatial cues was used for each delay condition, and a minimum of 2 d separated successive tests. Video tracking was performed by Ethovision (version 3.1, Noldus Technology).

Water maze

Experiment set I (fb-A_{2A}R KO and st-A_{2A}R KO mice)

The apparatus has been fully described before (Yee et al. 2007; Singer et al. 2009). Three tests were conducted: visible cue task, working memory task, and acquisition and reversal of reference memory. Each test was conducted in a distinct testing room with a unique set of distal spatial cues. In all tests, mice were given two trials per day to learn to escape from the water within 60 sec by climbing onto an escape platform (7-cm diameter) submerged 1 cm below the water surface, on which they remained for a 20-sec intertrial interval (ITI) before initiating the second trial. Platform placements were counterbalanced among subjects. A pseudorandomized, nonrepetitive, sequence of start positions was used. Video tracking was performed by Ethovision (version 3.1, Noldus Technology).

Visible cue task. This served to detect any confounding change in swimming or escape behavior. For two consecutive days, the location of the platform was made visible by a local cue placed directly above the escape platform and was varied between the two days.

Working memory task. Next, the escape platform stayed hidden and assumed a new position on each day, but remained unchanged across the two trials on the same day. Improvement from trial 1 to 2 provided a measure of working memory—the retrieval of the day-specific location of the escape platform learned from trial 1 (Hodges et al. 1995). St-A_{2A}R KO and st-WT mice were evaluated over a block of 4 d, with a minimal ITI (i.e., 20 sec), which proved to be sufficiently difficult for st-WT mice. On the

other hand, fb-A_{2A}R KO and fb-WT mice were tested for additional two 4-d blocks to accommodate more extended ITIs: 10 min for the second block and 15 min for the final block. These additional training blocks allowed us to achieve a level of difficulty that also reduced the performance of fb-WT mice to chance level. The daily platform locations and start positions within a block were counterbalanced with a pseudorandom sequence.

Acquisition and reversal of reference memory. This commenced 6 or 14 d after the working memory test for the forebrain or striatal cohort, respectively. The platform now assumed a constant location throughout the acquisition phase of the experiment, which lasted for 10 consecutive days. The platform position was counterbalanced across groups. A 60-sec probe test (with the escape platform removed) was performed on days 11 and 13. Next, reversal learning began and lasted for 4 or 8 d in the respective forebrain or striatal cohorts. In this phase, the escape platform assumed a new constant location that was diagonally opposite to its former location. An initial probe test, performed on day 5, revealed successful reversal learning in fb-A_{2A}R KO and fb-WT mice, which led to termination of the test. On the other hand, it was apparent that st-A_{2A}R KO and st-WT mice required further training. To monitor the progress of reversal learning across the additional training days, two more probe tests were performed: one at the beginning of day 7 and another at the beginning of day 9. Testing was then terminated in this cohort at the end of the final probe test on day 9. Performance was indexed by escape latency and path length. Probe test performance was evaluated by the proportion of time spent in the target quadrant, where the platform was previously located, relative to the other three quadrants.

Experiment set II (st-A_{2A} R KO mice)

As the performance enhancement in Experiment set I was observed against a background of weak st-WT control performance, additional visible and working memory tasks in Experiment set II were performed on a separate cohort of st-A_{2A}R KO and st-WT mice to solidify the original interpretation of enhanced working memory in st-A_{2A}R KO mice based on Experiment set I and to exclude the alternative interpretation that st-WT mice had failed to learn the matching rule. A four-trial-per-day training protocol was used to improve st-WT performance with the modifications described in the following.

Working memory task. Training: Mice were first trained for 4 d in the working memory task using four trials per day to facilitate learning and ensure that all mice could learn the procedures and the matching rule inherent to the working memory task.

Testing: Working memory testing proceeded for the next 8 d, as in Experiment set I, once effective working memory functioning (trial 1-to-2 improvement) during training was observed. Two delays (ITIs) of varying retention loads, 20 sec or 10 min, separated the two daily trials. Mice were tested for 4 d at each delay: 2 d at 20 sec, then 4 d at 10 min, and then 2 d more at 20 sec. This sequence ensured that comparison across the two delay conditions was not systemically confounded by overall training effects across the 8 d of testing.

Statistical analysis

Data from the two A_{2A}R KO mouse lines (i.e., fb-A_{2A}R KO or st-A_{2A}R KO) and from the two experiments (i.e., *Experimental sets I and II* in the st-A_{2A}R KO mouse line) were separately analyzed by parametric analysis of variance (ANOVA) of the appropriate design. Thus, all analyses involved comparisons between subjects that shared identical experimental histories. Between-subject factors included genotype (i.e., A_{2A}R KO vs. WT) and sex (i.e., male vs. female) if present. Within-subjects factors such as trials, days, delays, and arms were included as required by the experimental design. Post-hoc pair-wise comparisons or restricted ANOVAs

were performed to facilitate interpretation of higher-order interaction effects whenever appropriate. One-sample Student's *t*-tests were used to gauge performance against chance level. All analyses were conducted using SPSS (Version 16) with a Type I error rate of 0.05.

Acknowledgments

This work is supported by NIH grants R01MH083973, RO1NS48995, RO1NS41083-07, and RO1DA19362; Department of Defense grant W81XWH-071-1-0012; a grant from the Bumpus Foundation; and ETH Zurich.

References

- Aggleton JP, Vann SD, Oswald CJ, Good M. 2000. Identifying cortical inputs to the rat hippocampus that subservise allocentric spatial processes: A simple problem with a complex answer. *Hippocampus* **10**: 466–474.
- Bannerman DM, Deacon RM, Seeburg PH, Rawlins JN. 2003. GluR-A-deficient mice display normal acquisition of a hippocampus-dependent spatial reference memory task but are impaired during spatial reversal. *Behav Neurosci* **117**: 866–870.
- Bannerman DM, Niewoehner B, Lyon L, Romberg C, Schmitt WB, Taylor A, Sanderson DJ, Cottam J, Sprengel R, Seeburg PH, et al. 2008. NMDA receptor subunit NR2A is required for rapidly acquired spatial working memory but not incremental spatial reference memory. *J Neurosci* **28**: 3623–3630.
- Bastia E, Xu YH, Scibelli AC, Day YJ, Linden J, Chen JF, Schwarzschild MA. 2005. A crucial role for forebrain adenosine A(2A) receptors in amphetamine sensitization. *Neuropsychopharmacology* **30**: 891–900.
- Batista-Brito R, Machold R, Klein C, Fishell G. 2008. Gene expression in cortical interneuron precursors is prescient of their mature function. *Cereb Cortex* **18**: 2306–2317.
- Bellebaum C, Koch B, Schwarz M, Daum I. 2008. Focal basal ganglia lesions are associated with impairments in reward-based reversal learning. *Brain* **131** (Pt 3): 829–841.
- Berke JD, Breck JT, Eichenbaum H. 2009. Striatal versus hippocampal representations during win-stay maze performance. *J Neurophysiol* **101**: 1575–1587.
- Bliss TV, Collingridge GL. 1993. A synaptic model of memory: Long-term potentiation in the hippocampus. *Nature* **361**: 31–39.
- Boison D. 2007. Adenosine as a modulator of brain activity. *Drug News Perspect* **20**: 607–611.
- Canals M, Marcellino D, Fanelli F, Ciruela F, de Benedetti P, Goldberg SR, Neve K, Fuxe K, Agnati LF, Woods AS, et al. 2003. Adenosine A_{2A}-dopamine D₂ receptor-receptor heteromerization: Qualitative and quantitative assessment by fluorescence and bioluminescence energy transfer. *J Biol Chem* **278**: 46741–46749.
- Ciruela F, Casado V, Rodrigues RJ, Lujan R, Burgueno J, Canals M, Borycz J, Rebola N, Goldberg SR, Mallol J, et al. 2006. Presynaptic control of striatal glutamatergic neurotransmission by adenosine A₁-A_{2A} receptor heteromers. *J Neurosci* **26**: 2080–2087.
- Clarke HF, Robbins TW, Roberts AC. 2008. Lesions of the medial striatum in monkeys produce perseverative impairments during reversal learning similar to those produced by lesions of the orbitofrontal cortex. *J Neurosci* **28**: 10972–10982.
- Clatworthy PL, Lewis SJ, Brichard L, Hong YT, Izquierdo D, Clark L, Cools R, Aigbirhio FI, Baron JC, Fryer TD, et al. 2009. Dopamine release in dissociable striatal subregions predicts the different effects of oral methylphenidate on reversal learning and spatial working memory. *J Neurosci* **29**: 4690–4696.
- Cunha RA, Johansson B, van der Ploeg I, Sebastiao AM, Ribeiro JA, Fredholm BB. 1994. Evidence for functionally important adenosine A_{2A} receptors in the rat hippocampus. *Brain Res* **649**: 208–216.
- Cunha RA, Johansson B, Constantino MD, Sebastiao AM, Fredholm BB. 1996. Evidence for high-affinity binding sites for the adenosine A_{2A} receptor agonist [³H] CGS 21680 in the rat hippocampus and cerebral cortex that are different from striatal A_{2A} receptors. *Naunyn Schmiedebergs Arch Pharmacol* **353**: 261–271.
- d'Alcantara P, Ledent C, Swillens S, Schiffmann SN. 2001. Inactivation of adenosine A_{2A} receptor impairs long term potentiation in the accumbens nucleus without altering basal synaptic transmission. *Neuroscience* **107**: 455–464.
- Dalley JW, Cardinal RN, Robbins TW. 2004. Prefrontal executive and cognitive functions in rodents: Neural and neurochemical substrates. *Neurosci Biobehav Rev* **28**: 771–784.
- Dall'Igna OP, Fett P, Gomes MW, Souza DO, Cunha RA, Lara DR. 2007. Caffeine and adenosine A_{2A} receptor antagonists prevent beta-amyloid (25–35)-induced cognitive deficits in mice. *Exp Neurol* **203**: 241–245.
- de Mendonca A, Ribeiro JA. 1994. Endogenous adenosine modulates long-term potentiation in the hippocampus. *Neuroscience* **62**: 385–390.
- de Mendonca A, Ribeiro JA. 1997. Adenosine and neuronal plasticity. *Life Sci* **60**: 245–251.
- De Steno DA, Schmauss C. 2009. A role for dopamine D₂ receptors in reversal learning. *Neuroscience* **162**: 118–127.
- Dewar KM, Montreuil B, Grondin L, Reader TA. 1989. Dopamine D₂ receptors labeled with [³H]raclopride in rat and rabbit brains. Equilibrium binding, kinetics, distribution and selectivity. *J Pharmacol Exp Ther* **250**: 696–706.
- Dixon AK, Gubitza AK, Sirinathsinghji DJ, Richardson PJ, Freeman TC. 1996. Tissue distribution of adenosine receptor mRNAs in the rat. *Br J Pharmacol* **118**: 1461–1468.
- Dragatsis I, Zeitlin S. 2000. CaMKIIalpha-Cre transgene expression and recombination patterns in the mouse brain. *Genesis* **26**: 133–135.
- El-Ghundi M, O'Dowd BF, George SR. 2007. Insights into the role of dopamine receptor systems in learning and memory. *Rev Neurosci* **18**: 37–66.
- Ferre S, Fredholm BB, Morelli M, Popoli P, Fuxe K. 1997. Adenosine-dopamine receptor-receptor interactions as an integrative mechanism in the basal ganglia. *Trends Neurosci* **20**: 482–487.
- Ferre S, Karcz-Kubicha M, Hope BT, Popoli P, Burgueno J, Gutierrez MA, Casado V, Fuxe K, Goldberg SR, Lluís C, et al. 2002. Synergistic interaction between adenosine A_{2A} and glutamate mGlu5 receptors: Implications for striatal neuronal function. *Proc Natl Acad Sci* **99**: 11940–11945.
- Ferre S, Diamond I, Goldberg SR, Yao L, Hourani SM, Huang ZL, Urade Y, Kitchin I. 2007. Adenosine A_{2A} receptors in ventral striatum, hypothalamus and nociceptive circuitry implications for drug addiction, sleep and pain. *Prog Neurobiol* **83**: 332–347.
- Flajolet M, Wang Z, Futter M, Shen W, Nuangchamngong N, Bendor J, Wallach I, Nairn AC, Surmeier DJ, Greengard P. 2008. FGF acts as a co-transmitter through adenosine A_{2A} receptor to regulate synaptic plasticity. *Nat Neurosci* **11**: 1402–1409.
- Fontinha BM, Delgado-Garcia JM, Madronal N, Ribeiro JA, Sebastiao AM, Guart A. 2009. Adenosine A_{2A} receptor modulation of hippocampal CA₃-CA₁ synapse plasticity during associative learning in behaving mice. *Neuropsychopharmacology* **34**: 1865–1874.
- Fredholm BB, Chen JF, Cunha RA, Svenningsson P, Vaugeois JM. 2005. Adenosine and brain function. *Int Rev Neurobiol* **63**: 191–270.
- Fredholm BB, Chern Y, Franco R, Sitkovsky M. 2007. Aspects of the general biology of adenosine A_{2A} signaling. *Prog Neurobiol* **83**: 263–276.
- Gerevich Z, Wirkner K, Illes P. 2002. Adenosine A_{2A} receptors inhibit the N-methyl-D-aspartate component of excitatory synaptic currents in rat striatal neurons. *Eur J Pharmacol* **451**: 161–164.
- Ghanem N, Jarinova O, Amores A, Long Q, Hatch G, Park BK, Rubenstein JL, Ekker M. 2003. Regulatory roles of conserved intergenic domains in vertebrate *Dlx* bigene clusters. *Genome Res* **13**: 533–543.
- Gimenez-Llort L, Schiffmann SN, Shmidt T, Canela L, Camon L, Wassholm M, Canals M, Terasmaa A, Fernandez-Teruel A, Tobena A, et al. 2007. Working memory deficits in transgenic rats overexpressing human adenosine A_{2A} receptors in the brain. *Neurobiol Learn Mem* **87**: 42–56.
- Goldman-Rakic PS. 1995. Cellular basis of working memory. *Neuron* **14**: 477–485.
- Hill MN, Froese LM, Morrish AC, Sun JC, Floresco SB. 2006. Alterations in behavioral flexibility by cannabinoid CB1 receptor agonists and antagonists. *Psychopharmacology (Berl)* **187**: 245–259.
- Hillion J, Canals M, Torvinen M, Casado V, Scott R, Terasmaa A, Hansson A, Watson S, Olah ME, Mallol J, et al. 2002. Coaggregation, coinertalization, and codensitization of adenosine A_{2A} receptors and dopamine D₂ receptors. *J Biol Chem* **277**: 18091–18097.
- Hodges H, Sowinski P, Sinden JD, Netto CA, Fletcher A. 1995. The selective 5-HT₃ receptor antagonist, WAY100289, enhances spatial memory in rats with ibotenate lesions of the forebrain cholinergic projection system. *Psychopharmacology (Berl)* **117**: 318–332.
- Houchi H, Babovic D, Pierrefiche O, Ledent C, Daoust M, Naassila M. 2005. CB1 receptor knockout mice display reduced ethanol-induced conditioned place preference and increased striatal dopamine D₂ receptors. *Neuropsychopharmacology* **30**: 339–349.
- Kachroo A, Orlando LR, Grandy DK, Chen JF, Young AB, Schwarzschild MA. 2005. Interactions between metabotropic glutamate 5 and adenosine A_{2A} receptors in normal and parkinsonian mice. *J Neurosci* **25**: 10414–10419.
- Kellendonk C, Simpson EH, Polan HJ, Malleret G, Vronskaya S, Winiger V, Moore H, Kandel ER. 2006. Transient and selective overexpression of

- dopamine D₂ receptors in the striatum causes persistent abnormalities in prefrontal cortex functioning. *Neuron* **49**: 603–615.
- Kesner RP. 2009. The posterior parietal cortex and long-term memory representation of spatial information. *Neurobiol Learn Mem* **91**: 197–206.
- Kreitzer AC, Malenka RC. 2005. Dopamine modulation of state-dependent endocannabinoid release and long-term depression in the striatum. *J Neurosci* **25**: 10537–10545.
- Kreitzer AC, Malenka RC. 2007. Endocannabinoid-mediated rescue of striatal LTD and motor deficits in Parkinson's disease models. *Nature* **445**: 643–647.
- Kruzich PJ, Mitchell SH, Younkin A, Grandy DK. 2006. Dopamine D₂ receptors mediate reversal learning in male C57BL/6J mice. *Cogn Affect Behav Neurosci* **6**: 86–90.
- Lee B, Groman S, London ED, Jentsch JD. 2007. Dopamine D₂/D₃ receptors play a specific role in the reversal of a learned visual discrimination in monkeys. *Neuropsychopharmacology* **32**: 2125–2134.
- Lerner TN, Horne EA, Stella N, Kreitzer AC. 2010. Endocannabinoid signaling mediates psychomotor activation by adenosine A_{2A} antagonists. *J Neurosci* **30**: 2160–2164.
- Lidow MS, Goldman-Rakic PS, Rakic P, Innis RB. 1989. Dopamine D₂ receptors in the cerebral cortex: Distribution and pharmacological characterization with [³H]raclopride. *Proc Natl Acad Sci* **86**: 6412–6416.
- Lopes LV, Halldner L, Rebola N, Johansson B, Ledent C, Chen JF, Fredholm BB, Cunha RA. 2004. Binding of the prototypical adenosine A_{2A} receptor agonist CGS 21680 to the cerebral cortex of adenosine A₁ and A_{2A} receptor knockout mice. *Br J Pharmacol* **141**: 1006–1014.
- Lovinger DM. 2010. Neurotransmitter roles in synaptic modulation, plasticity and learning in the dorsal striatum. *Neuropharmacology* **58**: 951–961.
- Lynch MA. 2004. Long-term potentiation and memory. *Physiol Rev* **84**: 87–136.
- Marie RM, Defer GL. 2003. Working memory and dopamine: Clinical and experimental clues. *Curr Opin Neurol* **16** (Suppl 2): S29–S35.
- McDonald RJ, King AL, Foong N, Rizos Z, Hong NS. 2008. Neurotoxic lesions of the medial prefrontal cortex or medial striatum impair multiple-location place learning in the water task: Evidence for neural structures with complementary roles in behavioural flexibility. *Exp Brain Res* **187**: 419–427.
- Mehta MA, Montgomery AJ, Kitamura Y, Grasby PM. 2008. Dopamine D₂ receptor occupancy levels of acute sulpiride challenges that produce working memory and learning impairments in healthy volunteers. *Psychopharmacology (Berl)* **196**: 157–165.
- Naassila M, Ledent C, Daoust M. 2002. Low ethanol sensitivity and increased ethanol consumption in mice lacking adenosine A_{2A} receptors. *J Neurosci* **22**: 10487–10493.
- Norenberg W, Wirkner K, Assmann H, Richter M, Illes P. 1998. Adenosine A_{2A} receptors inhibit the conductance of NMDA receptor channels in rat neostriatal neurons. *Amino Acids* **14**: 33–39.
- Ohtsuka N, Tansky MF, Kuang H, Kourrich S, Thomas MJ, Rubenstein JL, Ekker M, Leeman SE, Tsien JZ. 2008. Functional disturbances in the striatum by region-specific ablation of NMDA receptors. *Proc Natl Acad Sci* **105**: 12961–12966.
- Packard MG, Knowlton BJ. 2002. Learning and memory functions of the basal ganglia. *Annu Rev Neurosci* **25**: 563–593.
- Packard MG, McGaugh JL. 1992. Double dissociation of fornix and caudate nucleus lesions on acquisition of two water maze tasks: Further evidence for multiple memory systems. *Behav Neurosci* **106**: 439–446.
- Pietro Paolo S, Sun Y, Li R, Brana C, Feldon J, Yee BK. 2009. Limited impact of social isolation on Alzheimer-like symptoms in a triple transgenic mouse model. *Behav Neurosci* **123**: 181–195.
- Prediger RD, Batista LC, Takahashi RN. 2005a. Caffeine reverses age-related deficits in olfactory discrimination and social recognition memory in rats. Involvement of adenosine A₁ and A_{2A} receptors. *Neurobiol Aging* **26**: 957–964.
- Prediger RD, Da Cunha C, Takahashi RN. 2005b. Antagonistic interaction between adenosine A_{2A} and dopamine D₂ receptors modulates the social recognition memory in reserpine-treated rats. *Behav Pharmacol* **16**: 209–218.
- Prediger RD, Fernandes D, Takahashi RN. 2005c. Blockade of adenosine A_{2A} receptors reverses short-term social memory impairments in spontaneously hypertensive rats. *Behav Brain Res* **159**: 197–205.
- Rebola N, Canas PM, Oliveira CR, Cunha RA. 2005. Different synaptic and subsynaptic localization of adenosine A_{2A} receptors in the hippocampus and striatum of the rat. *Neuroscience* **132**: 893–903.
- Rebola N, Lujan R, Cunha RA, Mulle C. 2008. Adenosine A_{2A} receptors are essential for long-term potentiation of NMDA-EPSCs at hippocampal mossy fiber synapses. *Neuron* **57**: 121–134.
- Ribeiro AC, Pfaff DW, Devidze N. 2009. Estradiol modulates behavioral arousal and induces changes in gene expression profiles in brain regions involved in the control of vigilance. *Eur J Neurosci* **29**: 795–801.
- Rodrigues RJ, Alfaro TM, Rebola N, Oliveira CR, Cunha RA. 2005. Co-localization and functional interaction between adenosine A_{2A} and metabotropic group 5 receptors in glutamatergic nerve terminals of the rat striatum. *J Neurochem* **92**: 433–441.
- Rosin DL, Robeva A, Woodard RL, Guyenet PG, Linden J. 1998. Immunohistochemical localization of adenosine A_{2A} receptors in the rat central nervous system. *J Comp Neurol* **40**: 163–186.
- Rosin DL, Hettinger BD, Lee A, Linden J. 2003. Anatomy of adenosine A_{2A} receptors in brain: Morphological substrates for integration of striatal function. *Neurology* **61** (11 Suppl 6): S12–S18.
- Rowe JB, Toni I, Josephs O, Frackowiak RS, Passingham RE. 2000. The prefrontal cortex: Response selection or maintenance within working memory? *Science* **288**: 1656–1660.
- Save E, Poucet B. 2009. Role of the parietal cortex in long-term representation of spatial information in the rat. *Neurobiol Learn Mem* **91**: 172–178.
- Schiffmann SN, Jacobs O, Vanderhaeghen JJ. 1991a. Striatal restricted adenosine A₂ receptor (RDC8) is expressed by enkephalin but not by substance P neurons: An in situ hybridization histochemistry study. *J Neurochem* **57**: 1062–1067.
- Schiffmann SN, Libert F, Vassart G, Vanderhaeghen JJ. 1991b. Distribution of adenosine A₂ receptor mRNA in the human brain. *Neurosci Lett* **130**: 177–181.
- Schiffmann SN, Fisone G, Moresco R, Cunha RA, Ferre S. 2007. Adenosine A_{2A} receptors and basal ganglia physiology. *Prog Neurobiol* **83**: 277–292.
- Sebastiao AM, Ribeiro JA. 1996. Adenosine A₂ receptor-mediated excitatory actions on the nervous system. *Prog Neurobiol* **48**: 167–189.
- Shen HY, Coelho JE, Ohtsuka N, Canas PM, Day YJ, Huang QY, Rebola N, Yu L, Boison D, Cunha RA, et al. 2008a. A critical role of the adenosine A_{2A} receptor in extrastriatal neurons in modulating psychomotor activity as revealed by opposite phenotypes of striatum and forebrain A_{2A} receptor knock-outs. *J Neurosci* **28**: 2970–2975.
- Shen W, Flajolet M, Greengard P, Surmeier DJ. 2008b. Dichotomous dopaminergic control of striatal synaptic plasticity. *Science* **321**: 848–851.
- Simpson EH, Kellendonk C, Kandel E. 2010. A possible role for the striatum in the pathogenesis of the cognitive symptoms of schizophrenia. *Neuron* **65**: 585–596.
- Singer P, Boison D, Mohler H, Feldon J, Yee BK. 2009. Deletion of glycine transporter 1 (GlyT1) in forebrain neurons facilitates reversal learning: Enhanced cognitive adaptability? *Behav Neurosci* **123**: 1012–1027.
- Tebano MT, Martire A, Rebola N, Pepponi R, Domenici MR, Gro MC, Schwarzschild MA, Chen JF, Cunha RA, Popoli P. 2005. Adenosine A_{2A} receptors and metabotropic glutamate 5 receptors are co-localized and functionally interact in the hippocampus: A possible key mechanism in the modulation of N-methyl-D-aspartate effects. *J Neurochem* **95**: 1188–1200.
- Tebano MT, Martire A, Chiodi V, Pepponi R, Ferrante A, Domenici MR, Frank C, Chen JF, Ledent C, Popoli P. 2009. Adenosine A_{2A} receptors enable the synaptic effects of cannabinoid CB1 receptors in the rodent striatum. *J Neurochem* **110**: 1921–1930.
- Varvel SA, Lichtman AH. 2002. Evaluation of CB1 receptor knockout mice in the Morris water maze. *J Pharmacol Exp Ther* **301**: 915–924.
- Wang JH, Ma YY, van den Buuse M. 2006. Improved spatial recognition memory in mice lacking adenosine A_{2A} receptors. *Exp Neurol* **199**: 438–445.
- Wei CJ, Li W, Chen JF. 2011. Normal and abnormal functions of adenosine receptors in the central nervous system revealed by genetic knockout studies. *Biochim Biophys Acta* **1808**: 1358–1379.
- Wirkner K, Assmann H, Koles L, Gerevich Z, Franke H, Norenberg W, Boehm R, Illes P. 2000. Inhibition by adenosine A_{2A} receptors of NMDA but not AMPA currents in rat neostriatal neurons. *Br J Pharmacol* **130**: 259–269.
- Wirkner K, Gerevich Z, Krause T, Gunther A, Koles L, Schneider D, Norenberg W, Illes P. 2004. Adenosine A_{2A} receptor-induced inhibition of NMDA and GABA_A receptor-mediated synaptic currents in a subpopulation of rat striatal neurons. *Neuropharmacology* **46**: 994–1007.
- Xu J, Zhu Y, Contractor A, Heinemann SF. 2009. mGluR5 has a critical role in inhibitory learning. *J Neurosci* **29**: 3676–3684.
- Yang JN, Chen JF, Fredholm BB. 2009. Physiological roles of A₁ and A_{2A} adenosine receptors in regulating heart rate, body temperature, and locomotion as revealed using knockout mice and caffeine. *Am J Physiol Heart Circ Physiol* **296**: H1141–H1149.
- Yee BK, Singer P, Chen JF, Feldon J, Boison D. 2007. Transgenic overexpression of adenosine kinase in brain leads to multiple learning

- impairments and altered sensitivity to psychomimetic drugs. *Eur J Neurosci* **26**: 3237–3252.
- Yu L, Shen HY, Coelho JE, Araujo IM, Huang QY, Day YJ, Rebola N, Canas PM, Rapp EK, Ferrara J, et al. 2008. Adenosine A_{2A} receptor antagonists exert motor and neuroprotective effects by distinct cellular mechanisms. *Ann Neurol* **63**: 338–346.
- Yu C, Gupta J, Chen JF, Yin HH. 2009. Genetic deletion of A_{2A} adenosine receptors in the striatum selectively impairs habit formation. *J Neurosci* **29**: 15100–15103.
- Zandbelt BB, Vink M. 2010. On the role of the striatum in response inhibition. *PLoS One* **5**: e13848. doi: 10.1371/journal.pone.0013848.
- Zerucha T, Stuhmer T, Hatch G, Park BK, Long Q, Yu G, Gambarotta A, Schultz JR, Rubenstein JL, Ekker M. 2000. A highly conserved enhancer in the *Dlx5/Dlx6* intergenic region is the site of cross-regulatory interactions between *Dlx* genes in the embryonic forebrain. *J Neurosci* **20**: 709–721.
- Zhou SJ, Zhu ME, Shu D, Du XP, Song XH, Wang XT, Zheng RY, Cai XH, Chen JF, He JC. 2009. Preferential enhancement of working memory in mice lacking adenosine A_{2A} receptors. *Brain Res* **1303**: 74–83.

Received January 6, 2011; accepted in revised form April 11, 2011.