

E2Fs regulate the expression of genes involved in differentiation, development, proliferation, and apoptosis

Heiko Müller,^{1,3} Adrian P. Bracken,^{1,5} Richard Vernell,^{1,5} M. Cristina Moroni,¹ Fred Christians,² Emanuela Grassilli,¹ Elena Prosperini,¹ Elena Vigo,¹ Jonathan D. Oliner,^{2,4} and Kristian Helin^{1,6}

¹Department of Experimental Oncology, European Institute of Oncology, 20141 Milan, Italy; ²Affymetrix Inc. Santa Clara, California 95051, USA

The retinoblastoma protein (pRB) and its two relatives, p107 and p130, regulate development and cell proliferation in part by inhibiting the activity of E2F-regulated promoters. We have used high-density oligonucleotide arrays to identify genes in which expression changed in response to activation of E2F1, E2F2, and E2F3. We show that the E2Fs control the expression of several genes that are involved in cell proliferation. We also show that the E2Fs regulate a number of genes involved in apoptosis, differentiation, and development. These results provide possible genetic explanations to the variety of phenotypes observed as a consequence of a deregulated pRB/E2F pathway.

[Key Words: E2F; retinoblastoma; microarray; target gene bias]

Received November 1, 2000; revised version accepted December 18, 2000.

One of the hallmarks of cancer cells is their insensitivity to antiproliferative signals, such as antigrowth factors (Hanahan and Weinberg 2000). Scientific results obtained in the last decade have pointed to a central role for the retinoblastoma protein (pRB), and its relatives, p107 and p130, in mediating these antiproliferative signals. Consistent with this role, the pRB pathway is deregulated in the majority of human cancers (for review, see Sherr 1996). In addition to the pRB family members, this pathway involves cyclin-dependent kinase (CDK) inhibitors of the INK4 family, which act as negative regulators of cell proliferation, and the positively acting D-type cyclins that, in association with CDK4 or CDK6, form active kinase complexes. The downstream effectors in the pathway are the E2F transcription factors (for review, see Dyson 1998; Nevins 1998). The E2Fs regulate the timely expression of a series of genes whose products are essential for cell proliferation (for review, see Helin 1998).

When associated with pRB family members, the E2Fs function as transcriptional repressors, whereas 'free' E2F activates transcription (for review, see Dyson 1998; Müller and Helin 2000). Repression of E2F-regulated promoters by pRB family members is alleviated on their

phosphorylation by a D-type cyclin-associated kinase, and this kinase activity is suppressed by the INK4 CDK inhibitors. Because all known mutations in this pathway lead to deregulated E2F activity, it is believed that most human tumors contain aberrant levels of E2F-regulated transcripts. In addition to controlling cell proliferation, the pRB family members also regulate apoptosis, differentiation, and development (for review, see Lipinski and Jacks 1999). In agreement with its role as a tumor suppressor, mice that carry one disrupted *Rb1* allele are predisposed to cancer. However, *Rb1* nullizygous embryos die between day 13 and day 15 of gestation. These embryos show signs of maturation defects in erythropoiesis, extensive apoptotic cell death, and hyperproliferation in the nervous system, liver, and ocular lens. In contrast to the strong phenotype that results from the lack of pRb, mice nullizygous for p107 or p130 do not have a discernible phenotype in the C57BL/6 genetic background. Moreover, mice that lack both p107 and p130 are defective in endochondrial bone development because of a deficiency in chondrocyte differentiation, which suggests that one of the physiological roles of p107 and p130 is to restrict chondrocyte proliferation and to induce terminal differentiation. pRB family members also play a role in the terminal differentiation of myoblasts, adipocytes, and cells of a variety of hematopoietic lineages (Lipinski and Jacks 1999). It has been suggested that the pRB family members regulate terminal differentiation through two distinct mechanisms. The first involves regulation of cell cycle progression and apoptosis through repression of E2F-dependent promoters. The

Present addresses: ³Pharmacia Corporation, 20014 Nerviano, Italy; ⁴Amgen Inc., Thousand Oaks, CA 91320, USA.

⁵These authors contributed equally to this work.

⁶Corresponding author.

E-MAIL khelin@ieo.it; FAX +39-02-5748-9851.

Article and publication are at www.genesdev.org/cgi/doi/10.1101/gad.864201.

second is related to the differentiation process and results from interaction of pRB family members with other transcriptional regulators, such as MyoD, C/EBPs and HBP1 (Lipinski and Jacks 1999).

There are six members of the E2F transcription factor family, E2F1–E2F6 (for review, see Dyson 1998; Müller and Helin 2000). All six E2Fs contain two highly conserved domains that are involved in sequence-specific DNA binding and dimerization with DP proteins. Association of the E2Fs with one of the two known DP proteins is required for high-affinity, sequence-specific DNA binding, and in the case of E2F1–E2F5, association with members of the pRB family. The E2Fs can be classified into three subgroups based on their structure, affinity for members of the pRB family, expression pattern, and putative function. The first group consists of E2F1, E2F2, and E2F3, whose expression is cell-growth regulated. At physiological levels these E2Fs associate exclusively with pRB, and ectopic expression of each of these three E2Fs is sufficient to induce S-phase in serum-starved tissue culture cells. This group of E2Fs are mainly involved in the regulation of cell proliferation and apoptosis (see, e.g., Leone et al. 1998; Tsai et al. 1998; Wang et al. 1998; Yamasaki et al. 1998; Humbert et al. 2000b). E2F4 and E2F5, members of the second group, bind to all three pRB family members, and, although they are expressed throughout the cell cycle, these E2Fs are unable to induce S-phase in quiescent cells. Mice that are nullizygous for *E2f4* and *E2f5* have specific terminal-differentiation defects. *E2f4*^{-/-} mice show abnormalities in the development of hematopoietic lineages and in gut epithelium and dramatic shortening and broadening of the snout (Humbert et al. 2000a; Rempel et al. 2000). *E2f5*^{-/-} mice develop hydrocephalus as a result of excessive cerebrospinal fluid production (Lindeman et al. 1998). E2F6, the only member of the third group, lacks the domains that are involved in transactivation and binding to the pRB family members. The physiological function of E2F6 is unknown. However, as its structure suggests, it is a repressor of E2F-dependent transcription (Müller and Helin 2000).

E2F DNA-binding sites have been identified in promoters of many genes that are central to the regulation of cell cycle progression. Among the putative targets for E2F regulation are genes whose products are essential for DNA replication and cell cycle control (Helin 1998; Nevins 1998). Expression of the majority of the known E2F target genes is regulated during cell growth, and they are activated just before S-phase. A notable exception is the p19/p14ARF gene, whose expression is not cell-growth regulated but nevertheless is controlled by E2F (Sherr 1998). Because the E2F transcription factors are essential for proliferation, identification of genes whose expression pattern changes as a consequence of deregulated E2F activity may ultimately lead to a better understanding of cell proliferation.

Thus, we have used oligonucleotide arrays to monitor changes in gene expression following activation of E2F1, E2F2, or E2F3.

Results

Generation and characterization of inducible E2F cell lines

We have shown previously that fusion proteins that contain the estrogen receptor ligand-binding domain (ER) and E2F1, E2F2, or E2F3 were inactive in the absence of 4-hydroxy tamoxifen (OHT) and rapidly activated by addition of OHT (Vigo et al. 1999). To identify genes whose expression is altered as a result of a deregulated pRB/E2F pathway, we generated a series of cell lines by infecting human U2OS cells with retroviruses that express N-terminally HA-tagged ER-E2F1, ER-E2F2, or ER-E2F3. Following puromycin selection, individual clones were isolated, and the expression of the E2F proteins was confirmed using antibodies that specifically recognize E2F1, E2F2, or E2F3. As can be seen in Figure 1A, ER-E2F fusion proteins were expressed at approximately five- to 20-fold higher levels as compared with the endogenous level of the respective E2Fs. The highest level of overexpression is observed for the clones carrying ER-E2F2.

Several experiments were performed to characterize the cell lines. In Figure 1B, HA-ER-E2F3 is shown to change its subcellular localization in response to addition of OHT. In untreated cells, ER-E2F3 was found almost exclusively in the cytoplasm, whereas 8 hr after the addition of OHT, the majority of the ER-E2F3 protein was detected in the nucleus (Figure 1B). Similar results were obtained for the E2F1 and E2F2 fusion proteins (data not shown).

To test the transcriptional activity of the ER-E2F fusion proteins, transactivation assays were performed using a sensitive synthetic E2F-dependent luciferase reporter construct (pGL3 TATA basic 6xE2F). As shown in Figure 1C, OHT treatment led to a significant increase in luciferase activity in all cell lines that were tested. These results showed that the ER-E2F fusion proteins are activated by OHT and suggested that the cell lines would be suitable for the identification of E2F target genes.

To identify E2F target genes and to assess changes in mRNA expression after E2F activation, we used hybridization to oligonucleotide microarrays (Lockhart et al. 1996). Using this technique it was possible to determine changes in the expression pattern of up to 35,000 different genes after activation of specific E2Fs. To determine the optimum duration of OHT treatment before isolation of RNA for hybridization, we investigated the kinetics of mRNA accumulation of a previously known E2F target gene, Cyclin E1 (*CCNE1*) (Dyson 1998; Helin 1998). We observed a strong induction of the Cyclin E1 transcript as early as 4 hr after addition of OHT by semi-quantitative PCR (Figure 1D), and by Northern blotting (data not shown). Although in most clones the level of Cyclin E1 induction had reached its maximum after 4 hr, in some clones the 8-hr timepoint showed slightly higher levels of Cyclin E1 transcript. We did not detect significant changes in cell cycle profiles in any of the cell lines 8 hr after addition of OHT (data not shown). There-




Figure 1. Characterization of cell lines. (A) Expression of estrogen receptor E2F (ER-E2F) fusion proteins. Western blot analysis of whole cell lysates using antibodies specific for E2F1, E2F2, or E2F3 show the expression levels of the ER-E2F fusion proteins as compared to endogenous E2Fs. (B) Nuclear translocation of ER-E2F3 fusion protein. Immunostaining of HA-tagged ER-E2F3 fusion protein was performed using HA-specific antibody 12CA5. The fusion protein was nuclear after the addition of 4-hydroxy tamoxifen (OHT). (C) ER-E2F fusion proteins are transcriptionally active. Fold changes in transcriptional activity were determined by dividing normalized luciferase activity from induced cells (300 nM OHT for 9 hr) by the normalized luciferase activity from uninduced cells. (D) Induction of Cyclin E1 transcription by activated fusion proteins. Semiquantitative RT-PCR of endogenous Cyclin E1 was performed on RNA isolated from cells at the indicated hours after induction of E2F activity by OHT. We used 27 PCR cycles to detect the Cyclin E1 transcript in the ER-E2F-expressing cell lines, whereas 30 PCR cycles were used to detect the transcript in the control (U2OS) samples.

fore, 8 hr was determined to be optimum duration for OHT treatment and was used for these studies.

To perform the microarray analysis, hybridization targets were prepared from total RNA isolated from the nine test clones and three independent control cultures of U2OS cells 8 hr after addition of OHT. Each target was hybridized to Hu6800, Hu35ksubA, Hu35ksubB, Hu35ksubC, and Hu35ksubD arrays containing oligonucleotide probe pairs for ~35,000 different genes.

Statistical analysis of the data

The McNemar test was used to determine the statistical quality of the data (see Materials and Methods). In this test we used the difference calls provided by the GeneChip® software, which were moderately induced (MI) or induced (I) for up-regulated genes or decreased (D) or moderately decreased (MD) for down-regulated genes, to calculate the statistical significance of the expression

data. By combining the statistically significant changes (>99% confidence) with the fold change in expression level for any given gene, we chose a minimal fold change (cutoff) to keep the number of false positives close to an acceptable minimum. The number of genes whose expression was modulated in response to induction of each of the ER-E2Fs as a function of fold change cutoff is shown in Figure 2. For one single experiment (Fig. 2A) it was necessary to use a cutoff of at least fivefold induction or repression to keep the number of false positives close to an acceptable minimum. For duplicate experiments (Fig. 2B), reproducible modulation of transcript levels were scored using a cutoff of threefold without significant risk of false positives. Figure 2C shows that triplicate measurements result in a further increase in sensitivity. Here, at a cutoff of twofold change, no genes were found to be modulated in the U2OS control, and the expression of almost no genes were modified in the randomized data lists.

Our statistical analyses showed that the E2Fs induced a significant change in the expression of a large number of genes in U2OS cells. Based on the analysis presented above, we used the following criteria for characterization of a gene as induced or repressed. For duplicate measurements, a minimum of threefold change and a difference call of I or MI for up-regulated genes or a cutoff of threefold decrease and D or MD difference calls for down-regulated genes were used. The cutoff was lowered to twofold when triplicate measurements were performed. Using these criteria, we found the expression of 913 genes to be modulated by E2F1, 292 by E2F2, and 652 by E2F3 (see Table 4 in supplementary information). The expression of a subset of 162 genes was altered by all three E2Fs. We characterized 498 genes as selectively modulated by E2F1, 55 genes by E2F2, and 232 genes by E2F3. The total number of genes regulated by at least one E2F was 1240. We estimate that this number represents ~7% of the genes expressed in U2OS cells (~19,000 of the ~35,000 unigenes represented on the microarrays were expressed at detectable levels). From this list, we selected 42 genes (representing the spectrum from poorly modulated to strongly modulated) for verification analysis (Table 1 and below).

Verification of microarray results by Northern blot analysis

Two potential sources of false positive signals in microarray studies are cross-hybridization between closely related genes and clonal selection. To confirm the array results, Northern blot analysis was performed using total (Fig. 3A) or poly A (Fig. 3B) RNA prepared from cell lines that express ER-E2F1, ER-E2F2, or ER-E2F3 that were harvested at 0 hr, 4 hr, and 8 hr after OHT treatment. The results shown in Figure 3, A and B do not only confirm the data from the microarray studies, they also indicate that the fold changes in gene expression calculated by the GeneChip® software underestimate the level of modulation as determined by Northern blot

analysis. As can be seen in Figure 3 and in Table 1, strong induction of mRNA levels was detected for *FST*, *EGR1*, *EZH2*, *CASP3*, *TEAD4*, *DYRK2*, *PTPNS1*, *PRKAR2B*, *RANBPM*, and *BMP2* after ER-E2F activation. The induction of *FST*, *EZH2*, *CASP3*, *TEAD4*, *DYRK2*, *PTPNS1*, *PRKAR2B*, and *RANBPM* was unaffected by addition of cycloheximide during the OHT treatment, which suggests that these genes are direct transcriptional targets of E2F. We also observed lower expression levels of certain genes as a result of ER-E2F activation (e.g., *PAI1*, *CTGF*, *EPLIN*, *TGFB2*, *BCL3*, and *INHBA*; Fig. 3; Table 1). In this case, however, cycloheximide treatment prevented the apparent repression of gene expression upon ER-E2F induction, which suggests that intermediate regulatory factors may be involved.

In 39 out of 40 cases using Northern blot analysis, increased E2F activity was confirmed to alter the expression of a particular gene identified by the microarray method (Table 2). In several cases, a gene that was predicted to be selectively modulated by one E2F transcription factor by the microarray analysis was shown to be regulated by the other E2Fs in Northern blotting studies (e.g., *DMRT1* by E2F1, *BMP2* by E2F2, *CTGF* by E2F3). We did not detect any discrepancies between the microarray and Northern blot analyses with respect to the direction of transcript modulation (induction vs. repression). Based on these experiments, we estimate the rate of false positives in the screen to be ~2%.

To ensure that the E2F-induced gene expression changes that were observed were not characteristic of only U2OS or transformed cells, we generated pools of human diploid fibroblasts (WI38) that were infected with retroviral constructs expressing ER-E2F1. Poly A RNA was isolated from infected WI38 cells at 0 hr, 4 hr, 8 hr, and 12 hr after addition of OHT, and Northern blots were performed using probes generated from a subset of the genes whose expression was modulated in U2OS cells after ER-E2F induction. As can be seen in Figure 4A, in WI38 cells the level of transcript modulation was equal to, and in some cases higher than (*CCNE1*, *APAF1*, and *CASP3*), that observed in U2OS cells. These results suggest that the set of E2F-regulated genes identified in U2OS cells is not specific for this cell line but, rather, that the E2Fs induce similar transcriptional changes in nontransformed cells.

We also were concerned that the differences in transcriptional activity detected in the microarray experiments might be caused by overexpression of the ER-E2F fusion proteins and that the genes identified were not regulated by physiological levels of E2F. We sought to address this concern by repressing endogenous E2F activity by overexpression of pRB. This assumes that genes that are up-regulated by induction of E2F should be down-regulated by expression of pRB. Conversely, genes that are down-regulated by induction of E2F should be up-regulated by expression of pRB. To test this possibility, we used a U2OS cell line expressing a pRB mutant (pRBΔCDK) in a tetracycline-dependent manner (Lukas




Figure 2. Analysis of gene expression changes. The columns represent numbers of regulated genes (hits) in one (A), two (B), and three (C) independent measurements. A gene is called regulated when its difference call is induced/moderately induced (I/MI) for up-regulated genes and decreased/moderately decreased (D/MD) for down-regulated genes in all replicates and the fold change (FC) is equal to or beyond the fold-change cutoff in all replicates ($FC \geq \text{cutoff}$ for up-regulated genes or $FC \leq \text{cutoff}$ for down-regulated genes). Checkered bars represent the data. Black bars represent U2OS noise lists (derived from comparing U2OS control chips to each other). White bars represent randomized data lists. See Materials and Methods for details.

et al. 1999). As shown in Figure 4B, two genes up-regulated by E2F (*CCNE1*, *EGR1*) were repressed by pRB Δ cdk expression.

Conversely, the levels of two E2F-repressed genes (*PAI1*, *CTGF*) were strongly induced as early as 4 hr after tetracycline removal at a time when the pRB Δ CDK pro-

Table 1. Fold change and difference for a selected number of DNA chip entries

Symbol	Acc. No.	Unigene	Fold change	D-call	E2F1	Fold change	E2F2	D-call	E2F2	Fold change	E2F3	D-call	E2F3	Description
APAF1	AA285132	77579	3/4.4/2.4	I/I/I	E2F1	1.2/1.6/1.0	E2F2	NC/NC/MD	E2F2	1.3/1.5/-1.4	E2F3	NC/NC/MD	E2F3	Apoptotic protease activating factor
ASH2L	AA397724	6856	6.6/14.3/8.2	I/I/I	E2F1	1.5/3.0/2.8	E2F2	NC/NC/I	E2F2	3.7/7.5/3.3	E2F3	I/I/I	E2F3	Ash2 (absent, small, or homeotic, Drosophila, homolog)-like
BCL3	AA293187	31210	-13/11.5/-7.3	D/NC/MD	E2F1	-7/-12.2/-1.5	E2F2	MD/NC/NC	E2F2	-12.4/-11.4/-6.1	E2F3	D/D/D	E2F3	B-cell CLL/lymphoma 3
BMP2	M22489	73853	6.8/6.4/13.2	I/I/I	E2F1	2.4/2.8/3.8	E2F2	NC/MI/I	E2F2	3.9/4.5/1.8	E2F3	I/I/NC	E2F3	Bone morphogenetic protein 2
BMPR1A	Z22535	2534	9.0/7.1/6.1	I/I/I	E2F1	3.7/1.1/1.3	E2F2	I/NC/NC	E2F2	3.5/3.4/1.4	E2F3	I/I/NC	E2F3	Bone morphogenetic protein receptor, type IA
CASP3	U13737	74552	14.5/12.8/15	I/I/I	E2F1	3.0/7.9/6.7	E2F2	NC/MI/I	E2F2	16.6/17.8/4	E2F3	I/MI/I	E2F3	Caspase 3, apoptosis-related cysteine protease
CASP7	U67319	9216	2.8/2.8/1.5	I/I/NC	E2F1	1.6/1.2/1.3	E2F2	NC/NC/I	E2F2	2.0/2.3/1.7	E2F3	I/I/I	E2F3	Caspase 7, apoptosis-related cysteine protease
CAV1	Z18951	74034	-2.0/-3.4/2.8	D/D/D	E2F1	-1.4/-3.7/-1.2	E2F2	NC/D/D	E2F2	-4.8/1.3/2.5	E2F3	D/NC/D	E2F3	Caveolin 1, caveolae protein, 22kD
CD58	Y00636	75626	4.6/5.8/3.8	I/I/I	E2F1	2.3/4.0/2.6	E2F2	I/I/I	E2F2	3.9/4.4/3.3	E2F3	I/I/I	E2F3	CD58 antigen, (lymphocyte function-associated antigen 3)
CHES1	U68723	211773	-9.9/-11.7/-4.0	MD/D/MD	E2F1	-4.9/-7.7/-1.4	E2F2	NC/NC/D	E2F2	-7.0/-1.8/-8.3	E2F3	NC/NC/D	E2F3	Checkpoint suppressor 1
CTGF	AA449789	75511	-1.8/-3.5/2.9	D/D/D	E2F1	1.2/-1.4/-1.3	E2F2	NC/D/NC	E2F2	1.3/-2.1/-1.1	E2F3	NC/D/NC	E2F3	Connective tissue growth factor
DMRT1	AA429597	98586	1.3/-1.4/-1.7	NC/NC/D	E2F1	5.1/3.4/2.9	E2F2	I/MI/I	E2F2	2.2/1.0/1.0	E2F3	NC/NC/NC	E2F3	Doublesex and mab-3 related transcription factor 1
DYRK1A	D86550	75842	3.7/5.3/3.9	I/I/I	E2F1	1.3/3.5/1.8	E2F2	NC/MI/I	E2F2	3.0/3.8/2.2	E2F3	I/I/I	E2F3	Dual-specificity tyrosine-(Y)-phosphorylation regulated kinase 1A
DYRK2	AA609151	173135	3.7/3.2/3.9	I/I/I	E2F1	3.6/4.8/3.7	E2F2	I/I/I	E2F2	6.4/4.6/5.0	E2F3	I/I/I	E2F3	Dual-specificity tyrosine-(Y)-phosphorylation regulated kinase 2
EGR1	X52541	738	4.5/5.9/4.9	I/MI/I	E2F1	2.0/2.6/3.1	E2F2	NC/NC/MI	E2F2	2.5/3.0/2.8	E2F3	I/MI/MI	E2F3	Early growth response 1
EPLIN	AA157836	10706	-2.7/-6.5/4.6	D/D/D	E2F1	-1.2/-3.0/-1.0	E2F2	NC/D/NC	E2F2	-3.0/-3.6/-2.4	E2F3	D/D/D	E2F3	Epithelial protein lost in neoplasm beta
EZH2	U61145	72256	4.5/6.2/4.7	I/I/I	E2F1	3.1/6.4/5.5	E2F2	I/I/I	E2F2	4.2/6.7/4.6	E2F3	I/I/I	E2F3	Enhancer of zeste (Drosophila) homolog 2
FGFR2	M87770	278581	2.9/5.3/3.1	I/NC/I	E2F1	1.2/1.4/1.4	E2F2	NC/NC/NC	E2F2	1.9/1.9/1.4	E2F3	I/MI/NC	E2F3	Fibroblast growth factor receptor 2
FGFR3	M64347	1420	8.1/7.2/6.0	I/I/I	E2F1	2.0/3.8/3.6	E2F2	NC/MI/I	E2F2	10.3/5.0/4.5	E2F3	I/I/I	E2F3	Fibroblast growth factor receptor 3
FJL20430	N50962	31714	6.9/25.8/22.7	I/I/I	E2F1	7.4/10.3/12.4	E2F2	I/I/I	E2F2	9.6/14.5/14.8	E2F3	I/I/I	E2F3	Homo sapiens mRNA, complete cds, WD-repeat like sequence
FST	AA236796	9914	10.7/7.4/9.6	I/I/I	E2F1	15.9/14.7/11.4	E2F2	I/I/I	E2F2	15.8/18.2/14.0	E2F3	I/I/I	E2F3	Follistatin
GCH1	U19523	86724	8.2/5.9/9.9	I/I/I	E2F1	2.9/2.8/3.8	E2F2	I/MI/I	E2F2	18.1/4.4/10.0	E2F3	I/I/I	E2F3	GTP cyclohydrolase 1 (dopa-responsive dystonia)
HEY1	R61374	234434	38.3/21.9/32.1	I/I/I	E2F1	9.9/5.3/5.3	E2F2	I/I/I	E2F2	26.8/10.8/10.3	E2F3	I/I/I	E2F3	Hairy/enhancer-of-split related with YRPW motif 1
HUNK	N66354	109437	14.2/7.3/3.3	I/I/I	E2F1	3.2/2.6/1.2	E2F2	I/I/NC	E2F2	8.4/4.3/3.3	E2F3	I/I/I	E2F3	Horizontally upregulated neu tumor-associated kinase
INHBA	N27159	727	-24.1/-42.5/-12.1	D/D/D	E2F1	-4.7/-3.4/4.4	E2F2	NC/D/MD	E2F2	-24.8/27.1/-11.0	E2F3	D/D/D	E2F3	Inhibin, beta A (activin A, activin AB alpha polypeptide)
KIAA1055	AA447707	126084	2.8/6.4/6.8	I/I/I	E2F1	3.7/3.3/2.7	E2F2	I/I/I	E2F2	4.3/5.8/4.1	E2F3	I/I/I	E2F3	KIAA1055 protein
MAP3K5	AA057432	151988	35.2/42.1/29.5	I/I/I	E2F1	5.3/7.0/4.4	E2F2	I/I/I	E2F2	32.0/11.9/10.4	E2F3	I/I/I	E2F3	Mitogen-activated protein kinase kinase kinase 5
No symbol	F03200	7886	5.8/8.3/7.6	I/I/I	E2F1	3.6/3.6/3.3	E2F2	I/I/I	E2F2	6.8/5.7/4.9	E2F3	I/I/I	E2F3	Weakly similar to Pelle associated protein Pellino [<i>Drosophila</i>]
No symbol	N63887	7155	8.8/12.0/9.0	I/I/I	E2F1	7.7/4.4/5.3	E2F2	I/I/I	E2F2	21.9/7.5/6.9	E2F3	I/I/I	E2F3	Weakly similar to TYKI protein [<i>M.musculus</i>]
OSMR	U15932	238648	-3.2/-2.1/-2.4	D/D/D	E2F1	-1.8/-2.9/2.5	E2F2	D/D/D	E2F2	-1.9/2.1/-2.0	E2F3	D/MD/D	E2F3	Oncostatin M receptor
PAGE	X17094	59242	3.0/2.5/6.7	I/NC/I	E2F1	-1.7/-1.1/1.3	E2F2	NC/NC/NC	E2F2	1.7/1.1/3.0	E2F3	NC/NC/NC	E2F3	Paired basic amino acid cleaving enzyme (turin)
PAI1	J03764	82085	-7.9/-5.5/-2.4	D/D/D	E2F1	-6.6/-5.7/-1.2	E2F2	D/D/NC	E2F2	-3.9/2.3/-1.7	E2F3	D/D/NC	E2F3	Plasminogen activator inhibitor, type I
PRKAR2B	M31158	77439	2.3/3.5/8.5	I/I/I	E2F1	1.5/1.7/3	E2F2	I/MI/I	E2F2	4.5/2.8/6.0	E2F3	I/I/I	E2F3	Protein kinase, cAMP-dependent, regulatory, type II, beta
PTPNS1	T16175	156114	9.4/5.9/5.6	I/I/I	E2F1	3.4/3.2/1.3	E2F2	I/I/NC	E2F2	19.1/2.8/6.5	E2F3	I/NC/I	E2F3	Protein tyrosine phosphatase, non-receptor type substrate 1
RANBP1	AA424865	106346	3.0/4.5/4.6	I/I/I	E2F1	2.0/2.9/3.2	E2F2	I/I/I	E2F2	4.5/4.3/4.2	E2F3	I/I/I	E2F3	Novel centrosomal protein RanBPM
SOX9	Z46629	2316	-3.1/-6.5/-5.7	D/D/D	E2F1	-2.0/-1.2/-1.3	E2F2	D/NC/NC	E2F2	-1.1/-1.4/1.0	E2F3	NC/NC/NC	E2F3	SRY (sex-determining region Y)-box 9
SPHK1	AA232646	68061	13.3/13/14.4	I/I/I	E2F1	4.3/6.8/3.2	E2F2	I/MI/NC	E2F2	10.1/7.6/6.0	E2F3	I/I/I	E2F3	Sphingosine kinase 1
TACC1	AA481427	173159	5.3/6.7/8.3	I/I/I	E2F1	2.6/2.9/5.0	E2F2	NC/NC/I	E2F2	6.8/6.9/2.1	E2F3	I/I/NC	E2F3	Transforming, acidic coiled-coil containing protein 1
TEAD4	U63824	94865	3.9/3.6/6.6	I/I/I	E2F1	4.4/4.6/6.6	E2F2	I/I/I	E2F2	3.7/4.0/6.1	E2F3	I/I/I	E2F3	TEA domain family member 4 (transcription factor RTEF-1)
TGFA	X70340	170009	2.6/7.7/7.6	NC/I/I	E2F1	-1.0/-1.4/1.1	E2F2	NC/NC/NC	E2F2	2.2/1.5/1.7	E2F3	NC/NC/NC	E2F3	Transforming growth factor, alpha
TGFB2	N48082	169300	-9.6/-24.6/-9.8	D/D/D	E2F1	-2.1/-2.5/-3.5	E2F2	D/D/D	E2F2	-4.4/-8.1/-4.7	E2F3	D/D/D	E2F3	Transforming growth factor, beta2
YY1	Z14077	97496	3.1/3.7/3.8	I/I/I	E2F1	1.7/1.5/1.6	E2F2	NC/NC/NC	E2F2	2.5/3.2/3.1	E2F3	I/I/NC	E2F3	YY1 transcription factor

Symbol for each gene as provided by Unigene <http://www.ncbi.nlm.nih.gov/UniGene/index.html>. Accession number refers to sequence used to construct the microarray. Fold changes and difference calls are provided by the GeneChip software. (I) Induced, (MI) moderately induced, (NC) no change, (MD) moderately decreased, (D) decreased.




Figure 3. Verification of microarray data by Northern blotting. (A) Northern blot analysis to verify target gene regulation using 10 μ g of total RNA. Cells were harvested after 0 hr, 4 hr, and 8 hr of exposure to 300 nM 4-hydroxy tamoxifen (OHT). (B) Northern blot analysis to verify target gene regulation using 4 μ g of poly A⁺ RNA. Cells were harvested after 0 hr, 4 hr, and 8 hr of exposure to 300 nM OHT.

tein is just starting to be detectable (data not shown). Although we cannot formally rule out the possibility that these effects are independent of endogenous E2F ac-

tivity, the most likely explanation is that nonphosphorylatable pRB binds to and inactivates endogenous E2F activity.

Table 2. *E2F-induced expression changes for selected genes: Summary call and verification*

Symbol	Summary call ($P < 0.01$)			Northern blot		
	E2F1	E2F2	E2F3	E2F1	E2F2	E2F3
APAF1	I/3.3	NC/1.3	NC/0.5	I	ND	ND
ASH2L	I/9.7	NC/2.4	I/4.8	I	I	I
BCL3	D/-10.6	NC/-6.9	D/-10.0	D	D	D
BMP2	I/8.8	NC/3.0	I/3.4	I	I	I
BMPR1A	I/8.0	NC/2.0	I/2.7	I	I	I
CASP3	I/14.1	I/5.9	I/10.7	I	I	I
CASP7	NC/2.4	NC/1.4	NC/2.0	I	I	I
CAV1	D/-2.7	NC/-2.1	NC/-2.1	D	NC	D
CD58	I/4.8	I/3.0	I/3.9	I	I	I
CHES1	D/-8.5	NC/-4.7	D/-5.7	NC	NC	NC
CTGF	D/-2.8	NC/-1.3	NC/-0.4	D	D	D
DMRT1	NC/-0.6	I/3.8	NC/1.4	I	I	I
DYRK1A	I/4.3	NC/2.2	I/3.0	I	I	I
DYRK2	I/3.6	I/4.0	I/5.3	I	I	I
EGR1	I/4.8	NC/2.7	I/2.8	I	NC	I
EPLIN	D/-4.6	NC/-1.7	D/-3.0	D	NC	D
EZH2	I/5.1	I/5.0	I/5.2	I	I	I
F03200	I/7.2	I/3.5	I/5.8	I	I	I
FGFR2	I/3.8	NC/1.3	NC/1.7	I	I	I
FGFR3	I/7.1	I/3.1	I/6.6	I	I	I
FLJ20430	I/18.5	I/10.0	I/13.0	I	I	I
FST	I/9.2	I/14.0	I/16.0	I	I	I
GCH1	I/8.0	I/3.2	I/10.8	I	I	I
HEY1	I/30.8	I/6.8	I/16.0	I	I	I
HUNK	I/8.3	NC/2.3	I/5.3	I	I	I
INHBA	D/-26.2	D/-4.2	D/-21.0	D	D	D
KIAA1055	I/5.3	I/3.2	I/4.7	I	I	I
MAP3K5	I/35.6	I/5.6	I/18.1	I	I	I
N63887	I/9.9	I/5.7	I/8.7	I	I	I
OSMR	D/-2.6	NC/-2.4	NC/-2.0	D	NC	D
PACE	I/4.1	NC/1.0	NC/1.9	I	NC	I
PAI1	D/-5.3	D/-4.5	NC/-2.6	D	D	D
PRKAR2B	I/5.1	NC/2.1	I/4.4	I	I	I
PTPNS1	I/7.0	I/2.6	I/9.5	I	I	I
RANBPM	I/4.0	I/2.7	I/4.3	I	I	I
SOX9	D/-5.1	NC/-1.5	NC/-0.5	D	NC	NC
SPHK1	I/13.6	I/4.8	I/7.9	I	I	I
TACC1	I/6.8	NC/3.5	I/5.3	I	ND	ND
TEAD4	I/4.7	I/5.2	I/4.6	I	I	I
TGFA	I/6.0	NC/-0.4	NC/1.4	I	I	I
TGFB2	D/-14.7	D/-2.7	D/-5.7	D	D	D
YY1	I/3.5	NC/1.6	I/2.6	I	I	I

Symbols are given for each gene as in Table 1. Two genes do not have an assigned symbol, and their accession numbers are given. The average fold change in expression for a gene is shown. The summary call ($P < 0.01$) is calculated using the McNemar test as described in the text. The verification of all genes have been performed by Northern blotting, and calls are based on these results (Figs. 3,4; data not shown).

(ND) Not determined.




Figure 4. E2F-induced changes in transcript levels are neither confined to transformed cells nor caused by overexpression of E2F fusion proteins. (A) Northern blot analysis of E2F-regulated genes in human diploid WI-38 ERE2F1 cells. 2 μ g poly A RNA was loaded. (B) Northern blot analysis of RNA prepared from U2OS pRb Δ cdk cells. 15 μ g of total RNA was loaded. Hours indicate the time after removal of tetracyclin. Note the increase in the levels of the faster migrating mouse Rb Δ CDK with time. Blots were probed using a human probe, which likely is the reason that mouse Rb Δ cdk mRNA produced a much weaker signal than endogenous human RB1.

Groups of genes regulated by E2F: Target gene bias analysis


The E2F transcription factors are best known for their involvement in the regulation of cell cycle and apoptosis. A considerable number of genes involved in these processes were found to be modulated by ER-E2F induction (Fig. 5; Tables 1, 2, and 3). In the case of apoptosis, E2F can induce apoptosis independent of p53 (Müller and Helin 2000). Based on our results, transcriptional regulation of *APAF1*, *CASP3*, *CASP7*, *MAP3K5*, and *BCL3* genes provides a plausible mechanism whereby E2Fs affect this process. Among the cell cycle regulators, cyclins were the preferred E2F targets. There were 16 different known cyclin genes represented on the microarrays used in our studies, of which E2F1 and E2F3 regulate four (*CCNE1*, *CCNE2*, *CCND3*, *CCNG2*), and E2F2 regulates three (*CCNE1*, *CCNE2*, *CCND3*).

However, most of the E2F target genes identified in the screen cannot be categorized as regulators of cell cycle or apoptosis. To better understand the complicated expression patterns induced by E2F, we performed target gene bias (TGB) analysis (Fig. 5; see Materials and Methods). This analysis is based on the hypothesis that a disproportionately high fraction of genes involved in a particular process will be regulated by a transcription factor if that transcription factor actually regulates that process. For example, if that fraction is significantly higher for TGF β signaling genes, we hypothesize that E2F activity influences TGF β signaling. In mathematical terms, the level of significance (*P* value) is given by the binomial distribution (Fig. 5A). The negative logarithm (base 10) of the *P* value is used as a quantitative measure of signifi-

cance. A bias value of 2 corresponds to a *P* value of 0.01. TGB analysis is not restricted to genes that are involved in signaling pathways but can be applied to arbitrary groups of genes that share a particular property of interest (see below).

Figure 5B shows the results of TGB analysis for genes grouped by function and/or structural similarities as defined by GeneCards (see Materials and Methods). As a positive control for our analysis, we included previously reported E2F target genes. Two randomly selected sets of genes similar in size to the analyzed gene groups are included as negative controls. We observed significant bias ($-\log(P) > 2$) for transcription factors (in particular homeobox genes), genes of the TGF β signaling pathway, and receptor tyrosine kinases. As expected the E2Fs also produced a significant bias toward cell cycle and apoptosis genes.

We also sought to determine if a significant overlap existed between the expression patterns found in other published gene expression screens and the patterns observed by induction of E2F activity. Figure 5C shows that there is indeed a significant overlap between E2F-modulated genes and those induced after addition of serum to starved fibroblasts (Iyer et al. 1999) as well as genes regulated by receptor tyrosine kinases (RTKs; Fambrough et al. 1999). We also observed significant bias toward genes regulated by p53 (Zhao et al. 2000), c-Myc (Coller et al. 2000), activated Ras (Zuber et al. 2000), and replicative senescence (Shelton et al. 1999). However, genes found to be regulated by aging (Lee et al. 2000) and caloric restriction (Lee et al. 1999) showed biases equal to those from randomly selected subsets of genes. The gene expression patterns for E2F1, E2F2, and E2F3 were also



compared to each other (Fig. 5D). As expected, the overlaps are highly significant and they show the relatedness of these three factors as well as the sensitivity of TGB analysis to reveal similarities in gene expression patterns.

E2F-regulated genes are widely expressed

The known functions of the E2F transcription factors suggest that they play a fundamental role in a number of cellular processes. It is therefore expected that E2F target genes are expressed in a variety of tissues and cell types. We tested this hypothesis by determining the distribution of E2F target genes in different cDNA libraries and compared this distribution to the profile of all genes expressed at detectable levels in U2OS cells. As shown in Figure 5E, most genes represented on the microarrays in our studies were isolated from a small number of cDNA libraries (Fig. 5E; "19K exp") and this profile does not change significantly for genes that are expressed in U2OS cells (genes called "present" by GeneChip® software; Fig. 5E; "19K obs").

Therefore, U2OS cells express a random subset of the genes that are present on the microarrays. A completely different profile is observed for E2F target genes. The results for E2F3 are shown (Fig. 5F), but very similar profiles were obtained for E2F1 and E2F2. Genes that are found in only a few libraries are strongly underrepresented, whereas genes with a broader library representation (7–31) are overrepresented. Although it may be a simplistic line of reasoning, these results suggest that E2F target genes serve fundamental functions required in a large number of cell types and tissues.

Discussion

Previous experiments have suggested that, in addition to having a role in cell proliferation, E2F also plays a role in regulating apoptosis, differentiation, and development. However, to date most known E2F-regulated genes are characterized as functioning in cell cycle control and/or DNA replication, which suggests that we only know of a

subset of E2F-regulated genes. The major findings of our experiments are that increased E2F activity leads to significant changes in the overall gene expression profile of a given cell and that E2F modulates the expression of a large number of transcripts. Furthermore, E2F controls, in addition to those involved in cell cycle control and DNA replication, genes that are directly involved in cell-fate decisions such as apoptosis, differentiation, and early development (see below).

Our results show that the activation of the E2Fs induced a significant change in the expression of 1240 genes present on the 35K microarray chips, or ~7% of the total 19,000 genes that were found to be expressed in U2OS cells. Of these, 718 or 58% were induced and the expression of 522 were decreased following E2F activation. Of the 1240 genes, ~60% were unnamed expressed sequence tags (ESTs).

There are several measures for the reliability of our gene expression data. Our statistical analysis has shown that the observed changes are highly significant. Biological and technical variability was significantly reduced by the use of three independently derived U2OS cell lines expressing each of the ER-E2F fusion proteins. In several cases, we also observed reproducible expression changes of the same gene represented more than once on the microarrays. Moreover, Northern blot analysis was used as an independent test for the observed expression changes and the results were extremely consistent with the microarray data. Of 40 genes determined to be significantly induced or repressed by E2F1, Northern blot analysis confirmed 39 of these changes. For E2F2 and E2F3 the corresponding numbers were 21 of 21 and 29 of 30. For the verification by Northern blotting, we deliberately chose several genes that showed only two- to threefold change in expression levels, as well as genes that fell just outside a statistically significant change. Several of these genes were found by Northern blot analysis to be regulated by E2F, clearly showing that our statistical test is conservative and that 1240 significant changes is a low estimate for the number of E2F targets. In addition, the Northern blot results also showed that the candidate genes are rapidly induced by E2F in the absence of de novo protein synthesis, which suggests

Figure 5. Bias analysis. (A) Principle of target gene bias analysis (TGB). TGB determines the significance of a particular class of genes being regulated based on the fraction of these genes being regulated in the total set of detectable genes on the microarrays. Applied to subsets of genes on a microarray (see panel B), it asks whether the fraction of regulated genes in this subset of genes is equal to the overall fraction of regulated genes. Applied to other microarray experiments (see panel C,D), TGB is sensitive to the relatedness of gene expression patterns. In both cases, significant deviation from the expected value (bias) is hypothesized to imply biological relevance. (B) TGB analysis of functional gene groups. Previously described E2F target genes were included as a positive control. (C) TGB analysis of published screens. Note that TGB is not confined to microarray-based screens. Ras transformation targets were identified by subtractive suppression hybridization (Zuber et al. 2000). (D) Comparison of E2F1, E2F2, and E2F3 expression patterns by TGB. (E) U2OS cells express a random subset of the genes present on the microarray chips. The distribution of unigenes on the chip found in a certain number of libraries is shown. The expected profile (19K exp) is the result of multiplying the number of genes in each category by the fraction of genes expressed in U2OS cells (i.e., the ~19,000 genes that have been called present in either control or test chips/28,000 unigenes found on the chips). The observed profile (19K obs) represents the genes expressed in U2OS cells. (F) E2F target genes tend to be widely expressed. E2F3 was shown to regulate significantly the expression of 633 genes. The expected profile presents a random expression of these genes, and is found by multiplying the number of genes in each category of genes expressed in U2OS cells (19K obs in panel E) by the fraction of regulated genes (i.e., 633/19,000). The observed profile (633 obs) represents the distribution of the 633 genes regulated by E2F3.

Table 3. Functional classification of selected E2F regulated genes

Name	Acc.	E2F1	E2F2	E2F3	Function	Name	Acc.	E2F1	E2F2	E2F3	Function
Protein regulated E2F genes											
CDK2	U05254	9.0	13.3	25.8	Transcription factor, transcription.	Transcription factor					
CDK2F1	U05251	-2.7	-2.1	-2.1	Transcription factor, transcription.	TEA domain family member 4	A443408	4.9	2.9	4.0	Differentiation, inhibition of, JH1, growth's
CDK2A	A007314	2.9	1.6	1.6	Cell cycle, proliferation.	MYC	L83828	4.7	5.2	4.8	Regulator of muscle specific genes
CDK2B	A443656	3.7	3.6	4.6	Cell cycle	MYC2	U05011	9.9	3.1	5.2	Regulates cyclin expression and T cell induced apoptosis
CDK2E1	T54121	5.8	5.3	5.9	Cell cycle, proliferation	MYC3	A443656	4.9	3.8	3.2	Regulates cyclin expression and T cell induced apoptosis
CDK2E2	N50327	5.0	3.2	4.6	Transcription factor, proliferation.	MYC4	A443656	5.4	-0.5	3.9	MYC family, development and differentiation.
CDK2E3	A443656	2.2	1.8	2.6	Transcription factor, proliferation.	MYC5	U17411	4.7	3.0	3.6	MYC family, development and differentiation.
CDK2E4	X51256	6.6	5.7	6.3	Cell cycle.	MYC6	U04291	3.3	3.1	3.4	MYC family, development and differentiation.
CDK2E5	A443656	6.6	4.8	6.5	Cell cycle, proliferation.	MYC7	U04291	4.8	2.9	3.6	MYC family, development and differentiation.
CDK2E6	U05252	-5.3	-6.6	-3.0	Transcription factor.	MYC8	U04291	4.8	3.0	6.6	MYC family, development and differentiation.
CDK2E7	X54918	8.3	4.1	7.4	Transcription factor.	MYC9	N21701	3.8	3.2	3.2	MYC family, development and differentiation.
CDK2E8	U05253	-3.7	-2.1	-2.3	Cell cycle, proliferation.	MYC10	X51341	3.8	2.5	1.5	MYC family, development and differentiation.
CDK2E9	U05254	-3.7	-2.1	-2.3	Cell cycle, proliferation.	MYC11	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDK2E10	A443656	13.6	8.8	14.4	Transcription factor, proliferation.	MYC12	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Cell cycle and DNA replication											
CDC2	A443656	11.5	8.7	12.3	Cell cycle, proliferation.	MYC13	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2A	U05251	-5.5	-6.6	-3.0	Transcription factor.	MYC14	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2B	F08876	4.6	-2.3	-3.1	Transcription factor.	MYC15	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2C	A443656	8.1	4.2	3.7	Cell cycle, proliferation.	MYC16	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2D	X40769	-2.7	-1.1	-2.0	Cell cycle, proliferation.	MYC17	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2E	N59175	-4.5	-2.0	-2.8	Cell cycle, proliferation.	MYC18	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2F	U05252	4.1	-1.4	1.6	Cell cycle.	MYC19	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2G	U11071	4.0	2.7	4.3	Transcription factor.	MYC20	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
CDC2H	M62486	2.3	1.6	3.7	DNA replication	MYC21	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Apoptosis											
Bcl-2	A443656	3.3	1.5	2.5	Cell cycle, proliferation.	MYC22	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Bcl-2L1	M14745	4.7	3.7	5.1	Protein, apoptosis.	MYC23	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Caspase-1	U11071	14.1	5.9	10.7	Transcription factor.	MYC24	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Caspase-4	A443656	-4.5	-2.9	-3.8	Transcription factor.	MYC25	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Caspase-9	X38172	1.5	-2.1	-2.1	Transcription factor.	MYC26	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Caspase-10	X11117	-2.2	-1.5	-2.6	Transcription factor.	MYC27	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Caspase-11	A443656	35.6	5.6	10.1	Transcription factor.	MYC28	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Caspase-12	A443656	-3.9	-8.0	-8.9	Transcription factor.	MYC29	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Caspase-13	A443656	2.5	4.4	1.8	Transcription factor.	MYC30	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Transcription											
Hemoglobin genes											
Actb	A443656	9.7	3.2	3.8	Transcription factor.	MYC31	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg1	R42307	3.8	5.9	4.8	Transcription factor.	MYC32	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg2	U05251	2.8	3.8	3.3	Transcription factor.	MYC33	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg3	U05251	5.1	5.0	5.3	Transcription factor.	MYC34	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg4	R81374	30.3	6.8	18.0	Transcription factor.	MYC35	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg5	A443656	6.0	0.7	1.4	Transcription factor.	MYC36	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg6	A443656	6.8	1.9	0.7	Transcription factor.	MYC37	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg7	A443656	2.1	-1.8	-0.5	Transcription factor.	MYC38	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg8	A443656	2.0	0.0	2.3	Transcription factor.	MYC39	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg9	A443656	2.7	0.0	2.3	Transcription factor.	MYC40	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg10	A443656	2.1	1.7	3.5	Transcription factor.	MYC41	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg11	A443656	2.1	1.7	3.5	Transcription factor.	MYC42	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg12	A443656	2.1	1.7	3.5	Transcription factor.	MYC43	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg13	A443656	2.1	1.7	3.5	Transcription factor.	MYC44	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg14	A443656	2.1	1.7	3.5	Transcription factor.	MYC45	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg15	A443656	2.1	1.7	3.5	Transcription factor.	MYC46	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg16	A443656	2.1	1.7	3.5	Transcription factor.	MYC47	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg17	A443656	2.1	1.7	3.5	Transcription factor.	MYC48	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg18	A443656	2.1	1.7	3.5	Transcription factor.	MYC49	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg19	A443656	2.1	1.7	3.5	Transcription factor.	MYC50	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg20	A443656	2.1	1.7	3.5	Transcription factor.	MYC51	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg21	A443656	2.1	1.7	3.5	Transcription factor.	MYC52	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg22	A443656	2.1	1.7	3.5	Transcription factor.	MYC53	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg23	A443656	2.1	1.7	3.5	Transcription factor.	MYC54	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg24	A443656	2.1	1.7	3.5	Transcription factor.	MYC55	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg25	A443656	2.1	1.7	3.5	Transcription factor.	MYC56	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg26	A443656	2.1	1.7	3.5	Transcription factor.	MYC57	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg27	A443656	2.1	1.7	3.5	Transcription factor.	MYC58	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg28	A443656	2.1	1.7	3.5	Transcription factor.	MYC59	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg29	A443656	2.1	1.7	3.5	Transcription factor.	MYC60	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg30	A443656	2.1	1.7	3.5	Transcription factor.	MYC61	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg31	A443656	2.1	1.7	3.5	Transcription factor.	MYC62	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg32	A443656	2.1	1.7	3.5	Transcription factor.	MYC63	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg33	A443656	2.1	1.7	3.5	Transcription factor.	MYC64	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg34	A443656	2.1	1.7	3.5	Transcription factor.	MYC65	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg35	A443656	2.1	1.7	3.5	Transcription factor.	MYC66	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg36	A443656	2.1	1.7	3.5	Transcription factor.	MYC67	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg37	A443656	2.1	1.7	3.5	Transcription factor.	MYC68	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg38	A443656	2.1	1.7	3.5	Transcription factor.	MYC69	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg39	A443656	2.1	1.7	3.5	Transcription factor.	MYC70	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg40	A443656	2.1	1.7	3.5	Transcription factor.	MYC71	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg41	A443656	2.1	1.7	3.5	Transcription factor.	MYC72	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg42	A443656	2.1	1.7	3.5	Transcription factor.	MYC73	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg43	A443656	2.1	1.7	3.5	Transcription factor.	MYC74	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg44	A443656	2.1	1.7	3.5	Transcription factor.	MYC75	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg45	A443656	2.1	1.7	3.5	Transcription factor.	MYC76	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg46	A443656	2.1	1.7	3.5	Transcription factor.	MYC77	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg47	A443656	2.1	1.7	3.5	Transcription factor.	MYC78	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg48	A443656	2.1	1.7	3.5	Transcription factor.	MYC79	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg49	A443656	2.1	1.7	3.5	Transcription factor.	MYC80	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg50	A443656	2.1	1.7	3.5	Transcription factor.	MYC81	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg51	A443656	2.1	1.7	3.5	Transcription factor.	MYC82	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg52	A443656	2.1	1.7	3.5	Transcription factor.	MYC83	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg53	A443656	2.1	1.7	3.5	Transcription factor.	MYC84	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg54	A443656	2.1	1.7	3.5	Transcription factor.	MYC85	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg55	A443656	2.1	1.7	3.5	Transcription factor.	MYC86	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg56	A443656	2.1	1.7	3.5	Transcription factor.	MYC87	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg57	A443656	2.1	1.7	3.5	Transcription factor.	MYC88	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg58	A443656	2.1	1.7	3.5	Transcription factor.	MYC89	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg59	A443656	2.1	1.7	3.5	Transcription factor.	MYC90	A443656	4.9	3.8	2.5	MYC family, development and differentiation.
Actg60	A443656	2.1	1.7	3.5	Transcription factor.	MYC91	A443656	4.9	3.8		

that the majority of the induced genes are regulated directly by E2F. In contrast, in all cases tested, the repression of a gene by the E2Fs required *de novo* protein synthesis. This result suggests that the E2Fs themselves do not function as transcriptional repressors, but may induce a repressor(s) of transcription.

In addition to verifying the expression changes observed on the microarrays by Northern blot analysis, we performed two other sets of experiments to determine the relevance of our data.

First, we have shown that increased E2F activity in primary human diploid cells leads to a similar change in expression of the tested genes as observed in U2OS cells. This shows that deregulation of E2F activity can lead to significant changes in gene expression in nontransformed cells in addition to tumor cells. This point is supported by our database analysis, which shows that the E2F-regulated genes are widely expressed and not restricted to specific tissues or certain tumor cell lines (see Fig. 5E,F). Second, we have shown that ectopic expression of a dominant form of pRB leads to down-regulation of genes that are induced by E2F and up-regulation of genes that are repressed by E2F activity. Importantly, we observed a change in expression of the E2F-regulated genes before the cells were arrested in G1 by pRB expression, which suggests that the changes in gene expression were not a consequence of pRB-induced pleiotropic cell cycle regulation but, rather, they resulted from pRB's ability to regulate E2F transcriptional activity. Because pRB is a repressor of E2F activity, these results suggest that endogenous levels of E2F regulate the expression of the tested genes.

Difference in target gene regulation by E2F family members

An open question is whether the E2F transcription factors regulate distinct genes or if they regulate a common set of genes. In the latter case, the differences in the biological function of the E2Fs could be explained by differences in expression levels and/or by tissue-specific expression patterns of the E2Fs. We addressed this question by analyzing the expression changes induced by three members of one of the E2F subgroups. Indeed, we observed that E2F1 activity led to significant changes in more genes than E2F2 and E2F3, which indicates that E2F1 was capable of specific regulation of certain genes. However, despite the fact that E2F1 was not overexpressed to the same extent as E2F2 and E2F3, our results show that the induction of E2F1 in the U2OS cells led to a stronger transactivation of a sensitive E2F reporter construct than did E2F2 and E2F3. Therefore, our results could be explained by a stronger transactivation potential of E2F1 in these cells. However, we also identified genes whose expression was selectively changed as a consequence of E2F2 or E2F3 activation, and which were not significantly modulated by either of the other E2Fs, thereby suggesting that some target-gene specificity indeed exists. However, we have not been able to confirm strict target-gene specificity by Northern blot analysis

and in the cases in which we observed significant induction by one of the E2Fs, we also detected a lower but consistent regulation of that gene by the other E2Fs. These results suggest that the E2Fs may not have strict target-gene specificity, but they may convey specificity in how efficiently they can activate a given gene (see e.g., DMRT1; Fig. 3B). Our findings are consistent with a recent report that showed that the different E2Fs bind to the promoter region in the same gene *in vivo*; however, they do so at distinct times during the cell cycle (Takahashi et al. 2000). Taken together, it is therefore likely that all E2Fs are candidates to regulate the expression of the E2F-induced genes identified in our analysis.

Comparison to published screens and TGB analysis

Based on the results of verification by Northern blots, we estimate that 2%, or ~20–30 genes out of 1240 in the list of regulated genes will be false positives. It is, therefore, difficult to draw conclusions based on changes observed for a single gene until they are verified by Northern blotting. It is, however, unlikely that a group of genes with a similar function that are characterized by our analysis as target genes are all false positives. We therefore sought to interpret the expression patterns induced by E2Fs based on changes of groups of genes. To this end, we analyzed the gene expression patterns for biases toward functional groups of genes and toward previously published expression patterns. We call this target gene bias (TGB) analysis.

TGB analysis of functional gene groups is problematic in the sense that the quality of results depends crucially on the quality of the functional classification that is applied. To minimize the probability of misleading conclusions, we used two independently assembled classifications. One was prepared by us based on information found in GeneCards and GenBank records and the other was published by Shelton et al. (1999). Although the numerical outcome was different, as expected, the main conclusions and statistical significance were surprisingly stable. Our analysis revealed significant bias toward genes that are involved in the regulation of cell cycle and apoptosis. The role of E2F in these processes is well documented and our data, which reveal several unknown cell cycle and apoptosis genes, provide some new vital insights as to the mechanisms of this regulation. Furthermore, significant bias toward gene groups with functions in processes that are known to be regulated by E2F validates our approach. Surprisingly, we observed highly significant bias toward homeobox and TGF β signaling genes. These genes are key regulators of development and differentiation. These findings suggest that E2Fs may play a role in coordinating fundamental biological processes such as proliferation, apoptosis, differentiation, and development.

We also compared the E2F-induced gene expression patterns to published expression patterns observed in discrete settings. This was done using TGB analysis by treating the previously published sets of regulated genes as individual functional gene groups. As compared to

functional gene groups, sets of regulated genes are well defined, which makes the analysis less sensitive to subjective judgements. We observed highly significant overlap between genes regulated by E2F and those regulated by serum. Serum acts largely by activation of tyrosine kinases and, interestingly, both receptor tyrosine kinases and receptor tyrosine kinase target genes also produced significant biases. Furthermore, tyrosine kinases are known to activate Ras and Ras transformation target genes also show strong TGB. One interpretation of this analysis is that E2F is a key mediator of the effects of serum, RTK, and Ras activation.

This interpretation has been supported previously by experiments showing that Ras is linked to cell cycle progression via the pRB/E2F pathway (Peeper et al. 1997). The spectrum of E2F target genes expressed at a certain point in time is therefore influenced by signals communicated by surrounding cells and the cellular environment and this may be the key to understanding the role of the E2Fs in integrating cell cycle, apoptosis, differentiation, and development. Because these processes are relevant to all cells, it would be expected that E2F target genes are expressed in a wide range of cell types. Our analysis of cDNA library distribution of E2F target genes suggests that this is indeed the case. The observed biases towards c-myc, p53, and replicative senescence genes may reflect the involvement of the E2Fs in these processes in ways that are not yet understood, but the investigation of commonly regulated target genes could provide some insight into this.

We suggest that TGB analysis can be useful in the analysis of complex gene expression patterns. As compared to hierarchical cluster analysis, TGB analysis can take advantage of prior knowledge about genes and, by doing so, makes the results easier to interpret. If applied to functional gene groups, the results depend on our current understanding of gene function, and as long as this understanding is incomplete, the bias analysis will be incomplete. If applied to published expression patterns, TGB analysis is conceptually similar to sequence similarity searches. As is the case also for sequence similarities, such similarities give rise to hypotheses that have to be tested by further experimentation. A particularly appealing application of this concept is the possibility to analyze signaling pathways as shown here in the case of the serum → RTK → Ras → E2F connection. It is tempting to speculate that it may be possible to use expression patterns observed in tumors or in virus-infected cells to identify the signaling pathways that are disturbed. This knowledge would be of tremendous value for the identification of drug targets whose manipulation results in selective cytotoxicity.

Compared to hierarchical clustering analysis (Eisen et al. 1998), TGB analysis includes supplementary information. This is of significant help in the process of interpreting observed gene expression patterns. In TGB analysis, clusters of genes (functional gene groups) are assembled beforehand (based on shared gene properties of interest) and then analyzed for statistical significance. The clusters in TGB analysis are therefore homoge-

neous, have clearly defined borders, and their significance is evaluated numerically rather than visually. On the other hand, TGB analysis is only as good as the gene groups that are available for analysis.

Hierarchical clustering and TGB analysis may therefore be used as complementary approaches to help in the interpretation of complex gene expression patterns.

Classification of E2F target genes

One of the most intriguing aspects of gene expression profiling is that the results can correlate gene expression patterns with specific phenotypes. Therefore, the profiling can provide several predictions and hypotheses that subsequently can be tested. Here, we identified a number of genes whose function could provide a molecular explanation as to how the proteins in the pRB/E2F-pathway control normal growth. We have listed 100 of these genes in Table 3. Although it will be too extensive to go into the specific function of each gene, we would like to highlight possible connections between the expression of some of these genes and the physiological function of the pRB/E2F pathway.

Cell cycle control and DNA replication As an internal control for our experiments, we performed a database search for genes that have been reported to be regulated by E2F or have an E2F DNA-binding site (or binding sites) in their promoters. A number of these genes were expressed at detectable levels in U2OS cells, and the expression of several of them was changed after E2F activation (e.g., Cyclin E1, Cyclin D3, and c-Myb; Table 3). Strikingly, we also detected down-regulation of the plasminogen activator inhibitor, type 1, *PAI-1*, the only gene so far reported to be repressed by E2F (Koziczak et al. 2000). In our screen, we did not detect regulation of several genes previously reported to be E2F targets. Most notably, we did not observe significant regulation of the MCM genes, *CDC6* and *B-Myb*. The reason for this lack of regulation is at present unclear, but may be because of the fact that the expression of these genes is strongly regulated during the transition from quiescence to growth and not in each cycle of exponentially growing cells. Because we performed our analysis using asynchronous populations of exponentially growing U2OS cells, the increased E2F activity may not have further affected change in the mRNA levels of these genes. An alternative, but less likely, explanation for the lack of detectable regulation could be the quality of the chosen oligonucleotide probes for these genes on the arrays.

Among the novel E2F target genes that are involved in cell cycle regulation, the expression of Cyclin E2 (*CCNE2*) was found to be strongly induced by the all three E2Fs.

Cyclin E2 expression is cell cycle regulated, and it is expressed with similar kinetics as Cyclin E1 (Lauper et al. 1998; Zariwala et al. 1998; Gudas et al. 1999). Our data indicate that, as for Cyclin E1, the cell cycle regulation of Cyclin E2 expression is controlled by the pRB/E2F pathway.

Centrosome duplication Recently, it was shown that E2F and Cyclin A/CDK2 activity are required for centrosome duplication (Meraldi et al. 1999). We identified RanBPM as a direct target for the E2F transcription factors (Figure 3; Table 3). RanBPM colocalizes with γ -tubulin and is required for microtubule nucleation (Nakamura et al. 1998). Interestingly, overexpression of this protein can induce ectopic microtubule nucleation, and these data could indicate that pRB restricts centrosome duplication through the transcriptional regulation of RanBPM. Furthermore, because the pRB/E2F pathway is abrogated in the majority of human tumors, it may provide an explanation for the increased number of centrosomes in many tumors.

Apoptosis Loss of pRB has been shown to lead to both p53-dependent and p53-independent apoptosis (Lipinski and Jacks 1999). It is widely believed that the apoptosis is a result of higher levels of E2F activity because ectopic expression of the E2Fs also can induce both p53-dependent and p53-independent apoptosis (Dyson 1998; Helin 1998). Although some E2F-regulated candidate genes have been suggested, none of these explain the strong apoptotic effect of deregulated E2F activity. Therefore, it is intriguing to find that the E2Fs induce the expression of several key regulators of apoptosis, such as the cytochrome-C-binding protein APAF1, and the crucial effector caspases, caspase 3 and caspase 7. Moreover, the MAP kinase kinase, kinase 5, also known as apoptosis-signal-regulating kinase 1 (ASK1), which is required for TNF α -induced apoptosis (Ichijo et al. 1997), was strongly induced. The E2Fs induced these four genes in the absence of de novo protein synthesis (Fig. 3), which shows that they are direct targets for the E2F transcription factors. Moreover, we observed a strong down-regulation of Bcl-3, a known coactivator of NF κ B, which is, therefore, an inhibitor of apoptosis (Rebollo et al. 2000). The alteration of expression in any of these five genes or a combination thereof could explain the strong apoptotic effects of E2F activity. However, as is often the case for signal transduction pathways, the E2Fs stimulate both positive and negative signals of proliferation. The decision of whether to proliferate, differentiate, or die becomes a balance between the intensity of the various signals. Although the E2Fs induce cell proliferation, they also induce p18^{INK4C}, an inhibitor of the cell cycle, and although the E2Fs induce apoptosis, they also induce growth factors and antiapoptotic genes, such as *BCL2*. However, because E2F expression induces apoptosis, the induction of *BCL2* and growth factors is apparently insufficient for counteracting the strong expression of the apoptosis inducing proteins.

Development and differentiation E2F activity is essential for early development in *Drosophila* and *Xenopus* (Duronio et al. 1995; Suzuki and Hemmati-Brivanlou 2000), and recent results have also suggested that E2F activity is required for early development in mice (L. Yamasaki, pers. comm.). More specifically, *Xenopus* and *Drosophila* E2F activity was shown to be required for

axis determination in early development. It has been suggested that the E2Fs regulate axis determination through homeobox-containing proteins in *Xenopus* (Suzuki and Hemmati-Brivanlou 2000) and an EGF-receptor ligand, Gurken in *Drosophila* (Myster et al. 2000).

Our results confirm and extend these previous data by showing that the E2Fs regulate the expression of several proteins that are involved in early development, including homeobox proteins, transcription factors involved in cell fate decisions, a number of proteins that determine homeotic gene transcription, and signaling pathways such as the TGF β and Wnt pathways that are essential for early development. As an example of the relevance of these findings, it has been reported that position-effect variegation (PEV) in *Drosophila* depends on E2F activity (Seum et al. 1996). Loss of E2F activity enhanced PEV, whereas overexpression of E2F activity suppressed PEV in *Drosophila* (Seum et al. 1996). These data suggested that the E2Fs themselves have an epigenetic effect by regulating chromatin structure or, more likely, that the E2Fs control PEV by regulating genes of the Polycomb group (PcG) family. In our screen, we have identified several PcG genes, like Enhancer of Zeste 2 (*EZH2*), Embryonic Ectoderm Development protein (*EED*) and Homolog of Polyhomeotic (*EDR2/HPH2*; see Table 3). The E2F-induced expression of these genes may provide an explanation for the role of E2F in the regulation of PEV and, more importantly, in development.

The pRB/E2F pathway is known to be central in the regulation of various types of cellular differentiation (see introduction; Lipinski and Jacks 1999). For example, pRB is required for erythroid, neuronal, eye, muscle, and adipocyte differentiation (Chen et al. 1996; Lipinski and Jacks 1999). Both p107 and p130 are required for normal endochondrial bone development (Cobrinik et al. 1996). In addition, E2F4 is known to contribute to hematopoietic lineage and to craniofacial development (Humbert et al. 2000a; Rempel et al. 2000), whereas loss of E2f5 leads to overproduction of cerebrospinal fluid and to hydrocephalus (Lindeman et al. 1998). Although these phenotypes are complex, our gene expression analysis may help to elucidate the role (or roles) played by the E2Fs in pRB/p107/p130 regulated differentiation. First, we found a number of transcription factors that are involved in cell fate decisions, such as Hairy/enhancer of split related (*HEY1*), Paired-like homeodomain (*PTX1*), *ID4*, MAF family members, and *Sox9*. These transcription factors have been associated with neurogenesis (*HEY1*; Leimeister et al. 1999), morphogenesis, hindlimb, and craniofacial development (*PTX1*; Crawford et al. 1997; Szeto et al. 1999), block of differentiation of various tissues (*ID4*; Kondo and Raff 2000), the regulation of early differentiation (MAF family; Blank and Andrews 1997), and cartilage formation (*Sox9*; Bi et al. 1999). The expression of transcription factors that are involved in cell fate decisions is exquisitely controlled, and in several cases it has been described that the over- or misexpression of such transcription factors leads to malformations and/or transformation. Second, E2F activation led to a dramatic change in the expression of genes in the TGF β pathway.

TGF β family members, which include TGF β s, activins, and bone morphogenetic proteins (BMPs), are secreted molecules that regulate a plethora of cellular responses, such as proliferation, differentiation, migration, and apoptosis (for review, see Massague and Chen 2000). E2F activation led to strong suppression of Inhibin β A and TGF β 2, and to a strong induction of Follistatin expression. Inhibin β A participates in the regulation of gametogenesis and craniofacial development, and mice that are homozygously null for Inhibin β A die within 24 h of birth (Matzuk et al. 1995).

Interestingly, Inhibin β A was also isolated as erythroid differentiation factor (Murata et al. 1988) and the gene has been shown to be an important regulator of photoreceptor differentiation in the developing retina (Davis et al. 2000). Together with the fact that Follistatin is an inhibitor of Activin A (the homodimer of inhibin β A that binds Activin receptors), our data suggest that the simultaneous induction of Follistatin and repression of inhibin β A could be involved in specific differentiation defects observed in the eye and in the erythrocytes of the Rb^{-/-} mice. For example, Rb^{-/-} erythroid cells develop normally in high percentage Rb^{-/-} chimeric mice (Maandag et al. 1994; Williams et al. 1994), which suggests that the Rb wild-type cells rescued the Rb^{-/-} cells, perhaps by secreting paracrine or endocrine erythroid differentiation factor (or factors) such as inhibin β A or Follistatin.

The last phenotype that we wish to discuss is the requirement for a normal pRB/E2F pathway in chondrocyte differentiation. We identified several genes that are involved in chondrocyte differentiation, such as the TGF β pathways genes, FGF receptor 2 and FGF receptor 3, Cartilage linking protein 1, and Connective Tissue Growth factor (Fig. 3; Table 3). Discrete changes in the expression of several of these genes as a result of deregulated E2F activity could lead to both the defects in chondrocyte differentiation observed in the p107/p130 double-knockout mice, and the craniofacial abnormalities observed in the p107/p130 and the E2f4 knockout mice. Moreover, it is striking that activating mutations of FGF receptor 3 causes achondroplasia, which is the most common genetic form of dwarfism in humans (Burke et al. 1998), considering the fact that p107/p130 knockout mice have short bones.

In summary, the experiments presented in this article have provided us with a list of genes whose expression levels are significantly altered after E2F activation. The E2F-regulated genes code for proteins whose activity control cell cycle progression, proliferation, apoptosis, differentiation, and development. The identification of these genes allows us to test a number of new hypotheses as to why the pRB/E2F pathway is so crucial for normal development and proliferation. The list of E2F-regulated genes could be important for diagnostic purposes as, for example, in determining the efficacy of targeting a deregulated pRB/E2F pathway by therapeutic drugs. Finally, 60% of the identified genes are without any known function (ESTs). Because the expression of these ESTs is regulated by the E2Fs, several of these may have important functions in cellular homeostasis.

Therefore, future experiments will pay special attention to identifying a possible function of some of these ESTs.

Materials and methods

Cell culture

U2OS cell lines that express HAER-E2F1, HAER-E2F2, or HAER-E2F3 were generated by infection of the human osteosarcoma cell line U2OS (which expresses functional pRB and p53, but does not express p16INK4A or p14ARF) with retroviruses that express the three fusion proteins (Vigo et al. 1999). Cells were selected in 2.5 μ g/mL puromycin. Individual clones were isolated after 3 weeks of selection and tested for the expression of the ER-fusion proteins by Western blotting. The cells were grown in DMEM supplemented with 10% fetal calf serum. Puromycin resistant clones were continuously grown in the presence of 2.5 μ g/mL puromycin.

Pools of early passage WI-38 HAER-E2F1 cells were generated by infection with the retroviral vector pBabePuro HAER-E2F1 (Vigo et al. 1999), and selection in 1.5 μ g/mL puromycin. Cells were grown in DMEM supplemented with 10% FCS. U2OS cells expressing a pRB mutant (pRB Δ CDK) in a tetracycline-dependent manner has been described previously (Lukas et al. 1999).

Western blotting

Western blots were performed using the mouse monoclonal antibodies KH95, TFE22, and TFE31 that specifically recognize E2F1, E2F2, and E2F3, respectively (Helin et al. 1993; K. Helin, unpubl.).

Immunostaining

Cells, treated with or without 300 nM 4-hydroxytamoxifen (OHT) for 6 hr, were stained for the expression of HA-tagged ER-E2F proteins using the 12CA5 antibody as previously described (Müller et al. 1997).

Transactivation assays

Transactivation assays were performed as described previously (Müller et al. 1997). Briefly, cells were transfected with 2 μ g pGL3TATAbasic-6x E2F luciferase reporter, 2 μ g pCMV- β Gal reporter, and 2 μ g pCMVneoBam carrier. To perform assays, lysates were prepared from cells exposed to 300 nM OHT for 9 hr and from control cells. Normalized luciferase activity was determined as previously described (Müller et al. 1997).

Semiquantitative RT-PCR

Reverse transcription PCR (RT-PCR) was performed essentially as described (Vigo et al. 1999), except that total RNA was prepared using the RNeasy extraction kit (Qiagen).

High density oligonucleotide microarrays

Total RNA was prepared from test and control cells using the RNeasy kit (Qiagen).

Targets for hybridization to the microarrays were prepared as described (Lockhart et al. 1996; Fambrough et al. 1999), except that hybridization was performed in 1 \times MES buffer (0.1 M 2-[N-Morpholino] ethanesulfonic acid [MES] pH6.7, 1 M NaCl, 0.01% Triton X-100) and chips were washed in 0.1 \times MES buffer (0.1 M 2-[N-Morpholino] ethanesulfonic acid [MES] pH6.7, 0.1

M NaCl, 0.01% Triton X-100). Target concentration was 30 μg fragmented cRNA in 200 μL hybridization solution. Images were scanned at 3- μm resolution using a Hewlett Packard GeneArray Scanner. The images were analyzed using Affymetrix's GeneChip® software.

Statistical analysis

The McNemar test was used to determine data quality and cutoff (Abell et al. 1999). A special purpose software was developed and is available on request or at <http://marray.lar.ieu.it>. The McNemar test is often used in clinical trials to assess the efficacy of drug treatment versus placebo controls. The test compares lists of yes/no values. A P value >0.01 indicates lack of efficacy (the lists are equivalent), whereas P values <0.01 suggest the presence of a therapeutic effect. Applied to our data, we used the results from pairwise chip comparisons to obtain these lists where yes means "gene regulated" and no means "gene not regulated". A gene was called regulated when the difference call provided by the GeneChip® software was MI or I for up-regulated genes, or D or MD for down-regulated genes, and the fold change was equal to or beyond a variable cutoff value.

Chip comparisons between two test chips or between two control chips were used to determine the noise level of the experiments (called noise lists or control lists here), whereas chip comparisons between a test and a control chip were used to determine the signal (called signal lists or data lists).

We estimated three parameters:

The equivalence of samples and chip performances (noise list versus noise list, $P > 0.01$).

The presence of differences in transcript levels (noise list versus signal list, $P < 0.01$).

The reproducibility of measurement of such differences (signal list versus signal list, $P > 0.01$. Note that this test does not depend on reproducibility of transcriptional regulation for the same gene in independent experiments.) Although all signal-to-noise comparisons were highly significant for the presence of signal (see below), they were not all equivalent. The following ranking was observed:

$E2F1_1 = E2F1_3 > E2F1_2$

$E2F2_1 = E2F2_2 > E2F2_3$

$E2F3_1 > E2F3_2 = E2F3_3$

Next, we asked if the detected signal reflected reproducible changes in gene expression.

Signal lists that resulted from single-chip comparisons were combined into three duplicate and one triplicate list using the logical AND operator (meaning that a gene was called regulated only when it was found to be regulated in each of the composing sublists, see above). Noise lists and randomized signal lists combined the same way were used as controls.

Randomization of signal lists was carried out using a pseudo-random number generator that reassigned a new position to each gene in the list before combining them. The numbers of genes that were found to be regulated in each of these lists as a function of fold-change cutoff is shown in Figure 2. For example, in Figure 2A it is evident that there are significantly ($P \sim 1.0E-150$, $E2F1_1$ at cutoff 3) more regulated genes in the data lists as compared with the U2OS control lists. The randomized data lists contain the same number of regulated genes as the data lists. In Figure 2B, the randomized lists contain significantly ($P = 1.6E-103$ at cutoff 3 for $E2F1$) fewer regulated genes, which indicates reproducible measurement of regulated transcript levels for a large proportion of genes. Even fewer genes are found to be regulated because of noise in the control lists. The fact that the randomized and nonrandomized U2OS derived duplicate lists contain similar numbers of regulated genes

($P = 0.43$ at cutoff 1) illustrate the stochastic nature of the noise lists.

Cloning of probes for Northern blots

Primers were designed using Oligo 4.1 (Primer Analysis software). Total RNA was prepared from cycling U2OS cells using the RNeasy kit (Qiagen). Double-stranded (ds) cDNA was synthesized using the ds cDNA synthesis system from Life Technologies. We used 10 ng of ds cDNA in a standard 50 μL PCR reaction using Taq polymerase. PCR fragments were cloned into pCR2.1 using a TA cloning kit (Invitrogen). The identity of the PCR fragments was verified by sequencing.

Northern blots

Nearly confluent cultures or U2OS clones were trypsinized and plated at 5×10^6 cells per 15-cm plate on the day before induction. The ER-E2F fusion proteins were activated by adding OHT to a final concentration of 300 nM, and samples were harvested at the indicated times. Cycloheximide was added where indicated to a final concentration of 10 $\mu\text{g}/\text{mL}$. RNA was isolated using either the RNeasy kit (Qiagen) for analysis of total RNA or by CsCl ultracentrifugation method as described (Ausubel et al. 1988) for analysis of poly A RNA. Poly A RNA was isolated with the Oligotex reagents from Qiagen using a batch protocol as described by the manufacturer.

Either 10 μg of total RNA or 4 μg of poly A RNA were separated on 1.25% formaldehyde-agarose gels, transferred to Hybond N⁺ membrane (Amersham) and probed as described (Ausubel et al. 1988).

TGB analysis

TGB analysis was performed as described in the results section using a special purpose software available on request or at <http://marray.lar.ieu.it>.

Acknowledgments

We gratefully acknowledge Elena Colli for help in generating the U2OS cell lines and Simona Ronzoni for FACS analysis. We thank Jiri Bartek for U2OS cells expressing pRB Δ CDK. We thank Ali R. Fattaey and Nick H. Heintz for discussions and critical reading of the manuscript and members of the Department of Experimental Oncology for discussions. This work was supported by grants from the TMR programme of the European Community, Ministero della Sanita, Associazione Italiana per la Ricerca sul Cancro (AIRC) and Fondazione Italiana per la Ricerca sul Cancro (FIRC).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Abell, M.L., Braselton, J.P., and Rafter, J.A. 1999. *Statistics with mathematical*. Academic Press, .
- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A., and Struhl, K. 1988. *Current protocols in molecular biology*. Greene Publishing Associates & Wiley-Interscience, New York.
- Bi, W., Deng, J.M., Zhang, Z., Behringer, R.R., and de Crombrughe, B. 1999. Sox9 is required for cartilage formation.

- Nat. Genet.* **22**: 85–89.
- Blank, V. and Andrews, N.C. 1997. The Maf transcription factors: Regulators of differentiation. *Trends Biochem. Sci.* **22**: 437–441.
- Burke, D., Wilkes, D., Blundell, T.L., and Malcolm, S. 1998. Fibroblast growth factor receptors: Lessons from the genes. *Trends Biochem. Sci.* **23**: 59–62.
- Chen, P.L., Riley, D.J., Chen, Y., and Lee, W.H. 1996. Retinoblastoma protein positively regulates terminal adipocyte differentiation through direct interaction with C/EBPs. *Genes & Dev.* **10**: 2794–2804.
- Cobrinik, D., Lee, M.-H., Hannon, G., Mulligan, G., Bronson, R.T., Dyson, N., Harlow, E., Beach, D., Weinberg, R.A., and Jacks, T. 1996. Shared role of the pRB-related p130 and p107 proteins in limb development. *Genes & Dev.* **10**: 1633–1644.
- Coller, H.A., Grandori, C., Tamayo, P., Colbert, T., Lander, E.S., Eisenman, R.N., and Golub, T.R. 2000. Expression analysis with oligonucleotide microarrays reveals that MYC regulates involved in growth, cell cycle, signaling, and adhesion. *Proc. Natl. Acad. Sci.* **97**: 3260–3265.
- Crawford, M.J., Lanctot, C., Tremblay, J.J., Jenkins, N., Gilbert, D., Copeland, N., Beatty, B., and Drouin, J. 1997. Human and murine PTX1/Ptx1 gene maps to the region for Treacher Collins syndrome. *Mamm. Genome* **8**: 841–845.
- Davis, A.A., Matzuk, M.M., and Reh, T.A. 2000. Activin A promotes progenitor differentiation into photoreceptors in rodent retina. *Mol. Cell. Neurosci.* **15**: 11–21.
- Duronio, R.J., O'Farrell, P.H., Xie, J.E., Brook, A., and Dyson, N. 1995. The transcription factor E2F is required for S-phase during *Drosophila* embryogenesis. *Genes & Dev.* **9**: 1445–1455.
- Dyson, N. 1998. The regulation of E2F by pRB-family proteins. *Genes & Dev.* **12**: 2245–2262.
- Eisen, M.B., Spellman, P.T., Brown, P.O., and Botstein, D. 1998. Cluster analysis and display of genome-wide expression patterns. *Proc. Natl. Acad. Sci.* **95**: 14863–14868.
- Fambrough, D., McClure, K., Kazlauskas, A., and Lander, E.S. 1999. Diverse signaling pathways activated by growth factor receptors induce broadly overlapping rather than independent sets of genes. *Cell* **97**: 727–741.
- Gudas, J.M., Payton, M., Thukral, S., Chen, E., Bass, M., Robinson, M.O., and Coats, S. 1999. Cyclin E2, a novel G1 cyclin that binds to CDK2 and is aberrantly expressed in human cancers. *Mol. Cell. Biol.* **19**: 612–622.
- Hanahan, D. and Weinberg, R.A. 2000. The hallmarks of cancer. *Cell* **100**: 57–70.
- Helin, K. 1998. Regulation of cell proliferation by the E2F transcription factors. *Curr. Opin. Genet. Dev.* **8**: 28–35.
- Helin, K., Wu, C.-L., Fattaey, A.R., Lees, J.A., Dynlacht, B.D., Ngwu, C., and Harlow, E. 1993. Heterodimerization of the transcription factors E2F-1 and DP-1 leads to cooperative transactivation. *Genes & Dev.* **7**: 1850–1861.
- Humbert, P.O., Rogers, C., Ganiatsas, S., Landsberg, R.L., Trimarchi, J.M., Dandapani, S., Brugnara, C., Erdman, S., Schrenzel, M., Bronson, R.T., et al. 2000a. E2F4 is essential for normal erythrocyte maturation and neonatal viability. *Mol. Cell* **6**: 281–291.
- Humbert, P.O., Verona, R., Trimarchi, J.M., Dandapani, S., and Lees, J.A. 2000b. E2f3 is critical for normal cellular proliferation. *Genes & Dev.* **14**: 690–703.
- Ichijo, H., Nishida, E., Irie, K., ten Dijke, P., Saitoh, M., Moriguchi, T., Takagi, M., Matsumoto, K., Miyazono, K., and Gotoh, Y. 1997. Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science* **275**: 90–94.
- Iyer, V.R., Eisen, M.B., Ross, D.T., Schuler, G., Moore, T., Lee, J.C.F., Trent, M., Staudt, L.M., Hudson Jr., J., Boguski, M.S., et al. 1999. The transcriptional program in the response of human fibroblasts to serum. *Science* **283**: 83–87.
- Kondo, T. and Raff, M. 2000. The Id4 HLH protein and the timing of oligodendrocyte differentiation. *EMBO J.* **19**: 1998–2007.
- Koziczak, M., Krek, W., and Nagamine, Y. 2000. Pocket protein-independent repression of urokinase-type plasminogen activator and plasminogen activator inhibitor 1 gene expression by E2F1. *Mol. Cell. Biol.* **20**: 2014–2022.
- Lauper, N., Beck, A.R.P., Cariou, S., Richman, L., Hofmann, K., Reith, W., Slingerland, J.M., and Amati, B. 1998. Cyclin E2: a novel CDK2 partner in the late G1 and S phases of the mammalian cell cycle. *Oncogene* **17**: 2637–2643.
- Lee, C.-K., Klopp, R.G., Weindruch, R., and Prolla, T.A. 1999. Gene expression profile of aging and its retardation by caloric restriction. *Science* **285**: 1390–1393.
- Lee, C.-K., Weindruch, R., and Prolla, T.A. 2000. Gene-expression profile of the aging brain in mice. *Nat. Genet.* **25**: 294–297.
- Leimeister, C., Externbrink, A., Klamt, B., and Gessler, M. 1999. Hey genes: A novel subfamily of hairy- and Enhancer of split related genes specifically expressed during mouse embryogenesis. *Mech. Dev.* **85**: 173–177.
- Leone, G., DeGregori, J., Yan, Z., Jakoi, L., Ishida, S., Williams, R.S., and Nevins, J.R. 1998. E2F3 activity is regulated during the cell cycle and is required for the induction of S phase. *Genes & Dev.* **12**: 2120–2130.
- Lindeman, G.J., Dagnino, L., Gaubatz, S., Xu, Y., Bronson, R.T., Warren, H.B., and Livingston, D.M. 1998. A specific, non-proliferative role for E2F-5 in choroid plexus function revealed by gene targeting. *Genes & Dev.* **12**: 1092–1098.
- Lipinski, M.M. and Jacks, T. 1999. The retinoblastoma gene family in differentiation and development. *Oncogene* **18**: 7873–7882.
- Lockhart, D.J., Dong, H., Byrne, M.C., Follettie, M.T., Gallo, M.V., Chee, M.S., Mittmann, M., Wang, C., Kobayashi, M., Horton, H., et al. 1996. Expression monitoring by hybridization to high-density oligonucleotide arrays. *Nat. Biotechnol.* **14**: 1675–1680.
- Lukas, J., Sørensen, C.S., Lukas, C., Santoni-Rugiu, E., and Bartek, J. 1999. p16INK4a, but not constitutively active pRb, can impose a sustained G1 arrest: molecular mechanisms and implications for oncogenesis. *Oncogene* **18**: 3930–3935.
- Maandag, E.C.R., van der Valk, M., Vlaar, M., Feltkamp, C., O'Brien, J., van Roon, M., van der Lugt, N., Berns, A., and te Riele, H. 1994. Developmental rescue of an embryonic-lethal mutation in the retinoblastoma gene in chimeric mice. *EMBO J.* **13**: 4260–4268.
- Massague, J. and Chen, Y.G. 2000. Controlling TGF- β signaling. *Genes & Dev.* **14**: 627–644.
- Matzuk, M.M., Kumar, T.R., Vassalli, A., Bickenbach, J.R., Roop, D.R., Jaenisch, R., and Bradley, A. 1995. Functional analysis of activins during mammalian development. *Nature* **374**: 354–356.
- Meraldi, P., Lukas, J., Fry, A.M., Bartek, J., and Nigg, E.A. 1999. Centrosome duplication in mammalian somatic cells requires E2F and Cdk2-Cyclin A. *Nat. Cell. Biol.* **1**: 88–93.
- Müller, H. and Helin, K. 2000. The E2F transcription factors: key regulators of cell proliferation. *Biochem. Biophys. Acta* **1470**: M1–M12.
- Müller, H., Moroni, M.C., Vigo, E., Petersen, B.O., Bartek, J., and Helin, K. 1997. Induction of S-phase entry by E2F transcription factors depends on their nuclear localization. *Mol. Cell. Biol.* **17**: 5508–5520.
- Murata, M., Eto, Y., Shibai, H., Sakai, M., and Muramatsu, M.

1988. Erythroid differentiation factor is encoded by the same mRNA as that of the inhibin β A chain. *Proc. Natl. Acad. Sci.* **85**: 2434–2438.
- Myster, D.L., Bonnette, P.C., and Duronio, R.J. 2000. A role for the DP subunit of the E2F transcription factor in axis determination during *Drosophila* oogenesis. *Development* **127**: 3249–3261.
- Nakamura, M., Masuda, H., Horii, J., Kuma, K.-i., Yokohama, N., Ohba, T., Nishitani, H., Miyata, T., Tanaka, M., and Nishimoto, T. 1998. When overexpressed, a novel centrosomal protein, RANBPM, causes ectopic microtubule nucleation similar to g-tubulin. *J. Cell Biol.* **143**: 1041–1052.
- Nevins, J.R. 1998. Toward an understanding of the functional complexity of the E2F and retinoblastoma families. *Cell Growth Differ.* **9**: 585–593.
- Peeper, D.S., Upton, T.M., Ladha, M.H., Neuman, E., Zalvide, J., Bernards, R., DeCaprio, J.A., and Ewen, M.E. 1997. Ras signaling linked to the cell-cycle machinery by the retinoblastoma protein. *Nature* **386**: 177–181.
- Rebollo, A., Dumoutier, L., Renauld, J.-C., Zaballos, A., Ayllon, V., and Martinez-A.C. 2000. Bcl-3 expression promotes cell survival following interleukin-4 deprivation and is controlled by AP1 and AP1-like transcription factors. *Mol. Cell Biol.* **20**: 3407–3416.
- Rempel, R.E., Saenz-Robles, M.T., Storms, R., Morham, S., Ishida, S., Engel, A., Jakoi, L., Melhem, M.F., Pipas, J.M., Smith, C., and Nevins, J.R. 2000. Loss of E2F4 activity leads to abnormal development and multiple cellular lineages. *Mol. Cell* **6**: 293–306.
- Seum, C., Spierer, A., Pauli, D., Szidonya, J., Reuter, G., and Spierer, P. 1996. Position-effect variegation in *Drosophila* depends on dose of the gene encoding the E2F transcriptional activator and cell cycle regulator. *Development* **122**: 1949–1956.
- Shelton, D.N., Chang, E., Whittier, P.S., Choi, D., and Funk, W.D. 1999. Microarray analysis of replicative senescence. *Curr. Biol.* **9**: 939–945.
- Sherr, C.J. 1996. Cancer Cell Cycles. *Science* **274**: 1672–1677.
- Sherr, C.J. 1998. Tumor surveillance via ARF-p53 pathway. *Genes & Dev.* **12**: 2984–2991.
- Suzuki, A. and Hemmati-Brivanlou, A. 2000. *Xenopus* embryonic E2F is required for the formation of ventral and posterior cell fates during early embryogenesis. *Mol. Cell* **5**: 217–229.
- Szeto, D.P., Rodriguez-Esteban, C., Ryan, A.K., O'Connell, S.M., Liu, F., Kiuoussi, C., Gleiberman, A.S., Izpisua-Belmonte, J.C., and Rosenfeld, M.G. 1999. Role of the Bicoid-related homeodomain factor Pitx1 in specifying hindlimb morphogenesis and pituitary development. *Genes & Dev.* **15**: 484–494.
- Takahashi, Y., Rayman, J.B., and Dynlacht, B.D. 2000. Analysis of promoter binding by the E2F and pRB families in vivo: distinct E2F proteins mediate activation and transactivation. *Genes & Dev.* **14**: 804–816.
- Tsai, K.Y., Hu, Y., Macleod, K.F., Crowley, D., Yamasaki, L., and Jacks, T. 1998. Mutation of E2F1 suppresses apoptosis and inappropriate S-phase entry and extends survival of Rb-deficient mouse embryos. *Mol. Cell* **2**: 293–304.
- Vigo, E., Müller, H., Prosperini, E., Hateboer, G., Cartwright, P., Moroni, M.C., and Helin, K. 1999. CDC25A phosphatase is a target of E2F and is required for efficient E2F-1 induced S phase. *Mol. Cell Biol.* **19**: 6379–6395.
- Wang, Z.M., Yang, H., and Livingston, D.M. 1998. Endogenous E2F-1 promotes timely G0 exit of resting mouse embryo fibroblasts. *Proc. Natl. Acad. Sci.* **95**: 15583–15586.
- Williams, B.O., Schmitt, E.M., Remington, L., Bronson, R.T., Albert, D.M., Weinberg, R.A., and Jacks, T. 1994. Extensive contribution of Rb-deficient cells to adult chimeric mice with limited histopathological consequences. *EMBO J.* **13**: 4251–4259.
- Yamasaki, L., Bronson, R., Williams, B.O., Dyson, N.J., Harlow, E., and Jacks, T. 1998. Loss of E2F-1 reduces tumorigenesis and extends the lifespan of Rb1(+/-)mice. *Nat. Genet.* **18**: 360–364.
- Zariwala, M., Liu, J., and Xiong, Y. 1998. Cyclin E2, a novel human G1 cyclin and activating partner of CDK2 and CDK3, is induced by viral oncoproteins. *Oncogene* **17**: 2787–2798.
- Zhao, R., Gish, K., Murphy, M., Yin, Y., Notterman, D., Hoffman, W.H., Tom, E., Mack, D.H., and Levine, A.J. 2000. Analysis of p53-regulated gene expression patterns using oligonucleotide arrays. *Genes & Dev.* **14**: 981–993.
- Zuber, J., Tchernitsa, O., Hinzmann, B., Schmitz, A.C., Grips, M., Hellriegel, M., Sers, C., Rosenthal, A., and Schafer, R. 2000. A genome-wide survey of RAS transformation targets. *Nat. Genet.* **24**: 144–152.