

The *Arabidopsis HOS1* gene negatively regulates cold signal transduction and encodes a RING finger protein that displays cold-regulated nucleo–cytoplasmic partitioning

Hojoung Lee, Liming Xiong, Zhizhong Gong, Manabu Ishitani, Becky Stevenson, and Jian-Kang Zhu¹

Department of Plant Sciences, University of Arizona, Tucson, Arizona 85721, USA

Low temperature is one of the most important environmental stimuli that control gene transcription programs and development in plants. In *Arabidopsis thaliana*, the *HOS1* locus is a key negative regulator of low temperature-responsive gene transcription. The recessive *hos1* mutation causes enhanced induction of the *CBF* transcription factors by low temperature as well as of their downstream cold-responsive genes. The *hos1* mutant plants flower early, and this correlates with a low level of *Flowering Locus C* gene expression. The *HOS1* gene was isolated through positional cloning. *HOS1* encodes a novel protein with a RING finger motif near the amino terminus. *HOS1* is ubiquitously expressed in all plant tissues. *HOS1*–GFP translational fusion studies reveal that *HOS1* protein resides in the cytoplasm at normal growth temperatures. However, in response to low temperature treatments, *HOS1* accumulates in the nucleus. Ectopic expression of *HOS1* in wild-type plants causes cosuppression of *HOS1* expression and mimics the *hos1* mutant phenotypes.

[Key Words: Low temperature; signal transduction; *HOS1*; *FLC*; RING finger]

Received November 15, 2000; revised version accepted February 2, 2001.

Plants and other poikilothermic organisms adapt to ambient temperature by adjusting membrane fluidity, metabolism, and gene expression profiles (Murata and Los 1997; Thomashow 1999). Low temperature is one of the most important environmental factors influencing plant distribution, growth, development, and survival. Plants from temperate regions can increase their freezing tolerance by being exposed to low nonfreezing temperatures, a process known as cold acclimation (Guy 1990). Low temperature-induced gene expression is key to cold acclimation (Guy et al. 1985; Thomashow 1999). For example, induction of lipid desaturase genes may contribute to freezing tolerance by altering lipid composition (Gibson et al. 1994). Cold-induced molecular chaperones may help prevent protein denaturation during freezing (Anderson et al. 1994). Many cold-induced genes encode novel proteins, some of which have been shown to discourage the formation of deleterious membrane structures during freezing (Steponkus et al. 1998). Ectopic expression of *CBF* transcription factors has been shown to

induce a set of cold-responsive genes at warm temperatures and to mimic cold acclimation in terms of increases in freezing tolerance (Jaglo-Ottosen et al. 1998; Liu et al. 1998).

Many of the cold-regulated genes can also be induced by the phytohormone abscisic acid (ABA) or by osmotic stress treatment (Gilmour et al. 1992; Kurkela and Borg-Franck 1992; Nordin et al. 1993). Low temperature and osmotic stresses appear to have several features in common. Freezing injury is largely a consequence of freezing-induced dehydration stress. Low nonfreezing temperatures can also lead to water stress by impairing water absorption by the roots and water transport to the shoot (Levitt 1980). ABA is known to play an important role in cold acclimation, and treatment with ABA at normal growth temperatures increases the freezing tolerance of a wide range of plants (Chen et al. 1983). Furthermore, exposure to low temperature transiently increases ABA levels (Chen et al. 1983).

Cold-responsive genes such as *RD29A* (also known as *COR78* or *LTI78*) and *COR15A* are known to contain two *cis*-regulatory elements in their promoters, that is, the ABRE and DRE/CRT elements (Yamaguchi-Shinozaki and Shinozaki 1994; Stockinger et al. 1997). The ABRE element is responsive to ABA, whereas the DRE/

¹Corresponding author.

E-MAIL jkzhu@ag.arizona.edu; FAX (520) 621-7186.

Article and publication are at www.genesdev.org/cgi/doi/10.1101/gad.866801.

CRT element mediates ABA-independent regulation of gene expression by cold or hyperosmotic stress. Transcription factors termed CBFs or DREBs that bind to DRE/CRT are induced early by cold or osmotic stress (Stockinger et al. 1997; Gilmour et al. 1998; Liu et al. 1998). Constitutive or inducible overexpression of some of the transcription factors leads to enhanced expression of downstream cold responsive genes and increased plant tolerance to freezing, drought, and salt stresses (Jaglo-Ottosen et al. 1998; Liu et al. 1998; Kasuga et al. 1999).

Besides the CBF transcription factors, other potential cold signal transduction components in plants include *SFR6*, *ESK1*, *HOS1*, and *HOS2*, which are defined by *Arabidopsis* mutations. The molecular identities of these genetic loci await molecular cloning. The *sfr6* mutation suppresses cold and ABA induction of genes with the CRT/DRE promoter element (Knight et al. 1999). The *esk1* mutants are constitutively freezing-tolerant but are apparently not affected in the expression of genes with the CRT/DRE element, suggesting that *ESK1* may be involved in a distinct pathway related to freezing tolerance (Xin and Browse 1998). *HOS1* and *HOS2* are both negative regulators of cold signaling pathways (Ishitani et al. 1998; Lee et al. 1999). Mutations in *HOS1* or *HOS2* lead to enhanced cold induction of genes such as *RD29A*, *COR15A*, *KIN1*, and *ADH*. Without cold acclimation, *hos1* mutant plants have reduced capacity for freezing tolerance. After cold acclimation, *hos1* plants acquire as much freezing tolerance as do wild-type plants (Ishitani et al. 1998).

hos1 mutant plants flower considerably earlier, which may be related to vernalization, another important plant process regulated by low temperature (Ishitani et al. 1998). *Vernalization* refers to the promotion of flowering by prolonged periods of low temperature exposure. The early flowering phenotype of *hos1* mutant plants appears to be a manifestation of constitutive vernalization (Ishitani et al. 1998). This is because (1) *hos1* plants have a reduced threshold of low temperature activation of cold-responsive genes, that is, cold signaling happens even at normal growth temperatures in the mutant plants (Ishitani et al. 1998); and (2) nonvernalized *hos1* mutant plants flower as early as vernalized wild-type plants (Ishitani et al. 1998). *hos1* mutant plants can be further vernalized by low temperature treatment to flower even earlier.

To address the mechanisms of cold-regulated gene expression and plant development, a necessary step is to determine the molecular nature of the key regulatory genes involved. Toward this goal, we report here the map-based cloning and molecular analysis of *HOS1*, the first such genetic locus involved in cold-regulated plant development and signaling to be characterized at the molecular level. The sequence data reveal that *HOS1* encodes a novel RING finger protein. Studies with *HOS1*-GFP fusion protein show that *HOS1* displays cold-regulated nucleo-cytoplasmic partitioning, suggesting an important role of *HOS1* may be to communicate cold-generated signals in the cytoplasm to the nucleus to regulate the level of gene transcription.

Results

Expression of CBF transcription factor genes in hos1 mutant plants

Transcripts for the *CBF/DREB1* transcription factors are rapidly induced by low temperature treatment (Gilmour et al. 1998; Liu et al. 1998; Medina et al. 1999). These transcription factors in turn activate the expression of downstream *COR/RD/KIN* genes such as *COR15A*, *COR47*, *RD29A*, and *KIN1*. We have shown previously that these *COR/RD/KIN* genes are superinduced by cold in *hos1* mutant plants (Ishitani et al. 1998). Northern analysis was carried out to determine whether *CBF* genes are also superinduced by cold in the *hos1* mutant. Both *CBF2* and *CBF3* were found to exhibit enhanced cold induction in *hos1* mutant plants (Fig. 1). At 3 h after cold treatment, *CBF2* expression in *hos1* mutants is approximately 50% higher than in wild-type plants. Increased *CBF2* and *CBF3* expression is more pronounced after longer cold treatment, for example, at 6, 12, 24, and 48 h (Fig. 1). We did not observe significant cold induction of *CBF1* in either the wild-type or *hos1* mutant plants (not shown).

The hos1 mutation reduces expression of the Flowering Locus C gene

The *Flowering Locus C (FLC)* gene that encodes a MADS-box protein is a central regulator of flowering induction by vernalization (Michaels and Amasino 1999; Sheldon et al. 2000). The level of *FLC* determines the extent of the vernalization response in the promotion of flowering: The duration of cold treatment correlates with the extent of down-regulation of *FLC* expression (Sheldon et al. 2000). Because the *hos1* mutant plants flower early (Ishitani et al. 1998) and appear to be constitutively vernalized (Fig. 2A), we carried out Northern analysis to determine whether *FLC* expression is altered in the mutant. Consistent with previous studies (Sheldon et al. 2000), the *FLC* transcript level is high in wild-




Figure 1. The *CBF* transcription factor genes are expressed at higher levels in *hos1* mutant plants in response to cold treatments. Total RNA (20 μ g per lane) was isolated from wild-type and *hos1* mutant seedlings treated with cold (0°C for the indicated time periods), ABA (100 μ M, 3 h), NaCl (300 mM, 5 h), or PEG (30%, 5 h). The RNA blot was probed with 32 P-labeled *RD29A*, *CBF2*, *CBF3*, and *actin* cDNA probes, with the *actin* used as a loading control.




Figure 2. The early flowering phenotype and reduced expression of the vernalization regulator *FLC* in *hos1* mutant plants. (A) Flowering time of C24 wild-type and *hos1* mutant plants with and without 2 or 4 wk of vernalization treatment at 0°C. Flowering time is indicated by the total number of rosette leaves at time of flower bud emergence. Shown are the average and standard deviation ($n = 15$). (B) *FLC* expression. Total RNA (20 μ g per lane) was isolated from wild-type and *hos1* mutant seedlings grown in agar plates that were incubated at 0°C for 0, 2, and 4 wk. The *tubulin* gene was used as a loading control.

type C24 plants, but it decreases significantly as the plants become vernalized by cold treatments (Fig. 2B). Interestingly, *hos1* mutant plants have a substantially lower level of *FLC* expression even in the absence of vernalization treatment (Fig. 2B). This lower level of *FLC* expression correlates with the early flowering phenotype of the *hos1* mutant plants (Fig. 2A). After 2 wk of vernalization treatment, the *FLC* level in *hos1* mutant plants is not reduced. After 4 wk of vernalization, the *FLC* level decreases slightly (Fig. 2B), as does the flowering time (Fig. 2A). As a control, the *tubulin* gene transcript levels are not different between *hos1* and the wild type, and are not changed by vernalization (Fig. 2B). These results suggest that wild-type *HOS1* is a negative regulator of vernalization and that it is required for a high level of *FLC* expression.

Positional cloning of *HOS1*

To map *hos1* genetically, homozygous *hos1* mutant plants in the C24 ecotype were crossed with wild-type plants of the Columbia ecotype. The resulting F_1 plants were selfed, and homozygous *hos1* mutant plants were selected from the segregating F_2 population by luciferase imaging after 2 d of treatment at 0°C. Surveying of representative molecular markers from each of the five *Arabidopsis* chromosomes located *hos1* to chromosome II. Further analysis showed that *hos1* is closely linked with the SSLP (simple sequence length polymorphism) marker nga168. Several new SSLP markers were devel-

oped based on the genomic sequences of BAC clones in the nga168 region. Fine mapping using these new markers delimited *HOS1* to the BAC clone T5I7 (Fig. 3). Candidate open reading frames on T5I7 were sequenced from wild-type as well as *hos1* mutant plants. The sequence analysis revealed a single nucleotide substitution in the hypothetical *T5I7.11* gene in the *hos1* mutant. This mutation is predicted to create a premature stop codon, resulting in a truncated protein.

To confirm that we have cloned the correct gene, the *T5I7.11* sequence including 2807 bp upstream of the predicted ATG start codon was amplified from wild-type plants and cloned into the binary vector pCIT20. The construct was introduced into *hos1* mutant plants via *Agrobacterium*-mediated transformation. Transformants were selected based on hygromycin resistance; the progenies of two transformants were tested by luciferase imaging after 2 d of cold treatment. Both hygromycin resistance and cold-induced *RD29A-LUC* expression segregated in the T_2 generation. All hygromycin-resistant plants exhibited the wild-type phenotype in terms of cold-induced *RD29A-LUC* expression and time to flower. Figure 4A–C shows an example of complemented *hos1* mutant plants in the T_3 generation. In response to cold treatment, *hos1* mutant plants exhibited brighter luminescence, indicating a higher level of *RD29A-LUC* expression compared with the wild type (Fig. 4B). In contrast, luminescence intensities from *hos1* mutant seedlings containing the introduced wild-type *HOS1* gene were similar to those in wild-type plants (Fig. 4B). Quantification of the luminescence intensities confirmed that the mutant was fully complemented by the wild-type *HOS1* gene (Fig. 4C). The wild-type *HOS1* transgene also rescued all the other phenotypes exhibited by *hos1* mutant plants, such as pale, small leaves and early flowering (data not shown).

HOS1 encodes a novel RING finger protein

The transcribed sequence of the *HOS1* gene was determined by sequencing cDNAs obtained by the reverse transcriptase polymerase chain reaction (RT-PCR). Comparison of the cDNA with the genomic sequence showed that the *HOS1* gene contains nine exons and eight introns (Fig. 3B). *HOS1* is predicted to encode a polypeptide of 915 amino acids, with an estimated molecular mass of 104 kD (Fig. 5). Database searches revealed that *HOS1* contains a short stretch of amino acids near the amino terminus that are similar to the RING finger domain found in a group of proteins known as IAPs (inhibitor of apoptosis) (Fig. 6; Miller 1999). The rest of *HOS1* does not show significant homology to any other sequences in the GenBank database. The *hos1* mutation, a G \rightarrow A substitution, is predicted to create a premature stop codon that truncates the protein after His 643. This suggests that the carboxy-terminal one-third of *HOS1* is functionally essential.

The RING finger is a small zinc-binding domain found in many proteins with diverse functions. Recent studies have shown that proteins containing a RING finger may




Figure 3. Positional cloning of the *HOS1* gene. (A) Physical mapping of *HOS1*. Genetic mapping delimited *HOS1* to the BAC clone T5I7. The *hos1* mutation was identified by sequencing and comparing all predicted genes on this BAC from *hos1* mutant and wild-type plants. (B) Structure of *HOS1* and the position of the *hos1* mutation. Positions are relative to the initiation codon. Filled boxes indicate the open reading frame, and lines between boxes indicate introns. The premature stop codon created by the *hos1* mutation is boxed.

act as E3 ubiquitin protein ligases, and that the RING finger can interact with E2 ubiquitin conjugating enzymes, thereby promoting ubiquitination of specific target proteins (for review, see Joazeiro and Weissman 2000). The RING finger motif is defined as a series of conserved cysteine and histidine residues: Cys-X₂-Cys-X₉₋₃₉-Cys-X₁₋₃-His-X₂₋₃-Cys/His-X²-Cys-X₄₋₄₈-Cys-X₂-Cys, where X can be any amino acid (Saurin et al. 1996). The RING finger of *HOS1* is most similar to that of IAPs, which can be classified as a C₃HC₄-type RING (Miller 1999). Interestingly, the first Cys in the RING finger of *HOS1* is replaced with Leu (Fig. 6). RING variants have also been seen in other well-known RING finger proteins (Saurin et al. 1996). For example, in mouse MDM2, Cys₃ is replaced by a Thr residue, whereas in CART1, Cys₇ is substituted for an Asp residue (Saurin et al. 1996).

HOS1 expression in *Arabidopsis* seedlings

Northern analysis was carried out to examine *HOS1* transcript abundance in *Arabidopsis* seedlings subjected to various treatments. In wild-type plants, *HOS1* mRNA was detected in unstressed control as well as cold-, ABA-, NaCl-, or PEG-treated samples (Fig. 7). *HOS1* transcript abundance was not substantially influenced by ABA, NaCl, or PEG treatment. In response to cold treatment, *HOS1* expression declined quickly and transiently. At 30 min after cold treatment, *HOS1* expres-

sion was almost undetectable (Fig. 7). However, by 1 h after cold treatment, *HOS1* expression recovered to pre-treatment levels. This rapid and transient reduction of *HOS1* message in response to brief cold treatment was consistently observed in independent experiments. After recovery from the decline, *HOS1* expression was then maintained at this level until 2 d after cold treatment, before declining again (Fig. 7).

Very low levels of *HOS1* transcript were detected in *hos1* mutant plants subjected to the same treatments as were applied to wild-type plants (Fig. 7). The low level of *HOS1* expression is probably attributable to active degradation of the mutant mRNA, since it is known that mRNA surveillance mechanisms degrade transcripts with premature stop codons (Hilleren and Parker 1999). To verify that the stress treatments were appropriate, the RNA blot was also probed with *RD29A* and *CBF2*. The results show that the two genes were up-regulated with the appropriate kinetics, that is, *CBF2* induction occurred earlier than *RD29A* (Fig. 7), although not as early as the *CBF2* induction seen in some other experiments (Fig. 1). The results also confirm that *CBF2* as well as *RD29A* are expressed at higher levels in *hos1* mutant plants under cold treatment. Hybridization with a β -*tubulin* cDNA probe showed RNA loading in stressed and in control unstressed sample lanes (Fig. 7).

To examine *HOS1* expression in various plant tissues, 2.78 kb of the genomic sequence upstream of the *HOS1*




Figure 4. Complementation of *hos1* mutant by the wild-type *HOS1* gene. Ten-day-old wild-type seedlings, *hos1*, and homozygous T_3 progenies of a *hos1* plant containing the wild-type *HOS1* transgene (designated as *hos1 + HOS1*) grown in agar medium were incubated at 0°C for 24 h before the bioluminescence image was taken. (A) A picture of wild-type, *hos1*, and T_3 progenies of *hos1* plants containing the *HOS1* complementation construct. (B) Bioluminescence image of plants shown in A. (C) Quantification of the luminescence intensities in B.

open reading frame (−2807 to −23 from ATG) was fused with the β -glucuronidase reporter gene (*GUS*), and the resulting construct was introduced into wild-type *Arabidopsis* plants. *GUS* expression was observed in all plant tissues examined (Fig. 8). In leaves, the expression was stronger in the vasculature (Fig. 8C), but could be seen in mesophyll tissues as well (Fig. 8D). Strong expression was observed in roots (Fig. 8A), with root tips having especially high levels (Fig. 8B).

HOS1 accumulates in the nucleus in response to cold treatment

Knowledge of the subcellular localization of a protein is crucial to the understanding of its cellular function. To determine the subcellular localization of *HOS1* gene product, *HOS1* cDNA was fused in-frame at its C terminus with the GFP (green fluorescence protein) marker. *HOS1*–GFP fusion protein was expressed under the CaMV 35S promoter in transgenic *Arabidopsis* plants. The subcellular localization of *HOS1*–GFP was determined by green fluorescence imaging under a confocal microscope. Without cold treatment, only very faint green fluorescence signals slightly above background (i.e., nontransformed plants) could be detected in the majority of cells (Fig. 9A). This suggests that *HOS1*–GFP was either expressed at a very low level or was being rapidly degraded at normal growth temperatures. Occasionally, a few cells were found to express a high level of

HOS1–GFP. In these cells, the green fluorescence was entirely cytoplasmically localized (Fig. 9A, indicated by arrow). Amazingly, when the *HOS1*–GFP transgenic plants were placed at 0°C for 1–2 d, green fluorescence could be seen in structures consistent with the nuclei (Fig. 9B). To verify that the structures emitting green fluorescence were nuclei, the seedling roots were stained with propidium iodide. The results show that the green fluorescence-emitting organelles were stained with propidium iodide, suggesting that the organelles are, indeed, nuclei (Fig. 9B, inserts). The occasional cells expressing high levels of cytoplasmic *HOS1*–GFP at normal temperatures also showed strong nuclear *HOS1*–GFP expression after cold treatment (Fig. 9C,D).

The nuclear accumulation of *HOS1*–GFP fusion protein was clearly visible after plants were subjected to 0°C treatment for 20 h. Longer cold treatment did not change the intensity of the nuclear green fluorescence. After the cold-treated plants were returned to normal growth temperatures (18°–23°C) for more than 12 h, the nuclear green fluorescence disappeared, suggesting that the fusion protein moved back to the cytoplasm and/or was rapidly degraded.

To determine whether the cold-induced nuclear accumulation is specific to *HOS1*–GFP, we treated transgenic *Arabidopsis* plants containing a known cytoplasmically localized small acidic ribosomal protein–GFP fusion (line Q1; Cutler et al. 2000) at 0°C for 48 h. Only cytoplasmically localized green fluorescence was detected in

CGCCAACACCAACCACGCTTTTTCTCTCTTTACCTCCATTTTTGTATTTCGTCACCGGAGTAGATGGATACGAGAGAAA
 TCACGGTTTTGCCTCCGCCGAGATCGATATCTCTTCCACGAGCCTAATTACAGTAGCAAGCCTGTTACAGTTTC
 TTCGCTTTATACCTTTTTATTCTCCACTGAATTAGTTTAGCTCAGGTCCTTAGGATTTCTAGGTTACAGTTAAAAA
 ATGGGTTTTGCAGAGCTTAGAGTTAATTCGAATTTATTGTTGTAACCTACAGGAGGCGTTGAAACATTTGGCATCT
 M G F A E L R V N F N F E F I V C N L O E A L K H L A S
 ATAAACCTCAGAGAATTATGTAATGAGGCAAAAGTTGAGCGATGTCGTCGCAACAGAGACTTAGCTAGCTGTGGAAGA
I N L R E L C N E A K V E R C R A T R D L A S C G R
 TTTGTGAACATATGTTGAATCCATGTGGTCATGCCCTCTGTGCACAGAGTGTGTGACAGGTTGTGATGTTTGTCTT
F V N Y V L N P C G H A S L C T E C C O R C D F V C P
 ATCTGTAGAAGTACGCTACCAAAATTTGGAGATAGACTTAGGCTTCGCTCTTATTATGAGTGTGTTGAAGCTGTTTGG
I C R S T L P K F G D R L R L R L Y Y E C V E A G L
 ATTTCCCGAACACATGAAGAGGCGTCACAGGATTCGGATGAGATGAACATCAATTTGGCAGCTGATGTTTCATCCGCTT
 I S R R T H E E A S Q D S D E D E H Q L A A D V H R L
 TATCTCTGTTGATGTGCTATGAACAATAATTTGATCTCGGTGGTTTGGCATTATATACCAATCTCTGTATGGAT
 Y S L F D V A M N N L I S V V C H Y I T N V C M D
 GAGACAGCTGTATCTAGTGATCCGGTCATTGCTTCTTGTCTGGATGAAGTTGTTGCAAGGATTGGGTCAGCGTACA
 E T A V S S D P V I A F L L D E V V K D W V K R T
 TTCCGGACCTTAGTGAACCTCAAGAAATCTATAATCTTGAAACAAAAGAGATGCAAGCGTGGTTAGATAAATC
 F R S T L A E L Q E I Y N L E T K E M Q A W L D K L
 CTAAGGTGTTCCAAACAGGTGGCTGGCATATGATGTGCTCGAAGTTATGGAATCAGCTTTAAGGGTCTGTGTCA
 L R C S K V A G I C S V L E V M E S A F K G S V S
 CCTCAGCTTCAAGATGTGCAGACGCTGAGAGAGAACATTGGTAAAACAAAGCAGCATTGGACATAATGGTCTGGTGC
 P Q L Q D V Q T L R E N I G K T K Q H L D I M V W C
 ATCAGACATGGATTTCTGGATGATGTGAGTCTCGATTCGAATTTACATCGTGAATGCATGGTAGGGGAAAGA
 I R H G F L D D V R S R Y S N F T S W N A L V G E R
 AAATCAAATGCAAGTTAAGCGTGCATGGCCTGATGCTGTAGACCACTTTCAGATTGCAGTGTACAGAGTCTCCCTC
 K S N A V K R A W P D A V D Q S V D C S V Q S A S L
 TTTATTGAGGATGCTTTAGAAAATCTTGAGAGAGAGCCAGAATACAGTCAGGAAATAGGTGCCGACCTAGAAGTTGGA
 F I E D A L E N L E R E P E Y S Q E I G A D L E V G
 CGTTTACAGAGGATAAGAGATCATTCTTAGGTCCAAATAGAGGGAACCTCAGGATCTTATCCATTGAGAACTT
 R L Q K D K R S F L R S K I E G T S G S Y P F E N L
 AGAAGTCTGCTGACATGCTCTTTTACATGGCGGTTGAGATTGTTGTTGCGAAGCAAGCAATTTTCTTTACTAC
 R T A A D M L F L H G G S D L V V A K Q A I F L Y Y
 CTGTTGATCGGCATTGGACTACCCCTGAAAAATTTGGAAGCACACTATAGATGATTTTGTCTACGCTTTGGCATA
 L F D R H W T T P E K Y W K H T I D D F A A T F G I
 ACGAGCATTTCATGCTGGAGTCTTTTGTATTTTATCTTCTCGATGATCACTCAGAGGAGGCACTCCAGGAAGTGT
 T R H S L L E S F V F Y L L D D H S E E A L Q E A C
 AGGATCTACCTGAAATATCGGTCAGAACTTACCCAAAGTTGCGCAAGTCTGTTAGAAAGGGATAACCCTGAA
 R I L P E I C G P E T Y P K V A Q V L L E R D N P E
 ACAGCTCTTATGGTCTTGGCTGGTCTGGTCTGGTGTATCAGAATTGGTTTCAATTGGTGAAGCTGTACAGCT
 T A L M V L R W S G R D G V S E L V S I G E A V T A
 CTTCTGTGAGAGTGAATGTGGTCTTTTGAAGTGAAGCATTACATACCAAAGGACACTCTGCTTGAAGTAAAGGAA
 L R V R V E C G L L S E A F T Y Q R T L C L K V K E
 AATAATTTGAAAAATGGAGCTGTGAAACATGCGTCTGATGATCTAGACATCTGGAGTTGGACGGAATGGATGGAGATA
 N N L K N G A V K H A S D D L D I W S W T E W M E I
 CTTGTCAATGAATCTGCTGTCTTCTATTCGGAGAACTTGGTTGATCGAATCATTGAGTTACCGTGAACCCAGAT
 L V N E F C C L S I R R N L V D R I I E L P G W N P D
 GAGGAGAAATATCTGCACAGATGCTTATTAGATTTCGCAACTGATGACCCCTCATCAGCTGTGGTAGTCTTGGTT
 E E K Y L H R C L L D S A T D D P S S A V G S L L V
 GTCCTTTATCTTACGCGTACCGTTACATCCAGGCATACCAAGTTCGATCTTAGGCTTTCAGAAAATCGAAGAGGCTTC
 V F Y L Q R Y R Y I Q A Y Q V D L R L Q K I E E A F
 GTATCCGACAATCAAATGGGGAAGAAGTTATGTTTCAAGATGCGATCACAGAGTCATTGGAGAAAAGAGTTAGTTGAT
 V S D N Q I G E E V M F R M R S Q S H W R K E L V D
 AGAGCATAATATCTCCAGTAATTCAGCAGCAGCAAGTTAGATCTGGACAATTTTCTGAGATGGAAAGATGCCTCT
 R A I D I L P V I Q Q Q Q V R S G Q F S E M E D A S
 GAAGGAGCAAAAAATCTGATCTTCTGATGACCTGACATGATTACTTCTCAGTTCCCTTTGCCACTACTAATTC
 E G A K K S D L P D A P D M I T S S V P F A A T N S
 GTTTTCTCCAAAGTGTAAACAATGCCAGAGCTAGAGAGCCAGTGGCTAATAATGGCAGTCTTTTCAACCTGGTCAT
 V F L Q S C A N N A R A R E P V A N N G S P F Q P G H
 ATGATAGGAAACGCGTCCCATGATCTTTTACATGGAAGGTTATTACAAACGCAAATAGAGGACAAAAGAGGAAAGTT
 M I G N A S H D L S H G R L F T N A N R G Q K S E V
 AGAAGCGTAACCAAGAAATCTGAAATTCGGTGAATGTCAACTCCGTTCAAAGATCTAAATAGAGCTCGTGGGAACAGC
 R S V T K N L K F G E M S T P F K D L N R A R G N S
 CAGCTCCAAGGAAAAGGACTGAAGAATCTCTCCAGAAGTAAATGTTGATAGATACATAGAGAATAATATGAGCTCT
 Q L Q G K R T E E S S P E V N V D R Y I E N N M S S
 CTTTATCTCCGCGAATCACAGCTAACAATCCCGTGACAGTGAATCTAGCAGCAACCATCTCAATGGATCTTCCAG
 P Y L R R I T A N N P V T V K S S S N H L N G S S Q
 AAACCGGAATCCACCTTTTGGCACAAGGATGCAACAGACAAAAGATAACTTTGTTGATTTGGATGATCCAATGGAC
 K P E S T F F G T R M Q P D K D N F V D L D P M D
 ATGTCTCGAGCTTAAAGGACAACAACAATGTCTTGGCCACGGAGAGCAGAAAATAACAGTGGTGGATTGAGATGG
 M S S S L K D N N N V L A T E S R N N S G G L R W
 CGGTCAGATGAAACAAGCGATGACGAAGATGAACACTAGTTTGGTTCCATGCCGTTGAAGGGAAGGAGGAGCGT
 R S D E T S D D E D E L T S F G S M P V K G R R R R
AGATTCGAGCAAGATGAAATGAAACAAACTCGAAAAGACCATCATCAATTATGTTTAAAGCATAGAGAAAGAGAG
R F A A R •
 AGAGAGGTAGAGCAAGTTGTGTAAGAGGGGAGAGAGACTGTTACAGTGAGCGTTGTTCTCTGTTATTTTTGTGCT
 GAATTTGTGTATTCTCTCTAGTCT

Figure 5. The *HOS1* cDNA sequence and the conceptual translational product of its longest ORF (GenBank accession no. AAB87130). The RING finger domain is underlined, and a putative nuclear localization signal is double-underlined.




Figure 6. Alignment of the RING finger domain of HOS1 with those of animal IAPs. Amino acid residues in black indicate identical matches, and those in gray indicate conserved substitutions. The conserved Cys and His residues of the RING finger are indicated by asterisk marks. A missing Cys residue in HOS1 is indicated by a question mark. GeneBank accession numbers for various IAPs are as follows: Human IAP, AAC83232; Pig IAP, AAC39271; Mouse IAP, AAC42078; Rat IAP, AAG22971; HOS1, AAB87130.

these control plants either at 23°C or 0°C (Fig. 9E,F). These results show that cold treatment does not induce a general nonspecific nuclear accumulation of GFP fusion proteins.

Cosuppression of HOS1 expression in wild-type plants mimics hos1 mutant phenotypes

The *HOS1* open reading frame was overexpressed in wild-type plants under the CaMV 35S promoter to determine whether increased *HOS1* expression would down-regulate *RD29A-LUC* regulation and affect plant tolerance to chilling or freezing stress. Among 25 independent transgenic plant lines tested in the T₂ and T₃ generations, none showed reduced *RD29A-LUC* induction by cold treatments compared to wild type (data not shown). We also did not find any transgenic lines with altered chilling or freezing tolerance. Interestingly, there were two transgenic lines that showed higher than wild-type levels of *RD29A-LUC* expression in response to cold treatment. This phenotype segregated in the T₂ generation, with an ~3:1 ratio of the high expression to regular expression plants. The *RD29A-LUC* phenotype of one of the lines, #4, is shown in Figure 10A–C. The high expression plants were all found to carry the 35S–


HOS1 transgene according to their hygromycin resistance.

The increased *RD29A-LUC* expression in these 35S–*HOS1* transgenic lines mimics the phenotype of the *hos1* mutant plants, although the luciferase expression level was not as high as in the mutant (Fig. 10C). The strong luminescence phenotype of these transgenic lines is probably a result of posttranscriptional cosuppression of *HOS1* expression, because semiquantitative RT-PCR tests revealed a decreased amount of *HOS1* transcript in the high expression plants derived from the #4 line (Fig. 10D).

Discussion

Low temperature is an important environmental signal for all living organisms. Two well-known plant physiological processes are associated with the low temperature cue, namely, cold acclimation and vernalization. Cold acclimation refers to the phenomenon that plants from temperate regions gain freezing tolerance by preexposure to low nonfreezing temperatures (Guy 1990; Thomashow 1994). Vernalization is a plant-specific process whereby prolonged exposure to low temperature accelerates flowering (Chandler et al. 1996). *HOS1* is the only

Figure 7. *HOS1* transcript levels in wild-type and *hos1* mutant plants subjected to various stress treatments. Wild-type and *hos1* mutant 3-week-old seedlings were incubated at 0°C for 0 min, 10 min, 30 min, 1 h, 3 h, 6 h, 12 h, 24 h, or 48 h. Seedlings were also treated with ABA (A: 100 μM, 3 h), NaCl (N: 300 mM, 5 h), and PEG (P: 30%, 5 h). Total RNA was extracted, and 20 μg RNA was loaded into each lane. The blots were hybridized with ³²P-labeled *HOS1*, *RD29A*, *CBF2*, and β -*tubulin* cDNA probes. The β -*tubulin* gene was used as a loading control.






Figure 8. Ubiquitous expression of *HOS1* in *Arabidopsis* seedlings. Fifteen-day-old transgenic plants containing the *HOS1* promoter-*GUS* construct were analyzed for GUS expression. (A) Whole seedling, (B) roots, (C) leaf, and (D) close-up of part of a leaf.


gene known to regulate both cold acclimation and vernalization (Ishitani et al. 1998). For cold acclimation, *HOS1* negatively regulates the expression of cold-responsive genes that are known to contribute to cold acclimation. *CBF/DREB* transcription factor genes are induced by cold at higher levels in the *hos1* mutant (Figs. 1 and 7). This is probably responsible for higher levels of cold induction of *RD29A*, *COR47*, *COR15A*, and *KIN1* in the mutant because these later genes are controlled by the *CBF/DREB* transcription factors (Stockinger et al. 1997; Liu et al. 1998). However, the *hos1* mutation seems to also affect genes outside the *CBF* regulon. This is suggested by the effect of the *hos1* mutation on cold induction of the alcohol dehydrogenase gene (Ishitani et al. 1998), which does not appear to be under control of *CBF* transcription factors because its promoter is not known to contain the *DRE/CRT* element. These observations indicate that *HOS1* functions early in cold signal transduction. An early role of *HOS1* in cold signaling is consistent with its involvement in vernalization, that is, *FLC* expression and flowering time control.

Molecular cloning of *HOS1* revealed that it encodes a novel protein with a modified RING finger domain. The RING finger motif was first identified in the human *Really Interesting New Gene 1* (*RING1*) product (for review, see Freemont 2000). Several hundred proteins from diverse eukaryotes have been found to contain this motif or its variants. RING finger proteins include some of the

most important regulators of human diseases and various cellular signaling pathways, for example, *BRCA1*, a product of a breast cancer-associated gene; the proto-oncogene products *Cbl*, *BMI-1*, and *PML*; and the p53-regulator *MDM2* (for reviews, see Freemont 2000; Jackson et al. 2000; Joazeiro and Weissman 2000). However, no specific function was ascribed to the RING finger until recently. In the past two years, a number of RING finger proteins have been found to have E3 ubiquitin ligase activities, that is, they mediate the transfer of ubiquitin to heterologous substrates and/or to the RING finger proteins themselves. The RING finger sequence of *HOS1* is most similar to those found in the family of antiapoptotic molecules in animals known as IAPs (Fig. 6). IAPs were recently shown to catalyze their own ubiquitination, and this ubiquitination activity is dependent on the RING domain (Yang et al. 2000).

It is now becoming abundantly clear that the stability of critical regulatory proteins in the cell is dynamically controlled in response to environmental or developmental stimuli (Joazeiro and Weissman 2000). Because protein degradation must be highly selective in order for the cell not to cannibalize itself, the substrate-recognition step mediated by the RING finger E3 ubiquitin ligases is of critical importance. The existence of a large number of RING finger proteins may thus reflect the need to selectively target many different substrate proteins for degradation.

Figure 9. Cold treatment alters HOS1-GFP localization in *Arabidopsis* seedlings. (A) Cytoplasmic localization of HOS1-GFP fusion protein in root cells without cold treatment. T₂ 10-day-old seedlings containing the *HOS1-GFP* translational fusion construct grown on MS agar plates in the dark were analyzed for GFP expression under confocal microscope. The arrow points to a cell with strong cytoplasmic green fluorescence. (B) Nuclear localization of HOS1-GFP after cold treatment. Seedlings were incubated at 0°C for 48 h before GFP analysis. The spots emitting green fluorescence correspond to nuclei as confirmed by propidium iodide staining (red stain in insert). Arrows point to nuclei. (C) Those cells expressing strong cytoplasmic GFP at normal growth temperatures show strong nuclear GFP after cold treatment. Arrows point to nuclei. (D) Cells in C were stained with propidium iodide. Arrows point to nuclei. (E,F) Cytoplasmic GFP localization in the roots of transgenic *Arabidopsis* expressing an acidic ribosomal protein-GFP fusion (Cutler et al. 2000) either before (E) or after 48 h of cold treatment (F).



No RING finger protein has been previously reported to function in low temperature signal transduction. Thus, our results provide the first evidence that a RING finger protein is also important in regulating gene transcription and plant development in response to cold environments. Future studies will determine whether HOS1 can serve as an E3 ubiquitin ligase and what types of substrate proteins might be specifically targeted by HOS1 for degradation. Because HOS1 negatively regulates CBF transcript levels, it is tempting to speculate that HOS1 might control the turnover of ICE, the proposed inducer of CBF expression (Thomashow 1999). The *Arabidopsis* COP1 is a well-known RING finger protein that negatively regulates light-responsive gene expression and photomorphogenesis (Deng et al. 1992). It has been shown recently that COP1 functions by controlling the degradation of HY5, a transcription factor that activates light-regulated gene expression (Osterlund et al. 2000).

The analogy with COP1 does not stop at the RING finger motif or negative regulation of gene transcription. The HOS1 and COP1 RING finger proteins both exhibit nucleo-cytoplasmic partitioning in response to environmental stimuli. COP1 is localized in the nucleus in the dark and translocates to the cytoplasm in response to light signals (von Arnim and Deng 1994). HOS1 accumulates in the nucleus in response to cold environments (Fig. 9). To our knowledge, HOS1 represents the first example of cold-induced nucleo-cytoplasmic partitioning of proteins. Our findings provide a connection be-

tween cold-triggered cytoplasmic signaling and nuclear gene transcription. Two histidine kinases and a response regulator were recently shown to contribute to cold perception and signaling in the cyanobacterium *Synechocystis* (Suzuki et al. 2000). It is known that the second-messenger calcium and protein phosphorylation are involved in early events of cold signaling in plant cells (Monroy and Dhindsa 1995; Jonak et al. 1996; Knight et al. 1996; Monroy et al. 1998). How these cytoplasmic events relay the cold signal to the nucleus, where cold-induced gene transcription occurs, is unknown. Perhaps the nucleo-cytoplasmic movement of HOS1 contributes to the communication between the cytoplasm and the nucleus under cold conditions.

It is a matter of speculation as to the mechanism of the nuclear accumulation of HOS1 in response to cold treatment. There is a putative nuclear localization sequence in HOS1 (Fig. 5). Whether this sequence is required or sufficient for HOS1 accumulation in the nucleus remains to be determined. It is possible that the nuclear translocation of HOS1 is triggered by phosphorylation changes in this protein. Alternatively, HOS1 may be co-translocated to the nucleus by binding to another protein that is destined to accumulate in the nuclear compartment in response to cold treatments.

It is interesting that *HOS1* transcript abundance is transiently down-regulated by cold shock (Fig. 7). Because the genetic role of *HOS1* is to attenuate cold signaling, perhaps the transient down-regulation of *HOS1* transcript is important to allow cellular cold signals to




Figure 10. Ectopic expression of *HOS1* causes silencing of *HOS1* expression and mimics the *hos1* mutant phenotypes. Ten-day-old wild-type seedlings (WT), *hos1*, and T₂ progenies of a wild-type plant containing the 35S-*HOS1* construct (designated as line #4) were incubated at 0°C for 24 h before the bioluminescence image was taken. (A) Picture of wild-type, *hos1*, and T₂ progenies of transgenic line #4. (B) Bioluminescence image of plants in A. (C) Quantification of the luminescence intensities in B. Shown are the averages and standard deviations ($n = 15$). T₂ progenies of the #4 line were segregated into two groups, which are designated #4-high and #4-low, corresponding to those exhibiting high expression and low expression of *RD29A-LUC*, respectively. (D) Reduced *HOS1* transcript abundance in the #4 cosuppression plants, as determined by quantitative RT-PCR.

be amplified. Our observation that cosuppression of *HOS1* mimics *hos1* mutant phenotypes (Fig. 10) indicates an inverse correlation between *HOS1* transcript level and the expression of cold-responsive *RD29A*. It will be of interest to determine whether natural plant variations in the extent of cold acclimation and vernalization might be correlated with their expression levels of *HOS1* or its orthologs.

Materials and methods

Plant materials and growth conditions

Arabidopsis thaliana plants of the C24 ecotype expressing the *RD29A* promoter-*Luciferase* (*RD29A-LUC*) transgene (referred as wild type herein) were obtained by *Agrobacterium* transformation (Ishitani et al. 1997). The methodology of mutant screening and characterization were described previously (Ishitani et al. 1997, 1998). For bioluminescence image analysis, wild-type and *hos1* mutant seeds were surface-sterilized and sown onto Petri plates containing Murashige and Skoog salts (MS salts) supplemented with 3% sucrose and solidified with 0.6% agar. Seedlings were grown at 23°C with cycles of 16 h light and 8 h darkness before stress treatments. For cold treatment, 1-wk-old seedlings in agar plate were incubated at 0°C ($\pm 0.1^\circ\text{C}$) for 48 h in the dark. For ABA treatment, 100 μM ABA (mixed isomers) in water was sprayed on leaves, and the plants were incubated at room temperature ($\sim 23^\circ\text{C}$) under light. For NaCl or PEG treatment, seedlings were placed on 3M filter paper saturated with MS salt solution supplemented with 300 mM NaCl or 30% PEG6000 and incubated under light at room temperature. After indicated times of incubation, bioluminescence images were taken as described (Ishitani et al. 1997).

For measuring flowering time in *hos1* and the wild-type, plants were grown in pots in a growth chamber under the conditions as described above. At the emergence of the first flower, the total number of rosette leaves was counted ($n = 15$).

Positional cloning

To generate the mapping population for *HOS1*, *hos1* mutant plants were crossed to wild-type *Arabidopsis* plants of the Columbia ecotype. A total of 586 *hos1* mutant plants were selected from the segregating F₂ population based on high luminescence expression of *RD29A-LUC* after cold treatment. Genomic DNA from these plants was extracted and analyzed for cosegregation with respect to SLP (simple sequence length polymorphism) markers. These markers were developed by searching for simple repeat sequences with the RepeatMasker program (<http://ftp.genome.washington.edu/cgi-bin/RepeatMasker>). Primer pairs covering the repeat regions were designed and used for PCR amplification of the genomic fragments from both C24 and Col ecotypes. Those that exhibited PCR fragment length polymorphisms were used as markers for mapping. Primer pairs for fine mapping of *HOS1* are as follows: F12L6-F, 5'-CACGCTTATTCTTAGCCTAG-3'; F12L6-R, 5'-TTGAAAGATCACGC CGGCGA-3'; F17A14-F, 5'-GGAGAATGAAAGACCGAATC-3'; F17A14-R, 5'-TGTAGAGACATGAGAATCT-3'; T28M21-F, 5'-TAGGGGGATATTCAAGTTTG-3'; T28M21-R, 5'-AATTTCCCCACTCCACTCCT-3'; T2P4-F, 5'-GTATTTCGATCTTCTTTAGGT-3'; T2P4-R, 5'-AGGAGTATACGTCCAATTGA-3'; T20B5-F, 5'-GAAGAGCAACGCATTACAGT-3'; T20B5-R, 5'-CTTCATAAATAACGAGGCCT-3'. In addition, nucleotide differences between C24 and Col ecotypes were also identified by direct sequencing of the open reading frames of *T517.2*, *T517.4*, and *T517.14*, and were used for fine mapping.

RNA analysis

Seedlings grown for 15 d on MS agar plates under continuous light were treated with low temperature, ABA, NaCl, or PEG as described above. Total RNA from control or stressed plants was extracted and analyzed as described by Liu and Zhu (1997). The *RD29A* gene-specific probe was from the 3' noncoding region (Liu and Zhu 1997). The *CBF2* and *CBF3* gene-specific probes were generated by PCR with the following primer pairs: CBF2-F, 5'-TTCGATTTTTATTCCATTTTTGG-3'; CBF2-R, 5'-CCAACGTCCTTGAGTCTTGAT-3'; CBF3-F, 5'-TAAACTCA GATTATTATTTCCATTT-3'; CBF3-R, 5'-GAGGAGCCACG TAGAGGGCC-3'. The *FLC* gene-specific probe was obtained by PCR with the following primer pair: FLC-F, 5'-CCGCTC GAGCTTAGTATCTCCGGCG-3'; FLC-R, 5'-GGACTAGTC GCCCTTATCAGCGGA-3'. The β -*tubulin* gene was amplified by PCR with the following primer pair: Tubulin-5', 5'CGTG GATCACAGCAATACAGAGCC-3'; Tubulin-3', 5'-CCTCCT GCACCTCCACTTCGTCTTC-3'. It was used for the loading control.

hos1 mutant complementation

For complementation of *hos1* mutant plants, T517 BAC (bacterial artificial chromosome) DNA was digested with *Xho*I, and the fragment corresponding to *HOS1* was cloned into the binary vector pCIT20 (Ma 1992). This and all other constructs described in this paper were completely sequenced to ensure that they did not contain PCR errors. The construct was transformed into *Agrobacterium* strain GV3101 by electroporation. *hos1* mutant plants were vacuum-infiltrated with the *Agrobacterium* containing the *HOS1* complementation construct. Hygromycin-resistant transgenic plants were selected, and their T₂ and T₃ progenies were subjected to luminescence imaging analysis for *RD29A-LUC* expression in response to cold, ABA, and NaCl treatments as described above.

Analysis of *HOS1* promoter-GUS expression

The promoter region (2.78 kb upstream from the initiation codon) of the *HOS1* gene was PCR-amplified with the following primer pair: *HOS1-GUS-F1*, 5'-CGCAAGCTTCTCGAGGTAA GACCGTAGATTCA-3'; *HOS1-GUS-R1*, 5'-CGGTCTAGAC CTAAGACCTGAGCTAAACTAAT-3'. The resulting fragment was digested with *Hind*III and *Xba*I and inserted into the pIG121 binary vector. This *HOS1* promoter-GUS construct was introduced into *Agrobacterium* strain GV3101 and transformed into wild-type *Arabidopsis*, and 20 independent lines of hygromycin-resistant transgenic plants were obtained. For GUS staining, T₂ seedlings grown on MS agar plate were incubated with X-Gluc for 12 h at 37°C and then washed 5 times with 70% (v/v) ethanol at 70°C to remove chlorophyll.

Expression and localization of *HOS1-GFP* fusion protein

The coding region of the *HOS1* gene was obtained by RT-PCR using the following primers: *HOS1BAMHF*, 5'-CGCGGATC CATGGGTTTTGCAGAGCTTAGAGTTAA-3'; *HOS1BAMHR*, 5'-CGCGGATCCGCGTTCTTGCTGCGAATCTACGTCTCC-3'. The resulting PCR fragment was digested with *Bam*HI and inserted into the binary vector pBIN35S-mGFP4 downstream from the CaMV 35S promoter. *Agrobacterium* strain GV3101 containing this *HOS1-GFP* translational fusion was introduced into *Arabidopsis* (Col ecotype), 40 lines of kanamycin-resistant transgenic plants were selected, and their T₂ were analyzed for GFP expression. To visualize the nucleus, root tissues were

stained with propidium iodide (1 μ g/mL). Green fluorescence (GFP expression) and red fluorescence (propidium iodide staining) analysis of transgenic plants were performed with a 1024 confocal laser-scanning microscope (Grebek et al. 1997).

HOS1 cosuppression and analysis

HOS1 cDNA was isolated by RT-PCR using primers *HOS1RTF1*, 5'-GATCCCGGGGCTCAGGTCTTAGGATTTCTAG-3'; *HOS1RTR1*, 5'-GCTCCCGGGTACCTCTCTCTCTTTCT CTATGC-3'. The PCR fragment was cloned into the pCR-BluntII-TOPO vector (Invitrogen) and sequenced. The coding sequence of the *HOS1* gene was PCR-amplified from the cDNA with primers *HOS1OVER-F1*, 5'-CGAGCTCGGCTCAGGTCT TAGGATTTCTAG-3'; *HOS1OVER-R1*, 5'-CCCCGGGGGT ACCTCTCTCTCTTTCTCTATGC-3'. The PCR fragment was digested with *Sst*I and *Sma*I and inserted into a pRTL104 vector downstream from the *CaMV* 35S promoter. The *HOS1* promoter-GUS fusion was removed from the construct by digestion with *Sph*I and cloned into pCAMBIA 1200. After transformation of wild-type *Arabidopsis* plants via *Agrobacterium* strain GV3101, 25 lines of hygromycin-resistant transgenic plants were obtained. Luciferase imaging was carried out on the T₂ plants. For analysis of the transcription level of *HOS1* in cosuppression lines, the following primer pair was used in quantitative RT-PCR: 5'-ACGGAATGGATGGAGATACT-3'; 5'-AAGAGACTACCCACAGCTGA-3' (resulting in a 170-bp PCR fragment). Total RNA from wild-type, *hos1*, and #4-high plants was extracted using the TRIzol (Gibco BRL) reagent system, and 1 μ g of RNA from each sample was used for RT-PCR reaction with the EZ *rTth* RNA PCR Kit (PERKIN ELMER, part no. N808-0179) according to the manufacturer's instructions. PCR products were then separated in a 4% agarose gel. As an internal control, the *ubiquitin* gene was amplified along with *HOS1* using the following primer pair: *UBQ11-F*, 5'-GAGGTATGCA GATCTTCGTG-3'; *UBQ11-R*, 5'-GTAGACTCCTTCTGGAT GTTG-3'.

Acknowledgments

We sincerely thank Cheol-Soo Kim, Joon-Hyun Park, and Suk-Whan Hong for assistance, and Sean Cutler, David Ehrhardt, and Chris Somerville for their generous gift of the Q1 GFP transgenic plants. This work was supported by National Science Foundation grant IBN-9808398 and USDA NRI grant 2000-00664.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Anderson, J.V., Li, Q.B., Haskell, D.W., and Guy, C.L. 1994. Structural organization of the spinach endoplasmic reticulum-luminal 70-kilodalton heat shock cognate gene and expression of 70-kilodalton heat shock genes during cold acclimation. *Plant Physiol.* **104**: 1359-1370.
- Chandler, J., Wilson, A., and Dean, C. 1996. *Arabidopsis* mutants showing an altered response to vernalization. *Plant J.* **10**: 637-644.
- Chen, H.-H., Li, P.H., and Brenner, M.L. 1983. Involvement of abscisic acid in potato cold acclimation. *Plant Physiol.*

- 71: 362–365.
- Cutler, S.R., Ehrhardt, D.W., Griffiths, J.S., and Somerville, C.R. 2000. Random GFP::cDNA fusions enable visualization of subcellular structures in cells of *Arabidopsis* at a high frequency. *Proc. Natl. Acad. Sci. USA* **97**: 3718–3723.
- Deng, X.-W., Matsui, M., Wei, N., Wagner, D., Chu, A.M., Feldmann, K.A., and Quail, P.H. 1992. *COP1*, an *Arabidopsis* regulatory gene, encodes a protein with both a zinc-binding motif and a G_β homologous domain. *Cell* **71**: 791–801.
- Freemont, P.S. 2000. Ubiquitination: RING for destruction? *Curr. Biol.* **10**: R84–R87.
- Gibson, S., Arondel, V., Iba, K., and Somerville, C. 1994. Cloning of a temperature-regulated gene encoding a chloroplast omega-3 desaturase from *Arabidopsis thaliana*. *Plant Physiol.* **106**: 1615–1621.
- Gilmour, S.J., Artus, N.N., and Thomashow, M.F. 1992. cDNA sequence analysis and expression of two cold-regulated genes of *Arabidopsis thaliana*. *Plant Mol. Biol.* **18**: 13–32.
- Gilmour, S.J., Zrka, D.G., Stockinger, E.J., Salazar, M.P., Houghton, J.M., and Thomashow, M.F. 1998. Low temperature regulation of the *Arabidopsis* CBF family of AP2 transcriptional activators as an early step in cold-induced *COR* gene expression. *Plant J.* **16**: 433–442.
- Grebenok, R.J., Pierson, E., Lambert, G.M., Gong, F.C., Afonso, C.L., Haldeman-Cahill, R., Carrington, J.C., and Galbraith, D.W. 1997. Green-fluorescent protein fusions for efficient characterization of nuclear targeting. *Plant J.* **11**: 573–586.
- Guy, C.L. 1990. Cold acclimation and freezing stress tolerance: Role of protein metabolism. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* **41**: 187–223.
- Guy, C.L., Niemi, K.L., and Brambl, R. 1985. Altered gene expression during cold acclimation of spinach. *Proc. Natl. Acad. Sci. USA* **82**: 3673–3677.
- Hilleren, P. and Parker, R. 1999. Mechanisms of mRNA surveillance in eukaryotes. *Annu. Rev. Genet.* **33**: 229–260.
- Ishitani, M., Xiong, L., Stevenson, B., and Zhu, J.-K. 1997. Genetic analysis of osmotic and cold stress signal transduction in *Arabidopsis*: Interactions and convergence of abscisic acid-dependent and abscisic acid-independent pathways. *Plant Cell* **9**: 1935–1949.
- Ishitani, M., Xiong, L., Lee, H., Stevenson, B., and Zhu, J.-K. 1998. *HOS1*, a genetic locus involved in cold responsive gene expression in *Arabidopsis thaliana*. *Plant Cell* **10**: 1151–1161.
- Jackson, P.K., Eldridge, A.G., Freed, E., Furstenthal, L., Hsu, J.Y., Kaiser, B.K., and Reimann, J.D.R. 2000. The lore of the RINGs: Substrate recognition and catalysis by ubiquitin ligases. *Trends Cell Biol.* **10**: 429–439.
- Jaglo-Ottosen, K.R., Gilmour, S.J., Zarka, D.G., Schabenberger, O., and Thomashow, M.F. 1998. *Arabidopsis CBF1* overexpression induces *COR* genes and enhances freezing tolerance. *Science* **280**: 104–106.
- Joazeiro, C.A.P. and Weissman, A.M. 2000. RING finger proteins: Mediators of ubiquitin ligase activity. *Cell* **102**: 549–552.
- Jonak, G., Kiegerl, S., Lighterink, W., Barker, P.J., and Huskisson, N.S. 1996. Stress signaling in plants: A mitogen-activated protein kinase pathway is activated by cold and drought. *Proc. Natl. Acad. Sci. USA* **93**: 11274–11279.
- Kasuga, M., Liu, Q., Miura, S., Yamaguchi-Shinozaki, K., and Shinozaki, K. 1999. Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nature Biotech.* **17**: 287–291.
- Knight, H., Trewas, A.J., and Knight, M.R. 1996. Cold calcium signaling in *Arabidopsis* involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell* **8**: 489–503.
- Knight, H., Veale, E.L., Warren, G.J., and Knight, M.R. 1999. The *sfr6* mutation in *Arabidopsis* suppresses low temperature induction of genes dependent on the CRT/DRE sequence motif. *Plant Cell* **11**: 875–886.
- Kurkela, S. and Borg-Franck, M. 1992. Structure and expression of *kin2*, one of two cold- and ABA-induced genes of *Arabidopsis thaliana*. *Plant Mol. Biol.* **19**: 689–692.
- Lee, H., Xiong, L., Ishitani, M., and Zhu, J.-K. 1999. Cold-regulated gene expression and freezing tolerance in an *Arabidopsis thaliana* mutant. *Plant J.* **17**: 301–308.
- Levitt, J. 1980. *Responses of plants to environmental stress*. Vol. 1: *Chilling, freezing, and high temperature stress*. Academic Press, New York.
- Liu, J. and Zhu, J.-K. 1997. Proline accumulation and salt-induced gene expression in a salt-hypersensitive mutant of *Arabidopsis*. *Plant Physiol.* **114**: 591–596.
- . 1998. A calcium sensor homolog required for plant salt tolerance. *Science* **280**: 1943–1945.
- Liu, Q., Kasuga, M., Sakuma, Y., Abe, H., Miura, S., Yamaguchi-Shinozaki, K., and Shinozaki, K. 1998. Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low temperature-responsive gene expression, respectively, in *Arabidopsis*. *Plant Cell* **10**: 1391–1406.
- Ma, H., Yanofsky, M.F., Klee, H.J., Bowman, J.L., and Meyerowitz, E.M. 1992. Vectors for plant transformation and cosmid libraries. *Gene* **117**: 161–167.
- Medina, J., Bargues, M., Terol, J., Perez-Alonso, M., and Salinas, J. 1999. The *Arabidopsis* CBF gene family is composed of three genes encoding AP2 domain-containing proteins whose expression is regulated by low temperature but not by abscisic acid or dehydration. *Plant Physiol.* **119**: 463–470.
- Michaels, S.D. and Amasino, R.M. 1999. *FLOWERING LOCUS C* encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell* **11**: 949–956.
- Miller, L.K. 1999. An exegesis of IAPs: Salvation and surprises from BIR motifs. *Trends Cell Biol.* **9**: 323–328.
- Monroy, A.F. and Dhindsa, R.S. 1995. Low-temperature signal transduction: Induction of cold acclimation-specific genes of alfalfa by calcium at 25°C. *Plant Cell* **7**: 321–331.
- Monroy, A.F., Sangwan, V., and Dhindsa, R.S. 1998. Low temperature signal transduction during cold acclimation: Protein phosphatase 2A as an early target for cold-inactivation. *Plant J.* **13**: 653–660.
- Murata, N. and Los D.A. 1997. Membrane fluidity and temperature perception. *Plant Physiol.* **115**: 875–879.
- Nordin, K., Vahala, T., and Palva, E.T. 1993. Differential expression of two related, low-temperature-induced genes in *Arabidopsis thaliana* (L.) Heynh. *Plant Mol. Biol.* **21**: 641–653.
- Osterlund, M., Hardtke, C., Wei, N., and Deng, X.-W. 2000. Targeted destabilization of HY5 during light regulated development of *Arabidopsis*. *Nature* **405**: 462–466.
- Saurin, A.J., Borden, K.L.B., Boddy, M.N., and Freemont, P.S. 1996. Does this have a familiar RING? *Trends Biochem. Sci.* **21**: 208–214.
- Sheldon, C.C., Rouse, D.T., Finnegan, E.J., Peacock, W.J., and Dennis, E.S. 2000. The molecular basis of vernalization: The central role of *FLOWERING LOCUS C* (*FLC*). *Proc. Natl. Acad. Sci. USA* **97**: 3753–3758.
- Steponkus, P.L., Uemura, M., Joseph, R.A., Gilmour, S.J., and Thomashow, M.F. 1998. Mode of action of the *COR15a* gene on the freezing tolerance of *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **95**: 14570–14575.
- Stockinger, E.J., Gilmour, S.J., and Thomashow, M.F. 1997. *Ar-*

- bidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a *cis*-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc. Natl. Acad. Sci. USA* **94**: 1035–1040.
- Suzuki, I., Los, D.A., Kanesaki, Y., Mikami, K., and Murata, N. 2000. The pathway for perception and transduction of low-temperature signals in *Synechocystis*. *EMBO J.* **19**: 1327–1334.
- Thomashow, M.F. 1994. *Arabidopsis thaliana* as a model for studying mechanisms of plant cold tolerance. In *Arabidopsis* (eds. E.M. Meyerowitz and C.R. Somerville), pp. 807–834. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Thomashow, M.F. 1999. Plant cold acclimation: Freezing tolerance genes and regulatory mechanisms. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **50**: 571–599.
- von Arnim, A.G. and Deng, X.-W. 1994. Light inactivation of *Arabidopsis* photomorphogenic COP1 involves a cell-specific regulation of its nucleo-cytoplasmic partitioning. *Cell* **79**: 1035–1045.
- Xin, Z. and Browse, J. 1998. *eskimo1* mutants of *Arabidopsis* are constitutively freezing-tolerant. *Proc. Natl. Acad. Sci. USA* **95**: 7799–7804.
- Yamaguchi-Shinozaki, K. and Shinozaki, K. 1994. A novel *cis*-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *Plant Cell* **6**: 251–264.
- Yang, Y., Fang, S., Jensen, J.P., Weissman, A.M., and Ashwell, J.D. 2000. Ubiquitin protein ligase activity of IAPs and their degradation in proteasomes in response to apoptotic stimuli. *Science* **288**: 874–877.