

# Acetogens and Acetoclastic *Methanosarcinales* Govern Methane Formation in Abandoned Coal Mines<sup>∇</sup>

Sabrina Beckmann,<sup>1</sup> Tillmann Lueders,<sup>2</sup> Martin Krüger,<sup>3</sup> Frederick von Netzer,<sup>2</sup>  
Bert Engelen,<sup>1</sup> and Heribert Cypionka<sup>1\*</sup>

*Institute for Chemistry and Biology of the Marine Environment (ICBM), University of Oldenburg, Carl-von-Ossietzky Straße 9-11, D-26129 Oldenburg,<sup>1</sup> Institute of Groundwater Ecology, Helmholtz Zentrum München, Ingolstädter Landstrasse 1, D-85764 Neuherberg,<sup>2</sup> and Federal Institute for Geosciences and Natural Resources (BGR), Stilleweg 2, D-30655 Hannover,<sup>3</sup> Germany*

Received 2 December 2010/Accepted 23 March 2011

**In abandoned coal mines, methanogenic archaea are responsible for the production of substantial amounts of methane. The present study aimed to directly unravel the active methanogens mediating methane release as well as active bacteria potentially involved in the trophic network. Therefore, the stable-isotope-labeled precursors of methane, [<sup>13</sup>C]acetate and H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub>, were fed to liquid cultures from hard coal and mine timber from a coal mine in Germany. Guided by methane production rates, samples for DNA stable-isotope probing (SIP) with subsequent quantitative PCR and denaturing gradient gel electrophoretic (DGGE) analyses were taken over 6 months. Surprisingly, the formation of [<sup>13</sup>C]methane was linked to acetoclastic methanogenesis in both the [<sup>13</sup>C]acetate- and the H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub>-amended cultures of coal and timber. H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub> was used mainly by acetogens related to *Pelobacter acetylenicus* and *Clostridium* species. Active methanogens, closely affiliated with *Methanosarcina barkeri*, utilized the readily available acetate rather than the thermodynamically more favorable hydrogen. Thus, the methanogenic microbial community appears to be highly adapted to the low-H<sub>2</sub> conditions found in coal mines.**

Worldwide, mine gas emissions from active and abandoned coal mines release substantial amounts of methane, contributing as much as 7% of global methane formation (4). Mine gas is a hazard but is also a potential source of methane for the industry. Stable carbon and hydrogen isotopic signatures indicate that methane in mine gas has mixed thermogenic and biogenic origins (31, 33). In abandoned coal mines, thermogenic methane is a remainder of geological processes, but biogenic formation of methane is still going on (14). Besides hard coal, possible sources for methane are large amounts of mine timber used for the construction of mines and left behind after the cessation of mining.

Generally, methane is produced from either acetate, hydrogen, or methylotrophic substrates as precursors. Recently, we showed that methylotrophic methanogenesis does not have a quantitative impact on *in situ* processes (2). However, while hydrogen is energetically favorable, acetate is the quantitatively more available substrate (38). In previous studies we have revealed that acetate is an important intermediate of the degradation processes and the main precursor of the biogenic methane in abandoned coal mines (14). We have also shown the presence of *Methanosarcina* spp. as the dominating archaea (2). *Methanosarcina* spp. are able to use acetate as well as H<sub>2</sub>-CO<sub>2</sub>. While other investigators reported that methanogenesis in coal mines is driven mainly by H<sub>2</sub>-utilizing archaea (7, 28), our studies indicated that acetoclastic methanogenesis

seems to be the main methanogenic process, at least in the abandoned coal mines we have investigated (2).

The activity of methanogens in different habitats can be studied by using stable-isotope probing (SIP) (16, 17). DNA-SIP allows the identification of specifically active members of microbial populations based on the incorporation of <sup>13</sup>C into the DNA of cells consuming labeled substrates (18). In this technique, labeled DNA is resolved after incubation under label addition by subsequent isopycnic gradient ultracentrifugation (22). Hence, guilds of methanogens that utilize <sup>13</sup>C-labeled methanogenic substrates such as [<sup>13</sup>C]acetate or H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub> can be recovered in <sup>13</sup>C-enriched DNA. However, both methanogenic precursors can also be utilized by syntrophic acetate-oxidizing and/or homoacetogenic bacteria, respectively (27, 32). Until now, only a few SIP studies have been performed in coal habitats. Han et al. (10) investigated the active methanotrophic community in a Chinese coal deposit. However, no studies have revealed the activity of microorganisms directly involved in methane production in abandoned coal mines. In our present study, we wanted to identify active methanogens and to test the hypothesis that acetate is the main precursor of methane, even if hydrogen is available as an energetically more favorable electron donor.

## MATERIALS AND METHODS

**Sample collection and enrichment cultures.** Samples were collected in May 2007 in sealed compartments of coal mines closed in the 1960s. The mines harbor hard coal ("rock coal") belonging to the fat coals according to the German classification (reference 33 and references therein). Briefly, large pieces of coal (2 to 15 cm in diameter) and mine timber (2 to 10 cm in diameter) were collected aseptically in glass bottles that were immediately flushed with N<sub>2</sub> and stored at 4°C until further processing. *In situ* temperatures were 35 to 36°C, with 100% air humidity. Coal and mine timber samples were processed in an anaerobic chamber under a nitrogen atmosphere to prevent oxidation. Samples were homoge-

\* Corresponding author. Mailing address: Institute for Chemistry and Biology of the Marine Environment (ICBM), University of Oldenburg, Carl-von-Ossietzky Straße, 9-11, D-26129 Oldenburg, Germany. Phone: 49 441-798 5360. Fax: 49 441-798 3404. E-mail: cypionka@icbm.de.

<sup>∇</sup> Published ahead of print on 1 April 2011.

nized and distributed into Hungate tubes containing 500 ml of sulfate-free mineral medium (36) with a salinity of 15 practical salinity units (PSU), according to *in situ* values. Controls were supplemented with 10 mM 2-bromoethanesulfonate (BES) to exclude abiotic degassing by inhibiting methanogenesis. Enrichment cultures were amended with 10 mM fully  $^{13}\text{C}$ -labeled acetate (Campro Scientific) or  $^{13}\text{C}$ -bicarbonate (Campro Scientific) plus  $\text{H}_2$ . All incubated samples were frozen at  $-80^\circ\text{C}$  after an incubation time of 6 months until further processing for DNA-SIP. All incubations were carried out at least in triplicate. Subsamples of all incubated samples were taken at the beginning and at month 3 of incubation. The increases in the amounts of methane and hydrogen in the headspace, as well as the stable isotopes of methane, were monitored continuously over 6 months and were analyzed by gas chromatography-mass spectrometry (GC-MS) as described previously by Krüger et al. (13). Concentrations of acetate were analyzed by high-performance liquid chromatography (Agilent Technologies) using a Zorbax Eclipse Plus  $\text{C}_8$  USP L7 column (Agilent Technologies) at  $60^\circ\text{C}$ . The eluent was a 5 mM  $\text{H}_2\text{SO}_4$ -methanol gradient at 1 ml/min. Acetate was detected by a diode array detector (DAD; Agilent Technologies).

**DNA extraction, isopycnic centrifugation, and gradient fractionation.** DNA was extracted from 0.5 g of incubation slurry after 0, 3, and 6 months using phenol-chloroform extraction as described by Lueders et al. (18). Three parallel extractions were carried out, and extracts were pooled for each incubation treatment. DNA was checked by standard agarose gel electrophoresis and was quantified using PicoGreen staining according to the method of Wilms et al. (37).

Gradient preparation, isopycnic centrifugation, and gradient fractionation were performed as described by Lueders et al. (17) with minor modifications. Each gradient consisted of 6.3 ml of  $\text{CsCl}$  (approximately  $1.72\text{ g ml}^{-1}$ ; Calbiochem) and ca. 1 ml of gradient buffer (100 mM Tris-HCl [pH 8.0]  $\text{liter}^{-1}$ , 100 mM KCl  $\text{liter}^{-1}$ , 1 mM EDTA  $\text{liter}^{-1}$ ) including 2,000 ng of DNA. Prior to centrifugation, the average density of the centrifugation medium was controlled refractometrically and was adjusted to  $1.84\text{ g cm}^{-3}$ . The samples were centrifuged in 6.3-ml Quick-Seal Polyallomer tubes (Beckman) in an NVT 65 near-vertical rotor (Beckman) using an LE-70 ultracentrifuge (Beckman Instruments). Centrifugation was performed at  $20^\circ\text{C}$  for 36 h at 44,500 rpm ( $184,000 \times g$ ). Gradients were fractionated as described previously by Neufeld et al. (21). Briefly, the gradients were fractionated from bottom to top into 12 equal fractions (400  $\mu\text{l}$ ). A precisely controlled flow rate was achieved by displacing the gradient medium with water at the top of the tube using a Graseby 3100 syringe pump at a flow rate of  $1\text{ ml min}^{-1}$ . The density of each collected fraction (a small aliquot of 100  $\mu\text{l}$ ) was measured by determining the refractive index using a digital refractometer (model AR-20; Reichert Analytical Instruments, Depew, NY). Subsequently, the DNA was precipitated using polyethylene glycol 6000 (Aldrich Chemistry). The DNA pellet was washed once with 70% ethanol and was dissolved in 25  $\mu\text{l}$  of elution buffer.

**Quantification of archaeal and bacterial 16S rRNA genes in density gradient fractions.** DNA was precipitated from gradient fractions and was quantified fluorometrically and by quantitative PCR (qPCR) using the Ar109f/Ar912rt and Ba27F/907R primer systems described by Lueders et al. (17). The qPCR mixtures contained 12.5  $\mu\text{l}$  of the premix solution of a DyNAmo HS SYBR Green qPCR kit (New England Biolabs, Inc., Hitchin, United Kingdom), 1  $\mu\text{l}$  of each primer, and 10  $\mu\text{l}$  of the standard or DNA extract as a template in a final reaction mixture volume of 25  $\mu\text{l}$ . The PCR was carried out in a Rotor-Gene 3000 cyclor (Corbett Research, Sydney, Australia). After initial denaturation at  $95^\circ\text{C}$  for 15 min, 50 cycles followed. Each cycle consisted of denaturation for 30 s at  $94^\circ\text{C}$ , annealing for 20 s at  $52^\circ\text{C}$ , elongation for 30 s at  $70^\circ\text{C}$ , and fluorescence measurement at  $70^\circ\text{C}$ . To check the amplification specificity, fluorescence was also measured at the end of each cycle for 20 s at  $80^\circ\text{C}$ . After the last cycle, a melting curve was recorded by increasing the temperature from  $50^\circ\text{C}$  to  $99^\circ\text{C}$  ( $1^\circ\text{C}$  every 10 s). The numbers of bacterial and archaeal 16S rRNA gene targets were calculated from the DNA concentrations according to the work of Süß et al. (30). DNA standards for quantitative (real-time) PCR were prepared as described by Wilms et al. (37) and Engelen et al. (6). Bacterial and archaeal targets were measured in at least three different dilutions of DNA extracts (1:10 to 1:1,000) and in triplicate.

**PCR and DGGE analysis.** For denaturing gradient gel electrophoresis (DGGE), an 803-bp fragment of the archaeal 16S rRNA gene was amplified by using primers Ar109f (5'-AC KGC TCA GTA ACA CGT-3') and Ar912rt (5'-GTG CTC CCC CGC CAA TTC CTT TA-3'). For the analysis of bacterial composition, primers BA27F (5'-AGA GTT TGA TCM TGG CTC AG-3') and 907R (5'-CCG TCA ATT CCT TTG AGT TT-3') were used to amplify an 880-bp fragment of the bacterial 16S rRNA gene. At the 5' end of each forward primer, an additional 40-nucleotide GC-rich sequence (GC-clamp) was added to achieve a stable melting point for the DNA fragments in the DGGE according to the method of Muyzer et al. (20). PCR amplification was performed using an

Eppendorf thermal cycler system (Mastercycler; Eppendorf, Hamburg, Germany) as follows: 2  $\mu\text{l}$  (1 to 100 ng) of template DNA, 1 U of *Taq* DNA polymerase, the manufacturer's recommended buffer supplied with the polymerase enzyme, 10 mM deoxynucleoside triphosphates (dNTPs), 50  $\mu\text{M}$  (each) appropriate primers, and 10 mM bovine serum albumin (BSA) were adjusted to a total volume of 50  $\mu\text{l}$  with PCR-grade water (Ampuwa; Fresenius, Bad Homburg, Germany). The PCR program for archaeal and bacterial DNA included an initial denaturation step for 5 min at  $94^\circ\text{C}$ , followed by 30 cycles of 30 s at  $94^\circ\text{C}$ , 30 s at  $52^\circ\text{C}$ , and 1 min at  $72^\circ\text{C}$ . Primer extension was carried out for 5 min at  $72^\circ\text{C}$ . Aliquots (5  $\mu\text{l}$ ) of the PCR products were analyzed by agarose gel electrophoresis in 1.5% (wt/vol) agarose gels and ethidium bromide (0.8 ng  $\text{ml}^{-1}$ ) staining for 20 min on a UV transilluminator as described previously (37).

DGGE was performed using an INGENYphorU-2 system (Ingeny, Goes, Netherlands). PCR products and loading buffer (40% [wt/vol] glycerol, 60% [wt/vol]  $1\times$  Tris-acetate-EDTA [TAE], and bromophenol blue) were included in a 1:2 ratio. The PCR amplicons were applied directly to 6% (wt/vol) polyacrylamide gels with a linear gradient of 30 to 80% denaturant for archaeal PCR products and 50 to 70% denaturant for bacterial PCR products (100% denaturant corresponds to 7 M urea and 40% [vol/vol] formamide). Electrophoresis was carried out in  $1\times$  TAE buffer (40 mM Tris-acetate [pH 7.4], 20 mM sodium acetate, 1 mM  $\text{Na}_2\text{EDTA}$ ) at a constant voltage of 100 V and a temperature of  $60^\circ\text{C}$  for 20 h. After electrophoresis, the gels were stained for 2 h in  $1\times$  SYBR Gold solution (Molecular Probes, Eugene, OR) in  $1\times$  TAE and were washed for 20 min with distilled water. The gel was digitized using a digital imaging system (BioDocAnalyze; Biometra, Göttingen, Germany) with UV transillumination (302 nm).

**Reamplification and sequencing of DGGE bands.** DGGE bands were excised for sequencing using a sterile scalpel and were treated as described by Del Panno et al. (3). Briefly, the bands were transferred into 50  $\mu\text{l}$  of PCR-grade water and were incubated for 48 h at  $4^\circ\text{C}$ . For reamplification, 2  $\mu\text{l}$  of the supernatant was taken as a template using the reaction mixture described above with a final volume of 50  $\mu\text{l}$ . The PCR protocol adjusted for the reamplification was the same as that described above, with the following minor modifications: the total number of cycles was 25, and the final elongation was carried out at  $72^\circ\text{C}$  for 10 min. PCR products were purified by using the QIAquick PCR purification kit (Qiagen GmbH) and were eluted in 30  $\mu\text{l}$  of PCR-grade water. DNA yields were estimated fluorometrically in a microtiter plate reader (FLUOstar Optima; BMG Labtechnologies, Offenburg, Germany) using a 1:200-diluted PicoGreen reagent according to a modified manufacturer's protocol (Molecular Probes, Eugene, OR) as described in detail by Wilms et al. (37). Sequence analyses were performed by GATC Biotech AG (Konstanz, Germany). Sequences were compared to those in GenBank using the BLAST tool of the National Center for Biotechnology Information server (1).

**Nucleotide sequence accession numbers.** The sequences determined in this study have been deposited in the GenBank nucleotide sequence database under accession no. FR838951 to FR838999.

## RESULTS

For stable-isotope probing (SIP), microcosms with coal or timber samples were amended with  $^{13}\text{C}$ -acetate or  $\text{H}_2$ - $^{13}\text{CO}_2$  and were incubated under *in situ* conditions. The accumulation of  $\text{CH}_4$  was followed over a period of 6 months. To identify active community members, samples for DNA extraction were taken at the beginning of incubation and after 3 and 6 months (Fig. 1A). The day zero incubation samples served as SIP controls.

**Acetate is the precursor of methane.**  $\text{CH}_4$  production was observed in all incubated samples (Fig. 1A), with higher activities in timber than in coal enrichment cultures. The highest  $\text{CH}_4$  formation rates (0.13  $\mu\text{mol per g}$  [wet weight] and day) were detected in the  $^{13}\text{C}$ -acetate enrichment cultures between months 3 and 6 of incubation. The addition of  $\text{H}_2$ - $^{13}\text{CO}_2$  resulted in less stimulation of methanogenesis (maximum rate, 0.05  $\mu\text{mol per g}$  [wet weight] and day). The isotopic signature of  $\text{CH}_4$  indicated that methane was formed from the labeled substrates added (data not shown). No methane formation was observed in control samples incubated with the methanogen-




FIG. 1. Long-term incubation of weathered hard coal and mine timber amended with <sup>13</sup>C-labeled acetate or H<sub>2</sub>-CO<sub>2</sub>. Shown are microbial methane formation (A) and acetate (B) and hydrogen (C) depletion/formation. Measurements were carried out in 5 replicates.

esis inhibitor 2-bromoethanesulfonate (BES). Therefore, the possibility of abiotic degassing of adsorbed methane from the incubated samples can be excluded.

While H<sub>2</sub> was largely used up after 3 months, acetate was completely depleted after 6 months (Fig. 1B and C). Interestingly, acetate formation was observed in the H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub> cultures, while there was no H<sub>2</sub> formation in the samples incubated with acetate. The fact that acetate formation was also detected in the unamended incubated samples indicates the central role of acetate in the process of methanogenesis. The isotopic signature of the acetate formed showed strong labeling, indicating its formation from <sup>13</sup>CO<sub>2</sub> (data not shown).

**DNA-SIP reveals that methanogenesis is mediated by *Methanosarcina* spp.** In order to identify active community members, samples from the <sup>13</sup>C-enriched incubated samples were analyzed by SIP. Density-resolved archaeal DNA was detected in gradient fractions using archaeal qPCR (Fig. 2 and 3). The amount of archaeal DNA detected in the heavy fractions increased over the incubation time between months 3 and 6. Only the timber cultures showed <sup>13</sup>C-labeled archaeal DNA after 3 months, with maximum quantities in DNA gradients from acetate enrichment cultures. After 6 months, “light” and “heavy” DNA fractions substantiated a clear labeling of archaeal DNA in timber and coal cultures amended with [<sup>13</sup>C]acetate or <sup>13</sup>CO<sub>2</sub> (Fig. 2 and 3).

Next, the archaeal community members detected in “light” and “heavy” gradient fractions were analyzed by denaturing gradient gel electrophoresis (DGGE) and subsequent band sequencing (Tables 1 and 2). In general, the compositions of the total archaeal communities in the [<sup>13</sup>C]acetate and H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub> cultures were similar, and these communities were dominated by relatives of *Methanosarcina* spp., *Methanosaeta* spp., and uncultured *Crenarchaeota* (Tables 1 and 2). However, clear <sup>13</sup>C labeling was evident mainly for relatives of *Methanosarcina* spp. Increased band intensities reflected larger amounts of <sup>13</sup>C-labeled DNA in acetate cultures (Fig. 2 and 3). To a minor extent, labeled DNA was also observed for a DGGE band related to *Methanosaeta* spp. in the acetate enrichment cultures. Members of the *Crenarchaeota* were abundant in the original coal and timber samples but showed no incorporation of <sup>13</sup>C-labeled substrates.

**Identification of active Bacteria.** Quantities of labeled bacterial 16S rRNA genes increased over the incubation time (Fig. 2 and 3), as found for the *Archaea*. While strongly labeled bacterial DNA was detected especially in the [<sup>13</sup>C]acetate-amended coal samples after 3 months, H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub> cultures showed substantial quantities of <sup>13</sup>C-labeled DNA only after 6 months of incubation.

In the acetate-amended coal samples, bacterial DNA labeled after 3 months was affiliated with a lineage of uncultured *Geobacteraceae* and *Pelobacter* spp., and bacterial DNA labeled after 6 months was additionally affiliated with *Pseudomonas* and *Clostridium* spp. Most surprisingly, however, abundant unlabeled and labeled DNA was detected for a relative of a *Pseudomonas* sp. after 6 months only. In H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub>-amended coal samples, clearly labeled DNA was evident from gradient fractions in DGGE analysis after 6 months only and was affiliated with *Clostridium*, *Desulfovibrio*, and *Pelobacter* spp., as well as, again, with uncultured *Geobacteraceae*.

In [<sup>13</sup>C]acetate-amended timber cultures, members of che-

moautotrophic bacteria (*Hydrogenophaga* and *Hyphomicrobium* spp.) as well as sulfate and sulfur reducers (*Desulfovibrio* and *Desulfuromonas* spp.) were detected primarily in “intermediate” and “heavy” gradient fractions. Surprisingly, the DGGE band dominating the highest-density DNA gradient fractions was related to *Burkholderia* spp. This could be a methodological bias due to the high-GC DNA content of *Burkholderia* spp. In H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub>-amended timber microcosms, *Hydrogenophaga* spp. clearly dominated “heavy” DNA after 6 months, while a relative of *Desulfovibrio* spp. was detected in “intermediate” gradient fractions after 3 months.

## DISCUSSION

In the present study, we have identified active microbes responsible for methane formation in samples taken from abandoned coal mines. We demonstrated that acetate is the main precursor of methane and identified the microbes involved in the processes leading to methane formation.


Within the past few years, methane release has also been observed and studied in other coal mines worldwide. Several of the community members we have detected in our samples have been found previously in other coal mine deposits (especially *Pelobacter acetylenicus*, *Clostridium* spp., *Pseudomonas* spp., uncultured *Geobacteraceae*, and *Methanosarcina* spp.), indicating their potential role in the process of methane release (8, 12, 25, 26). A predominance of acetoclastic *Methanosarcinales* was already shown in two other coal seams investigated recently (9, 34) and in comparable habitats, i.e., hydrocarbon-contaminated aquifers (5). However, in coal mine water and drainage liquids, hydrogenotrophic methanogens were prevalent (7, 29, 35). Hydrogenotrophic methanogenesis dominated in the drainage water of coal reservoirs. In the floating systems, easily degradable substrates that lead to intermediate hydrogen formation might be released.

**Methane release via acetoclastic methanogenesis.** The fact that *Methanosarcina* spp. were responsible for methane production in our enrichment cultures is in agreement with our earlier studies showing that the *Methanosarcinales* are most abundant in *in situ* coal and timber samples (2). Although *Methanosarcina* spp. are known to use either hydrogen or acetate, those identified here seem to be strictly adapted to the conditions in this habitat. In the mines studied, acetate seems to be quantitatively more available for the *Methanosarcinales*. Hydrogen might hardly be formed at low metabolic rates and therefore might not be available for methanogens. Even after an incubation time of 6 months with an adequate supply of hydrogen, coal mine methanogens did not make direct use of hydrogen for methane production. Hydrogen appeared rather to be used by acetogens producing acetate, which then, in turn, was utilized by the *Methanosarcinales*.

**Active *Geobacteraceae* predominate in coal.** Acetotrophic members of the *Geobacteraceae* were found to be labeled in the coal enrichment cultures. Their abundance could be increased by amendment with acetate. However, they were also active in the samples incubated with hydrogen. This could be explained by secondary cross-feeding processes, since labeled acetate was formed (15). In our cultures, it is not clear which electron acceptor is used by the *Geobacteraceae* for acetate oxidation. One possibility could be the utilization of electron



**A) Hard Coal +  $^{13}\text{C}$ -Acetate**



**B) Hard Coal +  $^{13}\text{CO}_2$  +  $\text{H}_2$**





FIG. 2. Quantitative PCR distribution and DGGE community profiles of density-resolved bacterial and archaeal DNA in SIP centrifugation gradients after 3 and 6 months of incubation of hard coal with  $^{13}\text{C}$ -acetate (A) or  $\text{H}_2$ - $^{13}\text{CO}_2$  (B). Each letter on the right of a DGGE profile indicates the same band throughout the gradient. Band letters correspond to those in Table 1. The same band letter indicates the same organism.

**A) Mine Timber +  $^{13}\text{C}$ -Acetate**



**B) Mine Timber +  $^{13}\text{CO}_2$  +  $\text{H}_2$**




FIG. 3. Quantitative PCR distribution and DGGE community profiles of density-resolved bacterial and archaeal DNA in SIP centrifugation gradients after 3 and 6 months of incubation of mine timber with  $^{13}\text{C}$ -acetate (A) or  $\text{H}_2$ - $^{13}\text{CO}_2$  (B). Each letter on the right of a DGGE profile indicates the same band throughout the gradient. Band letters correspond to those in Table 2. The same band letter indicates the same organism.

TABLE 1. Bacterial and archaeal diversity in incubated hard-coal samples amended with [<sup>13</sup>C]acetate or H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub><sup>a</sup>

Labeled substrate and closest cultured relative	Similarity (%)	Band letter in Fig. 2	GenBank accession no.
<b>[<sup>13</sup>C]acetate</b>			
<i>Bacteria</i>			
<i>Burkholderia cepacia</i>	100	A	FR838981
<i>Pseudoalteromonas</i> sp.	100	B	FR838963
<i>Clostridium</i> sp.	99	C	FR838991
<i>Pseudomonas stutzeri</i>	99	D	FR838972
<i>Desulfovibrio africanus</i>	100	E	FR838976
<i>Pelobacter acetylenicus</i>	95	F	FR838952
Uncultured <i>Geobacteraceae</i>	99	G	FR838962
<i>Archaea</i>			
<i>Methanosaeta</i> sp.	98	a	FR838958
Uncultured <i>Crenarchaeota</i>	98	b	FR838970
<i>Methanosarcina barkeri</i>	99	c	FR838992
<i>Methanosarcina</i> sp.	99	d	FR838971
<b>H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub></b>			
<i>Bacteria</i>			
<i>Burkholderia cepacia</i>	100	A	FR838986
<i>Desulfovibrio alkaliphilus</i>	92	B	FR838983
<i>Clostridium</i> sp.	99	C	FR838998
<i>Desulfovibrio africanus</i>	100	D	FR838979
<i>Pelobacter acetylenicus</i>	95	E	FR838951
Uncultured <i>Geobacteraceae</i>	98	F	FR838960
<i>Xanthomonas</i> sp.	100	G	FR838954
<i>Archaea</i>			
<i>Methanosaeta</i> sp.	98	a	FR838957
Uncultured <i>Crenarchaeota</i>	99	b	FR838966
<i>Methanosarcina barkeri</i>	93	c	FR838997

<sup>a</sup> Detected by denaturing gradient gel electrophoresis of density-resolved DNA gradient fractions. The phylogenetic affiliations of the 16S rRNA genes from microbes that incorporated <sup>13</sup>C-labeled substrates are shaded.

acceptors, such as sulfur, directly from the coal, since members of the family *Geobacteraceae* are reported to be elemental-sulfur reducers (11). Jones et al. (12) also obtained high numbers of *Geobacter* species from coal, but none of the known electron acceptors was present, suggesting that *Geobacter* might be capable of coupling the degradation of organics to an electron- or H<sub>2</sub>-accepting partner. This could also be proposed in our case. However, in contrast to our strictly anoxic enrichment cultures, coal mines showed low concentrations of oxygen in layers of coal near the surface, and *Geobacteraceae* constituted the bulk of the overall bacterial community in the original coal samples (F. Gründler, unpublished data).

**Coal and timber: two substrates, two different active bacterial communities.** The active bacterial communities in coal and timber differ. Amendment with acetate or H<sub>2</sub>-CO<sub>2</sub> did not have a strong effect on the composition of the active community. Besides the *Geobacteraceae*, the active community in the coal samples comprised *Pelobacter acetylenicus* and members of *Clostridium* and *Pseudomonas* species. In earlier studies, *Pseudomonas stutzeri* was isolated from coal samples; this species is potentially able to utilize polycyclic aromatic hydrocarbons (PAHs) (24). This suggestion can be supported by the fact that the first event of coal fragmentation is exoenzymatic hydrolysis into small PAHs (29). Moreover, *Pseudomonas stutzeri* is even more active when acetate is present as a second electron donor (19), and that could be one reason for its predom-

TABLE 2. Bacterial and archaeal diversity in incubated mine timber samples amended with [<sup>13</sup>C]acetate and H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub><sup>a</sup>

Labeled substrate and closest cultured relative	Similarity (%)	Band letter in Fig. 3	GenBank accession no.
<b>[<sup>13</sup>C]acetate</b>			
<i>Bacteria</i>			
<i>Acholeplasma</i> sp.	95	A	FR838980
Uncultured <i>Bacteroidetes</i>	99	B	FR838984
<i>Clostridium</i> sp.	99	C	FR838974
<i>Burkholderia cepacia</i>	100	D	FR838994
<i>Hyphomicrobium</i> sp.	99	E	FR838953
<i>Hydrogenophaga</i> sp.	94	F	FR838955
<i>Desulfovibrio</i> sp.	95	G	FR838961
<i>Desulfovibrio africanus</i>	100	H	FR838982
<i>Halochromatium</i> sp.	92	I	FR838989
<i>Desulfuromonas acetexigens</i>	95	J	FR838965
Uncultured <i>Geobacteraceae</i>	96	K	FR838964
<i>Archaea</i>			
Uncultured <i>Crenarchaeota</i>	98	a	FR838968
Uncultured <i>Crenarchaeota</i>	98	b	FR838969
<i>Methanosaeta</i> sp.	98	c	FR838959
<i>Methanosarcina barkeri</i>	99	d	FR838995
<i>Methanosarcina barkeri</i>	93	e	FR838996
<b>H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub></b>			
<i>Bacteria</i>			
<i>Acholeplasma</i> sp.	95	A	FR838990
Uncultured <i>Bacteroidetes</i>	99	B	FR838988
<i>Clostridium</i> sp.	99	C	FR838975
<i>Burkholderia cepacia</i>	99	D	FR838987
<i>Hyphomicrobium</i> sp.	99	E	FR838956
<i>Hydrogenophaga</i> sp.	94	F	FR838985
<i>Desulfovibrio</i> sp.	95	G	FR838973
<i>Desulfovibrio africanus</i>	95	H	FR838999
<i>Archaea</i>			
Uncultured <i>Crenarchaeota</i>	98	a	FR838967
<i>Methanosarcina</i> sp.	100	b	FR838977
<i>Methanosarcina</i> sp.	100	c	FR838978
<i>Methanosarcina barkeri</i>	99	d	FR838993

<sup>a</sup> Detected by denaturing gradient gel electrophoresis of density-resolved DNA gradient fractions. The phylogenetic affiliations of the 16S rRNA genes from microbes that incorporated <sup>13</sup>C-labeled substrates are shaded.

inance in the coal enrichment cultures amended with acetate. However, oxygen and nitrate were not available as electron acceptors.

Accumulation of acetate was detected in both supplemented and nonsupplemented cultures. The presence of isotopically heavy acetate in samples incubated with H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub> indicated its new formation from CO<sub>2</sub>. Relatives of *Pelobacter* and *Clostridium* species might be involved in this process. For *Pelobacter acetylenicus*, acetate formation from acetylene has been described as well (23). Which other coal and timber compounds could be feasible substrates for acetogenesis is not known. However, acetate formation appeared to continue, since the levels do not approach zero in most cases.

In the timber enrichment cultures, active bacteria similar to *Hydrogenophaga* and *Clostridium* species predominated, suggesting that they might use timber compounds or secondary fermentative products for metabolism. Only the H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub>-amended coal cultures showed distinct labeling of bacteria after 6 months. The slight heavy shift in the acetate-amended coal cultures might not be an indication of bacterial activity, but the result of an increase in the amount of bacterial DNA

with a high GC content, such as *Burkholderia* species (Fig. 3; Table 2). Like the coal cultures, the H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub>-enriched timber cultures also showed acetate formation. Obviously, timber provides acetogenic substrates other than those from coal. However, *Hydrogenophaga* as well as *Clostridium* species were recently also detected in coal samples from another mine (12).

**Acetate is the main precursor of methane.** In connecting the results from our earlier investigations (2, 14) with our new findings in the present study, we present the following arguments supporting the notion that acetoclastic methanogenesis is the main route for methane formation. First, the natural isotopic signatures of methane indicated an acetoclastic origin, supported by the isotopic signatures of acetate that was formed from <sup>13</sup>CO<sub>2</sub> (14). Second, the highest methane formation rates were observed in the acetate-amended enrichment cultures of coal and timber, while H<sub>2</sub> gave lower activities. Third, acetate was depleted in the acetate cultures but accumulated in the H<sub>2</sub>-CO<sub>2</sub>- and BES-treated enrichment cultures. Finally, DNA-SIP revealed that relatives of *Methanosarcina* spp. were responsible for methane production in both [<sup>13</sup>C]acetate- and H<sub>2</sub>-<sup>13</sup>CO<sub>2</sub>-amended cultures, and in the latter, their activity was coupled to the activity of bacteria related to acetogens.

#### ACKNOWLEDGMENTS

We thank Daniela Zoch (Federal Institute for Geosciences and Natural Resources [BGR], Hannover, Germany) and Katrin Hörmann (Helmholtz Zentrum, Munich, Germany) for technical assistance.

Financial support was provided by the BGR. Funding for M.K. was partially provided by the Deutsche Forschungsgemeinschaft (DFG) (grants KR 3311/5-1 and 6-1) and the Bundesministerium für Bildung und Forschung (grant 03G0697A). F.V.N. was financed by a grant from the DFG to T.L. within priority program SPP 1319 (grant LU 1188/4-1).

#### REFERENCES

- Altschul, S. F., et al. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Beckmann, S., M. Krüger, B. Engelen, A. A. Gorbushina, and H. Cypionka. 2011. Role of bacteria, archaea and fungi involved in methane release in abandoned coal mines. *Geomicrobiol. J.* **28**(4):UGMB-2010-0055.R1.
- Del Panno, M., I. Morelli, B. Engelen, and L. Berthe-Corti. 2005. Effect of petrochemical sludge concentration on microbial communities during a soil bioremediation process. *FEMS Microbiol. Ecol.* **53**:305–316.
- Denman, K., et al. 2007. Couplings between changes in the climate system and biogeochemistry, chapter 7. In S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, and H. L. Miller (ed.), *Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, England.
- Dojka, M. A., P. Hugenholz, S. K. Haack, and N. R. Pace. 1998. Microbial diversity in a hydrocarbon- and chlorinated-solvent-contaminated aquifer undergoing intrinsic bioremediation. *Appl. Environ. Microbiol.* **64**:3869–3877.
- Engelen, B., et al. 2008. Fluids from the oceanic crust support microbial activities within the deep biosphere. *Geomicrobiol. J.* **25**:56–66.
- Flores, R. M. 2008. Microbes, methanogenesis, and microbial gas in coal. *Int. J. Coal Geol.* **76**:1–2.
- Fry, C. J., et al. 2009. Prokaryotic populations and activities in an interbedded coal deposit, including a previously deeply buried section (1.6–2.3 km) above ~150 Ma basement rock. *Geomicrobiol. J.* **26**:163–178.
- Green, M. S., K. C. Flanagan, and P. C. Gilcrease. 2008. Characterization of a methanogenic consortium enriched from a coalbed methane well in the Powder River Basin, U.S.A. *Int. J. Coal Geol.* **76**:34–45.
- Han, B., et al. 2009. Diversity and activity of methanotrophs in alkaline soil from a Chinese coal mine. *FEMS Microbiol. Ecol.* **70**:40–51.
- Holmes, D. E., P. N. Kelly, and D. R. Lovley. 2004. Comparison of 16S rRNA, *nifD*, *recA*, *gvrB*, *rpoB*, and *fusA* genes within the family *Geobacteraceae* fam. nov. *Int. J. Syst. Evol. Microbiol.* **54**:1591–1599.
- Jones, E. J. P., M. A. Voytek, M. D. Corum, and W. H. Orem. 2010. Stimulation of methane generation from nonproductive coal by addition of nutrients or a microbial consortium. *Appl. Environ. Microbiol.* **76**:7013–7022.
- Krüger, M., P. Frenzel, and R. Conrad. 2001. Microbial processes influencing methane emission from rice fields. *Global Change Biol.* **7**:49–63.
- Krüger, M., et al. 2008. Microbial methane formation from hard coal and timber in an abandoned coal mine. *Geomicrobiol. J.* **25**:315–321.
- Lanoil, B. D., and S.-K. Han. 2003. Identification of microbes responsible for acetate consumption in soils under different wetting regimes. Kearney Foundation of Soil Science: Soil Carbon and California's Terrestrial Ecosystems, 2001–2006 Mission. Final Report: 2001028, 1/1/2002–12/31/2003. Kearney Foundation, Division of Agriculture and Natural Resources, University of California, Davis, CA.
- Lu, Y., T. Lueders, M. W. Friedrich, and R. Conrad. 2005. Detecting active methanogenic populations on rice roots using stable isotope probing. *Environ. Microbiol.* **7**:326–336.
- Lueders, T., B. Wagner, P. Claus, and M. W. Friedrich. 2004. Stable isotope probing of rRNA and DNA reveals a dynamic methylophilic community and trophic interactions with fungi and protozoa in oxic rice field soil. *Environ. Microbiol.* **6**:60–72.
- Lueders, T., M. Manfield, and M. W. Friedrich. 2004. Enhanced sensitivity of DNA- and rRNA-based stable isotope probing by fractionation and quantitative analysis of isopycnic centrifugation gradients. *Environ. Microbiol.* **6**:73–78.
- Mittal, M., and K. J. Rockne. 2008. Indole production by *Pseudomonas stutzeri* strain NAP-3 during anaerobic naphthalene biodegradation in the presence of dimethyl formamide. *J. Environ. Sci. Health A Tox. Hazard. Subst. Environ. Eng.* **43**:1027–1034.
- Muzyer, G., E. C. de Waal, and A. G. Uitterlinden. 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* **59**:695–700.
- Neufeld, J. D., et al. 2007. DNA stable-isotope probing. *Nat. Protoc.* **2**:860–866.
- Radajewski, S., P. Ineson, N. R. Parekh, and J. C. Murell. 2000. Stable-isotope probing as a tool in microbial ecology. *Nature* **403**:646–649.
- Schink, B. 1985. Fermentation of acetylene by an obligate anaerobe, *Pelobacter acetylenicus* sp. nov. *Arch. Microbiol.* **142**:295–301.
- Schreiber, A. F., and U. K. Winkler. 1983. Transformation of tetralin by whole cells of *Pseudomonas stutzeri* AS39\*. *Eur. J. Appl. Microbiol. Biotechnol.* **18**:6–10.
- Shimizu, S., et al. 2007. Molecular characterization of microbial communities in deep coal seam groundwater of northern Japan. *Geobiology* **5**:423–433.
- Shin, S. G., C. S. Lee, K. Hwang, J. H. Ahn, and S. Hwang. 2008. Use of order-specific primers to investigate the methanogenic diversity in acetate enrichment system. *J. Ind. Microbiol. Biotechnol.* **35**:1345–1352.
- Stams, A. J. M. 1994. Metabolic interactions between anaerobic bacteria in methanogenic environments. *Antonie Van Leeuwenhoek* **66**:271–294.
- Strapoc, D., M. Mastalerz, A. Schimmelmann, A. Drobnik, and S. Hedges. 2008. Variability of geochemical properties in a microbially dominated gas system from the eastern margin of the Illinois Basin, U. S. A. *Int. J. Coal Geol.* **76**:98–110.
- Strapoc, D., et al. 2008. Methane-producing microbial community in a coal bed of the Illinois Basin. *Appl. Environ. Microbiol.* **74**:2424–2432.
- Süss, J., et al. 2006. Widespread distribution and high abundance of *Rhizobium radiobacter* within Mediterranean subsurface sediments. *Environ. Microbiol.* **8**:1753–1763.
- Tao, M. X., et al. 2007. Secondary biological coalbed gas in the Xinji area, Anhui province, China: evidence from the geochemical features and secondary changes. *Int. J. Coal Geol.* **71**:358–370.
- Thauer, R. K., D. Möller-Zinkhan, and A. M. Spormann. 1989. Biochemistry of acetate catabolism in anaerobic chemotrophic bacteria. *Annu. Rev. Microbiol.* **43**:43–67.
- Thielemann, T., B. Cramer, and A. Schippers. 2004. Coalbed methane in the Ruhr Basins, Germany: a renewable energy resource? *Org. Geochem.* **35**:1537–1549.
- Ulrich, G., and S. Bower. 2008. Active methanogenesis and acetate utilization in Powder River Basin coals, United States. *Int. J. Coal Geol.* **76**:25–33.
- Warwick, P. D., F. C. Breland, and P. C. Hackley. 2008. Biogenic origin of coalbed gas in the northern Gulf of Mexico Coastal Plain, U.S.A. *Int. J. Coal Geol.* **76**:119–137.
- Widdel, F., and F. Bak. 1992. Gram-negative mesophilic sulfate-reducing bacteria, p. 3352–3378. In A. Balows, M. Dworkin, W. Harder, and K. H. Schleifer (ed.), *The prokaryotes*, 2nd ed. Springer-Verlag, New York, NY.
- Wilms, R., H. Sass, B. Köpke, H. Cypionka, and B. Engelen. 2007. Methane and sulfate profiles within the subsurface of a tidal flat are reflected by the distribution of sulfate-reducing bacteria and methanogenic archaea. *FEMS Microbiol. Ecol.* **59**:611–621.
- Zinder, S. H. 1993. Physiological ecology of methanogens, p. 128–206. In J. G. Ferry (ed.), *Methanogenesis: ecology, physiology, biochemistry and genetics*. Chapman & Hall, New York, NY.