

Peroxiredoxins and NADPH-Dependent Thioredoxin Systems in the Model Legume *Lotus japonicus*^{1[W][OA]}

Alejandro Tovar-Méndez², Manuel A. Matamoros, Pilar Bustos-Sanmamed, Karl-Josef Dietz, Francisco Javier Cejudo, Nicolas Rouhier, Shusei Sato, Satoshi Tabata, and Manuel Becana*

Departamento de Nutrición Vegetal, Estación Experimental de Aula Dei, Consejo Superior de Investigaciones Científicas, 50080 Zaragoza, Spain (A.T.-M., M.A.M., P.B.-S., M.B.); Biochemistry and Physiology of Plants, Bielefeld University, 33501 Bielefeld, Germany (K.-J.D.); Instituto de Bioquímica Vegetal y Fotosíntesis, Universidad de Sevilla y Consejo Superior de Investigaciones Científicas, 41092 Seville, Spain (F.J.C.); UMR Interactions Arbres Microorganismes 1136, Nancy Université, 54506 Vandoeuvre, France (N.R.); and Kazusa DNA Research Institute, Kisarazu, Chiba 292-0818, Japan (S.S., S.T.)

Peroxiredoxins (Prxs), thioredoxins (Trxs), and NADPH-thioredoxin reductases (NTRs) constitute central elements of the thiol-disulfide redox regulatory network of plant cells. This study provides a comprehensive survey of this network in the model legume *Lotus japonicus*. The aims were to identify and characterize these gene families and to assess whether the NTR-Trx systems are operative in nodules. Quantitative reverse transcription-polymerase chain reaction and immunological and proteomic approaches were used for expression profiling. We identified seven *Prx*, 14 *Trx*, and three *NTR* functional genes. The *PrxQ1* gene was found to be transcribed in two alternative spliced variants and to be expressed at high levels in leaves, stems, petals, pods, and seeds and at low levels in roots and nodules. The *1CPrx* gene showed very high expression in the seed embryos and low expression in vegetative tissues and was induced by nitric oxide and cytokinins. In sharp contrast, cytokinins down-regulated all other *Prx* genes, except *PrxQ1*, in roots and nodules, but only *2CPrxA* and *PrxQ1* in leaves. Gene-specific changes in *Prx* expression were also observed in response to ethylene, abscisic acid, and auxins. Nodules contain significant mRNA and protein amounts of cytosolic PrxIIB, *Trxh1*, and NTRA and of plastidic NTRC. Likewise, they express cytosolic *Trxh3*, *Trxh4*, *Trxh8*, and *Trxh9*, mitochondrial PrxIIF and *Trxo*, and plastidic *Trxm2*, *Trxm4*, and ferredoxin-Trx reductase. These findings reveal a complex regulation of Prxs that is dependent on the isoform, tissue, and signaling molecule and support that redox NTR-Trx systems are functional in the cytosol, mitochondria, and plastids of nodules.

In plants, reactive oxygen species (ROS), such as the superoxide radical and hydrogen peroxide (H₂O₂), are mainly formed in the chloroplasts, mitochondria, peroxisomes, and apoplast during photosynthesis, respiration, and other processes involving electron transfer (del Río et al., 2002; Mittler, 2002; Foyer and Noctor, 2005). Plant cells also produce reactive nitrogen species, such as nitric oxide (NO), S-nitrosoglutathione (GSNO), and peroxynitrite, under physiological conditions

(Lamattina et al., 2003; Valderrama et al., 2007; Neill et al., 2008). Overproduction of both types of reactive species is potentially deleterious, but, at tightly controlled concentrations, they fulfill essential functions in plant development, defense response, and redox signaling (Foyer and Noctor, 2005; Besson-Bard et al., 2008). Thus, antioxidant defenses are linked to cellular regulation through a complex network involving redox input elements, transmitters, targets, and sensory proteins, such as peroxiredoxins (Prxs), thioredoxins (Trxs), and glutaredoxins (Grxs; Meyer et al., 2009; Dietz and Pfannschmidt, 2011).

Prxs constitute a ubiquitous family of nonheme thiol peroxidases that catalyze the reduction of H₂O₂, alkylhydroperoxides, and peroxynitrite to water, alcohols, or nitrite, respectively (Rouhier and Jacquot, 2005; Tripathi et al., 2009). These enzymes contain one or two Cys residues at the active site and usually function as monomers or dimers. Their common catalytic mechanism involves the catalytic Cys (peroxidatic) thiol, which is oxidized by peroxides to sulfenic acid. In most Prxs, the sulfenic acid is then reduced by a second Cys (resolving) thiol forming an intra or intermolecular disulfide bond. A new catalytic cycle is allowed after the reduction of the disulfide bond using electron donors, such as Trxs, Grxs, or cyclophilins

¹ This work was funded by MICINN-FEDER (grant no. AGL2008-01298) and Gobierno de Aragón (group A53). K.-J.D.'s work on peroxiredoxins was supported by the Deutsch Forschungsgemeinschaft (grant no. Di346/11). F.J.C.'s laboratory was supported by MICINN-FEDER (grant no. BIO2010-15430) and Junta de Andalucía (grant nos. BIO-182 and CVI-5919).

² Present address: Division of Biochemistry, 117 Schweitzer Hall, University of Missouri, Columbia, MO 65211.

* Corresponding author; e-mail becana@eead.csic.es.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Manuel Becana (becana@eead.csic.es).

^[W] The online version of this article contains Web-only data.

^[OA] Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.111.177196

(Dietz et al., 2006). There are four types of Prxs in plants (1CPrx, 2CPrx, PrxII, and PrxQ), which play specific roles according to their spatio-temporal expression patterns and subcellular localizations. Plant Prxs protect the nuclei (1CPrx), plastids (2CPrxA, 2CPrxB, PrxQ, and PrxIIE), cytosol (PrxIIB, PrxIIC, and PrxIID), and mitochondria (PrxIIF) against excess ROS in stressful conditions but are also implicated in redox signaling (Romero-Puertas et al., 2007; Tripathi et al., 2009).

Unlike most other organisms, plants have a large number of *Trx* genes, at least 20 in the fully sequenced genomes of *Arabidopsis* (*Arabidopsis thaliana*) and rice (*Oryza sativa*), which are classified into seven types (for review, see Vieira Dos Santos and Rey, 2006; Meyer et al., 2009). The *Trxf*, *Trxm*, *Trxx*, *Trxy*, and *Trxz* are localized in the chloroplasts, the *Trxh* isoforms in the cytosol, and the *Trxo* in the mitochondria. However, some *Trxh* isoforms have been found also in the mitochondria, nuclei, phloem, and apoplast (Gelhay et al., 2004). Oxidized Trxs produced as a result of reactions with Prxs and other substrates are reduced back to the functional reduced state by NADPH-thioredoxin reductases (NTRA and NTRB) in the cytosol and mitochondria (Schürmann and Jacquot, 2000; Reichheld et al., 2005) or by ferredoxin-thioredoxin reductase (FTR) in the chloroplasts (Dai et al., 2004). Another NADPH-thioredoxin reductase (NTRC) has been recently found in green tissues (Serrato et al., 2004). This peculiar enzyme contains both NTR and Trx domains in the same polypeptide and may act as a complete NTR-Trx system, reactivating plastidic 2CPrx without the assistance of classical Trxs (Moon et al., 2006; Pérez-Ruiz et al., 2006; Alkhalifiou et al., 2007).

Legume root nodules are formed as a result of the molecular interaction between the roots and soil rhizobia. The bacteroids inside the nodules fix atmospheric N₂ into ammonia and in return host cells supply the bacteroids with carbon metabolites. Two model legumes, *Medicago truncatula* and *Lotus japonicus*, have been proposed for genetic analyses of indeterminate and determinate nodulation, respectively. The two types of nodules differ in some structural and biochemical features (Hirsch, 1992). The antioxidants of nodules, in particular the superoxide dismutase, catalase, and ascorbate-glutathione pathway enzymes, have been studied in some detail (for review, see Puppo et al., 2005; Becana et al., 2010), whereas there is a dearth of information concerning other antioxidant and redox sensor enzymes, such as Prxs, Trxs, and NTRs. The study of these enzymes in legumes, and particularly in nodules, is important because N₂ fixation requires a strict regulation of the redox state in the host cells and bacteroids. Thus, nodules contain abundant metalloproteins that are prone to oxidation, such as nitrogenase, ferredoxin, hydrogenase, and leghemoglobin, with a high potential for ROS generation (Dalton et al., 1998; Becana et al., 2010). Knowledge of the redox regulatory network of nodules is only slowly emerging and is still in a fragmentary state. In pea (*Pisum*

sativum), the content of PrxIIF in nodules is similar to that in roots and remains constant during nodule development (Groten et al., 2006), whereas in soybean (*Glycine max*), a *Trxh* isoform is essential for nodulation (Lee et al., 2005), and in *M. truncatula*, two isoforms of a new type of Trx, designated Trxs ("s" for symbiosis), are highly expressed in nodules (Alkhalifiou et al., 2008).

This study, designed to gain insights into the *Prx*, *Trx*, and *NTR* gene families of *L. japonicus*, is organized in two parts. First, we identified the *Prx* genes and determined their expression profiles in nodulated plants and in response to signaling compounds to better understand their functional diversity and regulation. Second, we focused on the expression of the Trx and NTR isoforms in nodules to identify possible Prx regenerating systems in these symbiotic organs.

RESULTS

Identification and Characterization of *LjPrx* Genes

The *L. japonicus Prx* (*LjPrx*) genes were identified by searching genomic and EST databases using the *Arabidopsis Prx* protein sequences as BLAST queries. The open reading frames of seven *LjPrx* genes were found to be complete based on their tentative consensus (TC) sequences (Table I), and the exon-intron structures were elucidated by comparison between the gene and TC sequences (Fig. 1). An additional gene, here termed *LjPrxQ2*, was detected in the selected genome assembly contig (Sato et al., 2008), but it is not transcribed or its expression is below detection limits, in agreement with the absence of ESTs for this gene. All the *LjPrx* genes, except *LjPrxQ2*, could be mapped (Table I). The two *Lj2CPrx* genes are highly homologous, with 93% (nucleotide) and 84% (amino acid) identities in their sequences and with 81% to 82% (nucleotide) and 90% to 93% (amino acid) identities with respect to the *2CPrxA* (*At3g11630*) and *2CPrxB* (*At5g06290*) genes of *Arabidopsis*. The *Lj2CPrx* genes were designated A and B based on the higher expression of the *Lj2CPrxA* gene in the leaves, as occurs for *Arabidopsis 2CPrxA*.

The number of exons and introns of the *LjPrx* genes (Fig. 1) is identical to that of the *Arabidopsis Prx* genes (Rouhier and Jacquot, 2005), with the exception of the *LjPrxQ1* gene. This single gene locus is transcribed in two mRNAs, *LjPrxQ1a* and *LjPrxQ1b*, by alternative splicing, using a different first exon but the same second and third exons (Fig. 1). The first exons display high homology, with identities of 92% (nucleotide) and 83% (amino acid). Although this high overall sequence identity precluded a separate analysis of each alternative spliced form, two sets of primers were designed that allowed us to quantify, respectively, the *LjPrxQ1b* mRNA and the sum of the *LjPrxQ1a* and *LjPrxQ1b* mRNAs. This quantitative reverse transcription (RT)-PCR experiment revealed that the *LjPrxQ1b* mRNA accounted for only <10% of the total *LjPrxQ1* mRNAs in leaves, stems, petals, and pods and that it was

Table 1. *Prx* genes and proteins of *L. japonicus*.

Gene	Clone ^a	Chr ^a	TC ^b	No. of ESTs ^b	Length ^c	Mol. Mass ^c	Localization ^d	Arabidopsis Ortholog ^e	Medicago Ortholog ^e
<i>Lj1CPrx</i>	LjT20M01	4	TC57452	4	219	24.7	Nucleus/cytosol	At1g48130	TC176842
<i>LjPrxQ1a</i>	LjT31N02	4	TC62358	26	224	24.4	Chloroplast	At3g26060	TC174754
<i>LjPrxQ1b</i>	LjT31N02	4	TC60736	6	226	24.6	Chloroplast	At3g26060	TC174754
<i>LjPrxQ2</i>	LjSGA_149250	ND	–	–	–	–	Chloroplast	At3g26060	TC174754
<i>Lj2CPrxA</i>	LjT18K22	1	TC75376	80	266	29.2	Chloroplast	At3g11630	TC179904
<i>Lj2CPrxB</i>	LjT04E07	5	TC76501	54	260	28.6	Chloroplast	At5g06290	TC174211
<i>LjPrxIIB</i>	LjT33L17	2	TC64422	22	162	17.5	Cytosol	At1g65980	TC182619
<i>LjPrxIIE</i>	LjT19I13	1	TC76090	20	218	23.2	Chloroplast	At3g52960	TC174129
<i>LjPrxIIF</i>	LjT27H06	6	TC60826	16	197	21.2	Mitochondrion	At3g06050	TC176989

^aDesignation of genomic clones and chromosome location (ND, not determined). ^bDesignation of TC sequences and number of ESTs according to the DFCI *Lotus* Gene Index (6.0). ^cPredicted number of amino acid residues and molecular mass (kD) of precursor proteins. ^dPredicted subcellular localizations of mature proteins. ^eOrtholog genes of Arabidopsis and *M. truncatula* according to The Arabidopsis Information Resource and DFCI *Medicago* Gene Index (11.0), respectively.

undetectable in the other tissues examined (data not shown).

The LjPrx family includes at least one member of each Prx type, as confirmed by phylogenetic analysis (Supplemental Fig. S1). Furthermore, in silico analyses predicted that Lj2CPrxs, LjPrxQs, and LjPrxIIE are targeted to plastids, LjPrxIIB to the cytosol, and LjPrxIIF to the mitochondria (Table 1). The Lj1CPrx sequence contains two highly conserved motifs: PVCTTE, which is thought to be the catalytic site of the enzyme, and KE (X₁₃)KK(X₂)LRFT, which is a putative nuclear localization signal (Mowla et al., 2002, and refs. therein). In addition, sequence analysis of Lj1CPrx with TargetP predicted that it is a cytosolic enzyme, consistent with the dual localization in the nucleus and cytosol of the 1CPrx proteins of barley (*Hordeum vulgare*), wheat (*Triticum aestivum*), and Arabidopsis (Stacy et al., 1999; Haslekaas et al., 2003; Pulido et al., 2009).

To gain insights into the functional diversification of LjPrxs, their expression profiles were determined in

plant tissues. Two genes, *Lj1CPrx* and *LjPrxIIB*, are highly expressed in specific tissues, as can be noted by the different scales used to represent their mRNA levels compared to those of the other genes (Fig. 2A). Thus, expression of *Lj1CPrx* is almost confined to the embryo and hence is also relatively high in whole seeds. Only low levels of *Lj1CPrx* mRNA could be detected in vegetative organs such as roots, nodules, and leaves. Likewise, *LjPrxIIB* shows very high expression in pollen, moderate expression in embryos and seeds, and low expression in other organs. By contrast, *LjPrxIIF* is expressed in all organs but at maximal levels in the embryo. The genes encoding the plastidic Prx isoforms, namely, *Lj2CPrxA*, *Lj2CPrxB*, *LjPrxIIE*, and *LjPrxQ1*, are also expressed throughout the plant, albeit for some of these genes the mRNA levels were close to detection limits in pollen and roots (Fig. 2A). The relative abundance of all *LjPrx* mRNAs within each plant tissue was determined (Fig. 2B). The leaves, stems, flowers, petals, and pods exhibit similar




Figure 1. Exon-intron organization of *LjPrx* genes. Exons are depicted in black boxes and intron in white boxes. Exon and intron sizes are indicated in numbers of base pairs and are drawn to scale.




Figure 2. Expression profiles of the *LjPrx* gene family. A, Relative *LjPrx* mRNA levels in roots (R), nodules (N), leaves (L), stems (St), flowers (F), petals (Pt), pollen (Pl), pods (Pd), seedless pods (Sp), seeds (S), and embryos (E). The mRNA levels were normalized with respect to *ubiquitin*, and the *LjPrxIIB* mRNA level in roots was arbitrarily assigned a value of 1. All data are means \pm SE of four to eight biological replicates. B, Relative *LjPrx* mRNA level within each plant organ, calculated from data in A.

expression profiles, with 68% to 82% of the transcripts encoding plastidic *LjPrxs*. The pollen and embryos show unique expression profiles. Thus, in pollen >99% of the transcripts correspond to *LjPrxIIB* mRNA, and in the embryo, the *Lj1CPrx* and *LjPrxIIB* mRNAs account for 75% and 15%, respectively, of the total transcripts (Fig. 2B).

The content of the *LjPrx* proteins in plant tissues was examined using immunoblots (Fig. 3A). The *Lj1CPrx* protein was found specifically in the embryo and was undetectable in any other tissues or even in seed extracts (Fig. 2A). Because seeds contain significant *Lj1CPrx* mRNA levels and the protein is present in the embryos, it may be below detection limits in the whole seed extracts due to a dilution effect. Similarly, the *LjPrxQ1* protein was found exclusively in leaves. The *Lj2CPrx* proteins accumulated in leaves and to a lower extent in flowers, pods, seeds, and embryos, whereas they were undetectable in roots, nodules, and pollen. In contrast, *LjPrxIIB* and *LjPrxIIF* were found in all organs, although the amount of *LjPrxIIB* was very low in roots. Also, it was necessary to load 5-fold more protein on immunoblots to unambiguously detect *PrxIIB* in nodules of *L. japonicus* and other legumes, such as pea, common bean (*Phaseolus vulgaris*), and cowpea (*Vigna unguiculata*; Fig. 3B).

Regulation of *LjPrx* Expression by Hormones and NO

The effect of several hormones and stress signaling molecules was also studied to gain information about their role in developmental and acclimatory regulation

of *LjPrx* gene expression. To this purpose, nodulated plants were grown in hydroponic medium supplemented with hormones for 48 h, and the expression levels of *LjPrx* genes were determined in the roots (Fig. 4). Treatment of plants with GA_3 , jasmonic acid (JA), or salicylic acid did not cause substantial changes in the *LjPrx* mRNA levels, whereas 1-aminocyclopropane-1-carboxylic acid (the immediate ethylene precursor) down-regulated *LjPrxQ1* and *LjPrxIIB*, abscisic acid (ABA) up-regulated *Lj2CPrxA* and *LjPrxIIB*, and indole-3-acetic acid up-regulated *Lj1CPrx*. However, the most marked effects were observed with cytokinins (CKs), which decreased the expression of all *LjPrx* genes, except *Lj1CPrx* and *LjPrxQ1*, in roots (Fig. 4). This finding led us to investigate the effects of CK on the expression of *LjPrx* genes also in nodules and leaves. The response of most *LjPrx* genes to CK in nodules was similar to that observed in roots, whereas in leaves, CK caused down-regulation of *LjPrxQ1* but had no effect on the *LjPrxII* genes. Although the *Lj1CPrx* mRNA levels were very low in vegetative organs, a strong induction of this gene in roots and leaves and a much weaker induction in nodules were detected (Fig. 4). Additional experiments showed that the increase of *Lj1CPrx* mRNA level in roots occurred after only 3 h and persisted for at least 48 h. This gene was induced with only 5 μ M CK and had maximal expression with 100 μ M CK.

The effects of CK on *LjPrx* expression suggest an important function of this hormone in the control of the cellular level of H_2O_2 . To test this possibility, the mRNA levels of important genes implicated in H_2O_2




Figure 3. Immunoblot analyses of LjPrx proteins in different organs. A, Relative abundance of LjPrx proteins in roots (R), nodules (N), leaves (L), flowers (F), pollen (PI), pods (Pd), seedless pods (Sp), seeds (S), and embryos (E). B, Detection of cytosolic PrxIIB proteins in nodules of *L. japonicus* (Lj), *P. sativum* (Ps), *P. vulgaris* (Pv), and *V. unguiculata* (Vu). Gels were loaded with 10 μg (A) or 50 μg (B) of protein per lane, and the apparent molecular masses (kD) of the proteins are indicated on the right. Blots are representative of two independent protein extractions.

metabolism were quantified. In roots, CK did not cause any significant change in the contents of transcripts encoding mitochondrial Mn-superoxide dismutase, cytosolic and plastidic Fe- and CuZn-superoxide dismutases, or thylakoidal, stromal, and peroxisomal ascorbate peroxidases (data not shown). However, in response to CK, the expression of catalase (*LjCAT*) increased in roots and nodules but not in leaves, whereas the expression of cytosolic ascorbate peroxidase (*LjAPXc*) decreased in nodules and increased in leaves (Fig. 4). The effect of NO on *Lj1CPrx* expression was also examined because this signal molecule has been implicated in the response of plant cells to CK (Tun et al., 2001; Carimi et al., 2005). Plants were supplied with two NO donors and *Lj1CPrx* expression levels were determined in the roots. Both S-nitroso-N-acetyl-DL-penicillamine (SNAP) and GSNO increased

gene expression, although maximal induction was achieved after 24 h with SNAP and after only 3 h with GSNO (Supplemental Fig. S2).

The stimulatory effect of CK on *Lj1CPrx* mRNA accumulation and the fact that this gene is almost exclusively expressed in the embryo (Fig. 2A) prompted us to investigate the effects of this hormone on *Lj1CPrx* expression in germinating seeds. In the absence of CK, the content of *Lj1CPrx* mRNA progressively decreased following germination and was hardly detectable after 48 h of imbibition (Fig. 5). In the presence of CK, the mRNA level was also reduced but, after 24 h, it was 65% of the initial value compared to 14% for the control seeds. These results suggest that the CK treatment induced *Lj1CPrx* expression in seeds and that CK was unable to completely overcome the down-regulation of the gene that takes place during germination, as can be seen at 48 h (Fig. 5).

Identification and Characterization of *LjTrx* and *LjNTR* Genes

Most Prx isoforms are reduced efficiently by Trxs, and, in turn, the nonplastidic Trxs are regenerated by NTRA and NTRB. To complete our study, we pursued the identification of the Trx and NTR proteins of *L. japonicus*. The search focused on the isoforms expressed in nodules to determine if the NTR-Trx system might be operative in these unique plant organs. Because of the complexity of the *Trx* gene family, we combined mRNA expression with immunoblot and proteomic analyses of nodules.

The *L. japonicus* EST and genomic databases were screened to identify Trxs using the Arabidopsis and *M. truncatula* protein sequences as BLAST queries. This analysis identified 14 *LjTrx* genes coding for six isoforms of *Trxh*, three isoforms of *Trxm*, and one isoform each of *Trxf*, *Trxx*, *Trxy*, *Trxz*, and *Trxo*, but it failed to detect any homologs of Trxs (Table II). All these gene sequences were already deposited in the data banks




Figure 4. Heat map of the hormone response of the expression of *LjPrx* genes in roots. Plants grown in hydroponic cultures were treated for 48 h with 50 μM of each hormone. The effects of CK on *LjPrx* expression in leaves and nodules, and on *LjCAT* and *LjAPXc* genes, are also shown. Transcript levels were normalized with *ubiquitin* and expressed relative to those found in control plants, which were arbitrarily given a value of 1. Values are means \pm SE of four to 10 biological replicates from at least two independent treatments. ACC, 1-Aminocyclopropane-1-carboxylic acid; IAA, indole-3-acetic acid; SA, salicylic acid.




Figure 5. Effect of CK on expression of the *Lj1CPrx* gene during seed germination. Seeds were stratified for 24 h at 4°C and then germinated in agar plates for up to 48 h in the absence (control) or presence of 50 μ M CK. Data are expressed with respect to the mRNA levels at time zero, which were arbitrarily given a value of 1, and are means \pm SE of two biological replicates, each of them corresponding to the total RNA from 10 germinating seeds.

except the *LjTrxh1* clone (LjT45J20), which was isolated using TC65928 sequence information. This clone was completely sequenced for this study (accession no. AP012058). The *LjTrx* genes were designated according to the similarities of their derived proteins with respect to those of *M. truncatula* (Alkhalifioui et al., 2008; Renard et al., 2011). An alignment of the *Trxh* sequences (Supplemental Fig. S3) and a phylogenetic analysis of the *Trxs* (Supplemental Fig. S4) of *L. japonicus* and other model plants were performed to verify protein assignments to the *Trx* types and the three *Trxh* subgroups. Thus, *LjTrxh1* and *LjTrxh3* belong to subgroup I, *LjTrxh4* and *LjTrxh6* to subgroup II, and *LjTrxh8* and *LjTrxh9* to subgroup III. *LjTrxh9* exhibits a very peculiar active site and may rather possess a protein disulfide isomerase activity that depends on glutathione instead of NTR (Gelhay et al., 2004; Serrato et al., 2008). Only one isoform of subgroup I, *LjTrxh1*, contains the N-terminal motif MAAEE (Supplemental Fig. S3) found in the *Trxh1* and *Trxh2* of *M. truncatula*, which might allow these proteins to be secreted to the phloem or apoplast in addition to being localized to the cytosol (Supplemental Fig. S3; Renard et al., 2011). The *LjTrxh3* gene was found to be transcribed (Table II), whereas no ESTs are available to date for the orthologous *Trxh3* gene of *M. truncatula* (Renard et al., 2011). The expression profiles of the *LjTrx* genes were determined in roots, nodules, and leaves (Fig. 6). Notably, *LjTrxh1* showed by far the greatest expression levels, whereas the *LjTrxh6* mRNA was virtually undetectable in the three plant organs. As expected, the *LjTrxf*, *LjTrxm1*, *LjTrxm4*, and *LjTrxx* genes, which encode plastidic isoforms, were highly expressed in leaves compared to roots or nodules. In fact, the amounts of *LjTrxf* mRNA in roots and nodules, or of *LjTrxm1* and *LjTrxx* mRNAs in roots, were near detection limit. Almost no expression of *LjTrxz* was observed in roots, nodules, and leaves. By contrast, in these three plant organs, the *LjTrxh3*, *LjTrxh8*, *LjTrxy*, and *LjTrxo* genes had low but significant expression, whereas *LjTrxh4* and *LjTrxh9* showed mod-

erate expression (Fig. 6). Proteomic analyses allowed the unambiguous identification of the cytosolic *Trxh1* isoform in nodules of *L. japonicus*, *M. truncatula*, and common bean (Table III). These analyses also identified in nodules several *Prxs* (*PrxII*B, *PrxIII*E, and *PrxIII*F) as well as two *Grxs* (*GrxC2* and *GrxC4*) that may act as putative electron donors of *Prxs* (Table III).

Similarly, a search of *L. japonicus* databases and genomic libraries allowed the identification of three *LjNTR* genes with their complete open reading frames (Table II). The sequence of one genomic clone (LjT16K13) was available from public databases, whereas two other clones (LjB04N17 and LjB24C14) were isolated from bacterial artificial chromosome (BAC) libraries using sequence information on TC63269 (accession no. AP012059) and TC57567 (accession no. AP012060), respectively. Phylogenetic analysis confirmed protein assignments to the NTRA/B and NTRC types (Supplemental Fig. S5). Designation of LjNTRA and LjNTRB was based on sequence identity (86% and 80%, respectively) to the single NTR isoform (NTRA) of *M. truncatula* (Alkhalifioui et al., 2007) rather than to the Arabidopsis *NTRA* and *NTRB* genes (approximately 80%). The motifs characteristic of NTRs, namely, catalytic Cys residues and FAD- and NADPH-binding sites, were fully conserved in the *L. japonicus* NTRA/B enzymes (Supplemental Fig. S6).

Because each of the Arabidopsis *NTRA* and *NTRB* genes can generate two types of transcripts, encoding cytosolic and mitochondrial NTR isoforms (Laloi et al., 2001; Reichheld et al., 2005), we investigated if this also occurred for the *LjNTR* genes by performing a semi-quantitative RT-PCR analysis (Fig. 7). Specific primers were designed so that one pair of primers amplified solely the long cDNA, whereas the second pair amplified both long and short cDNAs. The long *LjNTRA* mRNA was found in nodules and leaves but not in roots, whereas the long *LjNTRB* mRNA was only detected in nodules at very low levels (Fig. 7). Consistent with this, two ESTs are available for the long *LjNTRB* mRNA for nodules (accession no. CB829112) and nodulating roots (accession no. DC595411). Together, the data indicate that, under our plant growth conditions, the mitochondrial isoform of LjNTRA is produced only in nodules and leaves. On the other hand, LjNTRC is predicted to be localized in the chloroplasts, as occurs for rice NTRC (Serrato et al., 2004).

The expression of the *LjNTR* genes was investigated in roots, nodules, and leaves. Although the two alternative mRNAs for *LjNTRA* and *LjNTRB* were probably present, they could not be quantified separately. Instead, the total mRNA levels of *LjNTRA*, *LjNTRB*, and *LjNTRC* were determined (Fig. 8A). All these genes were expressed in the three plant organs, but the *LjNTRA* mRNA levels were considerably higher. As expected for a gene coding for a chloroplastic enzyme, the expression of *LjNTRC* was enhanced in leaves relative to roots or nodules. Additional experiments with other plant tissues indicated that expression

Table II. *Trx* and *NTR* genes and proteins of *L. japonicus*

Gene	Clone ^a	Chr ^a	TC ^b	No. of ESTs ^b	Length ^c	Mol. Mass ^c	Localization ^d	Arabidopsis Ortholog ^e	Medicago Ortholog ^e
<i>LjTrxh1</i>	LjT45J20	5	TC65928	34	121	13.3	Cytosol	At3g51030	TC197256
<i>LjTrxh3</i>	LjT17E09	2	TC68183	12	121	13.3	Cytosol	At3g51030	CR955005 ^f
<i>LjTrxh4</i>	LjT40C04	1	TC65406	6	131	14.4	Cytosol	At5g39950	TC177162
<i>LjTrxh6</i>	LjT58M11	2	TC65208	2	126	14.3	Cytosol	At1g69880	TC188623
<i>LjTrxh8</i>	LjSGA_031277	ND	TC58009	16	138	15.5	Cytosol	At3g08710	TC176865
<i>LjTrxh9</i>	LjSGA_132520	ND	TC63066	33	123	13.8	Cytosol	At1g11530	TC177045
<i>LjTrxf</i>	LjSGA_082631	ND	TC59402	23	179	19.3	Chloroplast	At5g16400	TC174294
<i>LjTrxm1</i>	LjSGA_017977	ND	TC60229	8	181	19.8	Chloroplast	At4g03520	TC180914
<i>LjTrxm2</i>	LjSGA_126827	ND	TC71331	15	183	19.8	Chloroplast	At3g15360	TC178205
<i>LjTrxm4</i>	LjSGA_126077	ND	TC67299	20	182	19.8	Chloroplast	At3g15360	TC193088
<i>LjTrxx</i>	LjT02A04	5	TC61897	9	185	20.2	Chloroplast	At1g50320	TC173902
<i>LjTrxy</i>	LjT08O18	5	TC61826	6	168	18.7	Chloroplast	At1g76760	TC175512
<i>LjTrxz</i>	LjSGA_025025	ND	TC62611	9	188	21.3	Chloroplast	At3g06730	NP7258770
<i>LjTrxo</i>	LjT11M06	4	TC61209	19	179	20.0	Mitochondrion	At2g35010	TC184686
<i>LjNTRA</i>	LjB04N17	ND	TC63269/73407	49	369	39.5 (35.6)	Mitochondrion (cytosol)	At2g17420	TC177239
<i>LjNTRB</i>	LjT16K13	ND	TC73044/80146	6	387	40.4 (30.7)	Mitochondrion (cytosol)	At2g17420	TC177239
<i>LjNTRC</i>	LjB24C14	ND	TC57567/68679	9	518	56.6	Chloroplast	At2g41680	TC189527

^aDesignation of genomic clones and chromosome location (ND, not determined). ^bTC sequences and number of ESTs according to the DFCI Lotus Gene Index (6.0). ^cPredicted number of amino acid residues and molecular mass (kD) of precursor proteins. The molecular mass in parentheses corresponds to the protein encoded by the putative alternative mRNA. ^dPredicted subcellular localizations of mature proteins. The localization in parentheses corresponds to the protein encoded by the putative alternative mRNA. ^eOrtholog genes of Arabidopsis and *M. truncatula* according to The Arabidopsis Information Resource and DFCI Medicago Gene Index (11.0), respectively. ^fGenomic clone (BAC number) is given because no ESTs are available (Renard et al., 2011).

levels of *LjNTRA* in pollen and *LjNTRB* in embryos were approximately 12- and 0.5-fold, respectively, those found in roots. The abundance of the *LjNTRA/B* and *LjNTRC* proteins in roots, nodules, and leaves was compared using immunoblots (Fig. 8B). A single *LjNTRA/B* protein (35 kD) was observed in all three organs, whereas the *LjNTRC* protein (51.6 kD) was very abundant in leaves, detectable in nodules, and undetectable in roots. The *NTRA/B* protein was also present in other legume nodules (Fig. 8C), and its identity was confirmed by proteomic analyses of *M. truncatula* nodules (Table III).

It is also worth noting that we detected significant mRNA levels not only of some plastidic Trxs but also of the *FTRB* gene (TC64844) in nodules of *L. japonicus*. This gene encodes the catalytic subunit of FTR, an essential component of the redox FTR-Trx system in the chloroplasts; hence, the *FTRB* mRNA levels were expected to be high in the leaves. However, the roots and nodules also contained significant amounts of transcript, approximately 7-fold lower than in the leaves (Supplemental Fig. S7).

DISCUSSION

In this work, we identified the Prx, Trx, and NTR multigenic families of *L. japonicus* and determined their expression profiles in plant tissues and, for Prxs, also in response to signaling compounds. Prxs play major roles in preventing oxidative damage and in maintaining redox homeostasis. These essential functions are con-

sistent with the presence of Prx isoforms in most, if not all, cellular compartments. Transcriptional regulation of the *Prx* genes depends on developmental and environmental factors (Dietz et al., 2006). In *L. japonicus*, the genes encoding plastidic Prx isoforms show high (*Lj2CPrxA* and *LjPrxQ1*) or moderate (*Lj2CPrxB* and *LjPrxIII*) expression in the leaves. The *Lj2CPrx* proteins, indistinguishable in immunoblots, were found to accumulate in leaves and, less abundantly, in flowers, pods, seeds, and embryos, which suggests that this type of Prxs is also present in nonphotosynthetic plastids. This was confirmed by the finding of *Lj2CPrxB* in the proteome of common bean nodules. A recent study linked the redox state of 2CPrx in animal and plant cells




Figure 6. Expression profiles of *LjTrx* genes in roots, nodules, and leaves. The mRNA levels were normalized with respect to *ubiquitin*, and the *LjTrxo* mRNA level in roots was arbitrarily assigned a value of 1. All data are means ± SE of four biological replicates.

Table III. Identification of Prxs and their putative physiological reductants by proteomic analyses of legume nodules

Protein	Legume	TC ^a	UniProt ^b	Peptides ^b
2CPrxB	<i>P. vulgaris</i>	TC32275	Q9FE12	ASSELPLVGNTPDFEAEAVFDQEFIK, SGGLGDLNYPLISDVTK, SYDVLIPDQGIAR
PrxIIB	<i>M. truncatula</i>	TC182619	B7FH22	YTHALGLELDLSDK, FALLVEDLK
PrxIIE	<i>M. truncatula</i>	TC174129	C6TFM7 ^c	AIGVELDLSKPKVGLGVR, LFNLEEGGAFTFSGADDILK
PrxIIF	<i>M. truncatula</i>	TC176989	B7FGM0	VATGSDIISAASNVSQK, SLELTTDLGALLGTR
	<i>P. vulgaris</i>	TC33229	Q6KBB1 ^c	VATGTDIVSAAPNVSLQK, SLELVTDLGALLGTR
Trxh1	<i>L. japonicus</i>	TC65928	Q6RJZ7 ^c	FIAPILAEIAK, TVAEWVNVEAMPTFLFK
Trxh1	<i>M. truncatula</i>	TC197256	A1BLP6	FIAPILAEIAK, TVAEWAVDAMPTFLFK
Trxh1	<i>P. vulgaris</i>	TC38652	A1BLP7 ^c	FIAPILAEIAK, WSIEMPTFLFK
GrxC4	<i>M. truncatula</i>	TC181572	C6TCR3 ^c	IQDVLVNIVGK, HLGGSDETVEAYESGLLAK
GrxC2	<i>P. vulgaris</i>	TC40141	B3F8F4 ^c	LIEMDVEPDGADIQAALLEWGTGQR, LVPLITSAGAITK
NTRA	<i>M. truncatula</i>	TC177239	A6XJ26	VSGLFFAIGHEPATK, TSVEGVFAAGDVQDKK

^aTC sequences according to the DFCI *Lotus* Gene Index (6.0), *Medicago* Gene Index (11.0), or Bean Gene Index (4.0). Data of *M. truncatula* nodules were taken from Larrainzar et al. (2007), stored in the ProMEX spectral library (<http://promex.pph.univie.ac.at/promex/>), and updated to the current DFCI version. ^bUniProt accessions (UniRef100) and peptides detected. ^cUniProt accessions of ortholog proteins showing best match hit.

to the circadian clock and described it as a mechanism that functions independently of transcription (O'Neill et al., 2011). Such a function in timing metabolism could be also important in nonphotosynthetic plant cells.

Interestingly, two alternative spliced variants of the *LjPrxQ1* gene and a putative *LjPrxQ2* pseudogene could be identified. The high similarity of the deduced LjPrxQ1a and LjPrxQ1b proteins suggests that they do not differ, at least substantially, in their catalytic properties. Rather, the reason for the occurrence of two *LjPrxQ1* spliced variants may reside on a different regulation because the *LjPrxQ1b* mRNA levels were 10-fold lower than those of *LjPrxQ1a*. The two PrxQ1 isoforms might be expressed in different types of leaf cells or under different environmental conditions. In *Arabidopsis*, the PrxQ protein is attached to thylakoids (Lamkemeyer et al., 2006), and its transcript is highly responsive to light, ascorbate, and compounds inducing oxidative stress (Horling et al., 2003). In poplar (*Populus* spp.), the *PrxQ* mRNA level increases following pathogen infection (Rouhier et al., 2004). Although the *LjPrxQ1* mRNA was present in leaves, flowers, and embryos, the protein was only detectable in the leaves, suggesting posttranscriptional regulation of the gene. This organ specificity was also observed for poplar PrxQ (Rouhier et al., 2004) and suggests that the protein may be exclusively implicated in chloroplast protection. Interestingly, the expression of the *LjPrxQ1* gene was down-regulated by two hormones, ethylene and JA, that play a major role in stress signaling. Unlike PrxQ, the chloroplastic PrxIIE of *Arabidopsis* is largely present as a soluble protein in the stroma (Dietz et al., 2006) and the gene is constitutively expressed (Bréhélin et al., 2003), which suggests that PrxIIE also plays a role in other types of plastids. Our finding of a high level of *LjPrxIIE* mRNA in seeds and particularly in the embryo is consistent with a function of its protein product in seed development and/or germination.

The *Arabidopsis* genome contains three genes encoding cytosolic PrxIIs: *PrxIIB* is ubiquitously expressed in plant tissues, whereas *PrxIIC* and *PrxIID* are expressed at high levels in pollen (Bréhélin et al., 2003) and at low levels in other tissues (Pena-Ahumada et al., 2006). We could identify only one homolog of such genes, *PrxIIB*, in the *L. japonicus* databases. The LjPrxIIB mRNA and protein were found in all organs, although at much higher levels in pollen, seeds, and embryos than in roots, nodules, and leaves. These observations point to a role of LjPrxIIB in the antioxidative protection of pollen grains in order to cope with oxidative stress during desiccation (Bréhélin et al., 2003). They also suggest that PrxIIB is important in seeds and maturing fruits (Matamoros et al., 2010) and that, in legumes, this single protein could fulfill the functions of the three cytosolic PrxIIs of *Arabidopsis*. Previous work failed to detect a typical cytosolic PrxII in pea nodules (Groten et al., 2006; Matamoros




Figure 7. Steady-state levels of alternative transcripts for the *LjNTRA* and *LjNTRB* genes in roots, nodules, and leaves. Semiquantitative RT-PCR analysis was carried out using gene-specific primers (Supplemental Table S1). One pair of primers amplified exclusively the long cDNA (L), and the second pair amplified both long and short cDNAs (LS). Numbers of base pairs are given on the right.




Figure 8. Expression of *LjNTR* genes in plant organs. **A**, Relative *LjNTR* mRNA levels in roots (R), nodules (N), and leaves (L). The mRNA levels were normalized with respect to *ubiquitin*, and the *LjNTRA* mRNA level in roots was arbitrarily assigned a value of 1. All data are means \pm SE of four biological replicates from two independent treatments. **B**, Immunoblots of *LjNTR* proteins in roots, nodules, and leaves. **C**, Immunoblots of NTRA/B proteins in nodules of *L. japonicus* (Lj), *P. sativum* (Ps), *P. vulgaris* (Pv), and *V. unguiculata* (Vu). Gels were loaded with 10 μ g (B) or 50 μ g (C) of protein per lane, and the apparent molecular mass (kD) of the proteins is indicated on the right. Blots are representative of at least two independent protein extractions.

et al., 2010), although the presence of a putative PrxIIA homolog (68 kD) was reported (Groten et al., 2006). This was probably due to the low abundance of PrxIIB in nodules because a genuine cytosolic PrxII (17 kD) was found here using higher protein loadings and its identity was verified by proteomic analyses of various legume nodules.

In contrast, the *LjPrxIIF* mRNA and protein were readily detected in all tissues, consistent with the hypothesis that PrxIIF is important in redox homeostasis and antioxidant defense of mitochondria (Finkemeier et al., 2005). This enzyme is widely distributed in all plant tissues and probably has a house-keeping function in mitochondria (Gama et al., 2007). In pea leaves, PrxIIF is induced by salt, cadmium, and cold stress (Barranco-Medina et al., 2007), whereas the poplar enzyme is relatively unresponsive (Gama et al., 2007). The high levels of *LjPrxIIF* mRNA and protein in the embryo may reflect an increased need for protection against ROS generated when respiration is resumed during imbibition, as proposed for cereal seeds (Stacy et al., 1999; Pulido et al., 2009). In fact, the production of superoxide radicals and H_2O_2 is enhanced in mitochondria from soybean embryonic axes during imbibition (Puntarulo et al., 1988), and PrxIIF could thus play a role in protecting mitochondrial DNA in seed cells. This protein was also found in nodules using immunoblots and proteomics. In pea nodules, the PrxIIF content remained unaffected with aging or after exogenous supply of ascorbate (Groten et al., 2006). In contrast, exogenous CK caused down-regulation of *LjPrxIIF* in roots and nodules, although this effect was not specific because the hormone also decreased the expression of most other *LjPrx* genes in both plant organs.

The strong effect of CK on expression of *Prx* genes has not been described so far and could be of physiological relevance by either linking Prxs to CK-dependent signal transduction or by adjusting the cellular redox

milieu in plants. The cell cycle is under control of CK and redox state (den Boer and Murray, 2000), and glutathione is recruited to the nucleus in proliferating cells (Díaz Vivancos et al., 2010). Thus, the up-regulation by CK of *Lj1CPrx*, which encodes a nuclear protein, supports the hypothesis that CK and Prx collaborate in tuning the proper redox state of the dividing cell. To understand whether the effects of CK are related to ROS metabolism, the expression of several other genes encoding H_2O_2 -scavenging enzymes was examined. Plant treatment with CK resulted in up-regulation of *LjCAT* and down-regulation of *LjAPXc* in roots and nodules and had the opposite effects in leaves. Also, external application of CK increases antioxidant enzyme activities and delays leaf senescence (Zavaleta-Mancera et al., 2007). Taken all these results together, we conclude that CK may affect H_2O_2 homeostasis in plant cells through changes in the regulation of critical antioxidant enzymes, such as Prxs, catalase, and ascorbate peroxidase. In this regard, a novel finding in this study is that *Lj1CPrx* is induced by CK in roots and leaves. This hormone promotes cell division (Romanov, 2009), which requires enhanced protection of DNA against ROS; therefore, the induction of 1CPrx would favor such a role, as has been proposed to occur during the desiccation and early imbibition of seeds (Aalen et al., 1994). The protective and regulatory functions proposed for 1CPrx (Pulido et al., 2009) would explain the presence of low levels of *Lj1CPrx* mRNA in vegetative tissues. The localization of 1CPrx is considered to be highly restricted to the nuclei and cytosol of the developing embryo and aleurone cells of seeds (Stacy et al., 1999; Hasleková et al., 2003; Pulido et al., 2009). The protein has nevertheless been recently detected in vegetative tissues of the resurrection plant (*Xerophyta viscosa*) under abiotic stress or following ABA application (Mowla et al., 2002). These results suggest that 1CPrx is also expressed, although at low levels, in some plant tissues or species, where the

protein may exert an antioxidant and/or signaling function in the nuclei. It is also worth mentioning that *Lj1CPrx* is induced by NO. This induction has not been reported to date for plant Prxs, but it was recently described for PrxI and PrxVI in murine macrophages and proposed to play a protective role against nitrosative stress and, indirectly, in H₂O₂ signaling (Diet et al., 2007). In any case, the NO-mediated induction of *Lj1CPrx*, should this occur also in seeds, would be consistent with the stimulating effect of NO on germination (Lamattina et al., 2003).

The presence of Prxs in nodules of *L. japonicus* shown in this work raised the question of whether the system most commonly used for Prx regeneration, consisting of Trx and NTR, is operative in these specialized organs. Consequently, the expression of *LjTrxs* and *LjNTRs*, particularly in nodules, was investigated. The genes encoding all Trx types, except the *s*-type, were identified in the *L. japonicus* genome and found to be transcribed. The *Trxs* genes were reported to be functional in *M. truncatula* (Alkhalfioui et al., 2008) but could not be found in the genomes of *L. japonicus* or soybean, which suggests that they are restricted to specific tribes or genera of legumes. In contrast, we could clearly detect a *Trxh1* isoform in nodules of *L. japonicus*, *M. truncatula*, and common bean. However, *LjTrxh4* rather than *LjTrxh1* is probably the ortholog of a soybean *Trxh* previously reported as being essential for ROS scavenging in nodules (Lee et al., 2005). This soybean *Trxh* isoform has higher amino acid identity with *LjTrxh4* (73%) than with *LjTrxh1* (53%), *LjTrxh3* (55%), or *LjTrxh8* (42%). Interestingly, *LjTrxh4* shows greater expression in nodules than in roots or leaves. Also, we could identify three functional *LjNTR* genes that are expressed in nodules. Of these, *LjNTRA* and *LjNTRB* produce long and short mRNAs presumably encoding the mitochondrial and cytosolic isoforms, as described for their *Arabidopsis* counterparts (Laloi et al., 2001; Reichheld et al., 2005).

The presence of NTR enzymes in nodules had not been previously reported and provides strong support to the functioning of redox NTR-Trx systems, in conjunction with Prxs, in the symbiotic tissue (Fig. 9). This is most evident by the finding of the PrxIIB, *Trxh1*, and NTRA proteins in the nodule cytosol. Such an NTR-Trx system requires a steady supply of NADPH, which is mainly produced by the enzymes Glc-6-P dehydrogenase, 6-phosphogluconate dehydrogenase, and isocitrate dehydrogenase, all of them very active in nodules (Marino et al., 2007). These dehydrogenases also may provide the NADPH needed for glutathione reductase and the ascorbate-glutathione pathway (Dalton et al., 1998), or for Grxs, which were also identified in nodules and are efficient reductants of PrxII (Rouhier et al., 2002). In this regard, a recent study has shown that a NTRA-*Trxh3* system can intervene as a functional backup for cytosolic glutathione reductase in *Arabidopsis* leaves (Marty et al., 2009).

Two other NTR-Trx pathways, localized in mitochondria and plastids, may be also functional in nodules




Figure 9. Model of the redox NTR-Trx systems and their putative Prx targets, which may be operating in the mitochondria, cytosol, and plastids of nodule host cells. A putative FTR-Trx system of plastids is also indicated in dashed lines. This model essentially has been built based on the known biochemical specificities observed *in vitro*. The NTRA/B, NTRC, FNR, Grx, *Trxh1*, PrxIIB, PrxIIE, PrxIIF, and 2CPrx isoforms were detected by immunoblots and/or proteomic analysis, whereas expression of FTR, *Trxo*, *Trxh4*, *Trxm*, *Trxx*, and *Trxy* was detected at the mRNA level. FNR, Ferredoxin-NADP reductase; Fdx, ferredoxin; GR, glutathione reductase; GSH, glutathione.

(Fig. 9). First, the expression of mitochondrial NTRA/B and *Trxo* described here, in addition to PrxIIF, points to the functioning of an NTR-Trx system in nodule mitochondria. This is supported by *in vitro* reconstitution systems with recombinant *Trxo* and PrxIIF from pea leaves, which have shown that *Trxo* strongly interacts with, and can act as an electron donor to, PrxIIF (Finkemeier et al., 2005; Barranco-Medina et al., 2008). Second, we could detect *LjNTRC* in nodules, albeit at the low levels expected for a nonphotosynthetic tissue. The identification of low levels of 2CPrxB and PrxIIE in nodules using proteomics would suggest that NTRC could act as an electron donor to those Prxs, lending support to the operation of such an antioxidant system in nodule proplastids or amyloplasts. Moreover, it is known that NTRC exerts functions that are independent of Prx reduction (Pulido et al., 2010), such as redox regulation of starch synthesis in photosynthetic and nonphotosynthetic tissues (Michalska et al., 2009). This also might be the case in nodules. Thus, it will be important to define the relative contribution of the NTR-Trx and ascorbate-glutathione pathways in peroxide removal and redox signaling, as both of them are likely to be operative in the cytosol, mitochondria, and plastids of legume nodules (Dalton et al., 1998; Iturbe-Ormaetxe et al., 2001). A comparison of the two pathways will need to consider the differences in abundance of ascorbate peroxidase (up to 0.9% of the total nodule soluble proteins; Dalton et al., 1998) and PrxIIB (low levels found here) in the nodule cytosol or in the responses of these enzymes to developmental or environmental cues. Besides the components of the NTR-Trx systems, we detected significant expression of the *FTRB* gene in nodules of *L. japonicus*. Considering that

FTR is an electron donor of some Trxs in the chloroplasts, an FTR-Trx system, comprising ferredoxin, FTR, and plastidic Trxs, might be also functional in nodules (Fig. 9). Although this system is beyond the scope of this work, it can be anticipated that FTR activity provides a means by which plastidic Trxs are regenerated in nodules because available information suggests that the NTRC enzyme is unable to reduce Trx in the chloroplasts and presumably in nonphotosynthetic plastids (Serrato et al., 2004; Traverso et al., 2008). The functionality of an FTR-Trx system in nodules is further supported by the presence of such a redox system in wheat endosperm amyloplasts (Balmer et al., 2006) and of ferredoxin-NADP reductase in *M. truncatula* nodules (Larrainzar et al., 2007).

MATERIALS AND METHODS

Plant Growth and Treatments

One-week-old seedlings of *Lotus japonicus* 'MG20' were inoculated with *Mesorhizobium loti* strain R7A, transferred to aerated hydroponic cultures lacking combined nitrogen (1:4 strength B&D nutrient solution), and grown under controlled environment conditions (Bustos-Sanmamed et al., 2011). Roots, leaves, nodules, and stems were harvested from 45-d-old plants (late vegetative stage), and flowers, pollen, pods, seeds, and embryos were collected from 60-d-old plants (pods of approximately 3.5 cm; late flowering-fruiting stage). Plant material was immediately flash-frozen in liquid nitrogen and stored at -85°C until use.

Plants grown in hydroponics were treated for 48 h with 50 μM ABA, GA₃, JA, indole-3-acetic acid, 1-aminocyclopropane-1-carboxylic acid, or CK (an equimolar mixture of kinetin and 6-benzyl-aminopurine) as described previously (Bustos-Sanmamed et al., 2011). To study the effects of CK on germination, seeds were surface-disinfected, stratified for 24 h in 0.5% agar plates at 4°C, and treated with the hormone for up to 48 h during germination. Control seeds were germinated on plates in the presence of 0.1 mM NaOH, which was also used to dissolve CK. The pH value was kept at 6.6 for both control and CK-treated seeds with 5 mM MES. The effect of NO on gene expression in roots of 15-d-old nonnodulated seedlings was studied by application of two NO-releasing compounds, SNAP (500 μM ; Sigma-Aldrich) and GSNO (250 μM ; Calbiochem), for up to 24 h. In the case of GSNO, a control treatment with 250 μM of glutathione was included because this physiological NO donor releases both NO and glutathione during the incubation period.

Gene Identification and Expression Profiling

Transformation-competent artificial chromosome and BAC genomic libraries of *L. japonicus* were screened with probes based on the cDNA sequences. The partial or full nucleotide sequences of the isolated transformation-competent artificial chromosome/BAC clones were determined according to the bridging shotgun method (Sato et al., 2008).

Total RNA was extracted from plant material with the RNAqueous isolation kit (Ambion), and cDNA was synthesized from 2 μg DNase-treated RNA with (dT)17 and Moloney murine leukemia virus reverse transcriptase (Promega). Quantitative RT-PCR analysis was performed with an iCycler iQ instrument using iQ SYBR-Green Supermix reagents (Bio-Rad) and gene-specific primers (Supplemental Table S1). The PCR program and other details were already described (Bustos-Sanmamed et al., 2011). The amplification efficiency of primers, calculated by serial dilutions of cDNAs, was $>80\%$. Gene expression levels were normalized with *ubiquitin*. The *PP2A* gene, encoding a subunit of the Ser/Thr protein phosphatase 2A (Czechowski et al., 2005), was used as an additional reference gene to verify that *ubiquitin* expression was not affected by any of the treatments.

Phylogenetic Analyses and Prediction of Subcellular Localization of Proteins

Multiple alignment of amino acid sequences was performed using the MegAlign-DNASTAR program (Lasergene) by the neighbor-joining ClustalW2

method, and the phylogenetic trees were built with 1,000 bootstrap replicates. Predictions of subcellular localizations were carried out using the MitoProt (<http://ihg.gsf.de/ihg/mitoprot.html>), TargetP v1.1 (<http://www.cbs.dtu.dk/services/TargetP/>), and PSORT (<http://psort.hgc.jp/>) programs. The nuclear localization of 1CPrx was predicted according to the presence of a bipartite nuclear localization signal in the C-terminal region of the amino acid sequence (Stacy et al., 1999).

Immunoblots

For protein extraction, frozen plant organs were pulverized with liquid nitrogen and homogenized in ice-cold extraction buffer (1 mL per 0.2 g fresh weight) containing 50 mM potassium phosphate (pH 7), 5 mM dithiothreitol, 1% Triton X-100, and complete protease inhibitor cocktail (Roche). The extracts were cleared by centrifugation and stored at -20°C if necessary. Total proteins were separated by 12% SDS-PAGE and blotted onto polyvinylidene difluoride membranes, and immunoblots were carried out as described elsewhere (Matamoros et al., 2010). The sources of the polyclonal antibodies were as follows: 2CPrx, PrxQ, PrxIIC, and PrxIIF of *Arabidopsis thaliana*; (Horling et al., 2003); 1CPrx of barley (*Hordeum vulgare*; Stacy et al., 1999); Trxh1 to Trxh5 of poplar (*Populus* spp.; Gelhaye et al., 2004); NTRA/B of wheat (*Triticum aestivum*; Serrato et al., 2002); and NTRC of rice (*Oryza sativa*; Serrato et al., 2004). The PrxIIC antibody was used to detect LjPrxIIB as this antibody recognizes the two cytosolic PrxII isoforms but does not cross-react with plastidic PrxIIE (Horling et al., 2003) or mitochondrial PrxIIF (Finkemeier et al., 2005).

Proteomic Analyses of Nodules

All proteomic analyses were performed at the University of Vienna using a gel-free protocol based on liquid chromatography and tandem mass spectrometry as outlined in detail by Larrainzar et al. (2007) and Hoehenwarter and Wienkoop (2010). After mass spectrometry analysis, raw files were searched against the Dana-Farber Cancer Institute (DFCI) *Lotus* Gene Index (6.0), *Medicago* Gene Index (11.0), or Bean Gene Index (4.0) databases using the Sequest algorithm. For identification and spectral count based data, the matrix generation Proteome Discoverer (version 1.1; Thermo Fisher Scientific) was used. A decoy database enabled false positive rate analysis. Only high confidence peptides (false positive rate $< 0.1\%$) better than 5 ppm precursor mass accuracy per protein passed criteria.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure S1. Phylogenetic analysis of Prx proteins from higher plants.

Supplemental Figure S2. Effect of NO donors on *Lj1CPrx* gene expression in roots.

Supplemental Figure S3. Sequence alignment of Trxh1 proteins of model legumes.

Supplemental Figure S4. Phylogenetic analysis of Trx proteins of model plants.

Supplemental Figure S5. Phylogenetic analysis of NTR proteins of higher plants.

Supplemental Figure S6. Sequence alignment of NTRA/B proteins of plants.

Supplemental Figure S7. Expression analysis of the *LjFTRB* gene.

Supplemental Table S1. Primers used for RT-PCR analyses.

ACKNOWLEDGMENTS

We thank Stefanie Wienkoop and Christiana Staudinger (University of Vienna) for proteomic analyses of nodules and two anonymous reviewers for helpful comments. A.T.M. was the recipient of a postdoctoral contract (JAE-CSIC program).

Received March 29, 2011; accepted May 6, 2011; published May 11, 2011.

LITERATURE CITED

- Aalen RB, Opsahl-Ferstad HG, Linnestad C, Olsen OA (1994) Transcripts encoding an oleosin and a dormancy related protein are present in both the aleurone layer and the embryo of developing barley (*Hordeum vulgare* L.) seeds. *Plant J* 5: 385–396
- Alkhalifiou F, Renard M, Frendo P, Keichinger C, Meyer Y, Gelhaye E, Hirasawa M, Knaff DB, Ritzenhaler C, Montrichard F (2008) A novel type of thioredoxin dedicated to symbiosis in legumes. *Plant Physiol* 148: 424–435
- Alkhalifiou F, Renard M, Montrichard F (2007) Unique properties of NADP-thioredoxin reductase C in legumes. *J Exp Bot* 58: 969–978
- Balmer Y, Vensel WH, Cai N, Manieri W, Schürmann P, Hurkman WJ, Buchanan BB (2006) A complete ferredoxin/thioredoxin system regulates fundamental processes in amyloplasts. *Proc Natl Acad Sci USA* 103: 2988–2993
- Barranco-Medina S, Krell T, Bernier-Villamor L, Sevilla F, Lázaro JJ, Dietz KJ (2008) Hexameric oligomerization of mitochondrial peroxiredoxin PrxIIIF and formation of an ultrahigh affinity complex with its electron donor thioredoxin Trx-o. *J Exp Bot* 59: 3259–3269
- Barranco-Medina S, Krell T, Finkemeier I, Sevilla F, Lázaro JJ, Dietz KJ (2007) Biochemical and molecular characterization of the mitochondrial peroxiredoxin PsPrxII F from *Pisum sativum*. *Plant Physiol Biochem* 45: 729–739
- Becana M, Matamoros MA, Udvardi M, Dalton DA (2010) Recent insights into antioxidant defenses of legume root nodules. *New Phytol* 188: 960–976
- Besson-Bard A, Pugin A, Wendehenne D (2008) New insights into nitric oxide signaling in plants. *Annu Rev Plant Biol* 59: 21–39
- Bréhélin C, Meyer EH, de Souris JP, Bonnard G, Meyer Y (2003) Resemblance and dissemblance of Arabidopsis type II peroxiredoxins: similar sequences for divergent gene expression, protein localization, and activity. *Plant Physiol* 132: 2045–2057
- Bustos-Sanmamed P, Tovar-Méndez A, Crespi M, Sato S, Tabata S, Becana M (2011) Regulation of nonsymbiotic and truncated hemoglobin genes of *Lotus japonicus* in plant organs and in response to nitric oxide and hormones. *New Phytol* 189: 765–776
- Carimi F, Zottini M, Costa A, Cattelan I, De Michele R, Terzi M, Lo Schiavo F (2005) NO signalling in cytokinin-induced programmed cell death. *Plant Cell Environ* 28: 1171–1178
- Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiol* 139: 5–17
- Dai S, Johansson K, Miginiac-Maslow M, Schürmann P, Eklund H (2004) Structural basis of redox signaling in photosynthesis: structure and function of ferredoxin:thioredoxin reductase and target enzymes. *Photosynth Res* 79: 233–248
- Dalton DA, Joyner SL, Becana M, Iturbe-Ormaetxe I, Chatfield JM (1998) Antioxidant defenses in the peripheral cell layers of legume root nodules. *Plant Physiol* 116: 37–43
- del Río LA, Corpas FJ, Sandalio LM, Palma JM, Gómez M, Barroso JB (2002) Reactive oxygen species, antioxidant systems and nitric oxide in peroxisomes. *J Exp Bot* 53: 1255–1272
- den Boer BGW, Murray JAH (2000) Triggering the cell cycle in plants. *Trends Cell Biol* 10: 245–250
- Díaz Vivancos P, Dong Y, Ziegler K, Markovic J, Pallardó FV, Pellny TK, Verrier PJ, Foyer CH (2010) Recruitment of glutathione into the nucleus during cell proliferation adjusts whole-cell redox homeostasis in *Arabidopsis thaliana* and lowers the oxidative defence shield. *Plant J* 64: 825–838
- Diet A, Abbas K, Bouton C, Guillon B, Tomasello F, Fourquet S, Toledano MB, Drapier JC (2007) Regulation of peroxiredoxins by nitric oxide in immunostimulated macrophages. *J Biol Chem* 282: 36199–36205
- Dietz KJ, Jacob S, Oelze ML, Laxa M, Tognetti V, de Miranda SM, Baier M, Finkemeier I (2006) The function of peroxiredoxins in plant organellar redox metabolism. *J Exp Bot* 57: 1697–1709
- Dietz KJ, Pfannschmidt T (2011) Novel regulators in photosynthetic redox control of plant metabolism and gene expression. *Plant Physiol* 155: 1477–1485
- Finkemeier I, Goodman M, Lamkemeyer P, Kandlbinder A, Sweetlove LJ, Dietz KJ (2005) The mitochondrial type II peroxiredoxin F is essential for redox homeostasis and root growth of *Arabidopsis thaliana* under stress. *J Biol Chem* 280: 12168–12180
- Foyer CH, Noctor G (2005) Oxidant and antioxidant signalling in plants: a re-evaluation of the concept of oxidative stress in a physiological context. *Plant Cell Environ* 28: 1056–1071
- Gama F, Keech O, Eymery F, Finkemeier I, Gelhaye E, Gardeström P, Dietz KJ, Rey P, Jacquot JP, Rouhier N (2007) The mitochondrial type II peroxiredoxin from poplar. *Physiol Plant* 129: 196–206
- Gelhaye E, Rouhier N, Jacquot JP (2004) The thioredoxin *h* system of higher plants. *Plant Physiol Biochem* 42: 265–271
- Groten K, Dutilleul C, van Heerden PDR, Vanacker H, Bernard S, Finkemeier I, Dietz KJ, Foyer CH (2006) Redox regulation of peroxiredoxin and proteinases by ascorbate and thiols during pea root nodule senescence. *FEBS Lett* 580: 1269–1276
- Haslekås C, Viken MK, Grini PE, Nygaard V, Nordgard SH, Meza TJ, Aalen RB (2003) Seed 1-cysteine peroxiredoxin antioxidants are not involved in dormancy, but contribute to inhibition of germination during stress. *Plant Physiol* 133: 1148–1157
- Hirsch AM (1992) Developmental biology of legume nodulation. *New Phytol* 122: 211–237
- Hoehenwarter W, Wienkoop S (2010) Spectral counting robust on high mass accuracy mass spectrometers. *Rapid Commun Mass Spectrom* 24: 3609–3614
- Horling F, Lamkemeyer P, König J, Finkemeier I, Kandlbinder A, Baier M, Dietz KJ (2003) Divergent light-, ascorbate-, and oxidative stress-dependent regulation of expression of the peroxiredoxin gene family in Arabidopsis. *Plant Physiol* 131: 317–325
- Iturbe-Ormaetxe I, Matamoros MA, Rubio MC, Dalton DA, Becana M (2001) The antioxidants of legume nodule mitochondria. *Mol Plant Microbe Interact* 14: 1189–1196
- Laloi C, Rayapuram N, Chartier Y, Grienberger JM, Bonnard G, Meyer Y (2001) Identification and characterization of a mitochondrial thioredoxin system in plants. *Proc Natl Acad Sci USA* 98: 14144–14149
- Lamattina L, García-Mata C, Graziano M, Pagnussat G (2003) Nitric oxide: the versatility of an extensive signal molecule. *Annu Rev Plant Biol* 54: 109–136
- Lamkemeyer P, Laxa M, Collin V, Li W, Finkemeier I, Schöttler MA, Holtkamp V, Tognetti VB, Issakidis-Bourguet E, Kandlbinder A, et al (2006) Peroxiredoxin Q of *Arabidopsis thaliana* is attached to the thylakoids and functions in context of photosynthesis. *Plant J* 45: 968–981
- Larrainzar E, Wienkoop S, Weckwerth W, Ladrera R, Arrese-Igor C, González EM (2007) *Medicago truncatula* root nodule proteome analysis reveals differential plant and bacteroid responses to drought stress. *Plant Physiol* 144: 1495–1507
- Lee MY, Shin KH, Kim YK, Suh JY, Gu YY, Kim MR, Hur YS, Son O, Kim JS, Song E, et al (2005) Induction of thioredoxin is required for nodule development to reduce reactive oxygen species levels in soybean roots. *Plant Physiol* 139: 1881–1889
- Marino D, González EM, Frendo P, Puppo A, Arrese-Igor C (2007) NADPH recycling systems in oxidative stressed pea nodules: a key role for the NADP⁺-dependent isocitrate dehydrogenase. *Planta* 225: 413–421
- Marty L, Siala W, Schwarzländer M, Fricker MD, Wirtz M, Sweetlove LJ, Meyer Y, Meyer AJ, Reichheld JP, Hell R (2009) The NADPH-dependent thioredoxin system constitutes a functional backup for cytosolic glutathione reductase in *Arabidopsis*. *Proc Natl Acad Sci USA* 106: 9109–9114
- Matamoros MA, Loscos J, Dietz KJ, Aparicio-Tejo PM, Becana M (2010) Function of antioxidant enzymes and metabolites during maturation of pea fruits. *J Exp Bot* 61: 87–97
- Meyer Y, Buchanan BB, Vignols F, Reichheld JP (2009) Thioredoxins and glutaredoxins: unifying elements in redox biology. *Annu Rev Genet* 43: 335–367
- Michalska J, Zauber H, Buchanan BB, Cejudo FJ, Geigenberger P (2009) NTRC links built in thioredoxin to light and sucrose in regulating starch synthesis in chloroplasts and amyloplasts. *Proc Natl Acad Sci USA* 106: 9908–9913
- Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7: 405–410
- Moon JC, Jang HH, Chae HB, Lee JR, Lee SY, Jung YJ, Shin MR, Lim HS, Chung WS, Yun DJ, et al (2006) The C-type *Arabidopsis* thioredoxin

- reductase ANTR-C acts as an electron donor to 2-Cys peroxiredoxins in chloroplasts. *Biochem Biophys Res Commun* **348**: 478–484
- Mowla SB, Thomson JA, Farrant JM, Mundree SG** (2002) A novel stress-inducible antioxidant enzyme identified from the resurrection plant *Xerophyta viscosa* Baker. *Planta* **215**: 716–726
- Neill S, Barros R, Bright J, Desikan R, Hancock J, Harrison J, Morris P, Ribeiro D, Wilson I** (2008) Nitric oxide, stomatal closure, and abiotic stress. *J Exp Bot* **59**: 165–176
- O'Neill JS, van Ooijen G, Dixon LE, Troein C, Corellou F, Bouget FY, Reddy AB, Millar AJ** (2011) Circadian rhythms persist without transcription in a eukaryote. *Nature* **469**: 554–558
- Pena-Ahumada A, Kahmann U, Dietz KJ, Baier M** (2006) Regulation of peroxiredoxin expression versus expression of Halliwell-Asada-Cycle enzymes during early seedling development of *Arabidopsis thaliana*. *Photosynth Res* **89**: 99–112
- Pérez-Ruiz JM, Spínola MC, Kirchsteiger K, Moreno J, Sahrawy M, Cejudo FJ** (2006) Rice NTRC is a high-efficiency redox system for chloroplast protection against oxidative damage. *Plant Cell* **18**: 2356–2368
- Pulido P, Cazalis R, Cejudo FJ** (2009) An antioxidant redox system in the nucleus of wheat seed cells suffering oxidative stress. *Plant J* **57**: 132–145
- Pulido P, Spínola MC, Kirchsteiger K, Guinea M, Pascual MB, Sahrawy M, Sandalio LM, Dietz KJ, González M, Cejudo FJ** (2010) Functional analysis of the pathways for 2-Cys peroxiredoxin reduction in *Arabidopsis thaliana* chloroplasts. *J Exp Bot* **61**: 4043–4054
- Puntarulo S, Sánchez RA, Boveris A** (1988) Hydrogen peroxide metabolism in soybean embryonic axes at the onset of germination. *Plant Physiol* **86**: 626–630
- Puppo A, Groten K, Bastian F, Carzaniga R, Soussi M, Lucas MM, de Felipe MR, Harrison J, Vanacker K, Foyer CH** (2005) Legume nodule senescence: roles for redox and hormone signalling in the orchestration of the natural aging process. *New Phytol* **165**: 683–701
- Reichheld JP, Meyer E, Khafif M, Bonnard G, Meyer Y** (2005) AtNTRB is the major mitochondrial thioredoxin reductase in *Arabidopsis thaliana*. *FEBS Lett* **579**: 337–342
- Renard M, Alkhalifioui F, Schmitt-Keichinger C, Ritzenthaler C, Montrichard F** (2011) Identification and characterization of thioredoxin h isoforms differentially expressed in germinating seeds of the model legume *Medicago truncatula*. *Plant Physiol* **155**: 1113–1126
- Romanov GA** (2009) How do cytokinins affect the cell? *Russ J Plant Physiol* **56**: 268–290
- Romero-Puertas MC, Laxa M, Mattè A, Zaninotto F, Finkemeier I, Jones AME, Perazzoli M, Vandelle E, Dietz KJ, Delledonne M** (2007) S-Nitrosylation of peroxiredoxin IIE promotes peroxynitrite-mediated tyrosine nitration. *Plant Cell* **19**: 4120–4130
- Rouhier N, Gelhaye E, Jacquot JP** (2002) Glutaredoxin-dependent peroxiredoxin from poplar: protein-protein interaction and catalytic mechanism. *J Biol Chem* **277**: 13609–13614
- Rouhier N, Gelhaye E, Gualberto JM, Jordy MN, De Fay E, Hirasawa M, Duplessis S, Lemaire SD, Frey P, Martin F, et al** (2004) Poplar peroxiredoxin Q. A thioredoxin-linked chloroplast antioxidant functional in pathogen defense. *Plant Physiol* **134**: 1027–1038
- Rouhier N, Jacquot JP** (2005) The plant multigenic family of thiol peroxidases. *Free Radic Biol Med* **38**: 1413–1421
- Sato S, Nakamura Y, Kaneko T, Asamizu E, Kato T, Nakao M, Sasamoto S, Watanabe A, Ono A, Kawashima K, et al** (2008) Genome structure of the legume *Lotus japonicus*. *DNA Res* **15**: 227–239
- Schürmann P, Jacquot JP** (2000) Plant thioredoxin systems revisited. *Annu Rev Plant Physiol Plant Mol Biol* **51**: 371–400
- Serrato AJ, Pérez-Ruiz JM, Cejudo FJ** (2002) Cloning of thioredoxin h reductase and characterization of the thioredoxin reductase-thioredoxin h system from wheat. *Biochem J* **367**: 491–497
- Serrato AJ, Pérez-Ruiz JM, Spínola MC, Cejudo FJ** (2004) A novel NADPH thioredoxin reductase, localized in the chloroplast, which deficiency causes hypersensitivity to abiotic stress in *Arabidopsis thaliana*. *J Biol Chem* **279**: 43821–43827
- Serrato AJ, Guilleminot J, Meyer Y, Vignols F** (2008) AtCXXS: atypical members of the *Arabidopsis thaliana* thioredoxin h family with a remarkably high disulfide isomerase activity. *Physiol Plant* **133**: 611–622
- Stacy RAP, Nordeng TW, Culiáñez-Macià FA, Aalen RB** (1999) The dormancy-related peroxiredoxin anti-oxidant, PER1, is localized to the nucleus of barley embryo and aleurone cells. *Plant J* **19**: 1–8
- Traverso JA, Vignols F, Cazalis R, Serrato AJ, Pulido P, Sahrawy M, Meyer Y, Cejudo FJ, Chueca A** (2008) Immunocytochemical localization of *Pisum sativum* TRXs f and m in non-photosynthetic tissues. *J Exp Bot* **59**: 1267–1277
- Tripathi BN, Bhatt I, Dietz KJ** (2009) Peroxiredoxins: a less studied component of hydrogen peroxide detoxification in photosynthetic organisms. *Protoplasma* **235**: 3–15
- Tun NN, Holk A, Scherer GFE** (2001) Rapid increase of NO release in plant cell cultures induced by cytokinin. *FEBS Lett* **509**: 174–176
- Valderrama R, Corpas FJ, Carreras A, Fernández-Ocaña A, Chaki M, Luque F, Gómez-Rodríguez MV, Colmenero-Varea P, del Río LA, Barroso JB** (2007) Nitrosative stress in plants. *FEBS Lett* **581**: 453–461
- Vieira Dos Santos C, Rey P** (2006) Plant thioredoxins are key actors in the oxidative stress response. *Trends Plant Sci* **11**: 329–334
- Zavaleta-Mancera HA, López-Delgado H, Loza-Tavera H, Mora-Herrera M, Trevilla-García C, Vargas-Suárez M, Ougham H** (2007) Cytokinin promotes catalase and ascorbate peroxidase activities and preserves the chloroplast integrity during dark-senescence. *J Plant Physiol* **164**: 1572–1582