

# Genomic Analysis of *Xanthomonas oryzae* Isolates from Rice Grown in the United States Reveals Substantial Divergence from Known *X. oryzae* Pathovars<sup>∇†</sup>

L. R. Triplett,<sup>1</sup> J. P. Hamilton,<sup>2</sup> C. R. Buell,<sup>2</sup> N. A. Tisserat,<sup>1</sup> V. Verdier,<sup>1,3</sup> F. Zink,<sup>1</sup> and J. E. Leach<sup>1\*</sup>

Department of Bioagricultural Sciences and Pest Management, Colorado State University, Fort Collins, Colorado 80523<sup>1</sup>;  
Department of Plant Biology, Michigan State University, East Lansing, Michigan 48824<sup>2</sup>; and  
UMR RPB, IRD-CIRAD-UM2, BP 64501, 34394 Montpellier Cedex 5, France<sup>3</sup>

Received 6 January 2011/Accepted 9 April 2011

**The species *Xanthomonas oryzae* is comprised of two designated pathovars, both of which cause economically significant diseases of rice in Asia and Africa. Although *X. oryzae* is not considered endemic in the United States, an *X. oryzae*-like bacterium was isolated from U.S. rice and southern cutgrass in the late 1980s. The U.S. strains were weakly pathogenic and genetically distinct from characterized *X. oryzae* pathovars. In the current study, a draft genome sequence from two U.S. *Xanthomonas* strains revealed that the U.S. strains form a novel clade within the *X. oryzae* species, distinct from all strains known to cause significant yield loss. Comparative genome analysis revealed several putative gene clusters specific to the U.S. strains and supported previous reports that the U.S. strains lack transcriptional activator-like (TAL) effectors. In addition to phylogenetic and comparative analyses, the genome sequence was used for designing robust U.S. strain-specific primers, demonstrating the usefulness of a draft genome sequence in the rapid development of diagnostic tools.**

The species *Xanthomonas oryzae* is comprised of pathovars *oryzae* and *oryzicola*, the causative agents of bacterial leaf blight (BLB) and bacterial leaf streak (BLS) on rice, respectively (27). Although the two pathovars are closely related, BLB is a vascular disease characterized by marginal leaf lesions, while BLS affects parenchyma cells and results in leaf streaking. Both pathovars can cause substantial losses to rice production (27). *X. oryzae* has been designated a USDA select agent in the United States, and movement is restricted by several international quarantines (25, 29).

The finished genomes of three Asian *X. oryzae* pv. *oryzae* strains and one *X. oryzae* pv. *oryzicola* strain are available, facilitating in-depth comparative genomic analyses (22, 28, 35) (GenBank accession no. AAQN01000001). Whole-genome alignment revealed that the sequenced *X. oryzae* pv. *oryzae* strains, MAFF 311018, KACC 10331, and PXO99A, are very closely related (24, 35). *X. oryzae* pv. *oryzicola* clusters with the *X. oryzae* group, forming a branch distinct from *X. oryzae* pv. *oryzae* strains (24, 35). All the *X. oryzae* genomes are characterized by large numbers of insertion sequence (IS) elements, the major contributors to sequence diversity within the species (28, 35), and by various numbers of secreted transcriptional activator-like (TAL) effectors required for full virulence (35, 42). The African *X. oryzae* pv. *oryzae* strains are different from Asian strains and more closely related to Asian *X. oryzae* pv. *oryzicola*. A specific and intriguing feature of African *X. oryzae* pv. *oryzae* strains is that the genome contains a smaller number

of TAL effector and IS elements than the Asian strains (8). Bacterial blight has also been observed in South America (12, 23); strains isolated in Colombia are closely related to Asian strains (15, 17).

Although *X. oryzae* is not historically considered indigenous to the United States (27), strains of a yellow bacterium causing mild BLB-like symptoms were collected from rice fields in Texas and Louisiana in 1987 (17). The bacterium was classified as *X. oryzae* based on serological tests and fatty acid profiling (17). However, the U.S. strains were weakly virulent on rice (17), and restriction enzyme length polymorphism (RFLP) fingerprint profiles suggested that they were genetically distinct from *X. oryzae* pv. *oryzae* and *oryzicola* (34). Most of the U.S. strains were shown to carry one or more plasmids (41). These findings led the authors to speculate that U.S. *Xanthomonas* strains were highly divergent from those found in Asia and had not entered the United States in the recent past. Although the *X. oryzae*-like strains were collected in subsequent years in U.S. fields of rice and the native southern cutgrass (*Leersia hexandra*) (9), there were no reports of any significant yield loss in the field, and there have been no subsequent reports of *X. oryzae*-related problems in the United States. However, due to the numerous global regulatory measures aimed at preventing the spread of bacterial pathogens, improved characterization of any U.S. strains of *X. oryzae* will be critical for development of diagnostic tools and crop protection.

Recent technological advances have substantially increased the speed and cost-efficiency of genome sequencing. However, the final steps of completion, such as gap closure and analysis of repetitive elements, remain time-consuming and expensive. As more bacterial species are sequenced, a draft sequence of genomes closely related to a finished reference genome will serve as a valuable resource for comparative analysis. Draft genome sequences have been used for rapid development of

\* Corresponding author. Mailing address: Department of Bioagricultural Sciences and Pest Management, Colorado State University, Campus Delivery 1177, Fort Collins, CO 80523-1177. Phone: (970) 491-2924. Fax: (970) 491-3862. E-mail: jan.leach@colostate.edu.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

∇ Published ahead of print on 22 April 2011.

TABLE 1. Strains used in this study

Species	Strain(s) <sup>a</sup>	Country
<i>X. oryzae</i> , unknown pathovar	X8-1A, X11-5A, Ru87-7, X4-2C, X7-2D, X212-3-1, X37-2, X1-8, X44-D, X211-2, X45-A1, X13-5C, X54-A1, X7-5A, X211-1 (C. Gonzalez), X207-A1 (C. Gonzalez)	United States
<i>X. oryzae</i> pv. <i>oryzae</i>	PXO99A, PXO86 Xoo475-304 Xoo94 Xoo228 CFBP 1947, CFBP 1948 MAI1, MAI2 NAI1, NAI8 BAI3, BAI4 CIAT 1185 XOO4 IXO51 MXO49	Philippines Japan China Korea Cameroon Mali Niger Burkina Faso Colombia Thailand India Malaysia
<i>X. oryzae</i> pv. <i>oryzicola</i>	BLS256, BLS177, BLS333 MAI8, MAI10	Philippines Mali
<i>X. campestris</i> pv. <i>leersiae</i>	NCPBP 4346 (N. Parkinson)	China
<i>Pseudomonas syringae</i> pv. <i>syringae</i>	ICPB PS296 (L. E. Claffin)	
<i>Ralstonia solanacearum</i>	K60 (B. A. Hetrick)	
<i>X. campestris</i> pv. <i>campestris</i>	Xcc X1g10 (L. E. Claffin)	
<i>X. axonopodis</i> pv. <i>vesicatoria</i>	85-10 (C. Gonzalez)	United States
<i>Xanthomonas translucens</i> pv. <i>hordei</i>	2181 (L. E. Claffin)	United States
<i>X. vasicola</i>		

<sup>a</sup> Strains are from the collections of J. E. Leach and V. Verdier unless otherwise noted.

diagnostic tools, comparison of gene repertoires, and identification of putative virulence factors (2, 4, 11).

In this work, we characterized the phylogeny and gene content of two U.S. *Xanthomonas* strains by analyzing high-quality draft genome sequence data. Comparative genomics and multilocus sequence typing revealed that the U.S. *Xanthomonas* strains, while part of the *X. oryzae* species, form a group substantially divergent from a clade formed by *X. oryzae* pv. *oryzae*, *X. oryzae* pv. *oryzicola*, and *Xanthomonas campestris* pv. *leersiae*. Genome-based diagnostic primers were developed and tested to identify this novel clade.

#### MATERIALS AND METHODS

**Bacterial strains, genome sequencing, and genome assembly methods.** Table 1 lists bacterial strains used for whole-genome and single-gene sequencing and for primer testing. DNA was extracted using the genomic DNA extraction kit (Qiagen, Valencia, CA). The DNA was sheared, a library was constructed (Illumina, San Diego, CA), and the DNA was end sequenced (76-bp paired-end reads) using an Illumina Genome Analyzer II (Illumina, San Diego, CA) at the Michigan State University Research Technology Support Facility. The paired-end reads were trimmed to 40 bp to remove low-quality regions at the 3' end of the reads. A custom PerlScript was used to clean and remove reads with low-complexity and low-quality regions. Low-quality regions were defined as an average quality score of  $\leq 20$  over a 10-bp window along the read or  $\geq 2$  N bases in the read. The low-complexity threshold was defined as  $>85\%$  of the base composition composed of a single nucleotide. In cases where one read passed the cleaning process and the paired end failed, the good read was retained in a single read file for use in the assembly.

The Velvet short read assembler (version 0.7.53; PubMed identifier 18349386) (43) was used to assemble the genomes in conjunction with the VelvetOptimiser.pl PerlScript provided with Velvet. VelvetOptimiser.pl automates the searching of hash length, expected coverage, and coverage cutoff parameter space to generate an optimal assembly. The program was provided the cleaned paired-end reads, single reads from the cleaning process described above, an estimated insert size of 300 bp, and a minimum contig length of 200 bp. The Velvet parameters used in the final assembly for X11-5A were a hash length of 31, expected coverage of 19.58-fold, and a k-mer coverage cutoff of 9.79-fold. The hash length for X8-1A was 31, the expected coverage was 18.02-fold, and the k-mer coverage cutoff was

9.01-fold. Sanger sequencing of individual *X. oryzae* genes was performed at the Proteomics and Metabolomics Facility at Colorado State University.

Contigs from the Velvet assembly were assembled into a pseudomolecule, or draft chromosome sequence, using the genome of PXO99A as a reference sequence; contigs in each strain not mapping to PXO99A were concatenated into a second pseudomolecule named "assembly 2." Open reading frames (ORFs) were predicted and annotated using the Institute for Genome Sciences annotation engine (<http://ae.igs.umaryland.edu/cgi/index.cgi>).

**Pathogenicity assays.** Seedlings of the rice *Oryza sativa* subsp. *japonica* cv. Lemont and the *O. sativa* subsp. *indica* cv. IR64 were grown for 21 days under controlled conditions (28°C, 80% humidity) in a growth chamber. Strains PXO99A, X11-5A, and X8-1A were grown for 48 h on plates of nutrient agar (BD, Franklin Lakes, NJ) at 28°C and suspended to  $10^9$  CFU/ml in water. Plants were inoculated using the clip inoculation method (18). Lesion length and length of leaf curling were measured at several time points over 10 to 16 days post-inoculation (dpi). PXO99A caused rapid leaf curling from clipped ends of leaves well in advance of lesion development, whereas X11-5A and X8-1A caused distinct lesions and very little leaf curling. Therefore, the extent of leaf curling caused by PXO99A was compared to lesion length caused by the two U.S. strains. Additionally four accessions of *Oryza glaberrima* (TOG5497, TOG5672, TOG6202, TOG6308) and four cultivars of *O. sativa* (*O. sativa* subsp. *japonica* cv. Azucena, Nipponbare, and Curinga and *O. sativa* subsp. *indica* cv. TN1) were grown for 45 days in a greenhouse and clip-inoculated with strains X8-1A and X11-5A and *X. oryzae* pv. *oryzae* strain BAI3 (Burkina Faso) as a control. Lesion length (mm) was measured 16 and 21 dpi on two leaves on each of five replicate plants.

**Multilocus sequence typing (MLST) and phylogenetic analyses.** The ORFs of the genes *gyrB*, *fusA*, *dnaK*, *glnA*, *groEL*, *atpD*, *gapA*, *recA*, and *efp* were identified in genomic sequences of X11-5A, X8-1A, MAI-1, NAI-8, and other *Xanthomonas* strains (GenBank accession numbers: *X. oryzae* pv. *oryzicola*, AAQN01000001; *X. oryzae* pv. *oryzae*, CP000967, AE013598, and AP008229; *Xanthomonas vasicola*, ACHT00000000; and *X. campestris* pv. *campestris*, CP000050). For MLST of unsequenced strains, *fusA*, *gyrB*, and *gapA* were amplified from selected *X. oryzae* strains using published primer sequences (1), and PCR products were sequenced. Other *Xanthomonas fusA*, *gyrB*, and *gapA* gene sequences were downloaded at the Plant Associated and Environmental Microbes Database (1). Sequences were aligned by ClustalW (21) with the Mega4 program (39). Phylogenetic analyses were performed using the MrBayes program for Bayesian analysis (16), using the general time-reversible model with inverse-gamma rates of evolution for 100,000 generations. Phylogenetic trees were drawn and formatted in Mega4.

TABLE 2. General characteristics of draft genomes of U.S. *X. oryzae* strains and completed *X. oryzae* genomes

Strain	Pathovar	Country of origin	Total sequence length (bp)	No. of predicted genes	% GC	% ANI with:		Source or reference
						PXO99A	BLS256	
X11-5A	Undesignated	United States (Texas)	4,641,765 <sup>a</sup>	4,655	64	97	97.11	This study
X8-1A	Undesignated	United States (Louisiana)	4,679,331 <sup>a</sup>	4,886	64	97.04	97.06	This study
PXO99A	<i>X. oryzae</i> pv. <i>oryzae</i>	Philippines	5,240,075 <sup>b</sup>	5,083	63.6	100	97.8	35
KACC	<i>X. oryzae</i> pv. <i>oryzae</i>	Korea	4,941,439 <sup>b</sup>	4,637	63.7	99.49	97.8	22
BLS256	<i>X. oryzae</i> pv. <i>oryzicola</i>	Philippines	4,831,739 <sup>b</sup>	4,614	64	97.76	100	GenBank accession no. AAQN00000000

<sup>a</sup> Metrics for genome reported in Table S1 in the supplemental material.

<sup>b</sup> Single contig representing the circular chromosome.

**Comparative genome analyses.** Average nucleotide identity (ANI) values were calculated using the SpeciesJ program with the default settings for BLAST-based analysis (33). Briefly, the program divides a query genome into 1,020-nucleotide fragments and then calculates the average percentage of identical nucleotides shared with the corresponding fragments in a reference genome. Unique fragments with no homology to the reference genome are not included in the analysis. Strains with ANI values of 95 to 96 or greater are typically considered to be within the same species (10, 33). PXO99A and BLS256 genomes were used as the query against the target *X. oryzae* genomes of strains KACC, MAFF, PXO99A, and BLS256 and the rough assemblies of U.S. strains X11-5A and X8-1A.

The number of common ORFs shared between species was determined using BLASTN (3) searches of predicted ORFs in the genomes of *X. oryzae* strains KACC, MAFF, PXO99A, and BLS256 and *X. campestris* strain 8004 against the draft assemblies of U.S. strains X11-5A and X8-1A. The criterion of 70% identity over 150 or more nucleotides was used to classify ORFs as present or absent from strains. Vmatch (www.vmatch.de) was used to search unassembled short reads of U.S. and African strains for evidence of TAL effector sequences.

**Primer design and testing.** The program ePrimer3 from the EMBOSS package (32) was used to design primers within U.S. strain-specific ORFs or at the junction of strain-specific regions with conserved regions. The PrimerSearch program (32) was used to confirm the specificity of the primers (cutoff of 20% mismatch) against the sequenced *X. oryzae* genomes named above, as well as the genome sequences of *Xanthomonas albilineans*, *Xanthomonas axonopodis* pv. citri, *X. axonopodis* pv. vesicatoria, *X. campestris* pv. campestris, *X. vasicola*, and *Xylella fastidiosa* 9a5c (GenBank accession numbers FP565176, AE008923, AM039952, CP000050, ACHT00000000, and AE003849, respectively). Fifteen primer pairs with different predicted locations and product sizes were chosen for testing. Each 25- $\mu$ l PCR mixture contained 1.5 mM MgCl<sub>2</sub>, 0.2  $\mu$ M forward and reverse primers, 1 $\times$  PCR buffer (Invitrogen, Carlsbad, CA), 0.2 mM each deoxynucleoside triphosphate (dNTP), 0.5 units *Taq* polymerase (Invitrogen, Carlsbad, CA), and 10 ng template DNA. After an initial denaturing step, PCR was conducted for 28 cycles of 30 s at 94°C, 30 s at 55°C, and 70 s at 72°C. Primers were tested against DNA from all strains listed in Table 2.

**Nucleotide sequence accession numbers.** This whole genome shotgun project was deposited at GenBank under accession numbers AFHK000000000 (X11-5A) and AFHL000000000 (X8-1A). Assemblies and annotation data can be downloaded from the Comprehensive Phytopathogen Genomics Resource at [http://cpgr.plantbiology.msu.edu/us\\_xo\\_anno\\_download.shtml](http://cpgr.plantbiology.msu.edu/us_xo_anno_download.shtml). Illumina reads were deposited in the NCBI Sequence Read Archive under accession number SRP00669. Novel housekeeping gene sequences used in this study have been deposited in GenBank under accession numbers JF830787 to JF830795.

## RESULTS

**Pathogenicity testing and selection of U.S. *X. oryzae* strains for sequence characterization.** U.S. strains of *Xanthomonas* were previously identified as *X. oryzae* and divided into groups based on biocin production and plasmid profiling (41). To determine if disease development varied between individual U.S. *X. oryzae* strains and to identify virulent strains for sequencing, 16 U.S. strains from various groups were tested for virulence on *Oryza sativa* subsp. *japonica* cv. Lemont. Lesion length was monitored over 16 days. Inoculation with U.S. *X.*

*oryzae* strains resulted in mean lesion lengths of between 2 and 10 cm for each individual strain (data not shown). The two strains causing the largest lesions in the initial screen were X11-5A and X8-1A, isolated from fields in Texas and in Louisiana, respectively (17). Based on this observation, X11-5A and X8-1A were selected for genome sequencing. Additional clip experiments confirmed that X11-5A and X8-1A caused lesions on *O. sativa* subsp. *japonica* cv. Lemont, although the lesions were smaller and slower to form than those caused by the highly virulent Asian *X. oryzae* pv. *oryzae* strain PXO99A (Fig. 1A). *O. sativa* subsp. *indica* cv. IR64 is a widely grown cultivar known to be susceptible to most Asian strains of *X. oryzae*. No lesions or other symptoms were observed on IR64




FIG. 1. Symptom development on leaves of the rice cultivars *Oryza sativa* subsp. *japonica* cv. Lemont (A) and *O. sativa* subsp. *indica* cv. IR64 (B) following inoculation with *X. oryzae* strains PXO99A, X11-5A, and X8-1A. The length of leaf curling caused by PXO99A was compared to lesion lengths in X11-5A and X8-1A. Curling caused by PXO99A extended the length of the leaf after 10 days on *O. sativa* subsp. *japonica* cv. Lemont. Error bars represent the means of six replicate plants  $\pm$  standard deviations (SD). The experiment was repeated once with similar results.




FIG. 2. Phylogenetic relationships between U.S., Asian, and African *Xanthomonas oryzae* strains. Bayesian analysis was performed for 100,000 generations using a GTR +1 model of evolution based on a concatenated data set of nine complete housekeeping genes totaling 13,425 bp. Bayesian probabilities are shown next to each branch. *X. vasicola* was among the closest relatives to the *X. oryzae* group in previous studies.

inoculated with U.S. *X. oryzae* strains (Fig. 1B). In contrast, *X. oryzae* pv. *oryzae* PXO99A caused long lesions on IR64.

Strains X8-1A and X11-5A did not cause lesions on four additional accessions of *O. sativa* or four accessions of *O. glaberrima* tested. Strain BAI3 caused lesions averaging  $205 \pm 14$  mm in length on the *O. glaberrima* and *O. sativa* accessions, except on accession TOG6202 ( $142 \pm 3$  mm).

**Genome sequence of U.S. *Xanthomonas* strains.** The U.S. *Xanthomonas* strains were sequenced to approximately 70-fold sequence coverage and assembled (see Table S1 in the supplemental material). Over 90% of the assembled sequence of each genome was covered by contigs greater than 10 kbp in length. Table 2 contains a summary of the characteristics of the X11-5A and X8-1A draft genomes compared with genome sequences of *X. oryzae* pv. *oryzae* and *X. oryzae* pv. *oryzicola*. ANI values among the U.S. strain assemblies and *X. oryzae* genomes were greater than 95% (Table 2), the minimum level of identity shared among members of the same species (10, 19). This result places the U.S. strains in the species *X. oryzae* according to widely adopted standards in prokaryotic taxonomy (33). Gene content was predicted and annotated as described in Materials and Methods.

**Phylogenetic analysis of *X. oryzae* housekeeping genes.** Concatenated housekeeping nucleotide sequences from X11-5A and X8-1A were aligned with corresponding sequences obtained from all published *X. oryzae* genome sequences (36) and sequences obtained from two draft genomes of *X. oryzae* strains NAI-8 and MAI-1 from West Africa. Bayesian phylogenetic analysis of the MLST alignments revealed that, although clustering with *X. oryzae*, the U.S. strains form a branch separate from that of *X. oryzae* pv. *oryzae* and *oryzicola* (Fig. 2). Phylogenetic analysis of sequences of the predicted *hrp*, *gum*, and *rpf* virulence clusters supported this finding (data not shown).

To determine whether the sequenced U.S. strains were closely related to other U.S. strains of *Xanthomonas* on rice, including those isolated in different years, we conducted a smaller-scale MLST experiment using additional field isolates. Fragments of the housekeeping genes *fusA*, *gapA*, and *gyrB* were amplified from three additional U.S. strains: 4-4C, isolated from Lemont rice in Bazorria County, TX, in 1987 (17),




FIG. 3. Phylogenetic tree based on a reduced sequence set, including additional *Xanthomonas* strains from rice and southern cutgrass. Bayesian analysis was performed based on a concatenated data set of three partial housekeeping genes totaling 1,440 bp.

and X207A1 and X211-2, isolated in 1990 from Lemont rice and southern cutgrass, respectively, in the same county (C. Gonzalez, personal communication). *X. campestris* pv. *leersiae* strain NCPPB4346, a pathogen of southern cutgrass, was previously shown to group distinctly from *X. oryzae* (30); this strain was also included in the analysis to investigate possible relatedness to the U.S. strains. Analysis of the alignment showed that the U.S. strains 4-4C, X207A1, and X211-2 are very closely related to X8-1A and X11-5A (Fig. 3). *X. campestris* pv. *leersiae* was grouped in the cluster of Asian and African strains. The African strains also grouped with *X. oryzicola* in this reduced data set, confirming previous findings (8). Although South American strains were not included in the phylogenetic analysis, we sequenced a fragment of *gyrB* from the Colombian *X. oryzae* strain CIAT1185 to confirm previous findings that Colombian strains group with the Asian *X. oryzae* pv. *oryzae* strains (data not shown) (15, 17). Together, these results demonstrate that the U.S. strains of *X. oryzae* form a closely related group that is distinct from the economically important *X. oryzae* pv. *oryzae* and *oryzicola* of Asia, Africa, and South America.

**Predicted gene content of U.S. *Xanthomonas* strains.** Predicted ORFs from U.S. *X. oryzae* strains were compared with those of sequenced *Xanthomonas* genomes using reciprocal BLASTN searches. Over 97% of predicted ORFs in the genome of U.S. strain X11-5A have predicted homologs in the genome of U.S. strain X8-1A (Fig. 4A) and vice versa. In contrast, 92% of predicted ORFs from *X. oryzae* pv. *oryzicola* strain BLS256, and 89 to 90% of those predicted in *X. oryzae* pv. *oryzae* strains KACC and PXO99A, have predicted homologs in the U.S. *X. oryzae* genomes (Fig. 4A). These results support the observation that the U.S. *X. oryzae* strains are more closely related to one another than to *X. oryzae* strains from Asia and suggest that the U.S. strains are slightly more similar to *X. oryzae* pv. *oryzicola* than to *X. oryzae* pv. *oryzae*.

**Genes present in Asian *X. oryzae* strains and absent in U.S. strains.** BLASTN analysis identified 200 ORFs in the genome of KACC with homologs in the genome of *X. oryzae* pv. *oryzae* PXO99A and *X. oryzae* pv. *oryzicola* BLS256 but not in the




FIG. 4. Comparison of predicted ORFs in the draft sequence of U.S. *X. oryzae* isolates with those in other previously sequenced *Xanthomonas* genomes. (A) Numbers and percentages of ORFs in the genomic sequences of *X. oryzae* strains X8-1A, KACC10331, PXO99A, and BLS256 and *X. campestris* strain 8004, with and without BLAST hits in the draft genome assembly of strain X11-5A. (B) Functional composition of predicted genes with BLAST hits among all *X. oryzae* genomes except the U.S. draft sequences (left column) and predicted genes shared among the U.S. draft sequences with no BLAST hits in other *X. oryzae* strains (right column).

draft genomic sequence of the U.S. strains. Most genes absent in the U.S. *X. oryzae* sequences are copies of insertion sequence (IS) elements or other transposable elements or are hypothetical genes (Fig. 4B). The lack of these transposable elements may partially result from the incomplete nature of the genomic sequence; comparison of annotated genes in the U.S. genomes with those in previously sequenced genomes suggest that transposable elements are likely underrepresented in the U.S. sequenced genomes (data not shown). Both genomes are predicted to encode intact copies of the major virulence clusters, including *hrp*, *gum*, *rax*, and *rpf* operons.

The U.S. strain genome sequences lack any evidence of genes associated with clustered regularly interspersed palindromic repeats (CRISPRs). CRISPR-associated genes are involved in a virus resistance mechanism (37); these genes are present in many bacterial species, including all three sequenced genomes of *X. oryzae* pv. *oryzae* (13, 35), but a search of the publicly available genome sequence of strain BLS256 revealed that they are lacking in *X. oryzae* pv. *oryzicola*. *X. oryzae* pv. *oryzae* and *oryzicola* are characterized by the presence of 8 to 26 copies of TAL effector genes, which encode secreted proteins required for full virulence (42). TAL effec-

tors are characterized by extensive central repeat regions that bind to promoter elements on host DNA, resulting in modification of host gene expression (6). No TAL effectors were detected in the genome assemblies of the two U.S. strains. Analysis of the unassembled short reads failed to find any evidence of TAL effectors in either U.S. strain genome, while the same analysis performed on sequences from non-U.S. *X. oryzae* strains demonstrated high coverage of TAL sequence (data not shown). In addition, two previous reports demonstrated that TAL effector probes do not hybridize to DNA blots from U.S. strains (7, 34). A BLASTN search revealed that the other predicted type III-secreted effectors reported in *X. oryzae* genomes (36, 40) were nearly all represented in both U.S. *X. oryzae* sequences, with the exception of those in the XopU and XopO families (see Table S2 in the supplemental material).

**U.S. strain-specific genes.** BLAST analysis identified 364 predicted ORFs in the genomic sequence of X11-5A, with close matches in the other U.S. strain sequenced, X8-1A, but no matches with previously sequenced *X. oryzae* genomes. While most of the genes missing from the U.S. strains had no predicted function, over half of the U.S. strain-specific genes were assigned a predicted function (Fig. 4B). The fragmented assemblies used here cannot be used to determine genome-scale gene arrangement; however, analysis of gene arrangement within individual contigs identified 175 unique predicted genes arranged in 47 clusters of two or more in both U.S. strains. The predicted composition of 9 U.S. strain-specific clusters, containing 70 predicted genes, is described in Table S3 in the supplemental material. Cluster 2 encodes a predicted homolog of the *bla* gene for ampicillin resistance. Accordingly, we observed that strains X11-5A and X8-1A both grow robustly on nutrient agar amended with 100  $\mu$ g/ml ampicillin, while growth of Asian strains PXO99 and PXO86 is completely suppressed.

Over 20 predicted ORFs in both U.S. genomes shared significant identity to predicted nonribosomal peptide synthase genes in the genome of *X. albilineans*. The genome of *X. albilineans* encodes multiple clusters of nonribosomal peptide synthases (NRPS) thought to function in antibiosis against bacterial competitors and chlorosis in plants (5, 31). The four predicted ORFs comprising one such *X. albilineans* cluster share 59 to 69% nucleotide identity with predicted NRPS genes in the U.S. *X. oryzae* strains (Fig. 5A). With the exception of the putative NRPS, the rest of the genome shares no more similarity to *X. albilineans* than with the *X. oryzae* pv. *oryzae* or *X. oryzae* pv. *oryzicola* genomes (30).

Due to the fragmented nature of the genome assembly, further study will be needed to determine the structure and genomic location of the NRPS genes. However, high coverage and nucleotide polymorphisms in some of the contigs strongly suggest the presence of multiple copies of NRPS clusters in the genome. Given that NRPS genes are often involved in the production of antibiotic peptides (14, 38), we hypothesized that the U.S. *X. oryzae* strains may have increased capacity for interspecies competition compared with that of *X. oryzae* pv. *oryzae*. When cultures of U.S. and Asian *X. oryzae* strains were incubated on a lawn of *Escherichia coli* DH5 $\alpha$ , the U.S. strains grew well, but two strains from the Philippines did not (Fig. 4B). There was no difference in the appearance of the strains




FIG. 5. U.S. strains of *Xanthomonas* harbor sequences similar to *X. albilineans* NRPS genes and thrive in coculture with *Escherichia coli*. (A) Map of *X. albilineans* genes from locus XAL\_1054 to XAL\_1059, a partially predicted NRPS cluster. Percent nucleotide identity shared with a predicted U.S. *X. oryzae* ORF is shown below each arrow. (B) Growth of U.S. *X. oryzae* strains (top) and common Asian strains (bottom) on a lawn of *E. coli* DH5α.

when incubated on nutrient agar without *E. coli* (data not shown). These results suggest that U.S. strains of *X. oryzae* have an increased ability to compete with bacteria in the environment, possibly as a result of antibiosis by NRPS genes present in the genome.

**Genomic differences between the U.S. *X. oryzae* strains.** Comparative analysis between the two *X. oryzae* assemblies revealed a large contig present in the X8-1A genome which is absent in the genome of X11-5A. The contig was identified as a 31-kb plasmid similar to *X. axonopodis* pv. *vesicatoria* plasmid pXAV38, sharing 97% identity with 74% of the pXAV38 genome. Electrophoresis of the X8-1A genomic DNA confirmed the presence of this plasmid. Despite previous reports of a plasmid in strain X11-5A (41), we found no evidence of a plasmid in the genomic sequence of strain X11-5A; it is possible that a plasmid may have been lost during lab culture.

There were 12 additional gene clusters predicted in X8-1A but not X11-5A, and six gene clusters were found in X11-5A but not X8-1A. These clusters, five of which are described in Table S4 in the supplemental material, include ORFs with predicted involvement in adhesion, modulation of active oxygen species, and O-antigen modification. Because we found no difference in virulence between the two strains, it is unlikely that any of these clusters has a significant role in virulence. The

genome of strain X11-5A contains a fragment corresponding to residues 130 to 257 of the *X. oryzae* pv. *oryzicola* effector protein AvrRxo1, although the putative gene product would lack the probable N-terminal signal necessary for secretion.

**Design of primers amplifying U.S. *X. oryzae* strains.** Previously, we identified primers specific to Asian pathovars of *X. oryzae* that did not amplify products from U.S. strains (20). Here, 15 primer sets were designed based on ORFs specific to U.S. strain genomes or based on the junction sites of predicted U.S.-specific genomic islands. Thirteen of the primer sets successfully differentiated X8-1A and X11-5A from the previously sequenced *X. oryzae* strains; seven primers with robust results in preliminary assays were tested against a panel of 16 U.S. *X. oryzae* strains, 25 strains of *X. oryzae* pv. *oryzae* and *oryzicola*, and six other *Xanthomonas*, *Pseudomonas*, and *Ralstonia* strains listed in Table 2. Five of these primer pairs, reported as USX1 to USX5 in Table 3, consistently amplified a product specific to all U.S. strains.

**DISCUSSION**

With the technological and fiscal advances in next-generation sequencing technologies, research can readily generate whole-genome sequences for an array of isolates of interest. These draft genome sequences are highly informative for comparative analyses of gene content, phylogenetic analyses, and development of diagnostic tests. Because sequencing and automated assembly can be performed in a short period of time, this strategy could be used to identify novel virulence islands and diagnostic markers of an outbreak pathogen well in advance of the following growing season. In this study, we generated a genome sequence by using next-generation sequencing technologies to characterize two U.S. strains of *X. oryzae*. Analysis of the draft sequences enabled placement of the U.S. strains within a novel subgroup of *X. oryzae* strains, identification of many genes missing or uniquely present in the U.S. strains, and development of robust diagnostic primers for identification of the U.S. strains.

Phylogenetic analyses placed the U.S. strains within a group of *X. oryzae* strains distinct from all known Asian and African strains, demonstrating that the diversity of the *X. oryzae* species extends far beyond the previously sequenced model organisms. To a lesser extent, *X. campestris* pv. *leersiae* strain NCPPB4346 and African strains of *X. oryzae* pv. *oryzae* were genetically distinct from Asian strains of *X. oryzae* pv. *oryzae*, supporting previous RFLP studies (7) (unpublished data). Further studies are being conducted to characterize the relationship between the African and Asian strains of *X. oryzae*. Little is known

TABLE 3. U.S. *Xanthomonas*-specific PCR primers

Primer	Forward sequence	Reverse sequence	Product length (bp)	Start position <sup>a</sup>
USX1	GCGCCTGCACAACAATATC	GTA CTGCACCACCGTCTGC	309	4127617
USX2	TCCTCAAAGTTCCCAAGTGC	GGCGTTGGTAAGACGAAGTC	302	1625928
USX3	ATGCAACACCTGCATTTACG	CGACACAGAAAACAGGCTCA	306	11089
USX4	TGGTGGCGAGCTTCTACTATG	GTAGGTCGTCCCAAGTTCAGC	327	2395182
USX5	AGTCGCGCTGTCTCTCAGT	AAGCAACAGCAGACCACCAT	310	1221500

<sup>a</sup> Position of forward primer in assembly 1 of the sequence of strain X11-5A.

about the *L. hexandra* pathogen *X. campestris* pv. *leersiae*, but it is likely a member of the species *X. oryzae*. Our phylogenetic analysis did not include strains isolated from South America (23), although *gyrB* sequencing and diagnostic primer amplification of one Colombian strain supports previous phenotypic and phylogenetic analyses grouping South American strains within the *X. oryzae* pv. *oryzae* (8, 12).

The phylogenetic separation of U.S. strains from other groups supports the hypothesis that *X. oryzae* has persisted in the United States for many years (17). Although bacterial blight is one of the oldest known diseases of rice (26), the existence of distantly related U.S. strains suggests that the divergence of *X. oryzae* pv. *oryzae* and *oryzicola*, and perhaps the entry of TAL effectors into *X. oryzae*, may have happened in the 300 years since the first importation of rice into the United States (assuming that the rice trade is a historically feasible means of *X. oryzae* spread). Given the extent of trade and travel between the United States, Asia, and Africa at that time, it is also possible that an *X. oryzae* progenitor could have entered the United States by way of Africa or traveled by way of other grassy hosts, such as *Leersia* spp. Several of the diagnostic primers tested in this study, initially assumed to be U.S. strain specific, amplified a product from African *X. oryzae* strains, suggesting that there may be some genomic similarities between the African and U.S. *X. oryzae* strains. Although we are currently limited to speculation in this area, the discovery of strains in Asia or Africa sharing U.S. strain-specific features could provide valuable clues regarding the origin and evolution of the *X. oryzae* species and TAL effectors.

Despite the weak pathogenicity and severely limited range of host cultivars of the U.S. strains compared to those of Asian and African *X. oryzae* pv. *oryzae*, we did not identify many ORFs common to the Asian pathovars that are lacking among the U.S. strains; TAL effectors are the only such genes with a known or suspected virulence function. This study, along with two previous reports that the U.S. strains lack TAL effectors (34), is compatible with speculation that the TAL effectors could have entered the *X. oryzae* genome in a single event (40). We hypothesize that an ancestor of *X. oryzae*, possibly a weak pathogen or commensal organism of rice, acquired increased survival on rice with the transfer of TAL effectors from another *Xanthomonas* species. Radiation and diversification of the TAL effectors may have conferred a selective virulence advantage to the descendant *X. oryzae* pv. *oryzae* and *oryzicola*. The U.S. strains of *X. oryzae* may provide a valuable platform for study of the individual or collective contributions of TAL effectors to virulence.

U.S. *X. oryzae* genomes are predicted to share several gene clusters highly similar to those in other environmental *Xanthomonas* spp., including *X. campestris*, *X. axonopodis* pv. *vesicatoria*, and *X. albilineans*. These findings underscore the extensive genetic variation and genomic plasticity among *X. oryzae* species and suggest that U.S. *X. oryzae* has shared a niche with other *Xanthomonas* spp. common in the United States. The U.S. strain-specific genes have suspected involvement in environmental survival and interspecies competition, including predicted genes for oxidative stress resistance, antibiotic resistance, and antibiosis. These gene clusters could equip these strains for survival in the U.S. environment, which

is drier and cooler than that encountered in the rice fields of Southeast Asia or West Africa.

This work demonstrates that U.S. strains of *X. oryzae* form a closely related group genetically and phenotypically distinct from both defined *X. oryzae* pathovars, as well as from *X. campestris* pv. *leersiae*. Given that the U.S. *X. oryzae* strains were isolated from both rice and asymptomatic native grasses in several seasons, it is very plausible that this group of *X. oryzae* still colonizes rice or *Leersia* fields in the southern United States. The clear distinctions between U.S. strains of *X. oryzae* and *X. oryzae* pv. *oryzae* should be taken into consideration in regulatory policy discussions related to bacterial leaf blight of rice. The U.S. strain-specific diagnostic primers developed here will serve as a valuable resource for regulatory programs to distinguish indigenous *X. oryzae* in U.S. rice fields from closely related *X. oryzae* of pathogenic importance. We are currently investigating whether these differences warrant the designation of a novel taxon within the *X. oryzae* species.

#### ACKNOWLEDGMENTS

This research was funded by grants USDA-CSREES-2006-55605-16645 and -2006-55605-04558 to C. R. Buell, N. A. Tisserat, and J. E. Leach and the CSU Infectious Diseases Supercluster. V. Verdier (IRD, CSU) is currently supported by a Marie Curie IOF Fellowship (EU grant PIOF-GA-2009-235457).

We thank C. Gonzales, R. Koebnik, and A. Bogdanove for thoughtful discussions and materials.

#### REFERENCES

- Almeida, N. F., et al. 2010. PAMDB, a multilocus sequence typing and analysis database and website for plant-associated microbes. *Phytopathology* **100**:208–215.
- Almeida, N. F., et al. 2009. A draft genome sequence of *Pseudomonas syringae* pv. *tomato* T1 reveals a type III effector repertoire significantly divergent from that of *Pseudomonas syringae* pv. *tomato* DC3000. *Mol. Plant Microbe Interact.* **22**:52–62.
- Altschul, S. F., et al. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Bhattacharyya, A., et al. 2002. Draft sequencing and comparative genomics of *Xylella fastidiosa* strains reveal novel biological insights. *Genome Res.* **12**:1556–1563.
- Birch, R. G. 2001. *Xanthomonas albilineans* and the antipathogenesis approach to disease control. *Mol. Plant Pathol.* **2**:1–11.
- Boch, J., et al. 2009. Breaking the code of DNA binding specificity of TAL-type III effectors. *Science* **326**:1509–1512.
- Choi, S. H. 1993. Restriction-modification systems and genetic variability of *Xanthomonas oryzae* pv. *oryzae*. Ph.D. dissertation. Kansas State University, Manhattan, KS.
- Gonzalez, C., et al. 2007. Molecular and pathotypic characterization of new *Xanthomonas oryzae* strains from West Africa. *Mol. Plant Microbe Interact.* **20**:534–546.
- Gonzalez, C. F., G. W. Xu, H. L. Li, and J. W. Cosper. 1991. *Leersia hexandra*, an alternative host for *Xanthomonas campestris* pv. *oryzae* in Texas. *Plant Dis.* **75**:159–162.
- Goris, J., et al. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int. J. Syst. Evol. Microbiol.* **57**:81–91.
- Greub, G., et al. 2009. High throughput sequencing and proteomics to identify immunogenic proteins of a new pathogen: the dirty genome approach. *PLoS One* **4**:e8423.
- Guevara, Y., and A. Maselli. 1999. El tizón bacteriano del arroz en Venezuela. *Agronomía Trop.* **49**:505–516.
- Haft, D. H., J. Selengut, E. F. Mongodin, and K. E. Nelson. 2005. A guild of 45 CRISPR-associated (Cas) protein families and multiple CRISPR/Cas subtypes exist in prokaryotic genomes. *PLoS Comput. Biol.* **1**:e60.
- Hashimi, S. M., M. K. Wall, A. B. Smith, A. Maxwell, and R. G. Birch. 2007. The phytotoxin albicidin is a novel inhibitor of DNA gyrase. *Antimicrob. Agents Chemother.* **51**:181–187.
- Hu, J., Y. Zhang, W. Qian, and C. He. 2007. Avirulence gene and insertion element-based RFLP as well as RAPD markers reveal high levels of genomic polymorphism in the rice pathogen *Xanthomonas oryzae* pv. *oryzae*. *Syst. Appl. Microbiol.* **30**:587–600.
- Huelsenbeck, J. P., and F. Ronquist. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* **17**:754–755.

17. Jones, R. K., et al. 1989. Identification of low-virulence strains of *Xanthomonas campestris* pv. *oryzae* from rice in the United States. *Phytopathology* **79**:984–990.
18. Kauffman, H. E., A. P. K. Reddy, S. P. Y. Hsieh, and S. D. Merca. 1973. An improved technique for evaluating resistance of rice varieties to *Xanthomonas oryzae*. *Plant Dis. Rep.* **57**:537–541.
19. Konstantinidis, K. T., A. Ramette, and J. M. Tiedje. 2006. The bacterial species definition in the genomic era. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **361**:1929–1940.
20. Lang, J. M., et al. 2010. Genomics based diagnostic marker development for *Xanthomonas oryzae* pv. *oryzae* and *X. oryzae* pv. *oryzicola*. *Plant Dis.* **94**: 311–319.
21. Larkin, M. A., et al. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* **23**:2947–2948.
22. Lee, B.-M., et al. 2005. The genome sequence of *Xanthomonas oryzae* pathovar *oryzae* KACC10331, the bacterial blight pathogen of rice. *Nucleic Acids Res.* **33**:577–586.
23. Lozano, J. C. 1977. Identification of bacterial leaf blight in rice, caused by *Xanthomonas oryzae*, in America [Central America, South America]. *Plant Dis. Rep.* **61**:644–648.
24. Lu, H., et al. 2008. Acquisition and evolution of plant pathogenesis-associated gene clusters and candidate determinants of tissue-specificity in *Xanthomonas*. *PLoS One* **3**:e3828.
25. Mew, T. W., J. Bridge, H. Hibino, J. M. Bonman, and S. D. Merca. 1988. Rice pathogens of quarantine importance, p. 101–105. *In* Rice seed health. The International Rice Research Institute, Manila, Philippines.
26. Mizukami, T., and S. Wakimoto. 1969. Epidemiology and control of bacterial leaf blight of rice. *Annu. Rev. Phytopathol.* **7**:51–72.
27. Nino-Liu, D. O., P. C. Ronald, and A. J. Bogdanove. 2006. *Xanthomonas oryzae* pathovars: model pathogens of a model crop. *Mol. Plant Pathol.* **7**:303.
28. Ochiai, H., Y. Inoue, M. Takeya, A. Sasaki, and H. Kaku. 2005. Genome sequence of *Xanthomonas oryzae* pv. *oryzae* suggests contribution of large numbers of effector genes and insertion sequences to its race diversity. *Jpn. Agric. Res.* **39**:275–287.
29. OEPP/EPPO. 1980. Data sheets on quarantine organisms no. 2, *Xanthomonas campestris* pv. *oryzae*. OEPP/EPPO bulletin 10. European and Mediterranean Plant Protection Organization, Paris, France.
30. Parkinson, N., C. Cowie, J. Heeney, and D. Stead. 2009. Phylogenetic structure of *Xanthomonas* determined by comparison of *gyrB* sequences. *Int. J. Syst. Evol. Microbiol.* **59**:264–274.
31. Pieretti, I., et al. 2009. The complete genome sequence of *Xanthomonas albilineans* provides new insights into the reductive genome evolution of the xylem-limited *Xanthomonadaceae*. *BMC Genomics* **10**:616.
32. Rice, P., I. Longden, and A. Bleasby. 2000. EMBOSS: the European molecular biology open software suite. *Trends Genet.* **16**:276–277.
33. Richter, M., and R. Rosselló-Móra. 2009. Shifting the genomic gold standard for the prokaryotic species definition. *Proc. Natl. Acad. Sci. U. S. A.* **106**: 19126–19131.
34. Ryba-White, M. 1995. Comparison of *Xanthomonas oryzae* pv. *oryzae* strains from Africa, North America, and Asia by restriction fragment length polymorphism analysis. *Int. Rice Res. Notes* **20**:25–26.
35. Salzberg, S., et al. 2008. Genome sequence and rapid evolution of the rice pathogen *Xanthomonas oryzae* pv. *oryzae* PXO99A. *BMC Genomics* **9**:204.
36. Song, C., and B. Yang. 2010. Mutagenesis of 18 type III effectors reveals virulence function of XopZ<sub>PXO99</sub> in *Xanthomonas oryzae* pv. *oryzae*. *Mol. Plant Microbe Interact.* **23**:893–902.
37. Sorek, R., V. Kunin, and P. Hugenholz. 2008. CRISPR—a widespread system that provides acquired resistance against phages in bacteria and archaea. *Nat. Rev. Microbiol.* **6**:181–186.
38. Stricker, M., A. Tanovic, and M. A. Marahiel. 2010. Nonribosomal peptide synthetases: structures and dynamics. *Curr. Opin. Struct. Biol.* **20**:234–240.
39. Tamura, K., J. Dudley, M. Nei, and S. Kumar. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**:1596–1599.
40. White, F. F., N. Potnis, J. B. Jones, and R. Koebnik. 2009. The type III effectors of *Xanthomonas*. *Mol. Plant Pathol.* **10**:749–766.
41. Xu, G. W., and C. F. Gonzalez. 1991. Plasmid, genomic, and bacteriocin diversity in U.S. strains of *Xanthomonas campestris* pv. *oryzae*. *Phytopathology* **81**:628–631.
42. Yang, B., and F. F. White. 2004. Diverse members of the AvrBs3/PthA family of type III effectors are major virulence determinants in bacterial blight disease of rice. *Mol. Plant Microbe Interact.* **17**:1192–1200.
43. Zerbino, D. R., and E. Birney. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* **18**:821–829.